

## Research Article

# On Some Solvable Difference Equations and Systems of Difference Equations

**Stevo Stević,<sup>1</sup> Josef Diblík,<sup>2,3</sup>  
Bratislav Iričanin,<sup>4</sup> and Zdeněk Šmarda<sup>3</sup>**

<sup>1</sup> *Mathematical Institute of the Serbian Academy of Sciences and Arts, Knez Mihailova 36/III, 11000 Beograd, Serbia*

<sup>2</sup> *Department of Mathematics and Descriptive Geometry, Faculty of Civil Engineering, Brno University of Technology, 60200 Brno, Czech Republic*

<sup>3</sup> *Department of Mathematics, Faculty of Electrical Engineering and Communication, Brno University of Technology, 61600 Brno, Czech Republic*

<sup>4</sup> *Faculty of Electrical Engineering, University of Belgrade, Bulevar Kralja Aleksandra 73, 11000 Beograd, Serbia*

Correspondence should be addressed to Stevo Stević, [sstevic@ptt.rs](mailto:sstevic@ptt.rs)

Received 8 August 2012; Accepted 27 September 2012

Academic Editor: Svatoslav Staněk

Copyright © 2012 Stevo Stević et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Here, we give explicit formulae for solutions of some systems of difference equations, which extend some very particular recent results in the literature and give natural explanations for them, which were omitted in the previous literature.

## 1. Introduction

Recently, there has been a great interest in difference equations and systems (see, e.g., [1–25]), and among them in those ones which can be solved explicitly (see, e.g., [1–5, 9–11, 15, 16, 18–24] and the related references therein). For some classical results in the topic see, for example, [7].

Beside the above-mentioned papers, there are some papers which give formulae of some very particular equations and systems which are proved by induction, but without any explanation how these formulae are obtained and how these authors came across the equations and systems. Our explanation of such a formula that we gave in [10] has re-attracted attention to solvable difference equations.

Our aim here is to give theoretical explanations for some of the formulae recently appearing in the literature, as well as to give some extensions of their equations.

Before we formulate our results, we would like to say that the system of difference equations

$$x_{n+1} = \frac{ax_{n-1}}{b + cy_n x_{n-1}}, \quad y_{n+1} = \frac{\alpha y_{n-1}}{\beta + \gamma x_n y_{n-1}}, \quad n \in \mathbb{N}_0, \quad (1.1)$$

where  $a, b, c, \alpha, \beta,$  and  $\gamma$  are real numbers, was completely solved in [15], that is, we found formulae for all well-defined solutions of system (1.1).

## 2. Scaling Indices

In the recent paper [25] were given some formulae for the solutions of the following systems of difference equations:

$$x_{n+1} = \frac{x_{n-3}}{\pm 1 \pm x_{n-3} y_{n-1}}, \quad y_{n+1} = \frac{y_{n-3}}{\pm 1 \pm y_{n-3} x_{n-1}}, \quad n \in \mathbb{N}_0. \quad (2.1)$$

Now we show that the results regarding system (2.1) easily follow from known ones. Indeed, if we use the change of variables

$$x_{2n+i} = u_n^{(i)}, \quad y_{2n+i} = v_n^{(i)}, \quad n \geq -2, \quad i = 1, 2, \quad (2.2)$$

then the systems in (2.1) are reduced to the next systems

$$u_{n+1}^{(i)} = \frac{u_{n-1}^{(i)}}{\pm 1 \pm u_{n-1}^{(i)} v_n^{(i)}}, \quad v_{n+1}^{(i)} = \frac{v_{n-1}^{(i)}}{\pm 1 \pm v_{n-1}^{(i)} u_n^{(i)}}, \quad n \geq -1, \quad i = 1, 2. \quad (2.3)$$

This means that  $(u_n^{(i)}, v_n^{(i)})_{n \geq -2}, i = 1, 2,$  are two (independent) solutions of the systems of difference equations

$$x_{n+1} = \frac{x_{n-1}}{\pm 1 \pm x_{n-1} y_n}, \quad y_{n+1} = \frac{y_{n-1}}{\pm 1 \pm y_{n-1} x_n}, \quad n \geq -1, \quad i = 1, 2. \quad (2.4)$$

However, all the systems of difference equations in (2.4) are particular cases of system (1.1). Hence, formulae for the solutions of systems (2.1) given in [25] follow directly from those in [15].

### 2.1. An Extension of Systems (2.1)

Systems (2.1) can be extended as follows:

$$x_{n+1} = \frac{ax_{n-2k+1}}{b + cx_{n-2k+1}y_{n-k+1}}, \quad y_{n+1} = \frac{\alpha y_{n-2k+1}}{\beta + \gamma y_{n-2k+1}x_{n-k+1}}, \quad n \in \mathbb{N}_0, \quad (2.5)$$

where  $k$  is a fixed natural number.

If we use the change of variables

$$x_{mk+i} = u_m^{(i)}, \quad y_{mk+i} = v_m^{(i)}, \quad m \geq -2, \quad i = 1, \dots, k, \quad (2.6)$$

then system (2.5) is reduced to the following  $k$  systems of difference equations:

$$u_{n+1}^{(i)} = \frac{au_{n-1}^{(i)}}{b + cu_{n-1}^{(i)}v_n^{(i)}}, \quad v_{n+1}^{(i)} = \frac{\alpha v_{n-1}^{(i)}}{\beta + \gamma v_{n-1}^{(i)}u_n^{(i)}}, \quad n \geq -1, \quad (2.7)$$

$i = 1, \dots, k$ . This means that  $(u_n^{(i)}, v_n^{(i)})_{n \geq -2}, i = 1, \dots, k$ , are  $k$  (independent) solutions of system (1.1), and solutions of system (2.5) are obtained by interlacing solutions of systems (2.7),  $i = 1, \dots, k$ .

For example, a natural extension of the systems in (2.1) is obtained for taking  $k = 3$ ,  $b/a = \pm 1$ ,  $c/a = \pm 1$ ,  $\beta/\alpha = \pm 1$ , and  $\gamma/\alpha = \pm 1$  in (2.5), that is, the system becomes

$$x_{n+1} = \frac{x_{n-5}}{\pm 1 \pm x_{n-5}y_{n-2}}, \quad y_{n+1} = \frac{y_{n-5}}{\pm 1 \pm y_{n-5}x_{n-2}}, \quad n \in \mathbb{N}_0. \quad (2.8)$$

In this way it can be obtained countable many, at first sight different, systems of difference equations. Systems of difference equations in (2.1) are artificially obtained in this way. This method can be applied to any equation or system of difference equations, and one can get papers with putative “new” results.

### 3. Some Third-Order Systems of Difference Equations Related to (1.1)

The following third-order systems of difference equations

$$x_{n+1} = \frac{x_{n-1}}{y_n x_{n-1} - 1}, \quad y_{n+1} = \frac{y_{n-1}}{x_n y_{n-1} - 1}, \quad z_{n+1} = \frac{1}{z_n y_n}, \quad (3.1)$$

$$x_{n+1} = \frac{x_{n-1}}{y_n x_{n-1} - 1}, \quad y_{n+1} = \frac{y_{n-1}}{x_n y_{n-1} - 1}, \quad z_{n+1} = \frac{z_{n-1}}{y_n z_{n-1} - 1}, \quad (3.2)$$

$$x_{n+1} = \frac{x_{n-1}}{y_n x_{n-1} - 1}, \quad y_{n+1} = \frac{y_{n-1}}{x_n y_{n-1} - 1}, \quad z_{n+1} = \frac{x_n}{z_{n-1} y_n}, \quad (3.3)$$

$n \in \mathbb{N}_0$ , have been studied recently (see [6] and the references therein).

As is directly seen, the first two equations in systems (3.1)–(3.3) are the same, and they form a particular case of system (1.1) which is solved in [15].

Since we know solutions for  $x_n$  and  $y_n$ , it is only needed to find explicit solutions for  $z_n$  in the third equations in systems (3.1)–(3.3), that is, in all three equations, the only unknown sequence is  $z_n$ . The joint feature for all three cases is that  $z_n$  can be solved in closed form.

Now we discuss systems of difference equations given in (3.1)–(3.3).

Case of System (3.1)

From the third equation in (3.1), we get

$$z_{n+1} = \frac{1}{z_n y_n} = \frac{y_{n-1}}{y_n} z_{n-1}, \quad n \in \mathbb{N}, \quad (3.4)$$

from which it follows that

$$z_{2n+i} = z_i \prod_{j=1}^n \frac{y_{2j+i-2}}{y_{2j+i-1}}, \quad n \in \mathbb{N}_0, \quad i = 0, 1. \quad (3.5)$$

Case of System (3.3)

From the third equation in (3.3), we get

$$z_{n+1} = \frac{x_n}{z_{n-1} y_n} = \frac{x_n y_{n-2}}{x_{n-2} y_n} z_{n-3}, \quad n \geq 2, \quad (3.6)$$

from which it follows that

$$z_{4n+i} = z_i \prod_{j=1}^n \frac{x_{4j+i-1} y_{4j+i-3}}{x_{4j+i-3} y_{4j+i-1}}, \quad n \in \mathbb{N}_0, \quad i = -1, 0, 1, 2. \quad (3.7)$$

*Remark 3.1.* The third equations in systems (3.1) and (3.3) are particular cases of the following difference equation (up to the shifting indices):

$$z_n = \frac{a_n}{z_{n-k}}, \quad n \in \mathbb{N}_0, \quad (3.8)$$

where  $k \in \mathbb{N}$ . From (3.8), it follows that

$$z_n = \frac{a_n}{a_{n-k}} z_{n-2k}, \quad n \geq k, \quad (3.9)$$

and consequently,

$$z_{2mk+i} = z_{i-2k} \prod_{j=0}^m \frac{a_{2kj+i}}{a_{2kj-k+i}}, \quad m \in \mathbb{N}_0, \quad i = k, k+1, \dots, 3k-1. \quad (3.10)$$

Case of System (3.2)

If we use the change of variables  $v_n = 1/z_n$ , the third equation in (3.2) becomes

$$v_{n+1} = -v_{n-1} + y_n, \quad n \in \mathbb{N}_0. \quad (3.11)$$

Hence,

$$v_{2n+i} = -v_{2(n-1)+i} + y_{2n+i-1}, \quad n \in \mathbb{N}_0, \quad i = 1, 2, \quad (3.12)$$

from which it follows that

$$v_{2n+i} = (-1)^{n+1} v_{i-2} + \sum_{j=0}^n (-1)^{n-j} y_{2j+i-1}, \quad (3.13)$$

so that

$$z_{2n+i} = \frac{z_{i-2}}{(-1)^{n+1} + z_{i-2} \sum_{j=0}^n (-1)^{n-j} y_{2j+i-1}}. \quad (3.14)$$

*Remark 3.2.* The third equation in system (3.2) is a particular case of the following difference equation:

$$z_n = \frac{a_n z_{n-k}}{b_n z_{n-k} + c_n}, \quad n \in \mathbb{N}_0, \quad (3.15)$$

which, by the change of variables  $z_n = 1/v_n$ , is transformed into

$$v_n = \frac{c_n}{a_n} v_{n-k} + \frac{b_n}{a_n}, \quad n \in \mathbb{N}_0, \quad (3.16)$$

from which it follows that

$$v_{km+i} = \frac{c_{km+i}}{a_{km+i}} v_{k(m-1)+i} + \frac{b_{km+i}}{a_{km+i}}, \quad m \in \mathbb{N}_0, \quad i = 0, 1, \dots, k-1, \quad (3.17)$$

so by a well-known formula, we have that

$$v_{km+i} = v_{i-k} \prod_{j=0}^m \frac{c_{kj+i}}{a_{kj+i}} + \sum_{j=0}^m \frac{b_{kj+i}}{a_{kj+i}} \prod_{l=j+1}^m \frac{c_{kl+i}}{a_{kl+i}}, \quad (3.18)$$

for  $m \in \mathbb{N}_0$  and  $i = 0, 1, \dots, k-1$ , and consequently,

$$z_{mk+i} = \frac{z_{i-k}}{\prod_{j=0}^m (c_{kj+i}/a_{kj+i}) + z_{i-k} \sum_{j=0}^m (b_{kj+i}/a_{kj+i}) \prod_{l=j+1}^m (c_{kl+i}/a_{kl+i})}, \quad (3.19)$$

for  $m \in \mathbb{N}_0$  and  $i = 0, 1, \dots, k-1$ .

*Remark 3.3.* Note that (3.16) suggests that the third equation in (3.2) can be also of the form

$$z_n = a_n z_{n-k} + b_n, \quad n \in \mathbb{N}_0, \quad (3.20)$$

where  $k \in \mathbb{N}$  is fixed, that is, to be an equation which consists of  $k$  (independent) linear first-order difference equation, which is solvable. In fact, the third equation in systems (3.1)–(3.3) can be any difference equation which can be solved in  $z_n$ , and in this way we can obtain numerous putative “new” results.

#### 4. A Generalization of System (1.1)

Consider the following system of difference equations:

$$x_{n+1} = g^{-1}\left(\frac{g(x_{n-1})}{ah(y_n)g(x_{n-1}) + b}\right), \quad y_{n+1} = h^{-1}\left(\frac{h(y_{n-1})}{cg(x_n)h(y_{n-1}) + d}\right), \quad n \in \mathbb{N}_0, \quad (4.1)$$

where  $g, h : \mathbb{R} \rightarrow \mathbb{R}$  are increasing functions such that

$$g(0) = h(0) = 0, \quad (4.2)$$

$$g(x)x > 0, \quad h(x)x > 0, \quad \text{for } x \neq 0. \quad (4.3)$$

Now we will find formulae for all well-defined solutions of system (4.1), that is, for the solutions  $(x_n, y_n)$ ,  $n \geq -1$ , such that

$$ah(y_n)g(x_{n-1}) + b \neq 0, \quad cg(x_n)h(y_{n-1}) + d \neq 0, \quad (4.4)$$

for every  $n \in \mathbb{N}_0$ .

If  $x_{-1} = 0$ , then from (4.1), (4.2), and (4.3), and by the method of induction, we get  $x_{2n+1} = 0$ ,  $n \in \mathbb{N}_0$ . Also, if  $x_0 = 0$ , then from (4.1), (4.2), and (4.3), and by the method of induction, we get  $x_{2n} = 0$ ,  $n \in \mathbb{N}_0$ . Similarly, if  $y_{-1} = 0$ , then we get  $y_{2n+1} = 0$ ,  $n \in \mathbb{N}_0$ , while if  $y_0 = 0$ , then we get  $y_{2n} = 0$ ,  $n \in \mathbb{N}_0$ .

If  $x_{n_0} = 0$  for some  $n_0 \in \mathbb{N}$ , then from (4.1)–(4.3) it follows that  $x_{n_0-2k} = 0$ , for each  $k \in \mathbb{N}_0$  such that  $n_0 - 2k \geq -1$ . Hence, in this case we have that  $x_{-1} = 0$  or  $x_0 = 0$ . Similarly, if  $y_{n_1} = 0$  for some  $n_1 \in \mathbb{N}$ , then from (4.1)–(4.3) it follows that  $y_{n_1-2k} = 0$ , for each  $k \in \mathbb{N}_0$  such that  $n_1 - 2k \geq -1$ . Hence, in this case we have that  $y_{-1} = 0$  or  $y_0 = 0$ . Thus, in both cases we arrive at a situation explained in the previous paragraph.

Hence, from now on, we assume that none of the initial values  $x_{-1}$ ,  $x_0$ ,  $y_{-1}$ , and  $y_0$  is equal to zero. Then, for every well-defined solution of system (4.1), we have that  $x_n \neq 0$  and  $y_n \neq 0$ , for every  $n \geq -1$ , and consequently  $g(x_n) \neq 0$  and  $h(y_n) \neq 0$ , for every  $n \geq -1$ .

Let

$$v_n = \frac{1}{g(x_n)h(y_{n-1})}, \quad u_n = \frac{1}{g(x_{n-1})h(y_n)}, \quad (4.5)$$

then by taking function  $g$  to the first equation in system (4.1) and function  $h$  to the second one, then multiplying the first equation in such obtained system by  $h(y_n)$  and the second by  $g(x_n)$ , system (4.1) is transformed into:

$$v_{n+1} = bu_n + a, \quad u_{n+1} = dv_n + c, \quad n \in \mathbb{N}_0, \quad (4.6)$$

from which it follows that

$$\begin{aligned}v_{n+1} &= bdv_{n-1} + bc + a, \quad n \in \mathbb{N}, \\u_{n+1} &= bdu_{n-1} + ad + c, \quad n \in \mathbb{N}.\end{aligned}\tag{4.7}$$

Hence, if  $bd \neq 1$ , we have that

$$\begin{aligned}v_{2n} &= \frac{(bd)^n}{g(x_0)h(y_{-1})} + (bc + a)\frac{1 - (bd)^n}{1 - bd}, \quad n \in \mathbb{N}_0, \\v_{2n+1} &= \frac{(bd)^n}{g(x_1)h(y_0)} + (bc + a)\frac{1 - (bd)^n}{1 - bd}, \quad n \in \mathbb{N}_0,\end{aligned}\tag{4.8}$$

while if  $bd = 1$ , we have that

$$\begin{aligned}v_{2n} &= \frac{1}{g(x_0)h(y_{-1})} + (bc + a)n, \quad n \in \mathbb{N}_0, \\v_{2n+1} &= \frac{1}{g(x_1)h(y_0)} + (bc + a)n, \quad n \in \mathbb{N}_0.\end{aligned}\tag{4.9}$$

We have also that

$$\begin{aligned}u_{2n} &= \frac{(bd)^n}{h(y_0)g(x_{-1})} + (ad + c)\frac{1 - (bd)^n}{1 - bd}, \quad n \in \mathbb{N}_0, \\u_{2n+1} &= \frac{(bd)^n}{h(y_1)g(x_0)} + (ad + c)\frac{1 - (bd)^n}{1 - bd}, \quad n \in \mathbb{N}_0,\end{aligned}\tag{4.10}$$

while if  $bd = 1$ , we have that

$$\begin{aligned}u_{2n} &= \frac{1}{h(y_0)g(x_{-1})} + (ad + c)n, \quad n \in \mathbb{N}_0, \\u_{2n+1} &= \frac{1}{h(y_1)g(x_0)} + (ad + c)n, \quad n \in \mathbb{N}_0.\end{aligned}\tag{4.11}$$

From (4.5), we have that

$$\begin{aligned}g(x_n) &= \frac{1}{v_n h(y_{n-1})} = \frac{u_{n-1}}{v_n} g(x_{n-2}), \quad n \in \mathbb{N}, \\h(y_n) &= \frac{1}{u_n g(x_{n-1})} = \frac{v_{n-1}}{u_n} h(y_{n-2}), \quad n \in \mathbb{N}.\end{aligned}\tag{4.12}$$

Using the relations (4.12), we get

$$\begin{aligned}
 x_{2n} &= g^{-1} \left( g(x_0) \prod_{j=1}^n \frac{u_{2j-1}}{v_{2j}} \right), \quad n \in \mathbb{N}_0, \\
 x_{2n+1} &= g^{-1} \left( g(x_{-1}) \prod_{j=0}^n \frac{u_{2j}}{v_{2j+1}} \right), \quad n \in \mathbb{N}_0, \\
 y_{2n} &= h^{-1} \left( h(y_0) \prod_{j=1}^n \frac{v_{2j-1}}{u_{2j}} \right), \quad n \in \mathbb{N}_0, \\
 y_{2n+1} &= h^{-1} \left( h(y_{-1}) \prod_{j=0}^n \frac{v_{2j}}{u_{2j+1}} \right), \quad n \in \mathbb{N}_0.
 \end{aligned} \tag{4.13}$$

*Example 4.1.* If we choose  $g(t) = t^{2k+1}$  and  $h(t) = t^{2l+1}$  for some  $k, l \in \mathbb{N}_0$ , then conditions (4.2) and (4.3) are obviously satisfied, and system (4.1) can be written in the form

$$x_{n+1} = \left( \frac{x_{n-1}^{2k+1}}{ay_n^{2l+1}x_{n-1}^{2k+1} + b} \right)^{1/(2k+1)}, \quad y_{n+1} = \left( \frac{y_{n-1}^{2l+1}}{cx_n^{2k+1}y_{n-1}^{2l+1} + d} \right)^{1/(2l+1)}, \quad n \in \mathbb{N}_0, \tag{4.14}$$

and from (4.13), we have that its solutions are given by

$$\begin{aligned}
 x_{2n} &= x_0 \prod_{j=1}^n \left( \frac{u_{2j-1}}{v_{2j}} \right)^{1/(2k+1)}, \quad n \in \mathbb{N}_0, \\
 x_{2n+1} &= x_{-1} \prod_{j=0}^n \left( \frac{u_{2j}}{v_{2j+1}} \right)^{1/(2k+1)}, \quad n \in \mathbb{N}_0, \\
 y_{2n} &= y_0 \prod_{j=1}^n \left( \frac{v_{2j-1}}{u_{2j}} \right)^{1/(2l+1)}, \quad n \in \mathbb{N}_0, \\
 y_{2n+1} &= y_{-1} \prod_{j=0}^n \left( \frac{v_{2j}}{u_{2j+1}} \right)^{1/(2l+1)}, \quad n \in \mathbb{N}_0.
 \end{aligned} \tag{4.15}$$

*Remark 4.2.* System (4.1) can be generalized by using the method of scaling indices from Section 2.1, that is, the following system is also solvable:

$$x_{n+1} = g^{-1} \left( \frac{g(x_{n-2k+1})}{ah(y_{n-k+1})g(x_{n-2k+1}) + b} \right), \quad y_{n+1} = h^{-1} \left( \frac{h(y_{n-2k+1})}{cg(x_{n-k+1})h(y_{n-2k+1}) + d} \right), \tag{4.16}$$



$n \in \mathbb{N}_0$ , where  $k \in \mathbb{N}$  and  $g, h : \mathbb{R} \rightarrow \mathbb{R}$  are increasing functions satisfying conditions (4.2) and (4.3).

It is easy to see that the change of variables in (2.6) leads to the following  $k$  systems of difference equations:

$$u_{n+1}^{(i)} = g^{-1} \left( \frac{g(u_{n-1}^{(i)})}{ah(v_n^{(i)})g(u_{n-1}^{(i)}) + b} \right), \quad v_{n+1}^{(i)} = h^{-1} \left( \frac{h(v_{n-1}^{(i)})}{cg(u_n^{(i)})h(v_{n-1}^{(i)}) + d} \right), \quad n \geq -1, \tag{4.17}$$

for  $i = 1, \dots, k$ .

*Remark 4.3.* Well-defined solutions of the following system of difference equations:

$$x_{n+1} = g^{-1} \left( \frac{g(x_{n-1})}{a_n h(y_n) g(x_{n-1}) + b_n} \right), \quad y_{n+1} = h^{-1} \left( \frac{h(y_{n-1})}{c_n g(x_n) h(y_{n-1}) + d_n} \right), \quad n \in \mathbb{N}_0, \tag{4.18}$$

where  $g, h : \mathbb{R} \rightarrow \mathbb{R}$  are increasing functions satisfying conditions (4.2) and (4.3) and  $a_n, b_n, c_n$ , and  $d_n, n \in \mathbb{N}_0$ , are real sequences, can be found similarly. We omit the details.

### 5. Solutions of a Generalization of a Recent Equation

Explaining some recent formulae appearing in the literature, in our recent paper [24], we have found formulae for well-defined solutions of the following difference equation:

$$x_{n+1} = \frac{x_n x_{n-k}}{x_{n-k+1} (a + b x_n x_{n-k})}, \quad n \in \mathbb{N}_0, \tag{5.1}$$

where  $k \in \mathbb{N}$  and the parameters  $a, b$  as well as initial values  $x_{-i}, i = \overline{0, k}$  are real numbers.

Equation (5.1) can be extended naturally in the following way:

$$x_{n+1} = \frac{x_n x_{n-k}}{x_{n-k+1} (a_n + b_n x_n x_{n-k})}, \quad n \in \mathbb{N}_0, \tag{5.2}$$

where  $k \in \mathbb{N}$  and the sequences  $(a_n)_{n \in \mathbb{N}_0}, (b_n)_{n \in \mathbb{N}_0}$ , as well as initial values  $x_{-i}, i = \overline{0, k}$ , are real numbers.

Employing the change of variables

$$y_n = \frac{1}{x_n x_{n-k}}, \quad n \in \mathbb{N}_0, \tag{5.3}$$

Equation (5.2) is transformed into the linear first-order difference equation

$$y_{n+1} = a_n y_n + b_n, \quad n \in \mathbb{N}_0, \tag{5.4}$$

whose general solution is

$$y_n = y_0 \prod_{l=0}^{n-1} a_l + \sum_{s=0}^{n-1} b_s \prod_{l=s+1}^{n-1} a_l, \quad n \in \mathbb{N}_0. \quad (5.5)$$

From (5.3), we have that

$$x_n = \frac{1}{y_n x_{n-k}} = \frac{y_{n-k}}{y_n} x_{n-2k}, \quad (5.6)$$

for  $n \geq k$ , which yields

$$x_{2km+i} = x_{i-2k} \prod_{j=0}^m \frac{y_{(2j-1)k+i}}{y_{2jk+i}}, \quad (5.7)$$

for every  $m \in \mathbb{N}_0$  and  $i \in \{k, k+1, \dots, 3k-1\}$ .

Using (5.5) in (5.7), we get

$$x_{2km+i} = x_{i-2k} \prod_{j=0}^m \frac{y_0 \prod_{l=0}^{(2j-1)k+i-1} a_l + \sum_{i=s}^{(2j-1)k+i-1} b_s \prod_{l=s+1}^{(2j-1)k+i-1} a_l}{y_0 \prod_{l=0}^{2jk+i-1} a_l + \sum_{s=0}^{2jk+i-1} b_s \prod_{l=s+1}^{2jk+i-1} a_l}, \quad (5.8)$$

for every  $m \in \mathbb{N}_0$  and  $i \in \{k, k+1, \dots, 3k-1\}$ .

Formula (5.8) generalizes the main formulae obtained in our paper [24].





## Acknowledgments

The second author is supported by the Grant P201/10/1032 of the Czech Grant Agency (Prague) and the "Operational Programme Research and Development for Innovations," no. CZ.1.05/2.1.00/03.0097, as an activity of the regional Centre AdMaS. The fourth author is supported by the Grant FEKT-S-11-2-921 of Faculty of Electrical Engineering and Communication, Brno University of Technology and by the Grant P201/11/0768 of the Czech Grant Agency (Prague). This paper is also supported by the Serbian Ministry of Science projects III 41025, III 44006, and OI 171007.

## References

- [1] A. Andruch-Sobiło and M. Migda, "On the rational recursive sequence  $x_{n+1} = ax_{n-1}/(b + cx_{n-1})$ ," *Tatra Mountains Mathematical Publications*, vol. 43, pp. 1–9, 2009.
- [2] I. Bajo and E. Liz, "Global behaviour of a second-order nonlinear difference equation," *Journal of Difference Equations and Applications*, vol. 17, no. 10, pp. 1471–1486, 2011.
- [3] L. Berg and S. Stević, "On difference equations with powers as solutions and their connection with invariant curves," *Applied Mathematics and Computation*, vol. 217, no. 17, pp. 7191–7196, 2011.
- [4] L. Berg and S. Stević, "On some systems of difference equations," *Applied Mathematics and Computation*, vol. 218, no. 5, pp. 1713–1718, 2011.
- [5] B. Iričanin and S. Stević, "On some rational difference equations," *Ars Combinatoria*, vol. 92, pp. 67–72, 2009.

- [6] A. S. Kurbanli, "On the behavior of solutions of the system of rational difference equations  $x_{n+1} = x_{n-1}/y_n x_{n-1} - 1$ ,  $y_{n+1} = y_{n-1}/(x_n y_{n-1} - 1)$  and  $z_{n+1} = z_{n-1}/(y_n z_{n-1} - 1)$ ," *Advances in Difference Equations*, vol. 2011, 40 pages, 2011.
- [7] H. Levy and F. Lessman, *Finite Difference Equations*, The Macmillan Company, New York, NY, USA, 1961.
- [8] G. Papaschinopoulos, C. J. Schinas, and G. Stefanidou, "On the nonautonomous difference equation  $x_{n+1} = A_n + (x_{n-1}^p/x_n^q)$ ," *Applied Mathematics and Computation*, vol. 217, no. 12, pp. 5573–5580, 2011.
- [9] G. Papaschinopoulos and G. Stefanidou, "Asymptotic behavior of the solutions of a class of rational difference equations," *International Journal of Difference Equations*, vol. 5, no. 2, pp. 233–249, 2010.
- [10] S. Stević, "More on a rational recurrence relation," *Applied Mathematics E-Notes*, vol. 4, pp. 80–85, 2004.
- [11] S. Stević, "A short proof of the Cushing-Henson conjecture," *Discrete Dynamics in Nature and Society*, vol. 2006, Article ID 37264, 5 pages, 2006.
- [12] S. Stević, "On the recursive sequence  $x_{n+1} = \max\{c, x_n^p/x_{n-1}^p\}$ ," *Applied Mathematics Letters*, vol. 21, no. 8, pp. 791–796, 2008.
- [13] S. Stević, "Global stability of a max-type difference equation," *Applied Mathematics and Computation*, vol. 216, no. 1, pp. 354–356, 2010.
- [14] S. Stević, "On a nonlinear generalized max-type difference equation," *Journal of Mathematical Analysis and Applications*, vol. 376, no. 1, pp. 317–328, 2011.
- [15] S. Stević, "On a system of difference equations," *Applied Mathematics and Computation*, vol. 218, no. 7, pp. 3372–3378, 2011.
- [16] S. Stević, "On the difference equation  $x_n = x_{n-2}/(b_n + c_n x_{n-1} x_{n-2})$ ," *Applied Mathematics and Computation*, vol. 218, no. 8, pp. 4507–4513, 2011.
- [17] S. Stević, "Periodicity of a class of nonautonomous max-type difference equations," *Applied Mathematics and Computation*, vol. 217, no. 23, pp. 9562–9566, 2011.
- [18] S. Stević, "On a system of difference equations with period two coefficients," *Applied Mathematics and Computation*, vol. 218, no. 8, pp. 4317–4324, 2011.
- [19] S. Stević, "On a third-order system of difference equations," *Applied Mathematics and Computation*, vol. 218, no. 14, pp. 7649–7654, 2012.
- [20] S. Stević, "On some solvable systems of difference equations," *Applied Mathematics and Computation*, vol. 218, no. 9, pp. 5010–5018, 2012.
- [21] S. Stević, "On the difference equation  $x_n = x_{n-k}/(b + c x_{n-1} \cdots x_{n-k})$ ," *Applied Mathematics and Computation*, vol. 218, no. 11, pp. 6291–6296, 2012.
- [22] S. Stević, J. Diblík, B. Iričanin, and Z. Šmarda, "On a third-order system of difference equations with variable coefficients," *Abstract and Applied Analysis*, vol. 2012, Article ID 508523, 22 pages, 2012.
- [23] S. Stević, J. Diblík, B. Iričanin, and Z. Šmarda, "On the difference equation  $x_n = a_n x_{n-k}/(b_n + c_n x_{n-1} x_{n-k})$ ," *Abstract and Applied Analysis*, vol. 2012, Article ID Article number409237, 20 pages, 2012.
- [24] S. Stević, J. Diblík, B. Iričanin, and Z. Šmarda, "On the difference equation  $x_{n+1} = x_n x_{n-k}/(x_{n-k+1}(a + b x_n x_{n-k}))$ ," *Abstract and Applied Analysis*, vol. 2012, Article ID Article number108047, 9 pages, 2012.
- [25] N. Touafek and E. M. Elsayed, "On the solutions of systems of rational difference equations," *Mathematical and Computer Modelling*, vol. 55, no. 7-8, pp. 1987–1997, 2012.



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

