

## ON STOCHASTIC INTEGRODIFFERENTIAL EQUATIONS VIA NON-LINEAR INTEGRAL CONTRACTORS I

Miljana Jovanović\* and Svetlana Janković†

### Abstract

The aim of this paper is to study the existence and uniqueness of solutions for a general stochastic integrodifferential equation of the Ito type, by using the concept of non-linear bounded random integral contractors, which includes the Lipschitz condition as a special case. The method applied in this consideration follows partially the basic ideas of the contractor theory introduced earlier by Altman [1, 2] and Kuo [6]. It is also shown that the Lipschitz condition and the condition based on a bounded random integral contractor for the coefficients of the considered equation, in general, cannot be compared.

## 1 Introduction

Several phenomena in life and sciences, especially in mechanics, engineering and, since recently, in finance, have been found to depend on random excitations. It therefore seems natural that current trend in describing and studying these phenomena is focused on the use of stochastic mathematical models rather than deterministic ones. Having in mind that in many cases random excitations are of the Gaussian white noise type, which is mathematically described as a formal derivative of the Brownian motion, all such phenomena are mathematically modelled and essentially represented by complex stochastic differential equations of the Ito type. Obviously, the interest of researchers is usually focused on conditions guaranteeing the existence, uniqueness and bifurcational behavior of solutions to these equations.

For example, the behavior of a non-linear dynamical system can be represented by the following differential equation

$$\dot{y} + f(t, \dot{y}, y) = g(t, \dot{y}, y) \cdot \xi(t, \omega), \quad t \geq 0,$$

---

\*Corresponding author

†Supported by Grant No 144003 of MNTRS.

2000 *Mathematics Subject Classifications*. 60H20.

*Key words and Phrases*. Stochastic integrodifferential equation, non-linear random integral contractor, existence and uniqueness of solutions.

Received: June 15, 2009

Communicated by Dragan S. Djordjević

where  $\xi(t, \omega)$  is a Gaussian white noise perturbation and  $\omega \in \Omega$  are random events. Since  $\xi(t, \omega) = \dot{w}(t, \omega)$ , where  $w(t, \omega)$  is a Brownian motion, this equation can be transformed into the following stochastic system,

$$\begin{aligned} dy(t) &= x(t) dt \\ dx(t) &= -f\left(t, x(t), c + \int_0^t x(s) ds\right) dt + g\left(t, x(t), c + \int_0^t x(s) ds\right) dw(t), \end{aligned}$$

where  $y(0) = c$  and  $\omega$  is usually omitted, as we will do throughout the paper. The second equation in this system is the stochastic integrodifferential equation of the Ito type, which is a special case of the equation considered in the present paper,

$$\begin{aligned} dx(t) &= F\left(t, x(t), \int_0^t f_1(t, s, x(s)) ds, \int_0^t f_2(t, s, x(s)) dw(s)\right) dt \\ &+ G\left(t, x(t), \int_0^t g_1(t, s, x(s)) ds, \int_0^t g_2(t, s, x(s)) dw(s)\right) dw(t), \\ &t \in [0, T], x(0) = x_0 \text{ a.s.} \end{aligned} \quad (1)$$

Here  $w = (w(t), t \geq 0)$  is a scalar Brownian motion defined on a complete probability space  $(\Omega, \mathcal{F}, \mathcal{P})$  with a natural filtration  $(\mathcal{F}_t, t \geq 0)$  of non-decreasing sub- $\sigma$ -algebras of  $\mathcal{F}$  ( $\mathcal{F}_t = \sigma\{w(s), 0 \leq s \leq t\}$ ),  $x_0$  is a random variable independent of  $w$ , the functions  $F : [0, T] \times R^3 \rightarrow R$ ,  $G : [0, T] \times R^3 \rightarrow R$ ,  $f_i : J \times R \rightarrow R$ ,  $g_i : J \times R \rightarrow R$ ,  $i = 1, 2$ , where  $J = \{(s, t) : 0 \leq s \leq t \leq T\}$ , are assumed to be Borel measurable on their domains. The process  $x = (x(t), t \in [0, T])$  is a strong solution to Eq. (1) provided it is adapted to  $(\mathcal{F}_t, t \geq 0)$ , all the Lebesgue and Ito integrals in the integral form of Eq. (1) are well defined, and Eq. (1) holds a.s. for each  $t \in [0, T]$ . We restricted ourselves to one-dimensional case; the multi-dimensional one is analogous and is not difficult by itself, but involves a complex notation.

Eq. (1) was studied earlier by many authors, first of all by Murge and Pachpatte [10, 11]. A somewhat simpler form of this equation, that is, linear with respect to Lebesgue and Ito integrals, was presented in paper [3] by Berger and Mizel. The basic existence-and-uniqueness theorem under classical conditions was proved in the above cited papers: Let  $E|x_0|^2 < \infty$  and the functions  $F, G, f_i, g_i$ ,  $i = 1, 2$  be globally Lipschitzian and satisfy the linear growth condition, i.e., let there exist a constant  $L > 0$  such that for all  $(t, s) \in J$  and  $(x, y, z), (x', y', z') \in R^3$ ,

$$\begin{aligned} |F(t, x, y, z) - F(t, x', y', z')| &\leq L(|x - x'| + |y - y'| + |z - z'|), \\ |f_i(t, s, x) - f_i(t, s, x')| &\leq L|x - x'|, \quad i = 1, 2 \end{aligned} \quad (2)$$

$$\begin{aligned} |F(t, x, y, z)| &\leq L(1 + |x| + |y| + |z|), \\ |f_i(t, s, x)| &\leq L(1 + |x|), \quad i = 1, 2, \end{aligned} \quad (3)$$

and analogously for  $G, g_1, g_2$ . Then, Eq. (1) has a unique a.s. continuous and  $\mathcal{F}_t$ -adapted solution  $x(t)$  satisfying  $E \sup_{t \in [0, T]} |x(t)|^2 < \infty$ .

The focus of our analysis in the present paper is to study the existence and uniqueness of the solution to Eq. (1) under some non-classical conditions, that is, by using the concept of a random integral contractor which includes the Lipschitz condition as a special case.

## 2 Formulation of the problem and main results

The concept of integral contractors was introduced by Altman [1, 2] for studying some different classes of deterministic equations in Banach spaces. Later, this approach was appropriately extended by Kuo [6] to the notion of random integral contractors for stochastic differential equations of the Ito type, and also for special classes of stochastic integral and integrodifferential equations in various functional spaces. In particular, we highlight papers [4, 5, 6, 8, 9, 13, 12, 14] and especially paper [8] by Mao treating stochastic differential-functional equations with semimartingales. The important fact is that in all these papers, with a partial exception of [9], the considered equations were linear with respect to Lebesgue and Ito integrals, so the regular integral contractor was defined as a solution to any linear functional equation. However, in the present paper we study the non-linear case, which makes it difficult for us to introduce notions and present conditions guaranteeing the existence and uniqueness of the solution to Eq. (1). For that reason, the main aim in this paper is to introduce the notion of a non-linear bounded random integral contractor, so that the non-linearity of the Lebesgue and Ito integrals in Eq. (1) could be exceeded, and then to prove the existence and uniqueness of the solution.

In the remainder, let us denote that  $\mathcal{C}$  is a collection of scalar stochastic processes, defined on  $[0, T]$ , continuous almost surely and adapted to the filtration  $(\mathcal{F}_t, t \geq 0)$ .

For reasons of notational simplicity, let us introduce the following operators: For each  $x \in \mathcal{C}$ ,

$$\begin{aligned} (A_1x)(t) &:= \int_0^t f_1(t, s, x(s)) ds, & (A_2x)(t) &:= \int_0^t f_2(t, s, x(s)) dw(s) \\ (B_1x)(t) &:= \int_0^t g_1(t, s, x(s)) ds, & (B_2x)(t) &:= \int_0^t g_2(t, s, x(s)) dw(s). \end{aligned}$$

Likewise, let us denote that

$$\begin{aligned} F[x(t)] &= F(t, x(t), (A_1x)(t), (A_2x)(t)), \\ G[x(t)] &= G(t, x(t), (B_1x)(t), (B_2x)(t)). \end{aligned}$$

In accordance with this, Eq. (1) can be written in a shorter integral form

$$x(t) = x_0 + \int_0^t F[x(s)] ds + \int_0^t G[x(s)] dw(s), \quad t \in [0, T]. \tag{4}$$

Let

$$\begin{aligned} \Phi : [0, T] \times R^3 &\rightarrow R, & \Gamma : [0, T] \times R^3 &\rightarrow R, \\ \Phi_i : J \times R &\rightarrow R, & \Gamma_i : J \times R &\rightarrow R, \quad i = 1, 2, \end{aligned}$$

be measurable mappings, bounded in the sense that there exist positive constants  $\alpha, \beta, \alpha_i, \beta_i$ ,  $i = 1, 2$ , such that for every  $(t, x, u, v) \in [0, T] \times R^3$ ,  $(t, s, x) \in J \times R$ ,  $y \in R$ ,

$$\begin{aligned} |\Phi(t, x, u, v) \cdot y| &\leq \alpha |y|, & |\Gamma(t, x, u, v) \cdot y| &\leq \beta |y|, \\ |\Phi_i(t, s, x) \cdot y| &\leq \alpha_i |y|, & |\Gamma_i(t, s, x) \cdot y| &\leq \beta_i |y|, \quad i = 1, 2. \end{aligned} \quad (5)$$

We also introduce the following operators: For every  $x, y \in \mathcal{C}$ ,

$$\begin{aligned} ((\tilde{\Phi}_1 x)y)(t) &:= \int_0^t \Phi_1(t, s, x(s)) y(s) ds, \\ ((\tilde{\Phi}_2 x)y)(t) &:= \int_0^t \Phi_2(t, s, x(s)) y(s) dw(s), \\ ((\tilde{\Gamma}_1 x)y)(t) &:= \int_0^t \Gamma_1(t, s, x(s)) y(s) ds, \\ ((\tilde{\Gamma}_2 x)y)(t) &:= \int_0^t \Gamma_2(t, s, x(s)) y(s) dw(s), \end{aligned} \quad (6)$$

and denote that

$$\begin{aligned} \Phi[x(t), y(t)] &= \Phi(t, x(t), ((\tilde{\Phi}_1 x)y)(t), ((\tilde{\Phi}_2 x)y)(t)), \\ \Gamma[x(t), y(t)] &= \Gamma(t, x(t), ((\tilde{\Gamma}_1 x)y)(t), ((\tilde{\Gamma}_2 x)y)(t)). \end{aligned}$$

We are now able to introduce the following non-linear operator  $A$ : For every  $x, y \in \mathcal{C}$ ,

$$\begin{aligned} ((Ax)y)(t) &:= y(t) + \int_0^t \Phi[x(s), y(s)] y(s) ds \\ &\quad + \int_0^t \Gamma[x(s), y(s)] y(s) dw(s), \quad t \in [0, T]. \end{aligned} \quad (7)$$

Clearly,  $(Ax)y \in \mathcal{C}$ .

**Definition 2.1** *Let there exist a positive constant  $K$  such that for every  $x, y \in \mathcal{C}$  the following inequalities hold almost surely:*

$$\begin{aligned} &|F[x(t) - ((Ax)y)(t)] - F[x(t)] + \Phi[x(t), y(t)] \cdot y(t)| \\ &\leq K [||y||_t + |(A_1(x - (Ax)y))(t) - (A_1x)(t) + ((\tilde{\Phi}_1 x)y)(t)| \\ &\quad + |(A_2(x - (Ax)y))(t) - (A_2x)(t) + ((\tilde{\Phi}_2 x)y)(t)|] \\ &|f_i(t, s, x(s) - ((Ax)y)(s)) - f_i(t, s, x(s)) + \Phi_i(t, s, x(s)) \cdot y(s)| \\ &\leq K ||y||_s, \quad i = 1, 2, \\ &|G[x(t) - ((Ax)y)(t)] - G[x(t)] + \Gamma[x(t), y(t)] \cdot y(t)| \\ &\leq K [||y||_t + |(B_1(x - (Ax)y))(t) - (B_1x)(t) + ((\tilde{\Gamma}_1 x)y)(t)| \\ &\quad + |(B_2(x - (Ax)y))(t) - (B_2x)(t) + ((\tilde{\Gamma}_2 x)y)(t)|], \\ &|g_i(t, s, x(s) - ((Ax)y)(s)) - g_i(t, s, x(s)) + \Gamma_i(t, s, x(s)) \cdot y(s)| \\ &\leq K ||y||_s, \quad i = 1, 2, \end{aligned} \quad (8)$$

where  $\|y\|_t = \sup_{0 \leq s \leq t} |y(s)|$ . Then the set of functions  $\{F, f_1, f_2, G, g_1, g_2\}$  has a bounded random integral contractor

$$\left\{ I + \int_0^t \Phi \left( s, x, \int_0^s \Phi_1 dr, \int_0^s \Phi_2 dw(r) \right) ds + \int_0^t \Gamma \left( s, x, \int_0^s \Gamma_1 dr, \int_0^s \Gamma_2 dw(r) \right) dw(s) \right\}. \quad (9)$$

**Definition 2.2** A bounded random integral contractor (9) is said to be regular if the equation

$$(Ax)y = z \quad (10)$$

has a solution  $y$  in  $\mathcal{C}$  for any  $x$  and  $z$  in  $\mathcal{C}$ .

Let  $L_2([0, T] \times \Omega)$  be a collection of stochastic processes in  $\mathcal{C}$  such that  $P\left\{\int_0^T |x(t)|^2 dt < \infty\right\} = 1$ .

**Definition 2.3** The functions  $F$  and  $G$  in Eq. (4) are said to be stochastically closed if for any  $x$  and  $x_n$  in  $\mathcal{C}$ , such that  $x_n \rightarrow x$  and  $F[x_n] \rightarrow y$ ,  $G[x_n] \rightarrow z$  in  $L_2([0, T] \times \Omega)$ , it follows that  $y = F[x]$  and  $z = G[x]$  almost surely, for every  $t \in [0, T]$ .

It is easy to check that if the functions  $F, f_1, f_2, G, g_1, g_2$  satisfy the global Lipschitz condition (2), then  $F$  and  $G$  are stochastically closed and the set  $\{F, f_1, f_2, G, g_1, g_2\}$  has a trivial integral contractor (9) for  $\Phi = \Gamma = \Phi_i = \Gamma_i \equiv 0, i = 1, 2$ . Obviously, the converse also holds. Moreover, if the global Lipschitz condition (2) is valid, let us prove that there exists a class of non-trivial bounded integral contractors, but that the converse assumption does not hold.

First, we can prove that

$$\left\{ I + \int_0^t \Phi \left( s, x, \int_0^s \Phi_1 dr, 0 \right) ds \right\} \quad (11)$$

is a bounded integral contractor for  $\Phi_2 = \Gamma = \Gamma_1 = \Gamma_2 \equiv 0$ . Since the Lipschitz condition (2) and the conditions (5) imply

$$\begin{aligned} & |F[x(t) - ((Ax)y)(t)] - F[x(t)] + \Phi[x(t), y(t)] \cdot y(t)| \\ & \leq |F[x(t) - ((Ax)y)(t)] - F[x(t)]| + |\Phi[x(t), y(t)] \cdot y(t)| \\ & \leq L[|((Ax)y)(t)| + |(A_1(x - (Ax)y))(t) - (A_1x)(t)| \\ & \quad + |(A_2(x - (Ax)y))(t) - (A_2x)(t)|] + \alpha |y(t)| \\ & \leq L[|((Ax)y)|_t + |(A_1(x - (Ax)y))(t) - (A_1x)(t) + ((\tilde{\Phi}_1x)y)(t)| \\ & \quad + |(A_2(x - (Ax)y))(t) - (A_2x)(t)|] + (\alpha + L\alpha_1T) \|y\|_t \quad \text{a.s.}, \end{aligned}$$

then

$$\begin{aligned} \|(Ax)y\|_t & \leq \sup_{0 \leq s \leq t} \left[ |y(s)| + \int_0^s |\Phi[x(r), y(r)] y(r)| dr \right] \\ & \leq (1 + \alpha T) \|y\|_t. \end{aligned} \quad (12)$$

Hence

$$\begin{aligned} & |F[x(t) - ((Ax)y)(t)] - F[x(t)] + \Phi[x(t), y(t)] \cdot y(t)| \\ & \leq K[||y||_t + |(A_1(x - (Ax)y))(t) - (A_1x)(t) + ((\tilde{\Phi}_1x)y)(t)| \\ & \quad + |(A_2(x - (Ax)y))(t) - (A_2x)(t)|]. \end{aligned}$$

that is,  $F$  satisfies (8) with the constant  $K = L(1 + \alpha_1T + \alpha T) + \alpha$ . However,

$$|((\tilde{\Phi}_1x)y)(t)| \leq \int_0^t |\Phi_1(t, s, x(s)) y(s)| ds \leq \alpha_1T ||y||_t \quad \text{a.s.,} \quad \Phi_2 \equiv 0,$$

and thus

$$\begin{aligned} & |f_1(t, s, x(s) - ((Ax)y)(s)) - f_1(t, s, x(s)) + \Phi_1(t, s, x(s)) \cdot y(s)| \\ & \leq L|((Ax)y)(s)| + \alpha_1 |y(s)| \leq [L(1 + \alpha T) + \alpha_1] \cdot ||y||_s \quad \text{a.s.} \end{aligned}$$

Since  $\Gamma = \Gamma_1 = \Gamma_2 \equiv 0$ , all the relations in (8) are satisfied and, therefore, the set of functions  $\{F, f_1, f_2, G, g_1, g_2\}$  has a class of bounded integral contractors (11).

Conversely, if there exists a regular bounded integral contractor (11), it follows from (7) and (10) that the equation

$$y(t) + \int_0^t \Phi[x(s), y(s)] y(s) ds = z(t), \quad t \in [0, T],$$

has a solution  $y \in \mathcal{C}$  for every  $x$  and  $z$  in  $\mathcal{C}$ . Then,

$$|z(t)| \leq |y(t)| + \int_0^t |\Phi[x(s), y(s)] y(s)| ds \leq (1 + \alpha T) ||y||_t \quad \text{a.s.,} \quad t \in [0, T]. \quad (13)$$

Since  $((Ax)y)(t) = z(t)$  a.s., from (8) we derive

$$\begin{aligned} & |F[x(t) - z(t)] - F[x(t)]| \\ & \leq |F[x(t) - z(t)] - F[x(t)] + \Phi[x(t), y(t)] y(t)| + |-\Phi[x(t), y(t)] y(t)| \\ & \leq K[||y||_t + |(A_1(x - z))(t) - (A_1x)(t) + ((\tilde{\Phi}_1x)y)(t)| \\ & \quad + |(A_2(x - z))(t) - (A_2x)(t)|] + \alpha |y(t)| \\ & \leq [K(1 + \alpha_1T) + \alpha][||y||_t + |(A_1(x - z))(t) - (A_1x)(t)| \\ & \quad + |(A_2(x - z))(t) - (A_2x)(t)|] \quad \text{a.s.,} \quad t \in [0, T]. \end{aligned}$$

However, from (13) we see that  $||y||_t$  does not have to be reduced with  $|z(t)|$  a.s., so that Eq. (4) can have a bounded integral contractor, although the Lipschitz condition, in general, does not have to be satisfied. Therefore, the Lipschitz condition and the one based on the integral contractor *cannot, in general, be mutually compared*. This fact could be a motivation to focus our future analysis on conditions and function spaces in order to obtain some alternative existence-and-uniqueness theorems, as well as to establish relations between them.

We now state the following existence-and-uniqueness theorems using the notion of a bounded random integral contractor.

**Theorem 2.1** *Let  $F$  and  $G$  be stochastically closed and  $\int_0^T |F[x_0]|^2 dt < \infty$ ,  $\int_0^T |G[x_0]|^2 dt < \infty$  a.s. Let also the set of functions  $\{F, f_1, f_2, G, g_1, g_2\}$  has a bounded random integral contractor (9). Then Eq. (4) has a solution  $x$  in  $\mathcal{C}$ .*

**Proof.** The proof is based on the following iteration procedures: We define the sequences  $\{x_n(t), n \geq 0\}$  and  $\{y_n(t), n \geq 0\}$  in  $\mathcal{C}$  such that

$$\begin{aligned} x_0(t) &= x_0 \text{ a.s.} \\ x_{n+1}(t) &= x_n(t) - ((Ax_n)y_n)(t) \end{aligned} \tag{14}$$

$$\begin{aligned} &= x_n(t) - y_n(t) - \int_0^t \Phi[x_n(s), y_n(s)] y_n(s) ds \\ &\quad - \int_0^t \Gamma[x_n(s), y_n(s)] y_n(s) dw(s), \\ y_n(t) &= x_n(t) - x_0 - \int_0^t F[x_n(s)] ds - \int_0^t G[x_n(s)] dw(s). \end{aligned} \tag{15}$$

For simplicity, we shall prove this assertion step by step.

Step 1.

$$E\|y\|_t^2 \leq a \frac{A^n t^n}{n!}, \quad 0 \leq t \leq T, \quad n \in N, \tag{16}$$

where  $a$  and  $A$  are some generic constants.

*Proof.* Let us denote that

$$\begin{aligned} a(t) &= F[x_n(t)] - \Phi[x_n(t), y_n(t)] y_n(t) - F[x_{n+1}(t)], \\ a_i(t, s) &= f_i(t, s, x_n(s)) - \Phi_i(t, s, x_n(s)) y_n(s) - f_i(t, s, x_{n+1}(s)), \quad i = 1, 2, \\ b(t) &= G[x_n(t)] - \Gamma[x_n(t), y_n(t)] y_n(t) - G[x_{n+1}(t)], \\ b_i(t, s) &= g_i(t, s, x_n(s)) - \Gamma_i(t, s, x_n(s)) y_n(s) - g_i(t, s, x_{n+1}(s)), \quad i = 1, 2. \end{aligned} \tag{17}$$

Then, from (14) and (15) we have

$$\begin{aligned} y_{n+1}(t) &= x_{n+1}(t) - x_0 - \int_0^t F[x_{n+1}(s)] ds - \int_0^t G[x_{n+1}(s)] dw(s) \\ &= x_n(t) - y_n(t) - \int_0^t \Phi[x_n(s), y_n(s)] y_n(s) ds \\ &\quad - \int_0^t \Gamma[x_n(s), y_n(s)] y_n(s) dw(s) \\ &\quad - x_0 - \int_0^t F[x_{n+1}(s)] ds - \int_0^t G[x_{n+1}(s)] dw(s) \\ &\equiv \int_0^t a(s) ds + \int_0^t b(s) dw(s). \end{aligned} \tag{18}$$

If we take  $x_n$  instead of  $x$  and  $y_n$  instead of  $y$  in (8), we obtain

$$\begin{aligned} |-a(t)| &= |F[x_n(t) - ((Ax_n)y_n)(t)] - F[x_n(t)] + \Phi[x_n(t), y_n(t)] y_n(t)| \\ &\leq K \left[ \|y_n\|_t + \left| -\int_0^t a_1(t, s) ds \right| + \left| -\int_0^t a_2(t, s) dw(s) \right| \right], \end{aligned}$$

and similarly,

$$|-b(t)| \leq K \left[ \|y_n\|_t + \left| -\int_0^t b_1(t, s) ds \right| + \left| -\int_0^t b_2(t, s) dw(s) \right| \right].$$

By applying the usual stochastic integral isometry, Schwarz inequality and Doob inequality [7], we find from (18) that

$$\begin{aligned} E \sup_{0 \leq s \leq t} |y_{n+1}(s)|^2 &\leq 2 \left[ t \int_0^t E|a(s)|^2 ds + 4 \int_0^t E|b(s)|^2 ds \right] \quad (19) \\ &\leq 2K^2 \left\{ 3t \int_0^t E \left[ \|y_n\|_s^2 + \left| -\int_0^s a_1(s, r) dr \right|^2 + \left| -\int_0^s a_2(s, r) dw(r) \right|^2 \right] ds \right. \\ &\quad \left. + 4 \cdot 3 \int_0^t E \left[ \|y_n\|_s^2 + \left| -\int_0^s b_1(s, r) dr \right|^2 + \left| -\int_0^s b_2(s, r) dw(r) \right|^2 \right] ds \right\}. \end{aligned}$$

We can estimate these integrals by using (8) and by applying integration by parts, which yields finally

$$E\|y_{n+1}\|_t^2 = E \sup_{0 \leq s \leq t} |y_{n+1}(s)|^2 \leq A \int_0^t E\|y_n\|_s^2 ds, \quad n \in N,$$

where  $A$  is a generic constant. By repeating integration, it follows that

$$E\|y_n\|_t^2 \leq \frac{A^n}{(n-1)!} \int_0^t (t-s)^{n-1} E\|y_0\|_s^2 ds, \quad n \in N.$$

Since  $y_0(t) = -\int_0^t F[x_0] ds - \int_0^t G[x_0] dw(s)$ , then

$$E\|y_0\|_T^2 \leq 2 \left[ T \int_0^T E|F[x_0]|^2 ds + 4 \int_0^T E|G[x_0]|^2 ds \right] = a,$$

so that

$$E\|y_n\|_t^2 \leq a \frac{A^n t^n}{n!}, \quad t \in [0, T], \quad n \in N,$$

which proves the first step.

Step 2.

$$P\{\|y_{n+1}\|_T > 2^{-n-1}\} \leq c \frac{c_1^n}{(n+1)!}, \quad (20)$$

where  $c$  and  $c_1$  are generic constants.



*Proof.* It follows from (18) that

$$\begin{aligned} & P\{\|y_{n+1}\|_T > 2^{-n-1}\} \\ & \leq P\left\{\int_0^T |a(s)| ds > 2^{-n-2}\right\} + P\left\{\sup_{0 \leq t \leq T} \left|\int_0^t b(s) dw(s)\right| > 2^{-n-2}\right\}. \end{aligned} \quad (21)$$

The application of Chebyshev's inequality and (19) yields

$$\begin{aligned} & P\left\{\int_0^T |a(s)| ds > 2^{-n-2}\right\} \leq 2^{2n+4} E\left(\int_0^T |a(s)| ds\right)^2 \\ & \leq 2^{2n+4} \cdot T \int_0^T E|a(s)|^2 ds \leq 3 \cdot 2^{2n+4} K^2 T \left[ \int_0^T E\|y_n\|_s^2 dt \right. \\ & \quad \left. + K^2 \int_0^T t \int_0^T E\|y_n\|_s^2 ds dt + K^2 \int_0^T \int_0^T E\|y_n\|_s^2 ds dt \right] \\ & \leq c \frac{(4AT)^n}{(n+1)!}, \end{aligned}$$

where  $c$  is a generic constant. The second term on the right-hand side in (21) can be estimated analogously and, therefore, (20) holds.

Step 3. The sequence  $\{x_n\}$  in  $\mathcal{C}$  converges almost surely, uniformly in  $[0, T]$ .

*Proof.* Let us start from (14) and derive that

$$\begin{aligned} \sup_{0 \leq t \leq T} |x_{n+1}(t) - x_n(t)| &= \|x_{n+1} - x_n\|_T \\ &\leq \|y_n\|_T + \int_0^T |\Phi[x_n(s), y_n(s)] y_n(s)| ds \\ &\quad + \left\| \int_0^t \Gamma[x_n(s), y_n(s)] y_n(s) dw(s) \right\|_T. \end{aligned}$$

From the boundedness of the mappings  $\Phi$  and  $\Gamma$  we find that

$$\begin{aligned} & P\left\{\int_0^T |\Phi[x_n(s), y_n(s)] y_n(s)| ds > 2^{-n}\right\} \\ & \leq 2^{2n} T \int_0^T E|\Phi[x_n(s), y_n(s)] y_n(s)|^2 ds \\ & \leq 2^{2n} T \alpha^2 \int_0^T E\|y_n\|_s^2 ds \\ & \leq 2^{2n} T \alpha^2 a \frac{A^n T^{n+1}}{(n+1)!}, \\ & P\left\{\left\| \int_0^t \Gamma[x_n(s), y_n(s)] y_n(s) dw(s) \right\|_T > 2^{-n}\right\} \leq 4 \cdot 2^{2n} \beta^2 a \frac{A^n T^{n+1}}{(n+1)!}. \end{aligned}$$

Thus,

$$\begin{aligned}
& P\{\|x_{n+1} - x_n\|_T > 3 \cdot 2^{-n}\} \\
& \leq P\{\|y_n\|_T > 2^{-n}\} + P\left\{\int_0^T |\Phi[x_n(s), y_n(s)] y_n(s)| ds > 2^{-n}\right\} \\
& \quad + P\left\{\left\|\int_0^t \Gamma[x_n(s), y_n(s)] y_n(s) dw(s)\right\|_T > 2^{-n}\right\} \\
& \leq c \cdot \frac{c_1^n}{n!},
\end{aligned}$$

where  $c$  and  $c_1$  are generic constants. Since

$$\sum_{n=1}^{\infty} P\{\|x_{n+1} - x_n\|_T > 3 \cdot 2^{-n}\} < \infty,$$

the application of Borel-Cantelli's lemma yields that for all large enough  $n$ ,

$$\sup_{0 \leq t \leq T} |x_{n+1}(t) - x_n(t)| \leq 3 \cdot 2^{-n} \text{ almost surely.}$$

Therefore, the sequence  $\{x_n(t)\}$  converges almost surely, uniformly in  $[0, T]$ .

Step 4. Let  $x_\infty$  be a limit of the sequence  $\{x_n\}$ . Then  $x_\infty \in \mathcal{C}$  and  $x_n \rightarrow x_\infty$  in  $\overline{L_2}([0, T] \times \Omega)$ .

*Proof.* It follows from Step 3 that  $x_\infty$  is in  $\mathcal{C}$  since the sequence  $\{x_n\}$  in  $\mathcal{C}$  converges to  $x_\infty$  almost surely. We deduce from (14) that

$$\begin{aligned}
|x_{n+1}(t) - x_n(t)|^2 & \leq 3|y_n(t)|^2 + 3 \left| \int_0^t \Phi[x_n(s), y_n(s)] y_n(s) ds \right|^2 \\
& \quad + 3 \left| \int_0^t \Gamma[x_n(s), y_n(s)] y_n(s) dw(s) \right|^2.
\end{aligned}$$

Then, (16) yields

$$\begin{aligned}
E\|x_{n+1} - x_n\|_t^2 & \leq 3 \left[ E\|y_n\|_t^2 + \alpha^2 t \int_0^t E\|y_n\|_s^2 ds + 4\beta^2 \int_0^t E\|y_n\|_s^2 ds \right] \\
& \leq 3aA^n \left[ \frac{t^n}{n!} + \alpha^2 \frac{t^{n+2}}{(n+1)!} + 4\beta^2 \frac{t^{n+1}}{(n+1)!} \right],
\end{aligned}$$

which implies that

$$\int_0^T E\|x_{n+1} - x_n\|_t^2 dt \rightarrow 0, \quad n \rightarrow \infty,$$

and, therefore,  $x_n \rightarrow x_\infty$  in  $L_2([0, T] \times \Omega)$ .

Step 5. Let  $U_n(t) = \int_0^t F[x_n(s)] ds$ ,  $V_n(t) = \int_0^t G[x_n(s)] dw(s)$ ,  $n \in N$ ,  
 $U(t) = \int_0^t F[x_\infty(s)] ds$ ,  $V(t) = \int_0^t G[x_\infty(s)] dw(s)$ .

Then  $U_n \rightarrow U, V_n \rightarrow V$  in  $L_2([0, T] \times \Omega)$ .

*Proof.* By applying (5) and (16), we find from (17) that

$$\begin{aligned} & E \int_0^T |F[x_{n+1}(t)] - F[x_n(t)]|^2 dt \\ & \leq 2 E \int_0^T |F[x_{n+1}(t)] - F[x_n(t)] + \Phi[x_n(t), y_n(t)] y_n(t)|^2 dt \\ & \quad + 2 E \int_0^T |\Phi[x_n(t), y_n(t)] y_n(t)|^2 dt \\ & \leq 2 \left[ \int_0^T E | - a(t) |^2 dt + \int_0^T \alpha E \|y_n\|_t^2 dt \right] \\ & \leq c \frac{A^n T^{n+1}}{(n+1)!} \rightarrow 0, \quad n \rightarrow \infty, \end{aligned}$$

where  $c$  is a generic constant. Therefore,  $\{F[x_n]\}$  is a Cauchy sequence in  $L_2([0, T] \times \Omega)$ , which implies that  $F[x_n] \rightarrow F[x_\infty]$  in  $L_2([0, T] \times \Omega)$  since  $x_n \rightarrow x_\infty$  in  $L_2([0, T] \times \Omega)$  and since the coefficients of Eq. (4) are stochastically closed in the sense of Definition 2.2. Hence

$$\begin{aligned} \int_0^T E |U_n(t) - U(t)|^2 dt &= \int_0^T E \left| \int_0^t (F[x_n(s)] - F[x_\infty(s)]) ds \right|^2 dt \\ &\leq T^2 \int_0^T E |F[x_n(t)] - F[x_\infty(t)]|^2 dt \rightarrow 0, \quad n \rightarrow \infty, \end{aligned}$$

and, therefore,  $U_n \rightarrow U$  in  $L_2([0, T] \times \Omega)$ .

Since  $V_n \rightarrow V$  in  $L_2([0, T] \times \Omega)$  is based on the same computation, the proof of Step 5 becomes complete.

Finally, we will complete the proof of Theorem 2.1. By taking  $L_2([0, T] \times \Omega)$  limits on the both sides in (15) and by applying the conclusions of Steps 1, 4 and 5, we observe for all  $t \in [0, T]$  that

$$x_\infty(t) = \int_0^t F[x_\infty(s)] ds + \int_0^t G[x_\infty(s)] dw(s) \quad \text{a.s.} \tag{22}$$

Consequently,  $x_\infty$  is the solution to Eq. (4) since the processes on the both sides of (22) are continuous almost surely for all  $t \in [0, T]$ , which completes the proof.  $\square$

**Theorem 2.2** *Let the functions  $F, f_1, f_2, G, g_1, g_2$  satisfy the assumptions of Theorem 2.1 and the bounded random integral contractor (9) is regular. Then the solution  $x$  to Eq. (4) in  $\mathcal{C}$  is unique.*

**Proof.** Let  $x_1$  and  $x_2$  be two solutions to Eq. (4). Since the bounded random integral contractor (9) is regular, the equation

$$(Ax_1)y = x_1 - x_2$$

has a solution  $y \in \mathcal{C}$ , that is,

$$\begin{aligned} y(t) + \int_0^t \Phi[x_1(s), y(s)] y(s) ds + \int_0^t \Gamma[x_1(s), y(s)] y(s) dw(s) \\ = x_1(t) - x_2(t) \\ = \int_0^t (F[x_1(s)] - F[x_2(s)]) ds + \int_0^t (G[x_1(s)] - G[x_2(s)]) dw(s). \end{aligned} \quad (23)$$

Therefore,

$$\begin{aligned} y(t) = - \int_0^t (F[x_2(s)] - F[x_1(s)] + \Phi[x_1(s), y(s)] y(s)) ds \\ - \int_0^t (G[x_2(s)] - G[x_1(s)] + \Gamma[x_1(s), y(s)] y(s)) dw(s). \end{aligned} \quad (24)$$

Since  $E\|y\|_t^2$  is generally not finite, we use the truncation:

$$I_N(t) = \begin{cases} 1, & \|y\|_s \leq N, \quad 0 \leq s \leq t, \\ 0, & \text{otherwise.} \end{cases}$$

Then  $I_N \in \mathcal{C}$  and  $I_N(t) = I_N(t) \cdot I_N(s) \cdot I_N(r)$  for  $0 \leq r \leq s \leq t \leq T$ . Now, (24) implies

$$\begin{aligned} EI_N(t)\|y\|_t^2 \\ \leq 2 \left[ t E \int_0^t I_N(s) |F[x_2(s)] - F[x_1(s)] + \Phi[x_1(s), y(s)] y(s)|^2 ds \right. \\ \left. + 4 E \int_0^t I_N(s) |G[x_2(s)] - G[x_1(s)] + \Gamma[x_1(s), y(s)] y(s)|^2 ds \right]. \end{aligned}$$

From (8) we find that

$$\begin{aligned} |F[x_2(s)] - F[x_1(s)] + \Phi[x_1(s), y(s)] y(s)| \\ \leq K [ \|y\|_s + |(A_1 x_2)(s) - (A_1 x_1)(s) + ((\tilde{\Phi}_1 x_1)y)(s)| \\ + |(A_2 x_2)(s) - (A_2 x_1)(s) + ((\tilde{\Phi}_2 x_1)y)(s)| ], \end{aligned} \quad (25)$$

and similarly for  $G$ . From now on, we observe that

$$\begin{aligned} EI_N(t)\|y\|_t^2 \\ \leq 6K^2 \left\{ t E \int_0^t I_N(s) [ \|y\|_s^2 + |(A_1 x_2)(s) - (A_1 x_1)(s) + ((\tilde{\Phi}_1 x_1)y)(s)|^2 \right. \\ \left. + |(A_2 x_2)(s) - (A_2 x_1)(s) + ((\tilde{\Phi}_2 x_1)y)(s)|^2 ] ds \right. \\ \left. + 4E \int_0^t I_N(s) [ \|y\|_s^2 + |(B_1 x_2)(s) - (B_1 x_1)(s) + ((\tilde{\Gamma}_1 x_1)y)(s)|^2 \right. \\ \left. + |(B_2 x_2)(s) - (B_2 x_1)(s) + ((\tilde{\Gamma}_2 x_1)y)(s)|^2 ] ds \right\}. \end{aligned}$$

To estimate the right-hand side in the previous relation, we will proceed analogously to (19). By omitting details, we obtain finally

$$\begin{aligned}
EI_N(t)\|y\|_t^2 &\leq 6K^2 \left\{ t \int_0^t EI_N(s)\|y\|_s^2 ds \right. \\
&\quad + 2K^2 t \int_0^t I_N(s) \int_0^s EI_N(r)\|y\|_r^2 dr ds \\
&\quad + 4 \int_0^t EI_N(s)\|y\|_s^2 ds \\
&\quad \left. + 8K^2 \int_0^t I_N(s) \int_0^s EI_N(r)\|y\|_r^2 dr ds \right\} \\
&\leq c \int_0^t (1+t-s)EI_N(s)\|y\|_s^2 ds \\
&\leq c(1+T) \int_0^t EI_N(s)\|y\|_s^2 ds,
\end{aligned}$$

where  $c$  is a constant. To close the proof, we apply the well-known the Gronwall-Bellman lemma and conclude that  $EI_N(t)\|y\|_t^2 = 0$  for all  $t \in [0, T]$ . The application of the Lebesgue monotone convergence theorem implies that  $E\|y\|_t^2 = \lim_{N \rightarrow \infty} EI_N(t)\|y\|_t^2 = 0$ ,  $t \in [0, T]$  and, therefore,  $y(t) = 0$  almost surely for all  $t \in [0, T]$ . Finally, from (23) it follows that  $x_1 = x_2$  almost surely, which completes the proof.  $\square$

## References

- [1] M. Altman, *Inverse differentiability contractors and equations in Banach space*, *Studia Mathematica* **46** 1–15, (1973).
- [2] M. Altman, *Contractors and contractor directions*, Marcel Dekker, New York, 1978.
- [3] M.A. Berger, and V.J. Mizel, *Volterra equations with Ito integrals I*, *J. of Integral Equations* 2 (1980), 187–245.
- [4] M. Jovanović, S. Janković, *On a class of nonlinear stochastic hereditary integrodifferential equations*, *Indian J. Pure Appl. Math.* **28** (8), 1062–1082, (1997).
- [5] M. Jovanović, S. Janković, *Existence and uniqueness problems for nonlinear stochastic hereditary integrodifferential equations*, *Indian J. Pure Appl. Math.*, **32** (5), 695-710, (2001).
- [6] H.H. Kuo, *On integral contractors*, *J. of Integral Equations* **1**, 35–46, (1979).
- [7] G. Ladde and V. Lakshmikantham, *Random Differential Inequalities*, Academic Press, New York (1980).

- [8] X. Mao, *Solutions of stochastic differential-functional equations via bounded stochastic integral constructors*, J. Theoretical Probability, 5(3) (1992), 487–502.
- [9] M.G. Murge, and B.G. Pachpatte, *Existence and uniqueness of solutions of nonlinear Ito type stochastic integral equations*, Math. Acta Scientia, 7(1987), 2, 207–216.
- [10] M.G. Murge, B.G. Pachpatte, *On generalized Itô type stochastic integral equation*, Yokohama Mathematical Journal, **34** (1986), 23–33.
- [11] M.G. Murge, B.G. Pachpatte, *Explosion and asymptotic behavior of nonlinear Ito type stochastic integrodifferential equations*, Kodai Math. J., **9** (1986), 1–18.
- [12] W.J. Padgett, A.N.V. Rao, *Solution of a stochastic integral equation using integral constructors*, Inform. and Control, 41, (1979), 56–66.
- [13] V. Popescu, and A. Rascanu, *On bounded deterministic and stochastic integral constructors*, Anale stiintifice ale Universitatii "Al. I. Cuza", Iasi, Tomul XXXIV. s. I a. Mathematica, (1988), f. 1, 37–51.
- [14] B.G. Zhang, and W.J. Padgett, *The existence and uniqueness of solutions to stochastic differential-difference equations*, Stochastic Analysis and Appl., 2(3)(1984), 335–345.

Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš, Serbia

*E-mail*

Miljana Jovanović: [mima@pmf.ni.ac.rs](mailto:mima@pmf.ni.ac.rs)

Svetlana Janković: [svjank@pmf.ni.ac.rs](mailto:svjank@pmf.ni.ac.rs)