

On Sufficient Degree Conditions for a Graph to be k -linked

KEN-ICHI KAWARABAYASHI¹,
ALEXANDR KOSTOCHKA² and GEXIN YU^{3†}

¹Graduate School of Information Sciences (GSIS), Tohoku University,
Aramaki aza Aoba 09, Aoba-ku Sendai, Miyagi, 980-8579, Japan
(e-mail: k.keniti@dais.is.tohoku.ac.jp)

²Department of Mathematics, University of Illinois, Urbana, IL 61801, USA
and
Institute of Mathematics, Novosibirsk 630090, Russia
(e-mail: kostochk@math.uiuc.edu)

³Department of Mathematics, University of Illinois, Urbana, IL 61801, USA
(e-mail: gexinyu@uiuc.edu)

Received 28 November 2003; revised 3 March 2004

A graph is k -linked if for every list of $2k$ vertices $\{s_1, \dots, s_k, t_1, \dots, t_k\}$, there exist internally disjoint paths P_1, \dots, P_k such that each P_i is an s_i, t_i -path. We consider degree conditions and connectivity conditions sufficient to force a graph to be k -linked.

Let $D(n, k)$ be the minimum positive integer d such that every n -vertex graph with minimum degree at least d is k -linked and let $R(n, k)$ be the minimum positive integer r such that every n -vertex graph in which the sum of degrees of each pair of non-adjacent vertices is at least r is k -linked. The main result of the paper is finding the exact values of $D(n, k)$ and $R(n, k)$ for every n and k .

Thomas and Wollan [14] used the bound $D(n, k) \leq (n + 3k)/2 - 2$ to give sufficient conditions for a graph to be k -linked in terms of connectivity. Our bound allows us to modify the Thomas–Wollan proof slightly to show that every $2k$ -connected graph with average degree at least $12k$ is k -linked.

1. Introduction

Dirac [2] proved that every n -vertex graph G with minimum degree at least $n/2$ is Hamiltonian, and Ore [12] observed that the condition $\delta(G) \geq n/2$ in Dirac's result can be replaced by ' $\sigma_2(G) \geq n$ ', where $\sigma_2(G)$ is the minimum value of the sum $\deg(u) + \deg(v)$ over all pairs $\{u, v\}$ of non-adjacent vertices in G .

[†] This work was partially supported by the NSF grant DMS-0099608 and by the Japan Society for the Promotion of Science for Young Scientists.

After Chartrand introduced the notion of *k-ordered graphs*, that is, graphs in which for every ordered sequence of *k* vertices there is a cycle that encounters the vertices of the sequence in the given order, several authors (see, e.g., [4, 11, 8, 6, 5]) studied the analogue of Dirac’s and Ore’s sufficient conditions for a graph to be *k-ordered*. Let $D_0(n, k)$ denote the minimum positive integer d such that every n -vertex graph with minimum degree at least d is *k-ordered*. Similarly, let $R_0(n, k)$ denote the minimum positive integer r such that every n -vertex graph G with $\sigma_2(G) \geq r$ is *k-ordered*. Improving on results in [4, 11], it was shown in [6] that $R_0(n, k) = n + \lceil (3k - 9)/2 \rceil$ for every $3 \leq k \leq n/2$. Furthermore, Kierstead, Sárkőzy and Selkow [8] showed that $D_0(n, k) = \lceil n/4 \rceil + \lfloor k/2 \rfloor - 1$ for $3 \leq k \leq (n + 3)/11$. These bounds demonstrate the interesting phenomenon that $R_0(n, k) > 2D_0(n, k)$ for k small with respect to n .

A graph is *k-linked* if, for every list of $2k$ vertices $\{s_1, \dots, s_k, t_1, \dots, t_k\}$, there exist internally disjoint paths P_1, \dots, P_k such that each P_i is an s_i, t_i -path. It is a folklore observation that if the number n of vertices of a graph G is at least $2k$, then in the definition of a *k-linked* graph it is enough to consider only the lists of *distinct* $s_1, \dots, s_k, t_1, \dots, t_k$. As in the previous paragraph, let $D(n, k)$ be the minimum positive integer d such that every n -vertex graph with minimum degree at least d is *k-linked*. Also, let $R(n, k)$ denote the minimum positive integer r such that every n -vertex graph G with $\sigma_2(G) \geq r$ is *k-linked*. Thomas and Wollan [14] used the bound $D(n, k) \leq (n + 3k)/2 - 2$ to give sufficient conditions for a graph to be *k-linked* in terms of connectivity. In this paper we determine the exact values of $D(n, k)$ and $R(n, k)$ for all n and k .

Theorem 1.1. *If $k \geq 2$, then*

$$R(n, k) = \begin{cases} 2n - 3, & n \leq 3k - 1, \\ \lfloor \frac{2(n+5k)}{3} \rfloor - 3, & 3k \leq n \leq 4k - 2, \\ n + 2k - 3, & n \geq 4k - 1, \end{cases} \tag{1.1}$$

and

$$D(n, k) = \left\lceil \frac{R(n, k)}{2} \right\rceil = \begin{cases} n - 1, & n \leq 3k - 1, \\ \lfloor \frac{n+5k}{3} \rfloor - 1, & 3k \leq n \leq 4k - 2, \\ \lceil \frac{n-3}{2} \rceil + k, & n \geq 4k - 1. \end{cases} \tag{1.2}$$

Note that $R(3k, k) < R(3k - 1, k)$. This is the only place for a fixed k where $R(n, k)$ decreases with growing n .

Egawa, Faudree, Győri, Ishigami, Schelp and Wang [3] proved the following very closely related result.

Theorem 1.2. ([3]) *Let $k \geq 2$ and $n \geq 3k$. Let $D_1(n, k)$ be the minimum positive integer d such that, for every n -vertex graph G with minimum degree at least d and every matching $M = \{s_i t_i \mid i = 1, \dots, m\}$ of size $m \leq k$ in G , there exist vertex-disjoint cycles C_1, \dots, C_m such that C_i contains $s_i t_i$ for each $i = 1, \dots, m$. Similarly, let $R_1(n, k)$ be the minimum positive*

integer r such that, for every n -vertex graph G with $\sigma_2(G) \geq r$ and every matching $M = \{s_i t_i \mid i = 1, \dots, m\}$ of size $m \leq k$ in G , there exist vertex-disjoint cycles C_1, \dots, C_m such that C_i contains $s_i t_i$ for each $i = 1, \dots, m$. Then

$$R_1(n, k) = \begin{cases} \lfloor \frac{n}{2} \rfloor + 4k - 2, & 3k \leq n \leq 4k - 2, \\ n + 2k - 2, & n \geq 4k - 1, \end{cases} \tag{1.3}$$

and

$$D_1(n, k) = \begin{cases} \lceil \frac{n+5k}{3} \rceil - 1, & 3k \leq n \leq 4k - 2, \\ \lceil \frac{n}{2} \rceil + k - 1, & n \geq 4k - 1. \end{cases} \tag{1.4}$$

This is closely related because, for a graph G and a matching $M = \{s_i t_i \mid i = 1, \dots, m\}$, the existence of cycles provided by Theorem 1.2 is equivalent to the existence in $G' = G - M$ of vertex-disjoint paths linking s_i with t_i provided by Theorem 1.1. Although the graphs G and G' differ only by a matching, the values of $R(n, k)$ and $R_1(n, k)$ for $3k < n < 4k$ differ significantly. On the other hand, the ideas of the proofs are similar. But neither of the bounds of Theorem 1.2 and Theorem 1.1 can be derived from the other. Also, in terms of linkages, Theorem 1.1 gives slightly better bounds for some parities of n and k , which could perhaps be used for extremal problems on linkages.

A very interesting problem is estimating $f(k)$ – the minimum positive integer f such that every f -connected graph is k -linked. After a series of papers by Jung [7], Larman and Mani [9], Mader [10], and Robertson and Seymour [13], the first linear upper bound for f , namely $f(k) \leq 22k$, was proved by Bollobás and Thomason [1]. Very recently, Thomas and Wollan [14] improved this bound to $f(k) \leq 16k$. If one were to use Theorem 1.1 in the Thomas–Wollan proof [14], then their sufficient condition for a graph to be k -linked could be relaxed.

Theorem 1.3. *Every $2k$ -connected graph $G = (V, E)$ with $|E| \geq 6k|V|$ is k -linked. In particular, every $12k$ -connected graph is k -linked.*

We note that applying Theorem 1.2 also would yield Theorem 1.3. In the next section we prove lower bounds for $D(n, k)$ and $R(n, k)$. Then, in Section 3, the upper bounds are established. In the final section, we show how to modify the Thomas–Wollan proof [14] in order to derive Theorem 1.3.

Using new ideas (in particular, ideas of this paper), Thomas and Wollan improved the upper bound on $f(k)$ further to $f(k) \leq 10k$.

2. Constructions

In this section we present examples giving lower bounds for $D(n, k)$ and $R(n, k)$. Consider several cases.

Case 1: $n \leq 3k - 1$. Let G be K_n with a deleted matching $\{(v_1, v_2), \dots, (v_{2\lfloor n/2 \rfloor - 1}, v_{2\lfloor n/2 \rfloor})\}$. Clearly, $\delta(G) = n - 2$ and $\sigma_2(G) = 2n - 4$. Let $m = \min\{k, \lfloor n/2 \rfloor\}$. For $i = 1, \dots, m$, let $s_i = v_{2i-1}$ and $t_i = v_{2i}$. Assume that there exist internally disjoint paths P_1, \dots, P_k such




Figure 1. Graph G for Case 2.

that each P_i is an s_i, t_i -path. Denote $S = \{s_1, \dots, s_m, t_1, \dots, t_m\}$. For every i , the path P_i contains a vertex $x_i \notin S$, since $s_i t_i \notin E(G)$. Therefore, $n \geq |S| + m = 3m$. This is impossible if $n \leq 3k - 1$, $k \geq 2$, and $m = \min\{k, \lfloor n/2 \rfloor\}$.

Case 2: $3k \leq n \leq 4k - 2$. Let $x = \lceil \frac{2n-5k}{3} \rceil + 1$. It is easier to describe the complement $\overline{G} = (V, \overline{E})$ of G . Fix six disjoint subsets $S_0, S_1, S_2, T_0, T_1, T_2$ of V with $|S_0| = |T_0| = x - 1$, $|S_1| = |T_1| = x$, and $|S_2| = |T_2| = k - x$. Let $V_0 = V - S_0 - S_1 - S_2 - T_0 - T_1 - T_2$.

Looking ahead, we may assume that

$$\begin{aligned} S_1 &= \{s_1, \dots, s_x\}, & S_2 &= \{s_{x+1}, \dots, s_k\}, \\ T_1 &= \{t_1, \dots, t_x\}, & T_2 &= \{t_{x+1}, \dots, t_k\}. \end{aligned}$$

The set \overline{E} of edges of \overline{G} is $E_1 \cup E_2 \cup E_3$, where

$$\begin{aligned} E_1 &= \{s_i t_i : i = 1, \dots, k\}, \\ E_2 &= \{vw : v \in S_0, w \in T_1\}, \\ \text{and } E_3 &= \{vw : v \in T_0, w \in S_1\}. \end{aligned}$$

The graph G itself is drawn in Figure 1.

Assume that there exist internally disjoint paths P_1, \dots, P_k such that each P_i is an s_i, t_i -path. As in Case 1, each P_i contains a vertex in $S_0 \cup T_0 \cup V_0$. Moreover, if $i \leq x$ (i.e., $s_i \in S_1$ and $t_i \in T_1$), then either P_i contains a vertex in V_0 , or it has at least two internal vertices, since no vertex in $S_0 \cup T_0$ is adjacent to both s_i and t_i . Therefore, $n \geq 2k + k + (x - |V_0|) = 5k - n - 2 + 3x$. By the definition of x , the last expression exceeds n , a contradiction.

The vertices in $S_0 \cup S_1 \cup T_0 \cup T_1$ have degree x in \overline{G} and all other vertices have degree at most 1. It follows that $\delta(G) = n - 1 - x = \lfloor \frac{n+5k}{3} \rfloor - 2$ and $\sigma_2(G) = 2n - 2 - 2x = 2 \lfloor \frac{n+5k}{3} \rfloor - 4$. This proves the lower bound on $D(n, k)$ for every $3k \leq n \leq 4k - 2$ and on $R(n, k)$ for $3k \leq n \leq 4k - 2$ such that $n + 5k \not\equiv 2 \pmod{3}$. For $n + 5k \equiv 2 \pmod{3}$, we slightly modify the construction: we change x to $x = \lceil \frac{2n-5k+2}{3} \rceil$ and move one vertex from V_0 to T_0 , so that $|T_0| = x$. Then, as in the previous paragraph, we obtain $n \geq 2k + k + (x - |V_0|) = 5k - n - 2 + 3x + 1$, which contradicts the new definition of x . On the other




Figure 2. Graph G for Case 3.

hand, $\sigma_2(G) = 2n - 3 - 2x = 2\lfloor \frac{n+5k-2}{3} \rfloor - 3$. Since we have $2\lfloor \frac{n+5k-2}{3} \rfloor - 3 = \lfloor \frac{2(n+5k)}{3} \rfloor - 4$, for $n + 5k \equiv 2 \pmod{3}$, this completes Case 2.

Case 3: $n \geq 4k - 1$. This is a standard example of a graph with connectivity $2k - 2$ (see Figure 2). Clearly, $\delta(G) = \lfloor n/2 \rfloor + k - 2$ and $\sigma_2(G) = n + 2k - 4$. If $s_1, \dots, s_{k-1}, t_1, \dots, t_{k-1}$ are the vertices of the central K_{2k-2} , s_k belongs to the left $K_{\lfloor n/2 \rfloor - k + 1}$, and t_k belongs to the right $K_{\lfloor n/2 \rfloor - k + 1}$, then there is no corresponding linkage: there is simply no room for an s_k, t_k -path.

3. Upper bounds in Theorem 1.1

Observe that it is enough to prove the upper bound for $R(n, k)$, since $D(n, k) \geq R(n, k)/2$. The case of $n \leq 3k - 1$ is obvious, because $\sigma_2(G) \geq 2n - 3$ means that $G = K_n$.

Remark 1. If $n \leq 4k - 1$, then $n + 2k - 4 \leq \frac{2n+10k-1}{3} - 4$.

Let $n \geq 3k$ and $G = (V, E)$ be a graph on n vertices satisfying the conditions of the theorem. Let $M_i = \{s_i, t_i\}$, $i = 1, \dots, k$, be arbitrary disjoint vertex pairs to be linked, and let $M = \cup_{i=1}^k M_i$. If, say, $s_k t_k \in E(G)$, then, for the graph $G' = G - s_k - t_k$ with $n' = n - 2$ vertices and for $k' = k - 1$, we have $\sigma_2(G') \geq R(n, k) - 4 \geq R(n', k')$. Therefore, if the theorem holds for G' , it also holds for G . Thus, we may assume that none of the pairs $s_i t_i$ is an edge in G .

We will find the linkage in 3 steps (resembling the steps of the main proof in [6]). On each of the steps, if we cannot perform this step, then $\sigma_2(G) < R(n, k)$.

Construct the auxiliary bipartite graph H with partite sets W_1 and W_2 as follows. Let $W_1 = \{M_1, \dots, M_k\}$, $W_2 = V - M$, and a pair (M_i, v) be an edge in H if $v \in N_G(s_i) \cap N_G(t_i)$. If H has a matching saturating W_1 , then this matching gives the required linkage. Otherwise, let m be the size of a maximum matching in H .

By the König–Egerváry theorem, there is a $Q \subseteq W_1$ with $k - m = |Q| - |N_H(Q)|$. Denote $R = N_H(Q)$ and $S = V(G) - M - R$. We may assume that $Q = \{M_i : i = 1, \dots, q\}$. Let $Q' = \cup_{i=1}^q \{s_i, t_i\}$ (the elements of Q are pairs, and the elements of Q' are all the elements of these pairs). Note that $|Q| = q$, $|Q'| = 2q$, $|R| = q - k + m$, and $|S| = n - k - q - m$.

Let P be a maximum matching in H . We may assume that only vertices in $D = \{M_1, \dots, M_{k-m}\}$ are not covered by this matching. Let $D' = \{s_i, t_i : 1 \leq i \leq k - m\}$.

Consider the linkage \mathcal{P} of $M - D$ corresponding to P . Let Z be the set of vertices not participating in the linkage.

Lemma 3.1. $n \geq 4k - m$.

Proof. We need to prove this lemma only for $n \leq 4k - 1$.

Assume $n \leq 4k - m - 1$. Let $M_i \in Q$ and $x \in S$. Then either s_i or t_i is not a neighbour of x . Therefore,

$$\deg(s_i) + \deg(t_i) \leq 2(n - 2) - |S| = n + k - 4 + q + m. \tag{3.1}$$

For the same reasons, x is not adjacent to at least q vertices in Q' . We may assume that $x s_1 \notin E(G)$. If $s_1 y \notin E(G)$ for every $y \in S$, then

$$\deg(s_1) + \deg(x) \leq (n - 2 - |S|) + (n - 1 - q) = n - 3 + k + m \leq n - 4 + 2k.$$

By Remark 1, this contradicts (1.1). Otherwise, there is a $y \in S$ with $y s_1 \in E(G)$ and, therefore, $y t_1 \notin E(G)$. Thus, by (3.1),

$$\begin{aligned} 3\sigma_2(G) &\leq (\deg(x) + \deg(s_1)) + (\deg(y) + \deg(t_1)) + (\deg(s_1) + \deg(t_1)) \\ &\leq 2(n + k - 4 + q + m) + 2(n - q - 1) \leq 2(n + 5k - 6). \end{aligned}$$

It follows that $\sigma_2(G) < R(n, k)$. □

By Lemma 3.1, for every $i = 1, \dots, k - m$, we can assign a vertex $z_i \in Z$ to s_i and a vertex $z'_i \in Z$ to t_i so that we assign distinct members of Z to distinct vertices. Also, for every $k - m + 1 \leq i \leq k$, let y_i be the common neighbour of s_i and t_i corresponding to the matching P above. Among such assignments, choose an assignment \mathcal{A} with as many edges $z_i s_i$ and $z'_i t_i$ as possible. Let $Z' = \bigcup_{i=1}^{k-m} \{z_i, z'_i\}$.

Lemma 3.2. *In \mathcal{A} , every z_i is adjacent to s_i and every z'_i is adjacent to t_i .*

Proof. Assume that $s_1 z_1 \notin E(G)$. We will prove that

$$|N(s_1) - M| \leq |M - N(z_1)|. \tag{3.2}$$

To do this, for every neighbour w of s_1 outside M we show a non-neighbour $f(w)$ of z_1 in M . First, observe that either $w \in Z'$ or w was used in the linkage \mathcal{P} , since otherwise we can assign w as z_1 .

Case 1. If w is used in a path s_i, w, t_i in the linkage \mathcal{P} and z_1 is adjacent to both s_i and t_i , then by swapping w with z_1 , we will get an assignment with new z_1 (former w) adjacent to s_1 . Thus, either s_i or t_i can be chosen as $f(w)$.

Case 2. If $w \in Z'$, say, $w = z'_i$ (possibly, $i = 1$), and z_1 is adjacent to t_i , then swapping z_1 with w produces a better assignment. Thus, $z_1 t_i \notin E$, and we let $f(w) = t_i$.

Since all s_i and t_i are disjoint, (3.2) holds, and therefore $\deg(s_1) + \deg(z_1) \leq 2(n - 2) - |V - M| = n - 4 + 2k$. By Remark 1, this yields $\sigma_2(G) < R(n, k)$ for each $n \geq 3k$. □

The last step in the proof is given by the next lemma.

Lemma 3.3. *The assignment \mathcal{A} in Lemma 3.2 can be chosen in such a way that every z_i is adjacent to every z'_i .*

Proof. Consider an assignment \mathcal{A} satisfying Lemma 3.2. For $i = 1, \dots, k - m$, let $X_i = \{s_i, t_i, z_i, z'_i\}$, and for $i = k - m + 1, \dots, k$, let $X_i = \{s_i, t_i, y_i\}$. Let $X = \bigcup_{i=1}^k X_i$. Choose \mathcal{A} in Lemma 3.2 so that as many as possible z_i are adjacent to corresponding z'_i . Suppose that the lemma does not hold. Then we may renumber (s_i, t_i) so that $z_1 z'_1 \notin E(G)$. Let $A = N(s_1) \cap (V(G) - X)$ and $B = N(t_1) \cap (V(G) - X)$.

Note that $N(A + z_1) \cap (B + z'_1) = \emptyset$. For $i = 1, \dots, k$, let k_i denote the number of neighbours of X_1 (with multiplicities) in X_i . Since each member of X_1 has exactly one neighbour in X_i , $k_1 = 4$.

Claim 3.4. *If $2 \leq i \leq k - m$, then $k_j \leq 12$. If $k - m + 1 \leq i \leq k$, then $k_j \leq 10$.*

Proof.

Case 1: $2 \leq i \leq k - m$. By the maximality of m , neither of z_1, z'_1 is a common neighbour of s_i and t_i and neither of z_i, z'_i is a common neighbour of s_1 and t_1 . Therefore, $k_i \leq |X_1| \cdot |X_i| - 4 = 12$.

Case 2: $k - m + 1 \leq i \leq k$.

Subcase 2.1: $s_1 y_i, t_1 y_i \in E(G)$. If some of z_1, z'_1 (say, z_1) is a common neighbour of s_i and t_i , then assigning z_1 as the new y_i and assigning the old y_i as the new y_1 will contradict the maximality of m . Otherwise, $k_i \leq |X_1| \cdot |X_i| - 2 = 3 \cdot 4 - 2 = 10$.

Subcase 2.2: $s_1 y_i \notin E(G)$. If all the four edges $z'_1 s_i, z'_1 t_i, z_1 y_i, t_1 y_i$ are in $E(G)$, then we can swap y_i with z'_1 and get a better assignment, since the new z'_1 is adjacent to z_1 . Otherwise, we again have $k_i \leq 10$. □

Claim 3.5. *For each $v \notin X$, $|N(v) \cap \{s_1, t_1, z_1, z'_1\}| \leq 2$.*

Proof. Otherwise, we can swap v with either z_1 or z'_1 so that the new assignment is better. □

Let $F = \deg(s_1) + \deg(t_1) + \deg(z_1) + \deg(z'_1)$. In view of the claims above, $F \leq 2(n - |X|) + 4 + 12(k - m - 1) + 10m$. Since $|X| = 4(k - m) + 3m$, we obtain

$$F \leq 2(n - 4k + m) + 4 + 12(k - m - 1) + 10m = 2n + 4k - 8.$$

Since $F \geq 2\sigma_2(G) \geq 4\delta(G)$, we obtain a contradiction to both (1.1) and (1.2). This proves the lemma, and thus the theorem as well. □

4. Connectivity conditions

Thomas and Wollan [14] showed that $f(k) \leq 16k$ by proving the following (stronger) result.

Theorem 4.1. ([14]) *Every $2k$ -connected graph G with $|E(G)| \geq 8k|V(G)|$ is k -linked.*

Most of the proof works under weaker restrictions on the average degree. The bottleneck for the bounds in Theorem 4.1 is the following claim.

Claim 4.2. ([14]) *For $\alpha = 8$, every graph H on at most $2\alpha k$ vertices with minimum degree at least αk contains a k -linked subgraph.*

If one proves Claim 4.2 for any $3 \leq \alpha < 8$, this would imply the strengthening of Theorem 4.1 with α in place of 8. Using Theorem 1.1, we prove below an analog of Claim 4.2 with $\alpha = 6$. This will make Theorem 4.1 work with 6 in place of 8, *i.e.*, will yield Theorem 1.3. The beginning of the proof is reminiscent of that for Claim 4.2, but for completeness, we present the full proof.

Lemma 4.3. *For $\alpha = 6$, every graph H on at most $2\alpha k$ vertices with minimum degree at least αk contains a k -linked subgraph.*

Proof. Consider an H satisfying the conditions of the lemma. If H itself is not k -linked, then there is a set $X = \{s_i, t_i : 1 \leq i \leq k\} \subseteq V(H)$ such that there are no disjoint paths P_1, \dots, P_k such that each P_i is an s_i, t_i -path and all s_i and t_i are distinct. Link as many as possible pairs (s_i, t_i) by paths of length at most 6 and, subject to this, minimize the sum of the lengths of these paths. Suppose that l_1 pairs are not linked and the number of paths of length i is l_{i+1} , $1 \leq i \leq 6$. We may assume that s_1 and t_1 are not linked.

Let S be the union of X with vertex sets of all the paths of the linkage. Let $A = N_H(s_1) - S$ and $B = N_H(t_1) - S$. By the choice, A and B are disjoint and are at distance at least 5 in $H - S$. Since the paths P_i are chosen to be of the minimum total length, we have

$$|N_H(v) \cap V(P_i)| \leq 3 \quad \forall v \in V(G) - S \quad \forall P_i, \tag{4.1}$$

$$|N_H(v) \cap \{s_1, t_1\}| \leq 1 \quad \forall v \in V(G) - S, \tag{4.2}$$

and

$$|N_H(s_1) \cap N_H(t_1) \cap V(P_i)| \leq 3 \quad \forall i, \tag{4.3}$$

Claim 4.4. *For each $v \in V(H) - S - A - B$, $|N(v) \cap (A \cup B)| \geq 2$ and either $N(v) \cap A = \emptyset$ or $N(v) \cap B = \emptyset$.*

Proof. Suppose that $|N(v) \cap (A \cup B)| \leq 1$ for some $v \in V(H) - A - B$. By (4.1), (4.2), and (4.3), $|N_H(v) - S| \geq \delta(H) - 3(k - 1) - 1$ and $|A| + |B| \geq 2\delta(H) - |S| - 3k$, we have

$$|V(H)| \geq |A| + |B| + |S| + (1 + \deg_{H-S}(v)) - 1 \geq 2 + 3\delta(H) - 6k > 12k \geq |V(H)|,$$

a contradiction. If both $N(v) \cap A \neq \emptyset$ and $N(v) \cap B \neq \emptyset$, then there is an s_1, t_1 -path of length 4 outside S , a contradiction. □

By Claim 4.4, every vertex in $V(H) - S$ has distance at most two from either s_1 or t_1 , but not both. Therefore, $H - S$ is the union of $H_1 = H[A \cup N(A) - S]$ and $H_2 = H[B \cup N(B) - S]$, and there are no edges connecting $V(H_1)$ with $V(H_2)$. Assume that $|V(H_1)| \leq |V(H_2)|$.

Observe that $\delta(H_1) \geq \delta(H) - 3(k - l_1) - l_1 \geq 6k - 3k + 2l_1 = 3k + 2l_1$ and $|V(H_1)| \geq 1 + \delta(H_1) > 3k$. If $|V(H_1)| \leq 4k$, then $3k + 2l_1 \geq \frac{|V(H_1)| + 5k}{3}$, and by Theorem 1.1, H_1 is k -linked. This proves the lemma when $|V(H_1)| \leq 4k$.

Now consider the case $|V(H_1)|, |V(H_2)| \geq 4k + 1$. If $\delta(H_1) \geq (|V(H_1)| - 3)/2 + k$ or $\delta(H_2) \geq (|V(H_2)| - 3)/2 + k$, then by Theorem 1.1, either H_1 or H_2 is k -linked. Thus we may assume that there exist $v_1 \in H_1$ and $v_2 \in H_2$ such that $\deg_{H_1}(v_1) \leq |V(H_1)|/2 + k - 2$ and $\deg_{H_2}(v_2) \leq |V(H_2)| + k - 2$. Hence,

$$\deg_{H_1}(v_1) + \deg_{H_2}(v_2) \leq (|V(H_1)| + |V(H_2)|)/2 + 2k - 4,$$

that is,

$$\deg_H(v_1) + \deg_H(v_2) - (|N_H(v_1) \cap S| + |N_H(v_2) \cap S|) \leq (|V(H)| - |S|)/2 + 2k - 4.$$

Therefore

$$\begin{aligned} |N_H(v_1) \cap S| + |N_H(v_2) \cap S| &\geq 2\delta(H) - |V(H)|/2 + |S|/2 - 2k + 4 \\ &\geq 12k - 6k + |S|/2 - 2k + 4 \geq 4k + 4 + |S|/2. \end{aligned} \tag{4.4}$$

Claim 4.5. $|N(v_1) \cap S| + |N(v_2) \cap S| \leq 4l_1 + 4l_2 + 5l_3 + 6(l_4 + l_5 + l_6 + l_7)$.

Proof. It is enough to prove that v_1 and v_2 together have at most:

- (a) 4 neighbours (counted with multiplicities) on each P_i of length 1 and each unlinked pair in S ,
- (b) 5 neighbours on each P_i of length 2,
- (c) 6 neighbours on each P_i of length at least 3.

Statement (a) is evident and (c) follows from (4.1). To prove (b), suppose that each of v_1 and v_2 has exactly 3 neighbours in a path $P_i = (s_i, w, t_i)$. Since $|S| \leq |V(H)| - |V(H_1)| - |V(H_2)| \leq 12k - 2(4k + 1) = 4k - 2$, every $v \in V(H)$ has at least $6k - |S| \geq 2k + 2$ neighbours outside S . Thus we may assume that $wu \in E(G)$ for some $u \in V(H_1)$, $u \neq v_1$.

Since no $u \in V(H_1) + s_1$ is adjacent to any $w \in V(H_2) + t_1$, each distinct pair $u_1, u_2 \in V(H_1) + s_1$ has at least $6k + 6k - (8k - 2) = 4k + 2 \geq |S| + 4$ common neighbours. Similarly, each distinct pair $w_1, w_2 \in V(H_2) + t_1$ has at least $|S| + 4$ common neighbours. Thus, the graph $H - (S - s_1 - t_1)$ contains an s_1, u -path Q_1 of length at most 2 avoiding v_1 and a t_1, v_2 -path Q_2 of length at most 2. Then we replace P_i by $s_i v_1 t_i$ and add the path $s_1 Q_1 u w v_2 Q_2 t_1$ of length at most 6. This contradicts the maximality of the linkage. \square

Now by Claim 4.5 we have

$$\begin{aligned} |N(v_1) \cap S| + |N(v_2) \cap S| &\leq 4l_1 + 4l_2 + 5l_3 + 6(l_4 + l_5 + l_6 + l_7) \\ &= 4 \sum_{i=1}^7 l_i + (l_3 + 2l_4 + 2l_5 + 2l_6 + 2l_7) \\ &= 4k + (l_3 + 2l_4 + 2l_5 + 2l_6 + 2l_7) \leq 4k + |S|/2, \end{aligned}$$

which contradicts (4.4). □

As we have mentioned above, proving Lemma 4.3 for smaller α would yield the corresponding improvement for $f(k)$.

Acknowledgement

We thank a referee for helpful comments.

References

- [1] Bollobás, B. and Thomason, A. (1996) Highly linked graphs. *Combinatorica* **16** 313–320.
- [2] Dirac, G. (1952) Some theorems on abstract graphs. *Proc. London Math. Soc.* **2** 69–81.
- [3] Egawa, Y., Faudree, R. J., Győri, E., Ishigami, Y., Schelp, R. H. and Wang, H. (2000) Vertex-disjoint cycles containing specified edges. *Graphs Combin.* **16** 81–92.
- [4] Faudree, J., Faudree, R., Gould, R., Jacobson, M. and Lesniak, L. (2000) On k -ordered graphs. *J. Graph Theory* **35** 69–82.
- [5] Gould, R. J. (2003) Advances on the Hamiltonian problem: A survey. *Graphs Combin.* **19** 7–52.
- [6] Gould, R. J., Kostochka, A. V., Lesniak, L., Schiermeyer, I. and Saito, A. (2003) Degree conditions for k -ordered Hamiltonian graphs. *J. Graph Theory* **42** 199–210.
- [7] Jung, H. A. (1970) Verallgemeinerung des n -fachen Zusammenhangs für Graphen. *Math. Annalen* **187** 95–103.
- [8] Kierstead, H. A., Sárközy, G. and Selkow, S. (1999) On k -ordered Hamiltonian graphs. *J. Graph Theory* **32** 17–25.
- [9] Larman, D. G. and Mani, P. (1974) On the existence of certain configurations within graphs and the 1-skeletons of polytopes. *Proc. London Math. Soc.* **20** 144–160.
- [10] Mader, W. (1967) Homomorphieeigenschaften und mittlere Kantendichte von Graphen. *Math. Annalen* **174** 265–268.
- [11] Ng, L. and Schultz, M. (1997) k -ordered Hamiltonian graphs. *J. Graph Theory* **2** 45–57.
- [12] Ore, O. (1960) Note on Hamilton circuits. *Amer. Math. Monthly* **67** 55.
- [13] Robertson, N. and Seymour, P. D. (1995) Graph minors XIII: The disjoint path problem. *J. Combin. Theory Ser. B*, **63** 65–110.
- [14] Thomas, R. and Wollan, P. (2003) An improved linear edge bound for graph linkage. Submitted.