

Review

On the Chemical Stabilities of Ionic Liquids

Subbiah Sowmiah, Venkatesan Srinivasadesikan, Ming-Chung Tseng and Yen-Ho Chu *

Department of Chemistry and Biochemistry, National Chung Cheng University, 168 University Road, Chia-Yi, 621 Taiwan

* Author to whom correspondence should be addressed; E-Mail: cheyhc@ccu.edu.tw;
Tel.: +886-5-242-8148; Fax: +886-5-272-1040.

Received: 1 July 2009; in revised form: 16 September 2009 / Accepted: 18 September 2009 /
Published: 25 September 2009

Abstract: Ionic liquids are novel solvents of interest as greener alternatives to conventional organic solvents aimed at facilitating sustainable chemistry. As a consequence of their unusual physical properties, reusability, and eco-friendly nature, ionic liquids have attracted the attention of organic chemists. Numerous reports have revealed that many catalysts and reagents were supported in the ionic liquid phase, resulting in enhanced reactivity and selectivity in various important reaction transformations. However, synthetic chemists cannot ignore the stability data and intermolecular interactions, or even reactions that are directly applicable to organic reactions in ionic liquids. It is becoming evident from the increasing number of reports on use of ionic liquids as solvents, catalysts, and reagents in organic synthesis that they are not totally inert under many reaction conditions. While in some cases, their unexpected reactivity has proven fortuitous and in others, it is imperative that when selecting an ionic liquid for a particular synthetic application, attention must be paid to its compatibility with the reaction conditions. Even though, more than 200 room temperature ionic liquids are known, only a few reports have commented their effects on reaction mechanisms or rate/stability. Therefore, rather than attempting to give a comprehensive overview of ionic liquid chemistry, this review focuses on the non-innocent nature of ionic liquids, with a decided emphasis to clearly illuminate the ability of ionic liquids to affect the mechanistic aspects of some organic reactions thereby affecting and promoting the yield and selectivity.

Keywords: ionic liquid; imidazolium; pyridinium; phosphonium; quaternary ammonium; chemical stability

1. Introduction

Ionic liquids (ILs) are low-melting molten salts composed entirely of ions, and many of them are liquids at room temperature [1-6]. These novel solvents are attracting interest as greener alternatives to conventional organic solvents with the aim of facilitating sustainable chemistry. Room temperature ionic liquids (RTILs), often referred to as ‘designer solvents’, have been the great focus of scientists in various fields since they can be tuned for specific applications [7-11]. Promising diverse applications of RTILs continue to expand significantly due to their unusual physical and chemical properties like high thermal stability, lack of inflammability, low volatility, chemical stability and excellent solubility with many organic compounds. Thus, these are considered to be emerging green solvents and potential alternatives to the classical volatile organic solvents. In addition to being “green” solvents [12-14], ILs have been used for a myriad of applications in diverse synthetic reactions [15-18], separations and extractions [19-22], and electrochemical [23,24], nanotechnological [25-28], biotechnological [29], and engineering [30-32] processes.

The perception that all ionic liquids are “green solvents” may lead to inappropriate experimental design and utilization of these chemicals. As reviewed by Song *et al.*, switching from an organic solvent to an ionic liquid often results in a significant improvement in catalytic performance *e.g.* rate acceleration, (enantio)selectivity improvement and an increase in catalyst stability [33-35], but synthetic chemists cannot ignore the stability data and intermolecular interactions (or even reactions) that are directly applicable to organic reactions in ionic liquids.


It is usually assumed that these ionic liquids are entirely innocent and non-coordinating solvents. However, such innocuous behavior was not always observed and often led to ‘undesirable’ transformations in reactions [36,37]. The lack of detailed knowledge on their structures and physical-chemical properties requires more attention and certain degree of caution should be exercised when ionic liquids are chosen as solvents.

Even though more than 200 room temperature ionic liquids are known, only a few reports evidence their effects on reaction mechanisms or rate/stability. It is important to note that although many IL-like organic salts have important industrial and commercial applications, the environmental fate and any potential toxicity issues for most ionic liquids are not known, initial data are only now being determined [38]. This review, rather than attempting to give a comprehensive overview of IL chemistry, is focused on the non-innocent nature of ILs, with a decided emphasis on clearly illuminating the ability of ILs to affect the mechanistic aspects of some organic reactions thereby affecting and promoting the yields as well as stereoselectivity.

2. Scope of This Review

ILs are advanced, technological solvents that can be tailor-synthesized to fit well to a particular application. This emerging field has been extensively reviewed by a number of renowned chemists, including Welton [7], Holbrey [39], and Seddon [40]. Recently, uses of ILs as solvents for chemical synthesis have also been carefully studied and thoroughly reviewed [1-11,41-44]. There is no doubt that this area of research has been a focal point of green chemistry for the past two decades (Figure 1).

Figure 1. IL publications (on May 27, 2009) determined from the ISI Web of Science in the last fourteen years.




The first room temperature ionic liquid [EtNH₃][NO₃] (m.p. 12 °C) was discovered in 1914 [45], but interest did not develop until the discovery of binary ionic liquids made from mixtures of aluminum(III) chloride and *N*-alkylpyridinium [46] or 1,3-dialkylimidazolium chloride [47]. A major drawback of all chloroaluminate(III) ionic liquids was their moisture sensitivity. In 1992, Wilkes and Zawarotko [48] prepared and characterized air and water stable 1-ethyl-3-methylimidazolium based ionic liquids incorporating different anions. Over the years that followed, new classes of cations and anions have been reported. The common cations and anions so far involved in ionic liquids are given below.


Possible Anions:

BF₄⁻; B(CN)₄⁻; CH₂CHBF₃⁻; CF₃BF₃⁻; C₂F₅BF₃⁻; nC₃H₇BF₃⁻; nC₄H₉BF₃⁻; PF₆⁻;
 CF₃CO₂⁻; CF₃SO₃⁻; N(SO₂CF₃)₂⁻; N(COCF₃)(SO₂CF₃)⁻; N(SO₂F)₂⁻; N(CN)₂⁻;
 C(CN)₃⁻; SCN⁻; SeCN⁻; CuCl₂⁻; AlCl₄⁻; OH⁻


Common Cations:




Tetraalkylammonium




Di, tri and tetraalkylimidazolium




[R-3C-im]




[R-4C-im]




Alkylpyridinium




Dialkylpyrrolidinium




Dialkylpiperidinium




Tetraalkylphosphonium



Trialkylsulfonium



Dialkylpyrazolium



N-Alkylthiazolium

[R-3C-im] = 1-alkyl-2,3-trimethyleneimidazolium

[R-4C-im] = 1-alkyl-2,3-tetramethyleneimidazolium

Table 1. Comparison of ionic liquids with organic solvents [49].

| Property | Organic Solvents | Ionic Liquids |
|----------------------------|---------------------------------------|---|
| Number of solvents | >1,000 | >1,000,000 |
| Applicability | Single function | Multifunction |
| Catalytic ability | Rare | Common and tunable |
| Chirality | Rare | Common and tunable |
| Vapour pressure | Obeys the Clausius-Clapeyron Equation | Negligible under normal conditions |
| Flammability | Usually flammable | Usually nonflammable |
| Solvation | Weakly solvating | Strongly solvating |
| Tunability | Limited range of solvents available | Unlimited range means 'designer solvents' |
| Polarity | Conventional polarity concepts apply | Polarity concept questionable |
| Cost | Normally inexpensive | 2 to 100 times the cost of organic solvents |
| Recyclability | Green imperative | Economic imperative |
| Viscosity/cP | 0.2-100 | 22-40,000 |
| Density/g cm ⁻³ | 0.6-1.7 | 0.8-3.3 |
| Refractive index | 1.3-1.6 | 1.5-2.2 |


One of the biggest and challenging industrial concerns is the replacement of volatile organic compounds (VOCs) [50], particularly those that are toxic, such as CH₂Cl₂, and those that are highly volatile and flammable, such as ethers (Table 1). The use of ILs [51] to replace or reduce VOCs is a move that could have significant positive environmental impact. There would also be safety benefits

resulting from the preferential use of ILs over VOCs, largely due to their low or non-flammability with normal use [52-54]. To date, the needed information on the environmental outcome, any potential toxicity, stability issues and others for most of the ionic liquids are not yet fully made available and thus they should be treated with caution during their applications.


The uniqueness of structural tunability of both the cations and anions suggests great flexibility in the potential applications of ionic liquids. For example, the melting points of 1-alkyl-3-methylimidazolium tetrafluoroborates [55] and hexafluorophosphates [40] are a function of the length of the 1-alkyl group, which forms a liquid crystalline phase for alkyl chain length over 12 carbon atoms. Another important property that changes with structure is the miscibility of water in these ionic liquids. For example, 1-alkyl-3-methylimidazolium tetrafluoroborate salts are miscible with water at 25 °C where the alkyl chain length is less than six, but at or above six carbon atoms, they form a separate phase when mixed with water. This behavior can be of substantial benefit when carrying out solvent extractions or product separations, as the relative solubilities of the ionic and extraction phase can be adjusted to make the separation as easy as possible. Hence, a tremendous amount of work has been carried out on applications of ionic liquids and has stimulated interest in both academia and industry [1-6,56].

Recent development of ILs turned on designing suitable ionic liquids with specific application repetition of ILs many times that can be used both as catalysts/promoters and solvents. Several innovative synthetic procedures on this task lead to successful results and emerge as a new field in ionic liquid chemistry. For example, Ranu *et al.* explored the influence of a new tailor-made, task-specific ionic liquid like [bmim][OH] on Michael addition reactions [57]. This IL, that acts as both catalyst and solvent, plays a dramatic role in the chemical transformation (Scheme 1). Several reviews related to this type ILs are available and, therefore, this subject is not included as a part of our review work.

Scheme 1. IL-mediated Michael addition reaction in ionic liquid.




Dyson *et al.* studied the multi-role of the hydroxyl group in hydroxyl-functionalized ILs (Scheme 2). This IL not only acts as a solvent but also can be viewed as a promoter and demonstrates that careful choice/design of the solvent can be used to tune catalytic reactions [58]. This type of designed functionalized ILs is beyond the scope of this review and will not be discussed here.

Scheme 2. Functionalized ILs used efficiently for the Suzuki coupling reaction.

3. Chemical Stability of Ionic Liquids

3.1. Imidazolium ionic liquids


Among RTILs, those derived from imidazole continue to attract increasing interest in many different areas of chemistry [7-11,56]. The imidazolium cation has earned a great attention among scientists due to its low melting points and viscosity, ease of synthesis, and good stability to oxidative and reductive conditions [59]. RTILs are, in general, viewed solely as solvents and, in recent days, its non-innocent nature however began to emerge (Scheme 3). Deprotonation affording *N*-heterocyclic carbenes (NHCs), high acidity at C2 position, and easier deuterium exchange have significant implications in the chemistry of this type of ILs [60].

Scheme 3. The non-innocent nature of imidazolium-based ILs.


Noting these properties, it becomes immediately obvious that imidazolium-based ILs are likely to be chemically unstable under basic conditions and caution must be exercised in interpreting reaction results obtained in such ILs, under basic conditions.

3.1.1. Acidity and deuterium exchange

In early 1964, Olofson *et al.* reported the kinetic evidence and study of the rates of deuterium incorporation in heavy water buffers of imidazole related compounds [61]. Their experiments clearly demonstrated the acidic nature of C2 proton of the 1,3-dialkylimidazolium cation, such as 1,3-dimethylimidazolium [dmim], that can be exchanged with deuterium under mild conditions (Scheme 4).


Scheme 4. H-D exchange at C2 position of [dmim] cation.

Although substitution at the C2 position of the imidazolium cation prevents the side reaction upon chemical synthesis, Handy and Okello later showed that even the C2 methyl substituted 1-butyl-2,3-dimethylimidazolium [bdmim] cation is not completely inert [62]. They found that the C2 methyl group underwent slow proton exchange even in the presence of a weak base such as triethylamine (Scheme 5).

Scheme 5. Deuterium isotope exchange at C2 methyl group of [bdmim] cation under basic conditions.

3.1.2. pKa of C2 hydrogen in imidazolium ions


Since pKa values can reveal the acidic nature of imidazolium cation, several reports determined these values by varying substituents (Table 2). The pKa values of the simple 1,3-dialkylimidazolium cations have been determined in both DMSO [63] and H₂O [64] and were found to be in the range of 21–24 [63-66].

Table 2. pKa values with different substituents on imidazolium cations.

| Entry | R | Solvent | pKa value |
|-------|-------------------------------|------------------|-----------|
| 1 | ^t Bu | DMSO | 22.7 |
| 2 | Me | H ₂ O | 23.0 |
| 3 | Me | DMSO | 21.1 |
| 4 | ^t Bu | DMSO | 22.6 |
| 5 | Ph | DMSO | 16.1 |
| 6 | ⁱ Pr(4,5-dimethyl) | DMSO | 24.0 |

Even though the pKa values of some of these imidazolium cations have been determined, much less attention has been directed towards the impact that the counteranion has on this acidity and exchange at C2 position. Handy and co-workers recently demonstrated that the anion had a strong influence on the conditions required for H-D exchange [62]. Their study revealed that more basic anions such as dicyanamide $N(CN)_2$ resulted in RTILs that would undergo deuterium exchange in the absence of any added base (entry 1, Table 3) while weakly coordinating and non-basic anions such as tetrafluoroborate BF_4 resulted in salts which required an external base for deuterium exchange to occur (entry 2, Table 3).


Table 3. Effect of anion on the H-D exchange rate of C2 hydrogen of [bmim] cation.



| Entry | Base | X | Rate (min^{-1}) |
|-------|------|-----------|----------------------------|
| 1 | None | $N(CN)_2$ | 41×10^{-3} |
| 2 | None | BF_4 | 0 |

As aforementioned, the C2 methyl substituted imidazolium ILs also underwent slow proton exchange in the presence of a base. Handy and Okello have shown that even the 2-methyl substituted imidazolium cation is not completely inert (Table 4) [62].

Table 4. Effect of anion on the H-D exchange rate of C2 methyl group of [bdmim] cation.




| Entry | Base | X | Rate (min^{-1}) |
|-------|---------|-----------|----------------------------|
| 1 | Et_3N | Cl | 0.04×10^{-3} |
| 2 | Et_3N | $N(CN)_2$ | 0.04×10^{-3} |
| 3 | KOH | Cl | 2.1×10^{-3} |
| 4 | KOH | $N(CN)_2$ | 1.0×10^{-3} |

A distinct influence of the anion on exchange process was reported by Schatz [67]. Very different H-D exchange rates (up to one order of magnitude) could be noted for chloride and bromide salts, depending upon the size of the substituents on the two imidazole nitrogens. The influence of the

counterion on the H-D exchange was investigated for compound **1**. In no case, similar rate constants for bromide and chloride were found in methanol- d_4 containing 3% water. The values differ by as much as a factor of 10 (Table 5). For bromides, the exchange rates dropped significantly compared to those of chlorides. However, in the case of bulky substituents, the bromide salts showed higher exchange rates [62].

Table 5. Rate constants of the H-D exchange in methanol- d_4 containing 3% water at 300 K.



| Entry | R | X | k (d ⁻¹) |
|-------|------------|----|----------------------|
| 1 | mesityl | Cl | 1.060 |
| 2 | isopropyl | Cl | 0.553 |
| 3 | tert-butyl | Cl | << 0.001 |
| 4 | mesityl | Br | 0.176 |
| 5 | isopropyl | Br | << 0.001 |
| 6 | tert-butyl | Br | 0.158 |

Our [b-3C-im][NTf₂] ionic liquid appears to provide sufficient chemical tolerance to bases so that many potential side reactions that are common in imidazolium-based ionic liquids may therefore be avoided [68]. Under neutral conditions (CD₃OD/D₂O = 1:1), ionic liquids such as [bdmim][NTf₂] and [bdmim][PF₆] produced less than 10% deuterium exchange at C2 methyl group at room temperature for one week time and only [bmim][PF₆] was found to be deuterium exchanged rapidly (50% in 1 h) and our [b-3C-im][NTf₂] gave no detectable exchanges even after one week at ambient temperature [68,69]. Under basic condition (0.1 M KOD in CD₃OD/D₂O = 1:1), only [b-3C-im][NTf₂] ionic liquid is chemically stable to base (t_{1/2} ~ 6 days). Among all tested ionic liquids, [b-3C-im][NTf₂] ionic liquid was most stable to solvent deuterium isotope exchange while the previously reported [bmim][NTf₂] and [bdmim][NTf₂] was deuterium exchangeable instantaneously and in 30 min, respectively, at ambient temperature.


From the results of solvent deuterium exchange experiments, we therefore concluded that the new ionic liquid [b-3C-im][NTf₂] is far more chemically stable than the previously reported [bmim][PF₆], [bdmim][NTf₂], and [bdmim][PF₆] (i.e., [b-3C-im][NTf₂] » [bdmim][NTf₂] ≈ [bdmim][PF₆] » [bmim][PF₆]).

3.1.3. Generation and stability of carbenes

The characteristic source of reactivity is the high acidity of the C2 hydrogen position in the imidazolium cation [36,37]. Deprotonation of C2 hydrogen results in a highly stabilized *N*-heterocyclic carbene (NHC) (Scheme 6). Initially, NHCs were first proposed by Wanzlick in the 1960s and were characterized by the formation of their adducts with a variety of other compounds [70]. For several years, no other evidence other than these adducts were proposed for NHC formation.


In 1991, Arduengo *et al.* achieved a breakthrough by isolating the stable NHC. By deprotonation of an imidazolium salt, they were able to characterize a stable singlet *N*-heterocyclic carbene, 1,3-di-1-adamantylimidazol-2-ylidene [71,72]. They demonstrated that imidazolium cations with larger groups on the two nitrogens afforded stable isolable NHCs.

Scheme 6. Preparation of a stable NHC.




Amyes *et al.* have calculated the equilibrium constant and ΔG° for the conversion of the singlet carbene to neutral imidazole *via* a 1,2-hydrogen shift in water to be 5×10^{16} and -22.7 kcal/mol, respectively [73]. The concerted 1,2-shift is however symmetry forbidden and hence the reaction to give imidazole occurs in two steps as shown in Scheme 7.

Scheme 7. Conversion of NHC to imidazole.




The first experimental measurements of gas phase proton affinity (251.3 ± 4 kcal/mol) of the *N*-heterocyclic carbene, 1-ethyl-3-methyl-imidazol-2-ylidene **2**, lead to greater insight into their reactivity [74].



Preparations of NHCs from imidazolium salts usually involve the use of strong bases such as potassium or sodium hydride, LDA or KHMDS (Scheme 8) [75,76].

Scheme 8. Common preparation of NHCs.




3.1.4. Reactions involving C-C bond formation

3.1.4.1. Baylis-Hillman reaction


In 2001, Afonso *et al.* anticipated that imidazolium-based ionic liquids would be a good choice of solvent for the Baylis–Hillman reaction [77]. As they assumed, the reaction was “found” to be 33 times faster in [bmim][PF₆] than in acetonitrile. The reactions were, however, low yielding. Soon after Afonso’s paper was published, Aggarwal *et al.* quickly pointed out that the ionic liquids used were all [bmim]-based and, in the presence of bases such as DABCO or 3-hydroxyquinuclidine, the imidazolium moiety in ionic liquids could be deprotonated at its C2 position and the resulting nucleophile directly reacted with and thereby consumed the aldehyde (Scheme 9), leading to the misinterpretation of both the apparent reaction rates and the lower yields [78,79].

Scheme 9. Nucleophilic addition of [bmim] cation onto benzaldehyde.



Further, they also showed that caution must be exercised when using ionic liquids recycled from one reaction to another. If the adduct formation is reversible as in Scheme 9, the ionic liquid recycled from one run to another of a Baylis–Hillman reaction provided a mixture of products (Scheme 10).

Scheme 10. Mixed Baylis-Hillman adducts result if [bmim]-based ionic liquid was recycled for new reaction with different starting aldehyde.



Our group made a comparative study on the use of [bmim][PF₆] and [bdmim][PF₆] in the Baylis–Hillman reaction and demonstrated that the product was contaminated with aldehyde-conjugated [bmim][PF₆] when this IL was used [80].


Itoh *et al.* also reported that no reaction took place when PhMgBr was reacted with benzaldehyde in conventional [bmim][NTf₂] ionic liquid and the desired product was obtained for the [bdmim][NTf₂]

solvent system [81]. These results clearly demonstrated once again that the strong acidity of the C2 proton of the imidazolium salt caused decomposition of the Grignard reagents [82,83].

Handy *et al.* carried out alkylation reactions of [bmim][Br] with various alkyl halides using the base NaH *via* generation of carbene intermediates [84]. Their results revealed that primary alkyl halides (butyl, hexyl, heptyl, decyl, and hexadecyl), except allyl bromide, provided 2-substituted imidazolium RTILs, while secondary alkyl halides failed to do so. They also showed that 2-substituted imidazolium could be obtained as a sole product by using excess NaH [62].

Most recently, Rosenau *et al.* demonstrated unambiguously that 1-alkyl-3-methylimidazolium ILs react at C2 with the reducing ends of cellulose and aldopyranose models [85]. They found the reaction was strongly catalyzed by bases such as imidazole and 1-methylimidazole, commonly present impurities in ILs. In solutions of cellulose in 1-alkyl-3-methylimidazolium ILs, both the ionic liquid and the cellulose are evidently not inert (Scheme 11). This again confirmed that C2 position of imidazolium was chemically unstable under basic conditions.

Scheme 11. Reaction of 1-methyl-3-(2-naphthylmethyl)imidazolium acetate ionic liquid with cellulose at its reducing end.



This might induce adverse effects in medical and biological applications if even minor impurities present in cellulose and cellulose derivatives. The application of alkylmethylimidazolium ILs in the processing of oxidized cellulose, such as TEMPO-oxidized or periodate-oxidized cellulose, seems to be rather problematic, for medical and physiological scenarios.

3.1.4.2. Palladium catalysed reactions

Heck Reaction: Carmichael *et al.* reported that ionic liquids provide a convenient medium for the Heck reaction, while allowing recycling of the catalyst [86]. Herrmann and Böhm found that the imidazolium-based ionic liquids gave less satisfactory results compared to quaternary ammonium salts (Scheme 12, Table 6) [87,88].

Scheme 12. Heck reaction forming palladacycle.





Table 6. Heck reaction of chlorobenzene and styrene in presence of palladacycle X^a.

| IL | Base | Time (h) | Yield (%) ^b |
|-----------------------------|-------|----------|------------------------|
| [N _{4,4,4,4}][Br] | NaOAc | 18 | 51 |
| [pmim][Br] | NaOAc | 19 | 22 |
| [pbim][Br] | NaOAc | 16 | 11 |
| [bbim][PF ₆] | NaOAc | 15 | 5 |

^a Conditions: 1.0 equiv of chlorobenzene, 1.5 equiv styrene, 1.2 equiv base; ^b GC yields.


This observation suggested that the Heck reaction when carried out in imidazolium ionic liquids could proceed through a different mechanism *via* formation of carbene species. Xu *et al.* obtained the first convincing evidence that the imidazolium ion can react with a catalyst precursor to form NHCs [89]. They also clearly showed that the carbene complex of palladium formed between Pd(OAc)₂ and the solvent [bmim][Br] is catalytically active for olefination reactions of aryl halides (C-C bond formation reaction) (Scheme 13) [90].

Scheme 13. Catalytic active [bmim][Br] for olefination of aryl halides.

In order to determine the effect of the ionic liquid, a control study on these reactions was performed. Like in the Heck reaction, Pd(OAc)₂ was heated in [bmim][Br] in the absence of the alkene and aryl halide, under reaction conditions. They observed rapid color changes from dark brown to red and then to yellow and the products formed were analyzed by NMR. The characteristic chemical shifts of the olefin protons (δ 7 in the ¹H-NMR spectrum) and the carbene carbons (δ 160–170 in the ¹³C-NMR spectrum) suggested the presence of a mixture of palladium carbene complexes. Supporting this formation of Pd carbene complexes, other species such as Ir carbene complexes (as well as hydrogenated species) are also obtained in reactions with imidazolium salts [91].


McGuinness *et al.* proposed a mechanism for the Heck reaction catalysed by Pd-carbene complexes [92]. The active species is the 14-electron Pd(0) complex Pd(tmiy)₂ to which the oxidative addition of an aryl halide occurs, giving the Pd(II) aryl complex. Dissociation of the halide ligand is accompanied by olefin coordination, insertion, and β -hydride elimination to release the product and give (after recoordination of the halide) Pd(tmiy)₂HX. Reductive elimination of HX in the presence of base then regenerates the active Pd(0) species. Srinivasan *et al.* have performed the Heck reaction at ambient temperature with considerably enhanced reaction rates by the combined use of ultrasonic irradiation and ionic liquids as solvents (Scheme 14) [93]. They confirmed the formation of the carbene complex **3** by ¹H-NMR analysis.

Scheme 14. Heck reaction *via* ultrasonic irradiation with IL.



Suzuki Coupling Reaction: Welton and co-workers reported that a mixed phosphine-NHC palladium species **4** was formed under Suzuki coupling conditions in RTILs (Scheme 15) [94,95]. Interestingly, the resulting catalyst was highly active in the Suzuki coupling reactions.

Scheme 15. Suzuki coupling reaction with RTILs.




Mathews *et al.* had previously demonstrated the *in situ* formation of a mixed phosphine-imidazolylidene palladium complex in the ionic liquid [bmim][BF₄] [94]. Wong *et al.* [96] carried out the Suzuki cross-coupling of 4-bromoacetophenone (BrAP) and phenylboronic acid (PBA) catalysed by tris(dibenzylideneacetone) dipalladium chloroform complex (Pd₂(dba)₃-CHCl₃) which showed the dramatic difference in yields in different solvents.

Negishi Cross-Coupling Reaction: Obviously, if deprotonation of the imidazolium cation with an amine base can be a problem for organic chemists, use of nucleophilic organometallic reagents such as


Grignard, organolithium, and even organozincs should certainly be more problematic. Indeed, the use of [bmim]-based RTILs in organozinc catalysed Negishi couplings met with failure; on the other hand, when a new ionic phosphine ligand **5** was incorporated for the reaction performed in a biphasic solvent system of toluene and [bdmim][BF₄], this palladium catalyzed cross-coupling reaction proceeded smoothly giving high isolated yields (Scheme 16) [97]. Chan and co-workers attempted a reaction of diethyl zinc with carbonyl compounds in the ionic liquid [bmim][Br], in which they observed gas evolution and formation of NHC-zinc complexes [98].

Scheme 16. Negishi cross-coupling of an organozinc in ionic liquid.



Sonogashira Reaction: Gholap *et al.* have demonstrated that the copper- and ligand-free Sonogashira reaction catalyzed by Pd(0) (or PdCl₂) nanoparticles proceeds under ultrasound irradiation in the ionic liquid [bbim][BF₄] (Scheme 17) [99]. The formation of Pd(0) nanoparticles was investigated and detected in this reaction. Control experiments showed that both ultrasound and the ionic liquid were necessary for the formation of these Pd(0) nanoparticles.

Scheme 17. Sonogashira reaction by Pd nanoparticles in IL.



Magna *et al.* investigated the telomerization of butadiene with methanol and clearly demonstrated the existence of reactivity between the Pd catalyst and 1,3-dialkylimidazolium ionic liquids [100]. They believed that this indicates a stoichiometric reactivity between 1,3-dialkylimidazolium salt and palladium, leading eventually to an inactive form. The putative formation of a “carbene-Pd” complex was evoked to explain the inactivity of systems using this kind of ionic liquids.

Aromatic Substitution Reactions: Earle *et al.* have recently reported that certain ionic liquids are chemically reactive and can greatly influence the outcome of the electrophilic aromatic substitution reactions [101]. Lancaster *et al.* investigated the aromatic nitration in various ionic liquids with a range of nitrate salts and explained the importance of cation choice among ILs that can influence the


selectivity, yield, and reaction time (Table 7) [102]. They observed low yields when using [bmim] and [bdmim] compared to [bmpy] ILs. This might be due to the stability of the ionic liquids towards the nitrating agent used. Their MS analysis revealed that one or two nitro group substitutions were observed in [bmim] and [bdmim].

Table 7. Electrophilic aromatic nitration of toluene by acyl nitrates in various solvents.

| Solvent | System | Time (h) | Yield (%) |
|---------------------------------|--------------------------------------|----------|-----------|
| CH ₂ Cl ₂ | HNO ₃ - Ac ₂ O | 1 | 35 |
| [bmim][NTf ₂] | HNO ₃ - Ac ₂ O | 24 | 42 |
| [bdmim][NTf ₂] | HNO ₃ - Ac ₂ O | 1 | 63 |
| [bmpy][NTf ₂] | HNO ₃ - Ac ₂ O | 1 | 93 |


Chiappe *et al.* have demonstrated the use of ionic liquid solvents for stereoselective halogenations of alkenes and alkynes [103]. Depending on the ionic liquid the addition may be stereospecific, *erythro*- (or *meso*-) bromochlorides or dibromides are obtained from *trans*-olefins while *cis*-olefins give the corresponding *threo*- (or *d,l*-) adducts (Scheme 18).

Scheme 18. Bromination of 1,3-cyclohexadiene in [bmim][Br] at room temperature.



Our group also found that the nucleophilic aromatic substitution reaction when conducted in the commonly used [bmim][PF₆] ionic liquid instead of [bdmim][PF₆], the desired arylamine products **6** were obtained with low yields and found to be contaminated with the 1-fluoro-2-nitrobenzene-conjugated [bmim][PF₆] adduct **7** (Scheme 19). The chemical nature of adduct could be readily verified and unambiguously confirmed by ¹H-NMR [104].

Scheme 19. Nucleophilic aromatic substitution in IL.




Being aware of this acidic incompatibility of imidazolium ionic liquids, our group synthesized and characterized a new class of room temperature ionic liquids [R-3C-im][NTf₂], which are chemically stable ionic liquids [68,105]. We have also performed high temperature organic reaction such as the Claisen rearrangement in the chemically and thermally stable and non-volatile room temperature ionic liquid [b-3C-im][NTf₂] under microwave conditions and successfully obtained high isolated yields [106]. Thus, these ionic liquids appear to fulfil practical requirement as an inert solvent for chemical reactions and these ionic liquids may open exciting perspectives of use in various synthetic applications of natural and non-natural products of biological significance. The [R-3C-im][NTf₂] are the most chemically stable imidazolium-based room temperature ionic liquids available today.

3.2. Phosphonium ionic liquids

Apart from imidazolium ionic liquids, phosphonium ILs have also been investigated with the view of exploring their general scope and their unique capabilities. Phosphonium-based ILs (PILs) are less expensive to manufacture on an industrial scale, and are also an efficient and recyclable media for Pd-mediated cross-coupling reactions [107,108]. PILs are stable under thermal conditions, and strongly basic reagents including Grignard reagents [109,110]. For these reasons they offer greater practicality and scope and deserve far more consideration as unique reaction media in the ILs field than they have received thus far. Clyburne *et al.* [111] found that highly basic NHCs (pK_a = 22-24) [63,64] are persistent in PILs [112] such as trihexyl(tetradecyl)phosphonium chloride **8** [107] and were surprised to observe that deprotonation of the IL **8** to produce a phosphorane **9** did not occur (Scheme 20).

Scheme 20. Expected phosphorane product on deprotonation of PIL cation.




Supporting this observation, they also examined whether stronger bases (Grignard reagents) would be persistent and their behaviour in PILs. Interestingly, these solutions again show no sign of degradation after one month, as shown by reactivity studies. The failure of deprotonation of **8** to produce a phosphorane **9** indicates that PIL **8** is capable of supporting reactions involving strong bases such as Grignard reagents without any degradation/reaction between the IL and the strongly basic reagents. They also examined the stability of PIL in the presence of several different nucleophilic reagents. The absence of phosphoranes and phosphoryl in the PILs suggests that they are a suitable choice as solvent for strongly basic, carbon-centered reagents. These observations open up the use of PILs as a reliable reaction media for a wide variety of basic reagents. They also reported that the stability of organometallic species in PILs are anion dependent and evidently shown that trihexyltetradecylphosphonium decanoate would act as solvents for bases such as Grignard reagents, isocyanides and Wittig reagents (phosphoranes) without any decomposition of the solvent [113]. Thus, for phosphonium ions, small bases are susceptible to react with the PIL through deprotonation reactions whereas large bases are resistant to this reaction.

Later, Itoh *et al.* designed new phosphonium salt ionic liquids that are applicable to various types of Grignard reagent mediated reactions [81]. In their study, they demonstrated that the introduction of an alkyl ether moiety on the side arm of phosphonium salt ionic liquid was quite effective in improving the capability of the phosphonium salt ionic liquids as a solvent for ether free Grignard reaction and Grignard reagent mediated reaction. Besides being reported to be sterically hindered at its central core [112,114], lack of acidic ring protons made a belief that commercially available trihexyl-(tetradecyl)phosphonium chloride **8** ionic liquid is chemically stable and remains inert towards Grignard reagents under strongly basic conditions [36,115].


Very recently, our group demonstrated the instability of tetraalkylphosphonium ionic liquids [116]. Under basic conditions, **8** was reactive and 50% deuterium exchanged on all four P-CH₂ methylene groups within 9 h (*t*_{1/2}) at ambient temperature. Further we investigated the direct carboxylate alkylation of organobromides with the presence of Hunig's base in ionic liquid **8** (Scheme 21).

Scheme 21. Reaction of benzoate salts with PIL **8**.



Our studies quickly revealed that the phosphonium cation reacted with sodium salts of substituted benzoates apparently through direct S_N2 carboxylate alkylation to form esters **10** and the resulting esters further converted, *via* Wittig reaction, to finally afford aryl ketones **11** by microwave irradiation at elevated temperature. We have also isolated and spectroscopically characterized these aryl ketones. A plausible mechanism was then proposed to account for the fact that aryl ketones **11** are reaction products and benzoate esters **10** are the most probable intermediates (Scheme 22).

Scheme 22. Mechanism proposed for benzoate reaction with PIL **8**.



Our results revealed for the first time that the commercially available tetraalkylphosphonium cation in ionic liquid **8** is acidic, electrophilic, and can be chemically reactive. Even though phosphonium-

based ionic liquids are less prone to thermal decomposition, the presence of O₂ may lead to the formation of phosphine oxides in many cases [117] and transfer of O from oxygen containing anions, such as HSO₄⁻, also leads to byproduct formation [118].

3.3. Quaternary ammonium ionic liquids


The date of discovery of the "first" ionic liquid starts with the quaternary ammonium ionic liquids. In 1888, ethanolanmonium nitrate (mp 52-55 °C) was reported by Gabriel [119]. Later in 1914, one of the earlier known room temperature ionic liquids was [EtNH₃][NO₃] (mp 12 °C), synthesized by Walden [120]. These ILs make some unique applications in various fields. Their special properties like low-melting points, low-viscosity, chemical and electrochemical stability facilitate development of ILs as possible safe electrolytes for high-energy density devices [121-126].

Calò *et al.* [127] made a comparative study of the effects exerted by different ILs (quaternary ammonium, pyridinium, imidazolium) on catalyst stability, reaction rates and regio- and stereoselectivity in carbon-carbon coupling (Heck, Suzuki, and Stille) reactions. The superiority of quaternary ammonium halides over other ILs might be due to the Coulombic interaction between the cations and anions in ILs as well as the nucleophilicity of the anions. The bulkiness of tetrahedral quaternary ammonium cations, which forces the anions away from the cation, renders these anions available for a good activity and stability of the palladium catalysts. On the contrary, the planar structures of imidazolium and pyridinium cations, due to a strong Coulombic interaction that binds the anions tightly, decrease their availability for stabilisation and activity of the catalysts.

In the Suzuki reaction, quaternary ammonium ionic liquid was less efficient in catalyzing the reaction in presence of inorganic bases. However, the presence of an aqueous base, which is necessary for the Suzuki reaction, accelerated the aggregation of Pd nanoparticles to afford catalytically inactive "Pd black", which inhibits the catalyst recycling. As previously reported [127,128], whatever the base, these reactions were devoid of stereoselectivity and, yet again, [bmim]-based ionic liquid was less efficient than quaternary ammonium ionic liquid. Similar to earlier reports, the influence of bases on the reaction between 4-bromotoluene and *trans*-ethyl cinnamate catalysed by Pd-benzothiazole carbene complex in quaternary ammonium ionic liquid [129], proved a dramatic role on the regio- and stereoselectivity regardless of the base used.

Two modes of thermal decomposition of quaternary ammonium salt are known [130]: the reverse Menshutkin (Scheme 23) and Hofmann (Scheme 24) types. One would expect reactivity in reverse Menshutkin reaction to follow established orders of X⁻ nucleophilicity toward saturated carbon (Scheme 23).


Scheme 23. Reverse Menshutkin decomposition in quaternary ammonium ILs.



One might expect reactivity in Hofmann elimination reaction to follow the order of X⁻ basicities (approximately ClO₄ < I < Br < NO₃, SCN, picrate) although it is not clear whether bases as weak as the anions can indeed bring about such E2 process [131,132]. However, the equilibrium condition of

$R_4N^+X^-$ with strongly nucleophilic X^- in the fused salt is evidently undergoing complete decomposition to olefin and $R_3NH^+X^-$ (Scheme 24).

Scheme 24. Hofmann elimination reaction in quaternary ammonium ILs.





Zhou *et al.* synthesized and characterized cyclic quaternary ammonium salts comprising *N*-alkyl-(alkyl ether)-*N*-methylpyrrolidinium, -oxazolidinium, -piperidinium, and -morpholinium (alkyl = *n*-C₄H₉, alkyl ether = CH₃OCH₂, CH₃OCH₂CH₂) with a series of weakly coordinating and electrochemically stable fluoro anions, namely perfluoroalkyltrifluoroborate [RFBF₃]⁻, (RF = CF₃, C₂F₅, *n*-C₃F₇, *n*-C₄F₉). Some of these salts are potential candidates for use as electrolytes in high-energy storage devices [126].

3.4. Pyridinium ionic liquids

The modern era of ionic liquids stems from the work on alkyipyridinium ionic liquids and has various applications. From the literature available for these ILs, it is understood that no systematic and detailed studies on their reactivity and stability have been undertaken.

Magna *et al.* observed a poor regioselectivity when using pyridinium salt in the telomerization of butadiene with methanol, compared to imidazolium salts [133]. This poor regioselectivity was due to the transfer of an alkyl group from pyridinium to triphenylphosphine [134,135], thus producing phosphonium salts, which cannot stabilize Pd(0), thus the appearance of the palladium black.

Strehmel *et al.* revealed that the use of *N*-butyl-4-methylpyridinium salts in free radical polymerization of *n*-butyl methacrylate in ionic liquids tends to result in relatively low degrees of polymerization and very high polydispersities, compared to the analogously substituted 1-alkyl-3-methyl imidazolium tetrafluoroborate and hexafluorophosphate, although the viscosities of **12** was higher [136].



12

4. Instability of Anions in Ionic Liquids


The stability of the anions involved in ionic liquids plays a critical role in certain applications or their toxicity. Typical ionic liquids consist of halogen containing anions such as [AlCl₄], [PF₆], [BF₄], [CF₃SO₃] or [(CF₃SO₂)₂N], which in some regard limits their 'greenness'. The presence of halogen atoms may cause serious concerns if the hydrolytic stability of the anion is poor (e.g., for [AlCl₄] and

[PF₆]) or if a thermal treatment of spent ionic liquids is required. In both cases, additional effort is needed to avoid the liberation of toxic and corrosive HF or HCl into the environment.


The lower homologues of alkylsulfate anions, namely methanesulfate and ethanesulfate, are known for their sensitivity towards hydrolysis. In the presence of water these anions form the corresponding alcohol and hydrogen sulfate at elevated temperatures. Obviously, this process is undesired for most applications since the ionic liquid system changes dramatically in its properties and an acidic proton is formed that may be a reactive species in many applications. For alkylsulfates with longer alkyl groups the sensitivity towards hydrolysis is much lower [137,138].

Anions present in ILs play a significant role in the property of ionic liquids. The assumption that all ionic liquids are non-coordinating solvents is incorrect. The coordinating property of ILs may also depend on the anion present in that ionic liquid. Ionic liquids containing PF₆⁻ ions are hydrolytically unstable, have the propensity to decompose and release HPO₂F₂, H₂PO₃F, H₃PO₄, and highly corrosive HF [139]. It is worth mentioning that several ‘uncatalyzed’ reactions reported in [cation][PF₆] ionic liquids that are, in fact, catalyzed by adventitious HF. Thus, a significant impact of IL anions is the release of HF containing compounds for which considerable care should be taken while using PF₆ containing ionic liquids. Thus, ILs should be treated with caution due to their unexplored toxicity and/or stability [140]. Rogers *et al.* identified 1-butyl-3-methylimidazolium fluoride hydrate crystallographically as a decomposition product that was obtained from hydrolytic degradation of [bmim][PF₆] during purification process [141]. They observed the evolution of acidic HF white fumes, which are colorless, highly toxic and corrosive.

Sundermeyer *et al.* designed and synthesised three new fluorinated anions BPFPA, PFTFSI, and PFNFSI related to the most prominent parent BTFSI *via* stepwise substitution of alkylsulfonyl groups by electron withdrawing pentafluorophenyl groups that attracts interests with respect to their use in highly hydrophobic, water immiscible and hydrolytically stable ILs or their use as electrolyte anions such as in lithium salts [142].




The thermal stability of this set of ILs is mainly dependent on the nature of the anion (Scheme 25). Whereas BPFPA derived ILs show a mass loss at temperatures between 170 °C and 190 °C, triflyl and nonafllyl derivatives with PFTFSI and PFNFSI anions exhibit higher thermal stabilities of up to 290 °C. Obviously, this type of ILs is undesired for most applications since the ionic liquid system changes dramatically in its properties and an acidic proton is formed that may be a reactive species in many applications.

Scheme 25. Thermal stability of ILs with fluorinated anions.

General procedure for the synthesis of the ILs ($X^- = Cl^-, Br^-$ or I^- ; HA = BPFPA-H, PFTFSI-H or PFNFSI-H)

5. Thermal Stability and Decomposition

One of the benefits of ionic liquids, when used as reaction solvents, is their wide liquid range and reportedly high thermal stability [143], which allows application at significantly elevated temperatures in various fields of science. Resistance to degradation (either by intramolecular reaction, reaction between cation and anion or reaction with the atmosphere) is always expected. Most recently, decomposition of some ionic liquids was reported to occur at much lower temperatures than previously reported [144]. A brief summary of the thermal stability of imidazolium and quaternary ammonium ionic liquids is given in this review. Almost in all ionic liquid cases, nucleophilic attack of the ionic liquid anion in a reverse Menshutkin-type reaction to yield neutral products is possible (Scheme 26) [130,145].

Scheme 26. Neutral products formation by nucleophilic attack *via* reverse Menshutkin decomposition.

It is understood from the literature that phosphonium salts are much more thermally stable than the corresponding ammonium salts and even have an edge on imidazolium salts. This is very important for the processes which operate at temperatures greater than 100 °C. Thermogravimetric analysis suggested that phosphonium ionic liquids were thermally stable up to nearly 400 °C [148]. Moreover, ionic liquids composed of cyano containing anions and cyclic quaternary ammonium cation are prone to polymerisation upon decomposition, while phosphonium-based ionic liquids give only volatile products [149]. Details regarding the thermal stability of phosphonium and pyridinium ionic liquids are not discussed in this review.

5.1. Imidazolium ionic liquids

In general, remarkable differences in T_d are observed by changing the anions, while a simple extension of alkyl chain hardly affects the T_d in imidazolium cation. ILs composed of BF_4^- , PF_6^- , NTf_2^- , are thermally more stable than corresponding halides. The relative anion stability follows the order: $\text{PF}_6^- > \text{BF}_4^- > \text{AsF}_6^- \gg \text{I}^-, \text{Br}^-, \text{Cl}^-$ (Table 8). Among anions, inorganic anions undergo endothermic thermal decomposition whereas organic anions like $(\text{C}_2\text{F}_5\text{SO}_2)_2\text{N}^-$ and $(\text{CF}_3\text{SO}_2)_2\text{N}^-$ generally restricts its exothermic decomposition up to 350 °C. This exothermicity of organic anions is likely to be a consequence of the sulfonyl groups.

Table 8. Range of T_d for some imidazolium ILs [146-150].

| Ionic liquids | T_d range (°C) |
|--|------------------|
| [bmim][dca], [bmim][Cl], [bmim][Br], [bmim][I], [Bnmim][Cl], [C ₃ mim][Cl], [bmim][Cl], [C ₅ mim][Cl], [eC ₃ mim][I], [mC ₂ mim][Cl], [mC ₃ mim][Cl], [mC ₄ mim][Cl], [mBnmim][Cl] | 240-280 |
| [bmim][PF ₆], [bmim][dca], [eC ₃ mim][I], [mBnmim][BF ₄] | 281-320 |
| [bmim][BF ₄], [bmim][OTf], [bdmim][PF ₆], [bdmim][BF ₄], [C ₃ mim][BF ₄], [bmim][BF ₄], [Bnmim][BF ₄], [decimim][BPh ₄], [eC ₃ mim][BF ₄], [emim][PF ₆], [mC ₂ mim][BF ₄], [mC ₃ mim][BF ₄], [mC ₄ mim][BF ₄] | 360-400 |
| [C ₅ mim][BF ₄], [bmim][methide], [bmim][NTf ₂], [emim][BF ₄], [pmmim][NTf ₂], [emim][NTf ₂] | 401-450 |
| [bdmim][N ₃], [C ₄ mim] ₂ [ZnBr ₂ Cl ₂], [pmim][NTf ₂] | 451-520 |

A thermolysis study of dialkylimidazolium salts reports that the decomposition is a $\text{S}_{\text{N}}2$ process that results in a mixture of *N*-alkylimidazoles and 1-alkylhalides [151]. Ohtani *et al.* studied the thermal decomposition behaviours of several imidazolium based ionic liquids by pyrolysis-gas chromatography [152]. They observed that the thermal decomposition proceeds through C-N bond cleavage. As for the imidazolium halides, corresponding haloalkanes and 1-alkylimidazoles were predominantly formed through nucleophilic attack of halide anions to the alkyl groups, which could preferably occur at methyl groups probably due to a steric effect. Meanwhile, imidazole rings did not decompose under the experimental conditions at around 550 °C.

Suslick *et al.* have investigated the stability of a variety of ionic liquids such as [bmim][Cl], [bmim][BF₄], [bmim][PF₆], urea ammonium nitrate (UAN), and decylmethylimidazolium tetraphenyl-

borate under ultrasound conditions and reported that, upon sonication, all of the imidazolium ionic liquids darkened from colorless to amber while UAN did not undergo a color change [153]. $^1\text{H-NMR}$ analysis of [bmim][Cl] and [bmim][BF₄] also indicated the appearance of new peaks in the imidazole region. Imidazolium ionic liquids produced gases/vapors that contained trace amounts of light hydrocarbons and nitriles consistent with the decomposition of imidazoles. Thus, the ionic liquids do decompose under ultrasonic conditions.

5.2. Quaternary ammonium ionic liquids

T_d of quaternary ammonium ILs give a wide range for a particular pair of cation and anion (Table 9).

Table 9. Range of T_d for some quaternary ammonium ILs [150].

| Entry | Ionic liquid | T_d range (° C) |
|-------|--------------------------------------|-------------------|
| 1 | [NH ₄][NO ₃] | 160 |
| 2 | [TMA][BF ₄] | 688-808 |
| 3 | [TEA][BF ₄] | 663-745 |
| 4 | [TPA][BF ₄] | 605-710 |
| 5 | [TBA][BF ₄] | 598-705 |

Previously, Sheikh studied the thermal decomposition of R₄NBX₄, where R is ethyl and X are Cl, Br, and phenyl dichloride [154]. Prasad *et al.* studied the onset temperatures of thermal decomposition and showed that, as the size of the substituted alkyl group increases, the thermal stability of these compounds decreases [150]. It is well accepted that quaternary ammonium compounds of the type R₄N⁺X⁻ decompose thermally to yield an amine (R₃N) and the corresponding alkyl compound. In a prelude to the study of the effects of R₄N⁺X⁻ on ammonium perchlorate (AP, an oxidant) decomposition, they interpreted that the amine (R₃N) formed by decomposition of IL can interact with the HClO₄ produced during the thermal decomposition of AP, thus forming an amine perchlorate [155,156].

5.3. Decomposition

As organic salts, ionic liquids are prone to oxidative degradation, particularly with vigorous oxidizing agents has been demonstrated by Pernak *et al.* with KMnO₄ [157] and O₃ [158]. They observed that the cations were ultimately oxidized to CO₂ and H₂O, while the anions were not oxidized. They observed that ILs undergo decomposition under the effect of ozone in aqueous solution more efficiently. Dynamically growing application of ILs induces misgivings as to their utilization when the real effects of cations and anions are unrevealed.

Shkrob *et al.* reported that the radiation stability from pulse radiolysis studies indicate that in ILs composed of aromatic cations (such as imidazolium and pyridinium), the electrons rapidly attach to the ring to form the corresponding neutral radicals [159]. In another intentional degradation study,

Stepnowski and Zaleska demonstrated that imidazolium and pyridinium ionic liquids are degraded by a combination of UV radiation and photocatalysis [160].

6. Conclusions

Our review emphasizes the urgent need for understanding the fundamental chemistry of ionic liquids as their incompatibility is now being revealed gradually. Most repeated justifications that ILs are inert and do not interact with the reagents is only an assumption until its properties are clearly studied. Unique and hidden chemistry of ILs limits their application in certain synthetic reactions. ILs, like all other solvents are chemicals which have intrinsic reactivity and thus the acidic nature, instability of anions, thermal stability, and degradation with oxidising agents has always to be kept in mind while choosing ionic liquids as reaction media.

Acknowledgements

We would like to thank V. Rajendiran and Mani Ramanathan for their gracious assistance in gathering literatures. This work was supported by the Taiwan Fellowship (to S.S.), the NCCU Predoctoral Scholarship (to V.S.), a multi-year Grant-in-Aid from ANT Technology (Taiwan, ROC) (to M.C.T. and Y.H.C.), and a multi-year grant from the National Science Council (Taiwan, ROC) (to Y.H.C.).

References and Notes

1. Xue, H.; Verma, R.; Shreeve, J.M. Review of ionic liquids with fluorine-containing anions. *J. Fluorine Chem.* **2006**, *127*, 159-176.
2. Singh, R.P.; Verma, R.D.; Meshri, D.T.; Shreeve, J.M. Energetic nitrogen-rich salts and ionic liquids. *Angew. Chem. Int. Ed.* **2006**, *45*, 3584-3601.
3. MacFarlane, D.R.; Pringle, J.M.; Johansson, K.M.; Forsyth, S.A.; Forsyth, M. Lewis base ionic liquids. *Chem. Commun.* **2006**, 1905-1917.
4. Canal, J.P.; Ramnial, T.; Dickie, D.A.; Clyburne, J.A.C. From the reactivity of *N*-heterocyclic carbenes to new chemistry in ionic liquids. *Chem. Commun.* **2006**, 1809-1818.
5. Abedin, S.Z.E.; Endres, F. Ionic liquids: The link to high-temperature molten salts? *Acc. Chem. Res.* **2007**, *40*, 1106-1113.
6. Rogers, R.D.; Voth, G.A. Ionic liquids. *Acc. Chem. Res.* **2007**, *40*, 1077-1078.
7. Welton, T. Room-temperature ionic liquids. Solvents for synthesis and catalysis. *Chem. Rev.* **1999**, *99*, 2071-2084.
8. Wasserscheid, P.; Keim, W. Ionic liquids - New "Solutions" for transition metal catalysis. *Angew. Chem. Int. Ed.* **2000**, *39*, 3772-3789.
9. Gordon, C.M. New developments in catalysis using ionic liquids. *Appl. Catal. A.* **2001**, *222*, 101-117.
10. Zhao, D.; Wu, M.; Kou, Y.; Min, E. Ionic liquids: applications in catalysis. *Catal. Today* **2002**, *74*, 157-189.

11. Dupont, J.; de Souza, R.F.; Suarez, P.A.Z. Ionic liquid (molten salt) phase organometallic catalysis. *Chem. Rev.* **2002**, *102*, 3667-3692.
12. Plechkova, N.V.; Seddon, K.R. Ionic liquids: "Designer" solvents for green chemistry. In *Methods and Reagents for Green Chemistry*; Tundo, P., Perosa, A., Zecchini, F., Eds.; Wiley: Hoboken, NJ, USA, 2007; pp. 103-130.
13. Rogers, R.D.; Seddon, K.R.; Volkov, S. In *Green Industrial Applications of Ionic Liquids*, 1st ed.; Springer: Norwell, MA, USA, 2003.
14. Earle, M.J.; Seddon, K.R. Ionic liquids. Green solvents for the future. *Pure Appl. Chem.* **2000**, *72*, 1391-1398.
15. Haumann, M.; Riisager, A. Hydroformylation in room temperature ionic liquids (RTILs): Catalyst and process developments. *Chem. Rev.* **2008**, *108*, 1474-1497.
16. Parvulescu, V.I.; Hardacre, C. Catalysis in ionic liquids. *Chem. Rev.* **2007**, *107*, 2615-2665.
17. Martins, M.A.P.; Frizzo, C.P.; Moreira, D.N.; Zanatta, N.; Bonacorso, H.G. Ionic liquids in heterocyclic synthesis. *Chem. Rev.* **2008**, *108*, 2015-2050.
18. Miao, W.; Chan, T.H. Ionic-liquid-supported synthesis: A novel liquid-phase strategy for organic synthesis. *Acc. Chem. Res.* **2006**, *39*, 897-908.
19. Han, X.; Armstrong, D.W. Ionic liquids in separations. *Acc. Chem. Res.* **2007**, *40*, 1079-1086.
20. Zhao, H.; Xia, S.; Ma, P. Use of ionic liquids as 'green' solvents for extractions. *J. Chem. Technol. Biotechnol.* **2005**, *80*, 1089-1096.
21. Visser, A.E.; Swatloski, R.P.; Reichert, W.M.; Mayton, R.; Sheff, S.; Wierzbicki, A.; Davis, J.H.; Rogers, R.D. Task-specific ionic liquids incorporating novel cations for the coordination and extraction of Hg²⁺ and Cd²⁺: Synthesis, characterization, and extraction studies. *Environ. Sci. Technol.* **2002**, *36*, 2523-2529.
22. Huddleston, J.G.; Rogers, R.D. Room temperature ionic liquids as novel media for 'clean' liquid-liquid extraction. *Chem. Commun.* **1998**, *16*, 1765-1766.
23. Hapiot, P.; Lagrost, C. Electrochemical reactivity in room-temperature ionic liquids. *Chem. Rev.* **2008**, *108*, 2238-2264.
24. MacFarlane, D.R.; Forsyth, M.; Howlett, P.C.; Pringle, J.M.; Sun, J.; Annat, G.; Neil, W.; Izgorodina, E.I. Ionic liquids in electrochemical devices and processes: Managing interfacial electrochemistry. *Acc. Chem. Res.* **2007**, *40*, 1165-1173.
25. Ichikawa, T.; Yoshio, M.; Hamasaki, A.; Mukai, T.; Ohno, H.; Kato, T. Self-organization of room-temperature ionic liquids exhibiting liquid-crystalline bicontinuous cubic phases: Formation of nano-ion channel networks. *J. Am. Chem. Soc.* **2007**, *129*, 10662-10663.
26. Biswas, K.; Rao, C.N.R. Use of ionic liquids in the synthesis of nanocrystals and nanorods of semiconducting metal chalcogenides. *Chem. Eur. J.* **2007**, *13*, 6123-6129.
27. Wang, Y.; Yang, H. Synthesis of CoPt nanorods in ionic liquids. *J. Am. Chem. Soc.* **2005**, *127*, 5316-5317.
28. Zhu, Y.; Wang, W.; Qi, R.; Hu, X. Microwave-assisted synthesis of single-crystalline Tellurium nanorods and nanowires in ionic liquids. *Angew. Chem. Int. Ed.* **2004**, *43*, 1410-1414.
29. Rantwijk, F.V.; Sheldon, R.A. Biocatalysis in ionic liquids. *Chem. Rev.* **2007**, *107*, 2757-2785.
30. Greaves, T.L.; Drummond, C.J. Protic ionic liquids: Properties and applications. *Chem. Rev.* **2008**, *108*, 206-237.

31. Schneider, S.; Hawkins, T.; Rosander, M.; Vaghjiani, G.; Chambreau, S.; Drake, G. Ionic liquids as hypergolic fuels. *Energy Fuels* **2008**, *22*, 2871-2872.
32. Zhao, H. Innovative applications of ionic liquids as "GREEN" engineering liquids. *Chem. Eng. Commun.* **2006**, *193*, 1660-1677.
33. Song, C.E. Immobilisation of chiral catalysts: easy recycling of catalyst and improvement of catalytic efficiencies. *Ann. Rep. Prog. Chem. Sect. C* **2005**, *101*, 143-173.
34. Song, C.E. *Methodologies in Asymmetric Catalysis*; Malhotra, S.V., Ed.; ACS Symposium Series 880; American Chemical Society: Washington, DC, USA, 2004; pp. 145-160.
35. Song, C.E. Enantioselective chemo- and bio-catalysis in ionic liquids. *Chem. Commun.* **2004**, 1033-1043.
36. Chowdhury, S.; Mohan, R.S.; Scott, J.L. Reactivity of ionic liquids. *Tetrahedron* **2007**, *63*, 2363-2389.
37. Ennis, E.; Handy, S.T. The chemistry of the C2 position of imidazolium room temperature ionic liquids. *Curr. Org. Syn.* **2007**, *4*, 381-389.
38. Hassoun, E.A.; Abraham, M.; Kini, V.; Al-Ghafri, M.; Abushaban, A. Cytotoxicity of the ionic liquid, 1-*N*-butyl-3-methyl imidazolium chloride. *Res. Commun. Pharmacol. Toxicol.* **2002**, *7*, 23-31.
39. Holbrey, J.D.; Seddon, K.R. ionic liquids. *Clean Technol. Environ. Policy* **1999**, *1*, 223-236.
40. Seddon, K.R. Ionic liquids for clean technology. *J. Chem. Tech. Biotech.* **1997**, *68*, 351-356.
41. Sheldon, R. Catalytic reactions in ionic liquids. *Chem. Commun.* **2001**, 2399-2407.
42. Gordon, C.M. New developments in catalysis using ionic liquids. *Appl. Catal. A.* **2001**, *222*, 101-117.
43. Helene, O.B.; Magna, L. Ionic liquids: Perspectives for organic and catalytic reactions. *J. Mol. Catal. A Chem.* **2002**, *182*, 419-437.
44. Dupont, J.; de Souza, R.F.; Suarez, P.A.Z. Ionic liquid (Molten Salt) phase organometallic catalysis. *Chem. Rev.* **2002**, *102*, 3667-3692.
45. Walden, P. Ueber die Molekulargrösse und elektrische Leitfähigkeit einiger geschmolzenen salze. *Bull. Acad. Imper. Sci. St. Petersburg* **1914**, *8*, 405-422.
46. Chum, H.L.; Koch, V.R.; Miller, L.L.; Osteryoung, R.A. Electrochemical scrutiny of organometallic iron complexes and hexamethylbenzene in a room temperature molten salt. *J. Am. Chem. Soc.* **1975**, *97*, 3264-3267.
47. Wilkes, J.S.; Levisky, J.A.; Wilson, R.A.; Hussey, C.L. Electron transfer. 52. Reactions of dihydriboflavin with metal-center oxidants. *Inorg. Chem.* **1982**, *21*, 1236-1264.
48. Wilkes, J.S.; Zaworotko, M.J. Air and water stable 1-ethyl-3-methylimidazolium based ionic liquids. *J. Chem. Soc. Chem. Commun.* **1992**, 965-967.
49. Natalia, V.P.; Seddon, K.R. Applications of ionic liquids in the chemical industry. *Chem. Soc. Rev.* **2008**, *37*, 123-150.
50. Lancaster, M. *Green Chemistry: An Introductory Text*; Royal Society of Chemistry: Cambridge, UK, 2002.
51. Wilkes, J.S. A short history of ionic liquids-from molten salts to neoteric solvents. *Green Chem.* **2002**, *4*, 73-80.
52. Freemantle, M. Ionic liquids in organic synthesis. *Chem. Eng. News* **2004**, *82*, 44-49.

53. It should be noted that under specific conditions, some ILs are, in fact, combustible. However, the bulk properties of an IL are dramatically different than, for example, fine mists which in many ways are comparable to normal solvent vapors.
54. Smiglak, M.; Reichert, W.M.; Holbrey, J.D.; Wilkes, J.S.; Sun, L.; Thrasher, J.S.; Kirichenko, K.; Singh, S.; Katritzky, A.R.; Rogers, R.D. Combustible ionic liquids by design: is laboratory safety another ionic liquid myth? *Chem. Commun.* **2006**, 2554-2556.
55. Holbrey, J.D.; Seddon, K.R. The phase behaviour of 1-alkyl-3-methylimidazolium tetrafluoroborates; Ionic liquids and ionic liquid crystals. *J. Chem. Soc. Dalton Trans.* **1999**, 2133-2139.
56. Wasserscheid, P., Welton, T., Eds. *Ionic Liquids in Synthesis*; Wiley-VCH, Weinheim, Germany, 2003.
57. Ranu, B.C.; Banerjee, S. Ionic liquid as catalyst and reaction medium. The dramatic influence of a task-specific ionic liquid, [bmIm]OH, in Michael addition of active methylene compounds to conjugated ketones, carboxylic esters, and nitriles. *Org. Lett.* **2005**, 7, 3049-3052.
58. Yan, N.; Yang, X.; Fei, Z.; Li, Y.; Kou, Y.; Dyson, P.J. Solvent-enhanced coupling of sterically hindered reagents and aryl chlorides using functionalized ionic liquids. *Organometallics* **2009**, 28, 937-939.
59. Handy, S.T. Room temperature ionic liquids: Different classes and physical properties. *Curr. Org. Chem.* **2005**, 9, 959-988.
60. Canal, J.P.; Ramnial, T.; Dickie, D.A.; Clyburne, J.A.C. From the reactivity of *N*-heterocyclic carbenes to new chemistry in ionic liquids. *Chem. Commun.* **2006**, 1809-1818.
61. Olofson, R.A.; Thompson, W.R.; Michelman, J.S. Heterocyclic nitrogen ylides. *J. Am. Chem. Soc.* **1964**, 86, 1865-1866.
62. Handy, S.T.; Okello, M. The 2-position of imidazolium ionic liquids: Substitution and exchange. *J. Org. Chem.* **2005**, 70, 1915-1918.
63. Alder, R.W.; Allen, P.R.; Williams, S.J. Stable carbenes as strong bases. *J. Chem. Soc. Chem. Commun.* **1995**, 1267-1268.
64. Amyes, T.L.; Diver, S.T.; Richard, J.P.; Rivas, F.M.; Toth, K. Formation and stability of *N*-heterocyclic carbenes in water: The carbon acid pKa of imidazolium cations in aqueous solution. *J. Am. Chem. Soc.* **2004**, 126, 4366-4374.
65. Kim, Y.J.; Streitwieser, A. Basicity of a stable carbene, 1,3-Di-tert-butylimidazol-2-ylidene, in THF. *J. Am. Chem. Soc.* **2002**, 124, 5757-5761.
66. Magill, A.M.; Cavell, K.J.; Yates, B.F. Basicity of nucleophilic carbenes in aqueous and nonaqueous solvents theoretical predictions. *J. Am. Chem. Soc.* **2004**, 126, 8717-8724.
67. Fahlbusch, T.; Frank, M.; Schatz, J.; Schuehle, D.T. Kinetic acidity of supramolecular imidazolium salts-effects of substituent, preorientation, and counterions on H/D exchange rates. *J. Org. Chem.* **2006**, 71, 1688-1691.
68. Cheng, J.Y.; Chu, Y.H. 1-Butyl-2,3-trimethyleneimidazolium bis(trifluoromethyl sulfonyl) imide ([b-3C-im][NTf₂]): a new, stable ionic liquid. *Tetrahedron Lett.* **2006**, 47, 1575-1579.
69. Fraser, R.R.; Mansour, T.S.; Savard, S. Acidity measurements in THF. V. Heteroaromatic compounds containing 5-membered rings. *Can. J. Chem.* **1985**, 63, 3505-3509.

70. Wanzlick, H.W.; Schikora, E. Ein neuer Zugang zur Carben-Chemie. *Angew. Chem.* **1960**, *72*, 494.
71. Arduengo, A.J., III; Harlow, R.L.; Kline, M. A stable crystalline carbene. *J. Am. Chem. Soc.* **1991**, *113*, 361-363.
72. Arduengo, A.J., III. Looking for stable carbenes: The difficulty in starting anew. *Acc. Chem. Res.* **1999**, *32*, 913-921.
73. Amyes, T.L.; Diver, S.T.; Richard, J.P.; Rivas, F.M.; Toth, K. Formation and stability of *N*-heterocyclic carbenes in water: The carbon acid pK_a of imidazolium cations in aqueous solution. *J. Am. Chem. Soc.* **2004**, *126*, 4366-4374.
74. Chen, H.; Justes, D.R.; Cooks, G.R. Proton affinities of *N*-heterocyclic carbene super bases. *Org. Lett.* **2005**, *7*, 3949-3952.
75. Arduengo, A.J.; Goerlich, J.R.; Marshall, W.J. A stable diaminocarbene. *J. Am. Chem. Soc.* **1995**, *117*, 11027-11028.
76. Alcarazo, M.; Roseblade, S.J.; Alonso, E.; Fernandez, R.; Alvarez, E.; Lahoz, F.J.; Lassaletta, J.M. 1,3-Bis(*N,N*-dialkylamino)imidazolin-2-ylidenes: Synthesis and reactivity of a new family of stable *N*-heterocyclic carbenes. *J. Am. Chem. Soc.* **2004**, *126*, 13242-13243.
77. Rosa, J.N.; Afonso, C.A.M.; Santos, A.G. Ionic liquids as a recyclable reaction medium for the Baylis-Hillman reaction. *Tetrahedron* **2001**, *57*, 4189-4193.
78. Aggarwal, V.K.; Emme, I.; Mereu, A. Unexpected side reactions of imidazolium-based ionic liquids in the base-catalysed Baylis-Hillman reaction. *Chem. Commun.* **2002**, 1612-1613.
79. Rosa, J.N.; Afonso, C.A.M.; Santos, A.G. Ionic liquids as a recyclable reaction medium for the Baylis-Hillman reaction. *Tetrahedron* **2001**, *57*, 4189-4193.
80. Hsu, J.C.; Yen, Y.H.; Chu, Y.H. Baylis-Hillman reaction in [bdmim][PF₆] ionic liquid. *Tetrahedron Lett.* **2004**, *45*, 4673-4676.
81. Itoh, T.; Kude, K.; Hayase, S.; Kawatsura, M. Design of ionic liquids as a medium for the Grignard reaction. *Tetrahedron Lett.* **2007**, *48*, 7774-7777.
82. Jurcik, V.; Wilhelm, R. An imidazolinium salt as ionic liquid for medium and strong bases. *Green Chem.* **2005**, *7*, 844-848.
83. Handy, S.T. Grignard reactions in imidazolium ionic liquids. *J. Org. Chem.* **2006**, *71*, 4659-4662.
84. Ennis, E.; Handy, S.T. The chemistry of the C2 position of imidazolium room temperature ionic liquids. *Curr. Org. Synt.* **2007**, *4*, 381-389.
85. Ebner, G.; Schiehser, S.; Potthast, A.; Rosenau, T. Side reaction of cellulose with common 1-alkyl-3-methylimidazolium-based ionic liquids. *Tetrahedron Lett.* **2008**, *49*, 7322-7324.
86. Carmichael, A.J.; Earle, M.J.; Holbrey, J.D.; McCormac, P.B.; Seddon, K.R. The Heck Reaction in ionic liquids: A multiphasic catalyst system. *Org. Lett.* **1999**, *1*, 997-1000.
87. Herrmann, W.A.; Böhm, V.P.W. Heck reaction catalyzed by phospho-palladacycles in non-aqueous ionic liquids. *J. Organomet. Chem.* **1999**, *572*, 141-145.
88. Böhm, V.P.W.; Herrmann, W.A. Nonaqueous ionic liquids: Superior reaction media for the catalytic Heck-vinylation of chloroarenes. *Chem. Eur. J.* **2000**, *6*, 1017-1025.
89. Xu, L.; Chen, W.; Xiao, J. Heck reaction in ionic liquids and the *in situ* identification of *N*-heterocyclic carbene complexes of Palladium. *Organometallics* **2000**, *19*, 1123-1127.

90. Herrmann, W.A.; Bohm, V.P.W.; Reisinger, C.P. Application of palladacycles in Heck type reactions. *J. Organomet. Chem.* **1999**, *576*, 23-41.
91. Gründeman, S.; Kovacevic, A.; Albrecht, M.; Faller, J.W.; Crabtree, R.H. Abnormal ligand binding and reversible ring hydrogenation in the reaction of imidazolium salts with IrH₅(PPh₃)₂. *J. Am. Chem. Soc.* **2002**, *124*, 10473-10481.
92. McGuinness, D.S.; Cavell, K.J.; Skelton, B.W.; White, A.H. Zerovalent palladium and nickel complexes of heterocyclic carbenes: Oxidative addition of organic halides, carbon-carbon coupling processes, and the Heck reaction. *Organometallics* **1999**, *18*, 1596-1605.
93. Deshmukh, R.R.; Rajagopal, R.; Srinivasan, K.V. Ultrasound promoted C-C bond formation: Heck reaction at ambient conditions in room temperature ionic liquids. *Chem. Commun.* **2001**, 1544-1545.
94. Mathews, C.J.; Smith, P.J.; Welton, T.; White, A.J.P.; Williams, D.J. *In situ* formation of mixed phosphine-imidazolylidene Palladium complexes in room-temperature ionic liquids. *Organometallics* **2001**, *20*, 3848-3850.
95. McLachlan, F.; Mathews, C.J.; Smith, P.J.; Welton, T. Palladium-catalyzed Suzuki cross-coupling reactions in ambient temperature ionic liquids: Evidence for the importance of palladium imidazolylidene complexes. *Organometallics* **2003**, *22*, 5350-5357.
96. Wong, H.; Han, S.; Livingston, A.G. The effect of ionic liquids on product yield and catalyst stability. *Chem. Eng. Sci.* **2006**, *61*, 1338-1341.
97. Sirieix, J.; Ossberger, M.; Betzemeier, B.; Knochel, P. Palladium catalyzed cross-couplings of organozincs in ionic liquids. *Synlett.* **2000**, 1613-1615.
98. Law, M.C.; Wong, K.Y.; Chan, T.H. Organometallic reactions in ionic liquids. Alkylation of aldehydes with diethylzinc. *Green Chem.* **2004**, *6*, 241-244.
99. Gholap, A.R.; Venkatesan, K.; Pasricha, R.; Daniel, T.; Lahoti, R.; Srinivasan, K.V. Copper- and ligand-free Sonogashira reaction catalyzed by Pd(0) nanoparticles at ambient conditions under ultrasound irradiation. *J. Org. Chem.* **2005**, *70*, 4869-4872.
100. Magna, L.; Chauvin, Y.; Niccolai, G.P.; Basset, J.M. The importance of imidazolium substituents in the use of imidazolium-based room-temperature ionic liquids as solvents for Palladium-catalyzed telomerization of butadiene with methanol. *Organometallics* **2003**, *22*, 4418-4425.
101. Earle, M.J.; Katdare, S.P.; Seddon, K.R. Paradigm confirmed: The first use of ionic liquids to dramatically influence the outcome of chemical reactions. *Org. Lett.* **2004**, *6*, 707-710.
102. Lancaster, N.L.; Llopis-Mestre, V. Aromatic nitrations in ionic liquids: The importance of cation choice. *Chem. Commun.* **2003**, 2812-2813.
103. Chiappe, C.; Capraro, D.; Conte, V.; Pieraccini, D. Stereoselective halogenations of alkenes and alkynes in ionic liquids. *Org. Lett.* **2001**, *3*, 1061-1063.
104. Tseng, M.C.; Liang, Y.M.; Chu, Y.H. Synthesis of fused tetrahydro- β -carbolinequinoxalinones in 1-*n*-butyl-2,3-dimethylimidazolium bis(trifluoromethyl sulfonyl) imide ([bdmim][NTf₂]) and 1-*n*-butyl-2,3-dimethylimidazolium perfluorobutylsulfonate ([bdmim][PF₄SO₃]) ionic liquids. *Tetrahedron Lett.* **2005**, *46*, 6131-6136.
105. Kan, H.C.; Tseng, M.C.; Chu, Y.H. Bicyclic imidazolium-based ionic liquids: synthesis and characterization. *Tetrahedron* **2007**, *63*, 1644-1653.

106. Lin, Y.L.; Cheng, J.Y.; Chu, Y.H. Microwave-accelerated Claisen rearrangement in bicyclic imidazolium [b-3C-im][NTf₂] ionic liquid. *Tetrahedron* **2007**, *63*, 10949-10957.
107. McNulty, J.; Capretta, A.; Wilson, J.; Dyck, J.; Adjabeng, G.; Robertson, A.J. Suzuki cross-coupling reactions of aryl halides in phosphonium salt ionic liquid under mild conditions. *Chem. Commun.* **2002**, 1986-1987.
108. Gerritsma, D.A.; Robertson, A.; McNulty, A.; Capretta, A. Heck reactions of aryl halides in phosphonium salt ionic liquids: library screening and applications. *Tetrahedron Lett.* **2004**, *45*, 7629-7631.
109. Bradaric, C.J.; Downard, A.; Kennedy, C.; Robertson, A.J.; Zhou, Y. Industrial preparation of phosphonium ionic liquids. *Green Chem.* **2003**, *5*, 143-152.
110. Kim, Y.J.; Varma, R.S. Tetrahaloindate(III)-based ionic liquids in the coupling reaction of carbon dioxide and epoxides to generate cyclic carbonates: H-bonding and mechanistic studies. *J. Org. Chem.* **2005**, *70*, 7882-7891.
111. Ramnial, T.; Daisuke, D.I.; Clyburne, J.A.C. Phosphonium ionic liquids as reaction media for strong bases. *Chem. Commun.* **2005**, 325-327.
112. Gorodetsky, B.; Ramnial, T.; Branda, N.R.; Clyburne, J.A.C. Electrochemical reduction of an imidazolium cation: a convenient preparation of imidazol-2-ylidenes and their observation in an ionic liquid. *Chem. Commun.* **2004**, 1972-1973.
113. Ramnial, T.; Taylor, S.A.; Bender, M.L.; Gorodetsky, B.; Lee, P.T.K.; Dickie, D.A.; McCollum, B.M.; Pye, C.C.; Walsby, C.J.; Clyburne, J.A.C. Carbon-centered strong bases in phosphonium ionic liquids. *J. Org. Chem.* **2008**, *73*, 801-812.
114. Ramnial, T.; Ino, D.D.; Clyburne, J.A.C. Phosphonium ionic liquids as reaction media for strong bases. *Chem. Commun.* **2005**, 325-327.
115. This ionic liquid is available from Cytec (CYPHOS-IL 101) and Fluka/Riedel-de Haën. According to Cytec product description, the ionic liquid typically assays 96 to 97% trihexyl(tetradecyl)phosphonium chloride (1) with trace contaminants of tetradecene isomers (0.1-0.4%), hydrochloric acid (0.1-0.5%), and trihexylphosphonium hydrochloride (0.1-1.2%).
116. Tseng, M.C.; Kan, H.C.; Chu, Y.H. Reactivity of trihexyl(tetradecyl)phosphonium chloride, a room-temperature phosphonium ionic liquid. *Tetrahedron Lett.* **2007**, *48*, 9085-9089.
117. Abraham, S.J.; Criddle, W.J. Pyrolysis-gas chromatography of quaternary phosphonium compounds. *J. Anal. Appl. Pyrolysis* **1985**, *7*, 337-349.
118. Abraham, S.J.; Criddle, W.J. Anion effects in the pyrolysis-gas chromatography of quaternary phosphonium compounds. *J. Anal. Appl. Pyrolysis* **1985**, *9*, 65-74.
119. Gabriel, S.; Weiner, J. Ueber einige Abkömmlinge des Propylamins. *Ber.* **1888**, *21*, 2669-2679.
120. Walden, P. Ueber die Molekulargrösse und elektrische Leitfähigkeit einiger geschmolzenen salze. *Bull. Acad. Sci. St. Petersburg* **1914**, 405-422.
121. Sato, T.; Maruo, T.; Marukane, S.; Takagi, K. Ionic liquids containing carbonate solvent as electrolytes for lithium ion cells. *J. Power Sources* **2004**, *138*, 253-261.
122. Matsumoto, H.; Sakaebe, H.; Tatsumi, K. Preparation of room temperature ionic liquids based on aliphatic onium cations and asymmetric amide anions and their electrochemical properties as a lithium battery electrolyte. *J. Power Sources* **2005**, *146*, 45-50.

123. Zhou, Z.B.; Takeda, M.; Ue, M. New hydrophobic ionic liquids based on perfluoroalkyltrifluoroborate anions. *J. Fluor. Chem.* **2004**, *125*, 471-476.
124. Zhou, Z.B.; Matsumoto, H.; Tatsumi, K. Low-melting, low-viscous, hydrophobic ionic liquids: Aliphatic quaternary ammonium salts with perfluoroalkyltrifluoroborates. *Chem. Eur. J.* **2005**, *11*, 752-766.
125. Zhou, Z.B.; Matsumoto, H.; Tatsumi, K. A new class of hydrophobic ionic liquids: Trialkyl(2-methoxyethyl)ammonium perfluoroethyltrifluoroborate. *Chem. Lett.* **2004**, *33*, 886-887.
126. Zhou, Z.B.; Matsumoto, H.; Tatsumi, K. Cyclic quaternary ammonium ionic liquids with perfluoroalkyltrifluoroborates: Synthesis, characterization, and properties. *Chem. Eur. J.* **2006**, *12*, 2196-2212.
127. Calò, V.; Nacci, A.; Monopoli, A. Effects of ionic liquids on Pd-catalysed carbon-carbon bond formation. *Eur. J. Org. Chem.* **2006**, 3791-3802.
128. Calò, V.; Nacci, A.; Monopoli, A.; Lopez, L.; di Cosmo, A. Heck reaction of β -substituted acrylates in ionic liquids catalyzed by a Pd-benzothiazole carbene complex. *Tetrahedron* **2001**, *57*, 6071-6077.
129. Calò, V.; Nacci, A.; Monopoli, A.; Laera, S.; Cioffi, N. Pd nanoparticles catalyzed stereospecific synthesis of β -aryl cinnamic esters in ionic liquids. *J. Org. Chem.* **2003**, *68*, 2929-2933.
130. Gordon, J. E. Fused organic salts. III. Chemical stability of molten tetra-*n*-alkylammonium salts. Medium effects on thermal $R_4N^+X^-$ decomposition. $RBr + \Gamma = RI + Br^-$ equilibrium constant in fused salt medium. *J. Org. Chem.* **1965**, *30*, 2760-2763.
131. Kolthoff, I.M.; Bruckenstein, S.; Chantooni, M.K. Acid-base equilibria in acetonitrile. spectrophotometric and conductometric determination of the dissociation of various acids. *J. Am. Chem. Soc.* **1961**, *83*, 3927-3935.
132. Edwards, J.O. Correlation of relative rates and equilibria with a double basicity scale. *J. Am. Chem. Soc.* **1954**, *76*, 1540-1547.
133. Magna, L.; Chauvin, Y.; Niccolai, G.P.; Basset, J.M. The importance of imidazolium substituents in the use of imidazolium-based room-temperature ionic liquids as solvents for Palladium-catalyzed telomerization of butadiene with methanol. *Organometallics* **2003**, *22*, 4418-4425.
134. Kutney, J.P.; Greenhouse, R. The protection and deprotection of the pyridine nitrogen. *Synth. Commun.* **1975**, *5*, 119-124.
135. Deady, L.W.; Finlayson, W.L.; Korytky, O.L. Steric effects in forward and reverse Menschutkin reactions of some pyridines, quinolines and thiazoles. *Aust. J. Chem.* **1979**, *32*, 1735-1742.
136. Strehmel, V.; Laschewsky, A.; Wetzel, H.; Görnitz, E. Free radical polymerization of *n*-butyl methacrylate in ionic liquids. *Macromolecules* **2006**, *39*, 923-930.
137. Holbrey, J.D.; Reichert, W.M.; Swatloski, R.P.; Broker, G.A.; Pitner, W.R.; Seddon, K.R.; Rogers, R.D. Efficient, halide free synthesis of new, low cost ionic liquids: 1,3-dialkylimidazolium salts containing methyl and ethyl sulfate anions. *Green Chem.* **2002**, *4*, 407-413.
138. Wasserscheid, P.; Hal, V.R.; Bösmann, A. 1-*n*-Butyl-3-methylimidazolium ([bmim]) octylsulfate-an even 'greener' ionic liquid. *Green Chem.* **2002**, *4*, 400-404.
139. Huddleston, J.G.; Visser, A.E.; Reichert, W.M.; Willauer, H.D.; Broker, G.A.; Rogers, R.D. Characterization and comparison of hydrophilic and hydrophobic room temperature ionic liquids incorporating the imidazolium cation. *Green Chem.* **2001**, *3*, 156-164.

140. Almost without exception, current commercially available ionic liquids have no supporting toxicological data available within MSDS information. For example, in the case of [C₄mim][PF₆] most pertinent to this study, most suppliers indicate that the hazards have not been fully investigated. CHEMADA Fine Chemicals (HaNegrev, Israel) report LD₅₀ data as 300–500 mg Kg⁻¹ (rat/oral) and > 2000 mg Kg⁻¹ (rat/dermal) but indicate that the ionic liquid is stable under normal conditions and that there is no available data on hazardous decomposition products.
141. Swatloski, R.P.; Holbrey, J.D.; Rogers, R.D. Ionic liquids are not always green: hydrolysis of 1-butyl-3-methylimidazolium hexafluorophosphate. *Green Chem.* **2003**, *5*, 361-363.
142. Linder, T.; Sundermeyer, J. Three novel anions based on pentafluorophenyl amine combined with two new synthetic strategies for the synthesis of highly lipophilic ionic liquid. *Chem. Commun.* **2009**, 2914-2916.
143. Scammells, P.J.; Scott, J.L.; Singer, R.D. Ionic liquids: The neglected issues. *Aust. J. Chem.* **2005**, *58*, 155-169.
144. Del Sesto, R. E.; McCleskey, T.M.; Macomber, C.; Ott, K.C.; Koppisch, A.T.; Baker, G.A.; Burrell, A.K. Limited thermal stability of imidazolium and pyrrolidinium ionic liquids. *Thermochim. Acta* **2009**, *491*, 118-120.
145. Chan, B.K.M.; Chang, N.H.; Grimmett, M.R. The synthesis and thermolysis of imidazole quaternary salts. *Aust. J. Chem.* **1977**, *30*, 2005-2013.
146. Busi, S.; Lahtinen, M.; Kärnä, M.; Valkonen, J.; Kolehmainen, E.; Rissanen, K. Synthesis, characterization and thermal properties of nine quaternary dialkyldiaralkylammonium chlorides. *J. Mol. Struct.* **2006**, *787*, 18-30.
147. Fredlake, C.P.; Crosthwaite, J.M.; Hert, D.G.; Aki, S.N.V.K.; Brennecke, J.F. Thermophysical properties of imidazolium-based ionic liquids. *J. Chem. Eng. Data* **2004**, *49*, 954-964.
148. Kim, Y.J.; Varma, R.S. Tetrahaloindate(III)-based ionic liquids in the coupling reaction of carbon dioxide and epoxides to generate cyclic carbonates: H-bonding and mechanistic studies. *J. Org. Chem.* **2005**, *70*, 7882-7891
149. Wooster, T.J.; Johanson, K.M.; Fraser, K.J.; MacFarlane, D.R.; Scott, J.L. Thermal degradation of cyano containing ionic liquids. *Green Chem.* **2006**, *8*, 691-696.
150. Prasad, M.R.R.; Krishnan, K.; Ninan, K.N.; Krishnamurthy, V.N. Thermal decomposition of tetraalkyl ammonium tetrafluoroborates. *Thermochim. Acta* **1997**, *297*, 207-210.
151. Chan, B.K.M.; Chang, N.; Grimmett, M.R. The synthesis and thermolysis of imidazole quaternary salts. *Aust. J. Chem.* **1977**, *30*, 2005-2013.
152. Ohtani, H.; Ishimura, S.; Kumai, M. Thermal decomposition behaviors of imidazolium-type ionic liquids studied by pyrolysis-gas chromatography. *Anal. Sci.* **2008**, *24*, 1335-1340.
153. Oxley, J.D.; Prozorov, T.; Suslick, K.S. Sonochemistry and sonoluminescence of room-temperature ionic liquids. *J. Am. Chem. Soc.* **2003**, *125*, 11138-11139.
154. Sheikh, S.U. A systematic study of the thermal decomposition of tetraalkylammonium haloborates. *J. Thermal Anal.* **1980**, *18*, 299-306.
155. Nambiar, P.R.; Jain, S.R. Thermal characterization of tetramethylphosphonium perchlorate, nitrate and picrate. *Thermochim. Acta* **1974**, *9*, 295-302.
156. Nambiar, P.R.; Verneker, V.R.P.; Jain, S.R. Explosive sensitivity of methylammonium perchlorates. *J. Thermal Anal.* **1975**, *8*, 15-26.

157. Pernak, J.; Sobaszkiewicz, K.; Foksowicz-Flaczyk, J. Ionic liquids with symmetrical dialkoxymethyl-substituted imidazolium cations. *Chem. Eur. J.* **2004**, *10*, 3479-3485.
158. Pernak, J.; Goc, I.; Mirska, I. Anti-microbial activities of protic ionic liquids with lactate anion. *Green Chem.* **2004**, *6*, 323-329.
159. Shkrob, I.A.; Chemerisov, S.D. The initial stages of radiation damage in ionic liquids and ionic liquid-based extraction systems. *J. Phys. Chem. B* **2007**, *111*, 11786-11793.
160. Stepnowski, P.; Zaleska, A. Comparison of different advanced oxidation processes for the degradation of room temperature ionic liquids. *J. Photochem. Photobiol. A* **2004**, *170*, 45-50.

Sample availability: Not available.

© 2009 by the authors; licensee Molecular Diversity Preservation International, Basel, Switzerland. This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).