

ANNALI DELLA
SCUOLA NORMALE SUPERIORE DI PISA
Classe di Scienze

DANIELE BOFFI

FRANCO BREZZI

LUCIA GASTALDI

On the convergence of eigenvalues for mixed formulations

Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 4^e série, tome 25, n° 1-2 (1997), p. 131-154

http://www.numdam.org/item?id=ASNSP_1997_4_25_1-2_131_0

© Scuola Normale Superiore, Pisa, 1997, tous droits réservés.

L'accès aux archives de la revue « Annali della Scuola Normale Superiore di Pisa, Classe di Scienze » (<http://www.sns.it/it/edizioni/riviste/annaliscienze/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

On the Convergence of Eigenvalues for Mixed Formulations

DANIELE BOFFI – FRANCO BREZZI – LUCIA GASTALDI

Abstract. Eigenvalue problems for mixed formulation show peculiar features that make them substantially different from the corresponding mixed direct problems. In this paper we analyze, in an abstract framework, necessary and sufficient conditions for their convergence.

1. – Introduction

In a general way, we say that a variational problem is written in *mixed form* if it fits the following abstract setting. We assume that

- (1) Φ and Ξ are Hilbert spaces,
- (2) $a(\psi, \varphi)$ and $b(\psi, \xi)$ are bilinear forms on $\Phi \times \Phi$ and $\Phi \times \Xi$ respectively,

$a(\cdot, \cdot)$ and $b(\cdot, \cdot)$ are continuous, that is

- (3) $\exists M_a > 0 \quad \forall \psi, \varphi \in \Phi \quad a(\psi, \varphi) \leq M_a \|\psi\|_\Phi \|\varphi\|_\Phi$
 $\exists M_b > 0 \quad \forall \varphi \in \Phi, \forall \xi \in \Xi \quad b(\varphi, \xi) \leq M_b \|\varphi\|_\Phi \|\xi\|_\Xi,$

and, to simplify the presentation, we also assume that

- (4) $a(\cdot, \cdot)$ is symmetric and positive semidefinite.

Setting $\|\varphi\|_a := (a(\varphi, \varphi))^{1/2}$ (which in general will only be a seminorm on Φ) this immediately gives

- (5) $\forall \psi, \varphi \in \Phi \quad a(\psi, \varphi) \leq \|\psi\|_a \|\varphi\|_a.$

Properties (1) to (4) will be assumed to hold throughout all the paper.

For any given pair (f, g) in $\Phi' \times \Xi'$ we consider now the problem

- find (ψ, χ) in $\Phi \times \Xi$ such that
- (6)
$$\begin{cases} a(\psi, \varphi) + b(\varphi, \chi) = \langle f, \varphi \rangle & \forall \varphi \in \Phi \\ b(\psi, \xi) = \langle g, \xi \rangle & \forall \xi \in \Xi. \end{cases}$$

It is known that, in order to have existence, uniqueness and continuous dependence from the data for problem (6) it is necessary and sufficient that the bilinear forms $a(\cdot, \cdot)$ and $b(\cdot, \cdot)$ satisfy the following conditions

$$(7) \quad \text{(IS)} \quad \begin{array}{l} \text{there exists } \beta > 0, \text{ such that} \\ \inf_{\xi \in \Xi} \sup_{\varphi \in \Phi} \frac{b(\varphi, \xi)}{\|\varphi\|_{\Phi} \|\xi\|_{\Xi}} \geq \beta \end{array}$$

$$(8) \quad \text{(EK)} \quad \begin{array}{l} \text{there exists } \alpha > 0, \text{ such that} \\ a(\varphi, \varphi) \geq \alpha \|\varphi\|_{\Phi}^2 \quad \forall \varphi \in \mathbb{K} \end{array}$$

where the kernel \mathbb{K} is defined as:

$$\mathbb{K} = \{\varphi \in \Phi \text{ such that } b(\varphi, \xi) = 0 \quad \forall \xi \in \Xi\}.$$

EXAMPLE 1. Stokes problem. We take $\Phi = (H_0^1(\Omega))^2$, $\Xi = L^2(\Omega)/_{\mathbb{R}}$, $a(\psi, \varphi) = (\nabla \psi, \nabla \varphi)$, $b(\psi, \xi) = -(\operatorname{div} \psi, \xi)$ where, as usual, (\cdot, \cdot) is the inner product in $L^2(\Omega)$ or in $(L^2(\Omega))^2$. It is easy to see that (7) and (8) are satisfied. Moreover if we take $g = 0$ the solution of (6) is related to the solution of the Stokes problem

$$(9) \quad \begin{array}{ll} -\Delta \underline{u} + \nabla p = f & \text{in } \Omega \\ \operatorname{div} \underline{u} = 0 & \text{in } \Omega \\ \underline{u} = 0 & \text{on } \partial\Omega \end{array}$$

by the relations $\psi = \underline{u}$ and $\chi = p$. Approximations based on this approach are classical and are usually called approximations in the primitive variables.

EXAMPLE 2. Dirichlet problem with Lagrange multipliers. We take $\Phi = H^1(\Omega)$, $\Xi = H^{-1/2}(\partial\Omega)$, $a(\psi, \varphi) = (\nabla \psi, \nabla \varphi)$ and $b(\psi, \xi) = \langle \psi, \xi \rangle_{\partial\Omega}$ (duality between $H^{1/2}(\partial\Omega)$ and Ξ). It is easy to see that (7) and (8) are satisfied. Moreover, for every $f \in H^{-1}(\Omega)$ and $g \in H^{1/2}(\partial\Omega)$ the unique solution of (6) is related to the solution of

$$(10) \quad \begin{array}{ll} -\Delta u = f & \text{in } \Omega \\ u = g & \text{on } \partial\Omega, \end{array}$$

by the relations $\psi = u$ and $\chi = -\frac{\partial u}{\partial n}$. Approximations based on this approach where first introduced by Babuška [2].

EXAMPLE 3. Mixed formulation of second order linear elliptic problems. We take $\Phi = H(\operatorname{div}; \Omega)$, $\Xi = L^2(\Omega)$, $a(\psi, \varphi) = (\psi, \varphi)$, $b(\psi, \xi) = (\operatorname{div} \psi, \xi)$. It is easy to see that (7) and (8) are satisfied. Moreover if we take $f = 0$ the solution of (6) is related to the solution of the problem

$$(11) \quad \begin{array}{ll} \Delta u = g & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{array}$$

by the relations $\chi = u$ and $\psi = \nabla u$. Approximations based on this approach where first introduced by Raviart-Thomas [24].

EXAMPLE 4. *Biharmonic problem.* We take $\Phi = H^1(\Omega)$, $\Xi = H_0^1(\Omega)$, $a(\psi, \varphi) = (\psi, \varphi)$ and $b(\psi, \xi) = -(\nabla\psi, \nabla\xi)$. It is easy to see that (7) is satisfied, but (8) is not. However, if Ω is smooth enough and we take $f = 0$ then (6) has a unique solution, related to the solution of

$$(12) \quad \begin{aligned} \Delta^2 u &= -g && \text{in } \Omega \\ u &= \frac{\partial u}{\partial n} = 0 && \text{on } \partial\Omega, \end{aligned}$$

by the relations $\chi = u$ and $\psi = -\Delta u$. Approximations based on this approach where first introduced by Glowinski [16] and analyzed by Ciarlet-Raviart [13] and Mercier [20].

For many other examples of mixed formulations of boundary value problems related to various applications in fluidmechanics and in continuous mechanics we refer, for instance, to [9].

Let us now consider the problem of discretization. Assume that we are given two families of finite dimensional subspaces Φ_h and Ξ_h of Φ and Ξ , respectively. We consider the discretized problem:

$$(13) \quad \begin{aligned} &\text{find } (\psi_h, \chi_h) \text{ in } \Phi_h \times \Xi_h \text{ such that} \\ &\begin{cases} a(\psi_h, \varphi_h) + b(\varphi_h, \chi_h) = \langle f, \varphi_h \rangle & \forall \varphi_h \in \Phi_h \\ b(\psi_h, \xi_h) = \langle g, \xi_h \rangle & \forall \xi_h \in \Xi_h. \end{cases} \end{aligned}$$

It is known that discrete analogues of (7) and (8) are sufficient to ensure solvability of the discrete problem together with optimal error bounds. More precisely if the spaces Φ_h and Ξ_h satisfy the following conditions

$$(14) \quad \text{(DEK)} \quad \begin{aligned} &\text{there exists } \alpha > 0, \text{ independent of } h, \text{ such that} \\ &a(\varphi_h, \varphi_h) \geq \alpha \|\varphi_h\|_{\Phi}^2 \quad \forall \varphi_h \in \mathbb{K}_h \end{aligned}$$

where the discrete kernel \mathbb{K}_h is defined as

$$\mathbb{K}_h = \{\varphi_h \in \Phi_h \text{ such that } b(\varphi_h, \xi_h) = 0 \quad \forall \xi_h \in \Xi_h\}$$

and

$$(15) \quad \text{(DIS)} \quad \begin{aligned} &\text{there exists } \beta > 0, \text{ independent of } h, \text{ such that} \\ &\inf_{\xi_h \in \Xi_h} \sup_{\varphi_h \in \Phi_h} \frac{b(\varphi_h, \xi_h)}{\|\varphi_h\|_{\Phi} \|\xi_h\|_{\Xi}} \geq \beta, \end{aligned}$$

then we have unique solvability of (13) and the following error estimate

$$(16) \quad \|\psi - \psi_h\|_{\Phi} + \|\chi - \chi_h\|_{\Xi} \leq C \left(\inf_{\varphi \in \Phi_h} \|\psi - \varphi\|_{\Phi} + \inf_{\xi \in \Xi_h} \|\chi - \xi\|_{\Xi} \right).$$

As we shall see, conditions (14) and (15) are also necessary for having (16), in a suitable sense.

We turn now to the eigenvalue problems. As we can see from the examples above, the eigenvalue problem which is naturally associated with the corresponding boundary value problem in strong form (namely (9), (10), (11) or (12)) does not correspond to taking $(\lambda\psi, \lambda\chi)$ as right-hand side of (6). Instead, according with the different cases, the *natural* eigenvalue problem is obtained by taking $(\lambda\psi, 0)$ or $(0, -\lambda\chi)$ as right-hand side of (6). One expects, as for instance in [21], that (14) and (15), together with suitable compactness properties, are sufficient to ensure good convergence of the eigenvalues. However, when the problem is set in mixed variational form, compactness is more delicate to deal with. In a previous paper [5] we showed that, for the particular case of Example 3, even if the operator mapping g into u in (11) is clearly compact, assumptions (14) and (15) are not sufficient to avoid, for instance, the presence of spurious eigenvalues in the discrete spectrum. Here we address a more general problem, in abstract form, and we look for sufficient (and, possibly, necessary) conditions in order to have good approximation properties for the eigenvalue problems having either $(\lambda\psi, 0)$ or $(0, -\lambda\chi)$ at the right-hand side. As we shall see, in each of the two cases, (14) and (15) might be neither necessary nor sufficient for that.

Our approach turns out to be more similar to the one of [14] rather than the one of [8] or [1]. Important references for the study of eigenvalue problems in mixed form are [21], [3], [23]. As far as the *sufficient* conditions are concerned, we have only little improvements over the previous papers. For instance, our bilinear form $a(\cdot, \cdot)$ is not supposed to be positive definite as in the previous literature. Moreover, previous related papers deal mostly with cases in which the two components of the solution of the direct problem are both convergent, while we accept discretizations that can produce singular global matrices. On the other hand, having assumed symmetry of $a(\cdot, \cdot)$, we do not have to consider adjoint problems as in [14]. However, in practical cases, the actual gain is negligible. The major interest of the paper, in our opinion, consists in showing that our sufficient conditions are, mostly, also *necessary*, thus providing a severe test for assessing whether a given discretization is suitable for computing eigenvalues or not. This justifies, in our opinion, the apparently excessive generality of our abstract approach. Indeed, as we shall see, convergence of discrete eigenvalues does not even imply, for mixed formulations, the nonsingularity of the corresponding global matrices.

Finally we point out that in this paper we do not look for *a priori estimates* for eigenvalues and eigenvectors, but only deal with convergence. This is somehow in agreement with the fact that necessary conditions are a major issue here. However, in most cases, a priori error estimates can be readily deduced checking the last step in the proofs of sufficient conditions and/or applying the general instruments of, say, [7], [21], [3].

An outline of the paper is the following. In Section 2 we state the problem and relate the convergence of the spectrum with the uniform convergence of the resolvent operators. Moreover we point out the role of the discrete conditions (DEK) and (DIS) in order to have existence and boundedness of the different components of the solution of (13).

Section 3 and 4 are devoted to the analysis of the eigenvalue problems associated to (6) when the right-hand side is of the type $(\lambda\psi, 0)$ or $(0, -\lambda\chi)$, respectively. In both cases we state sufficient and necessary conditions for the good approximation of the spectrum. At the end of each section we will show how the known good approximations of the problems in the examples above satisfy our sufficient conditions for convergence of eigenvalues and eigenvectors, and more generally we discuss the validity of other possible approximations in light of our conditions.

2. – Statement of the problems

Let H be a Hilbert space and $T : H \rightarrow H$ a selfadjoint compact operator. To simplify the presentation we assume that T is nonnegative.

We are interested in the eigenvalues $\lambda \in \mathbb{R}$ defined by

$$(17) \quad \lambda T u = u, \quad \text{with } u \in H \setminus \{0\}.$$

In the above assumptions it is well-known that there exists a sequence $\{\lambda_i\}$ and an associated orthonormal basis $\{u_i\}$ such that

$$(18) \quad \begin{aligned} \lambda_i T u_i &= u_i, \\ 0 &\leq \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_i \leq \dots, \\ \lim_{i \rightarrow \infty} \lambda_i &= +\infty. \end{aligned}$$

We also set, for $i \in \mathbb{N}$, $E_i = \text{span}(u_i)$.

The following mapping will be useful. Let $m : \mathbb{N} \rightarrow \mathbb{N}$ be the application which to every N associates the dimension of the space generated by the eigenspaces of the first N distinct eigenvalues; that is

$$(19) \quad \begin{aligned} m(1) &= \dim \{\oplus_i E_i : \lambda_i = \lambda_1\}, \\ m(N+1) &= m(N) + \dim \{\oplus_i E_i : \lambda_i = \lambda_{m(N)+1}\}. \end{aligned}$$

Clearly, $\lambda_{m(1)}, \dots, \lambda_{m(N)}$ ($N \in \mathbb{N}$) will now be the first N *distinct* eigenvalues of (17).

Assume that we are given, for every $h > 0$, a selfadjoint nonnegative operator $T_h : H \rightarrow H$ with finite range. We denote by $\lambda_i^h \in \mathbb{R}$ the eigenvalues of the problem

$$(20) \quad \lambda T_h u = u, \quad \text{with } u \in H \setminus \{0\}.$$

Let H_h be the finite-dimensional range of T_h and $\dim H_h =: N(h)$; then T_h admits $N(h)$ real eigenvalues denoted λ_i^h such that

$$(21) \quad 0 \leq \lambda_1^h \leq \dots \leq \lambda_i^h \leq \dots \leq \lambda_{N(h)}^h.$$

The associated discrete eigenfuntions u_i^h , $i = 1, \dots, N(h)$, give rise to an orthonormal basis of H_h with respect to the scalar product of H . Let $E_i^h := \text{span}(u_i^h)$.

We assume that

$$(22) \quad \lim_{h \rightarrow 0} \|T - T_h\|_{\mathcal{L}(H)} = 0.$$

It is a classical result in spectrum perturbation theory that (22) implies the following convergence property for eigenvalues and eigenvectors:

$$(23) \quad \begin{aligned} & \forall \epsilon > 0, \quad \forall N \in \mathbb{N} \quad \exists h_0 > 0 \text{ such that } \forall h \leq h_0 \\ & \max_{i=1, \dots, m(N)} |\lambda_i - \lambda_i^h| \leq \epsilon, \\ & \hat{\delta}(\oplus_{i=1}^{m(N)} E_i, \oplus_{i=1}^{m(N)} E_i^h) \leq \epsilon, \end{aligned}$$

where $\hat{\delta}(E, F)$, for E and F linear subspaces of H , represents the gap between E and F and is defined by

$$(24) \quad \begin{aligned} \hat{\delta}(E, F) &= \max[\delta(E, F), \delta(F, E)], \\ \delta(E, F) &= \sup_{u \in E, \|u\|_H=1} \inf_{v \in F} \|u - v\|_H. \end{aligned}$$

Viceversa, it is not difficult to prove that (23) is a sufficient condition for (22).

We are interested in having (23) for eigenvalue problems in mixed form.

Let us therefore go back to the abstract framework already used in the introduction, with the assumptions therein. In particular assume, for the moment, that (7) and (8) are satisfied and that (13) has a solution for every (f, g) in $\Phi' \times \Xi'$. Problems (6) and (13) define then, in a natural way, two operators $S(f, g) = (\psi, \chi)$ (solution of (6)) and $S_h(f, g) = (\psi_h, \chi_h)$ (solution of (13)).

It is well-known (see [9]) that (DIS) and (DEK) (cfr. equations (15) and (14)) imply that the discrete operator S_h is bounded from $\Phi'_h \times \Xi'_h$ to $\Phi \times \Xi$, uniformly in h (see (16)). Moreover, the converse holds true, as it is proved in the following Lemma 1. Before it we introduce the following notation: for every $h > 0$ we define

$$(25) \quad \|f\|_{\Phi'_h} = \sup_{\varphi_h \in \Phi_h} \frac{\langle f, \varphi_h \rangle}{\|\varphi_h\|_{\Phi}} \quad \|g\|_{\Xi'_h} = \sup_{\xi_h \in \Xi_h} \frac{\langle g, \xi_h \rangle}{\|\xi_h\|_{\Xi}}.$$

LEMMA 1. *If there exists a constant C such that for all $f \in \Phi'$ and $g \in \Xi'$*

$$(26) \quad \|S_h(f, g)\|_{\Phi \times \Xi} \leq C(\|f\|_{\Phi'_h} + \|g\|_{\Xi'_h})$$

for all $h > 0$, then (DIS) and (DEK) are verified.

PROOF. Let ψ_h belong to \mathbb{K}_h , then $(\psi_h, 0, \tilde{f}, 0)$ satisfies (13) with $\langle \tilde{f}, \varphi_h \rangle := a(\psi_h, \varphi_h)$, for all $\varphi_h \in \Phi_h$. Hence the inequality (26) gives (DEK) with $\alpha = 1/(C^2 M_a)$, M_a being the continuity constant of a (see (3)).

If $\chi_h \in \Xi_h$, then $(0, \chi_h, \tilde{f}, 0)$ satisfies (13), with $\langle \tilde{f}, \varphi_h \rangle := b(\varphi_h, \chi_h)$ for all $\varphi_h \in \Phi_h$. Hence the inequality (26) yields (DIS) with $\beta = 1/C$. \square

REMARK 1. In the statement of Lemma 1 we implicitly assumed that the operator S_h was defined for every f and g . However, as it can be clearly seen in the proof, this was not really necessary. Indeed it is sufficient to assume that there exists a constant $C > 0$ such that for every $h > 0$ and for every quadruplet $(\psi_h, \chi_h, f, g) \in \Phi_h \times \Xi_h \times \Phi' \times \Xi'$ satisfying (13), one has

$$(27) \quad \|\psi_h\|_{\Phi} + \|\chi_h\|_{\Xi} \leq C(\|f\|_{\Phi'_h} + \|g\|_{\Xi'_h}).$$

This should not surprise, as (13) is always a linear system with a square matrix. \square

Consider now the eigenvalue problem. For the sake of simplicity, let us assume for the moment that there exist two Hilbert spaces H_{Φ} and H_{Ξ} such that we can identify

$$(28) \quad \begin{aligned} H_{\Phi} &\equiv H'_{\Phi}, \\ H_{\Xi} &\equiv H'_{\Xi} \end{aligned}$$

and such that

$$(29) \quad \begin{aligned} \Phi &\subseteq H_{\Phi} \subseteq \Phi' \\ \Xi &\subseteq H_{\Xi} \subseteq \Xi' \end{aligned}$$

hold with dense and continuous embedding, in a compatible way.

The restrictions of S and S_h to $H_{\Phi} \times H_{\Xi}$ define now two operators from $H_{\Phi} \times H_{\Xi}$ into itself.

As a consequence of (16) and Lemma 1, it is immediate to prove the following proposition.

PROPOSITION 1. *Assume that (DIS) and (DEK) hold. Then S_h converges uniformly to S in $\mathcal{L}(H_{\Phi} \times H_{\Xi})$ if and only if S (from $H_{\Phi} \times H_{\Xi}$ into itself) is compact.*

This proposition concludes our convergence analysis for the eigenvalue problems associated to (6) and (13). However in the applications one finds more often eigenvalue problems associated to (6) and (13) when one of the two components of the datum is zero. Let us set these eigenvalue problems in their appropriate abstract framework introducing the following operators:

$$(30) \quad \begin{aligned} C_{\Phi} : \Phi' &\rightarrow \Phi' \times \Xi' & C_{\Xi} : \Xi' &\rightarrow \Phi' \times \Xi' \\ C_{\Phi}(f) &= (f, 0) & C_{\Xi}(g) &= (0, g) \end{aligned}$$

and their adjoints

$$(31) \quad \begin{aligned} C_{\Phi}^* : \Phi \times \Xi \rightarrow \Phi & & C_{\Xi}^* : \Phi \times \Xi \rightarrow \Xi \\ C_{\Phi}^*(\varphi, \xi) = \varphi & & C_{\Xi}^*(\varphi, \xi) = \xi. \end{aligned}$$

We shall say that (6) is a *problem of the type* $\begin{pmatrix} f \\ 0 \end{pmatrix}$ if the right-hand side in (6) satisfies $g = 0$. Similarly, we shall say that (6) is a *problem of the type* $\begin{pmatrix} 0 \\ g \end{pmatrix}$ if the right-hand side in (6) satisfies $f = 0$. Correspondingly, we shall study the approximation of the eigenvalues of the following operators:

$$(32) \quad \begin{aligned} T_{\Phi} &= C_{\Phi}^* \circ S \circ C_{\Phi} : \Phi' \rightarrow \Phi, & \text{for problems of the type } \begin{pmatrix} f \\ 0 \end{pmatrix}, \\ T_{\Xi} &= C_{\Xi}^* \circ S \circ C_{\Xi} : \Xi' \rightarrow \Xi, & \text{for problems of the type } \begin{pmatrix} 0 \\ g \end{pmatrix}. \end{aligned}$$

Whenever the associated discrete problems are solvable, we can introduce the discrete counterparts of T_{Φ} and T_{Ξ} as:

$$(33) \quad \begin{aligned} T_{\Phi}^h &= C_{\Phi}^* \circ S_h \circ C_{\Phi} : \Phi' \rightarrow \Phi, & \text{for problems of the type } \begin{pmatrix} f \\ 0 \end{pmatrix}, \\ T_{\Xi}^h &= C_{\Xi}^* \circ S_h \circ C_{\Xi} : \Xi' \rightarrow \Xi, & \text{for problems of the type } \begin{pmatrix} 0 \\ g \end{pmatrix}. \end{aligned}$$

In the remaining part of this section we are going to relate the solvability and boundedness of the discrete operators with either (DIS) or (DEK).

PROPOSITION 2. *If (DEK) (see (14)) holds and $g = 0$, then problem (13) has at least one solution (ψ_h, χ_h) . Moreover ψ_h is uniquely determined by f and*

$$(34) \quad \|\psi_h\|_{\Phi} \leq \frac{1}{\alpha} \|f\|_{\Phi'_h},$$

(where α is the constant appearing in (14)).

PROOF. Let ψ_h be the unique solution of $a(\psi_h, \varphi_h) = \langle f, \varphi_h \rangle$ for all φ_h in \mathbb{K}_h . Clearly ψ_h exists, is unique and satisfies (34). Now look for χ_h in Ξ_h such that $b(\varphi_h, \chi_h) = \langle f, \varphi_h \rangle - a(\psi_h, \varphi_h)$ for all $\varphi_h \in \Phi_h$. As the right-hand side is in the polar set of \mathbb{K}_h , the system is compatible and hence has at least one solution. \square

PROPOSITION 3. *Assume that there exists a constant $C > 0$ such that for every $h > 0$ and for every quadruplet $(\psi_h, \chi_h, f, 0) \in \Phi_h \times \Xi_h \times \Phi' \times \Xi'$ satisfying (13) one has*

$$(35) \quad \|\psi_h\|_{\Phi} \leq C \|f\|_{\Phi'_h},$$

then the operator T_{Φ}^h is defined in all Φ' and (DEK) holds with $\alpha = 1/(C^2 M_a)$, M_a being the continuity constant of a (see (3)).

PROOF. With the same proof as in Lemma 1 we see that (DEK) holds true. The solvability of (13) is now a consequence of Proposition 2. \square

PROPOSITION 4. *Assume that the following weak discrete inf-sup condition holds: for every $h > 0$, there exists a constant $\beta_h > 0$ such that*

$$(36) \quad (\text{DIS}_h) \quad \inf_{\xi_h \in \Xi_h} \sup_{\varphi_h \in \Phi_h} \frac{b(\varphi_h, \xi_h)}{\|\varphi_h\|_{\Phi} \|\xi_h\|_{\Xi}} \geq \beta_h.$$

Then for every $g \in \Xi'$ and $f = 0$ problem (13) has at least one solution (ψ_h, χ_h) and χ_h is uniquely determined by g .

PROOF. The assumption (36) implies that, with obvious notation, B_h is surjective. Hence for $g \in \Xi'$ there exists at least one $\psi_g \in \Phi_h$ such that $B_h \psi_g = g$. Then find $\psi_k \in \mathbb{K}_h$ such that $a(\psi_k, \varphi_h) = -a(\psi_g, \varphi_h) \forall \varphi_h \in \mathbb{K}_h$. Finally, take $\chi_h \in \Xi_h$ such that $b(\varphi_h, \chi_h) = -a(\psi_g, \varphi_h) - a(\psi_k, \varphi_h)$ for all $\varphi_h \in \Phi_h$. Such a χ_h exists, by the same argument used in the proof of Proposition 2. Finally observe that $(\psi_g + \psi_k, \chi_h)$ solves (13) with $(0, g)$ as right-hand side.

To see the uniqueness, assume that (ψ_h^i, χ_h^i) ($i = 1, 2$) are two solutions. Clearly $a(\psi_h^1 - \psi_h^2, \varphi_h) = 0$ for all $\varphi_h \in \mathbb{K}_h$. Taking $\varphi_h = \psi_h^1 - \psi_h^2$ one obtains that $a(\psi_h^1 - \psi_h^2, \psi_h^1 - \psi_h^2) = 0$ and hence as a is symmetric and positive semidefinite, $a(\psi_h^1 - \psi_h^2, \varphi_h) = 0$ for all $\varphi_h \in \Phi_h$ (use (5)). Now $b(\varphi_h, \chi_h^1 - \chi_h^2) = 0$ for all φ_h in Φ_h and (DIS_h) implies $\chi_h^1 = \chi_h^2$. \square

PROPOSITION 5. *Assume that there exists a constant $C > 0$ such that for every $h > 0$ and for every quadruplet $(\psi_h, \chi_h, 0, g) \in \Phi_h \times \Xi_h \times \Phi' \times \Xi'$ satisfying (13) one has*

$$(37) \quad \|\chi_h\|_{\Xi} \leq C \|g\|_{\Xi'_h},$$

then the operator T_{Ξ}^h is defined in all Ξ' and the weak discrete inf-sup condition (DIS_h) holds. In general, (37) does not imply (DIS).

PROOF. Remark first that the assumption (37) implies that, with obvious notation, B_h' is injective, therefore B_h will be surjective and this implies (DIS_h) .

In order to see that (DIS) cannot be deduced in general, consider the case when $a \equiv 0$, $\Phi_h = \Xi_h$ and b is h times the scalar product in Φ_h . \square

PROPOSITION 6. *Assume that there exists a constant $C > 0$ such that for every $h > 0$ and for every quadruplet $(\psi_h, \chi_h, 0, g) \in \Phi_h \times \Xi_h \times \Phi' \times \Xi'$ satisfying (13) one has*

$$(38) \quad \|\psi_h\|_{\Phi} + \|\chi_h\|_{\Xi} \leq C \|g\|_{\Xi'_h},$$

then both T_{Ξ}^h and $C_{\Phi}^ \circ S_h \circ C_{\Xi}$ are defined on Ξ' and (DIS) holds with $\beta = 1/C$.*

PROOF. Remark first that, from Proposition 4, problem (13) has at least one solution for every $g \in \Xi'$, but now the estimate (38) ensures that such solution is unique. Hence $C_\Phi^* \circ S_h \circ C_\Xi$ is also well-defined in Ξ' . Let now ξ_h be an element of Ξ_h , and let $g \in \Xi'$ be such that $\|g\|_{\Xi'_h} = 1$ and $\langle g, \xi_h \rangle = \|\xi_h\|_\Xi$. Taking $\psi_h^* = C_\Phi^* \circ S_h \circ C_\Xi g$ we have

$$(39) \quad \frac{b(\psi_h^*, \xi_h)}{\|\psi_h^*\|_\Phi} = \frac{\langle g, \xi_h \rangle}{\|\psi_h^*\|_\Phi} = \frac{\|\xi_h\|_\Xi}{\|\psi_h^*\|_\Phi} \geq \frac{1}{C} \|\xi_h\|_\Xi. \quad \square$$

PROPOSITION 7. *If there exists $C > 0$ such that*

$$(40) \quad \|C_\Xi^* \circ S_h \circ C_\Phi\|_{\mathcal{L}(\Phi'_h, \Xi_h)} \leq C$$

for every $h > 0$, then (DIS) holds with $\beta = 1/C$.

PROOF. The same proof as in Lemma 1. □

We see from Propositions 3 and 7, that for problems of the type $\begin{pmatrix} f \\ 0 \end{pmatrix}$ the estimate (35) on ψ_h implies (DEK) and the estimate (40) on χ_h implies (DIS). Analogue properties do not entirely hold for problems of the type $\begin{pmatrix} 0 \\ g \end{pmatrix}$.

3. – Problems of the type $\begin{pmatrix} f \\ 0 \end{pmatrix}$

In this section, together with (1)-(4), we assume that (EK) and (IS) are verified. We also assume that we are given a Hilbert space H_Φ (that we shall identify with its dual space H'_Φ) such that

$$(41) \quad \Phi \subseteq H_\Phi \subseteq \Phi'$$

with continuous and dense embeddings. We consider the eigenvalue problem

$$(42) \quad \begin{aligned} &\text{find } (\lambda, \psi) \text{ in } \mathbb{R} \times \Phi, \text{ with } \psi \neq 0, \\ &\text{such that there exists } \chi \in \Xi \text{ verifying} \\ &\begin{cases} a(\psi, \varphi) + b(\varphi, \chi) = \lambda(\psi, \varphi)_{H_\Phi} & \forall \varphi \in \Phi \\ b(\psi, \xi) = 0 & \forall \xi \in \Xi, \end{cases} \end{aligned}$$

which in the formalism of the previous section can be written

$$(43) \quad \lambda T_\Phi \psi = \psi.$$

We assume that the operator T_Φ is compact from H_Φ to Φ .

Suppose now that we are given two finite dimensional subspaces Φ_h and Ξ_h of Φ and Ξ , respectively. Then the approximation of (42) reads

$$(44) \quad \begin{aligned} & \text{find } (\lambda_h, \psi_h) \text{ in } \mathbb{R} \times \Phi_h, \text{ with } \psi_h \neq 0 \text{ such that} \\ & \text{there exists } \chi_h \in \Xi_h \text{ verifying} \\ & \begin{cases} a(\psi_h, \varphi_h) + b(\varphi_h, \chi_h) = \lambda_h(\psi_h, \varphi_h)_{H_\Phi} & \forall \varphi_h \in \Phi_h \\ b(\psi_h, \xi_h) = 0 & \forall \xi_h \in \Xi_h, \end{cases} \end{aligned}$$

that is

$$(45) \quad \lambda_h T_\Phi^h \psi_h = \psi_h.$$

We are now looking for necessary and sufficient conditions that ensure the uniform convergence of T_Φ^h to T_Φ in $\mathcal{L}(H_\Phi, \Phi)$ which, as we have seen, implies the convergence of eigenvalues and eigenvectors (see (23)).

To start with, we look for sufficient conditions.

We introduce some notation. Let Φ_0^H and Ξ_0^H be the subspaces of Φ and Ξ , respectively, containing all the solutions $\psi \in \Phi$ and $\chi \in \Xi$, respectively, of problem (6) when $g = 0$; that is, with the formalism of the previous section,

$$(46) \quad \begin{aligned} \Phi_0^H &= C_\Phi^* \circ S \circ C_\Phi(H_\Phi) = T_\Phi(H_\Phi) \\ \Xi_0^H &= C_\Xi^* \circ S \circ C_\Phi(H_\Phi). \end{aligned}$$

Notice that the following inclusion holds true:

$$\Phi_0^H \subseteq \mathbb{K}.$$

The spaces Φ_0^H and Ξ_0^H will be endowed with the *natural* norm: that is, for instance,

$$(47) \quad \begin{aligned} \|\varphi\|_{\Phi_0^H} &:= \inf\{\|\eta\|_{H_\Phi}, T_\Phi \eta = \varphi\}; \\ \|\xi\|_{\Xi_0^H} &:= \inf\{\|\eta\|_{H_\Phi}, C_\Xi^* \circ S \circ C_\Phi \eta = \xi\}. \end{aligned}$$

DEFINITION 1. We say that the *weak approximability* of Ξ_0^H is verified if there exists $\omega_1(h)$, tending to zero as h goes to zero, such that for every $\chi \in \Xi_0^H$

$$(48) \quad \sup_{\varphi_h \in \mathbb{K}_h} \frac{b(\varphi_h, \chi)}{\|\varphi_h\|_\Phi} \leq \omega_1(h) \|\chi\|_{\Xi_0^H}.$$

Notice that, in spite of its appearance, (48) is indeed an approximability property. Actually as $\varphi_h \in \mathbb{K}_h$, we have $b(\varphi_h, \chi) = b(\varphi_h, \chi - \chi^I)$ for every $\chi^I \in \Xi_h$, which has, usually, to be used to verify (48).

DEFINITION 2. We say that the *strong approximability* of Φ_0^H is verified if there exists $\omega_2(h)$, tending to zero as h goes to zero, such that for every $\psi \in \Phi_0^H$ there exists $\psi^I \in \mathbb{K}_h$ such that

$$(49) \quad \|\psi - \psi^I\|_\Phi \leq \omega_2(h) \|\psi\|_{\Phi_0^H}.$$

THEOREM 1. *Let us assume that (DEK) is verified (see (14)). Assume moreover the weak approximability of Ξ_0^H and the strong approximability of Φ_0^H . Then the sequence T_Φ^h converges uniformly to T_Φ in $\mathcal{L}(H_\Phi, \Phi)$, that is there exists $\omega_3(h)$, tending to zero as h goes to zero, such that*

$$(50) \quad \|T_\Phi f - T_\Phi^h f\|_\Phi \leq \omega_3(h) \|f\|_{H_\Phi}, \quad \text{for all } f \in H_\Phi.$$

PROOF. Let $f \in H_\Phi$ and let $(\psi, \chi) \in \Phi_0^H \times \Xi_0^H$ be solution of (6): $(\psi, \chi) = S(f, 0)$. As we assumed (DEK) Proposition 2 ensures that T_Φ^h is well defined on Φ' . Recall that $\psi := T_\Phi(f)$. Let $\psi_h := T_\Phi^h(f)$ and let χ^I be such that (ψ_h, χ^I) is a solution of (13) (such χ^I might not be unique). In order to prove the uniform convergence of T_Φ^h to T_Φ , we have to estimate the difference $\|\psi - \psi_h\|_\Phi$. We do it by bounding the term $\|\psi^I - \psi_h\|_\Phi$, where ψ^I is given by (49), and then by using the triangular inequality. We have

$$(51) \quad \begin{aligned} \alpha \|\psi^I - \psi_h\|_\Phi^2 &\leq a(\psi^I - \psi_h, \psi^I - \psi_h) \\ &= a(\psi^I - \psi, \psi^I - \psi_h) + a(\psi - \psi_h, \psi^I - \psi_h) \\ &\leq M_a \|\psi^I - \psi\|_\Phi \|\psi^I - \psi_h\|_\Phi - b(\psi^I - \psi_h, \chi - \chi_h) \\ &\leq \left(M_a \|\psi^I - \psi\|_\Phi + \sup_{\varphi_h \in \mathbb{K}_h} \frac{b(\varphi_h, \chi - \chi_h)}{\|\varphi_h\|_\Phi} \right) \|\psi^I - \psi_h\|_\Phi \\ &= \left(M_a \|\psi^I - \psi\|_\Phi + \sup_{\varphi_h \in \mathbb{K}_h} \frac{b(\varphi_h, \chi)}{\|\varphi_h\|_\Phi} \right) \|\psi^I - \psi_h\|_\Phi. \end{aligned}$$

The result then follows immediately from the *strong approximability* of Φ_0^H and the *weak approximability* of Ξ_0^H . In particular we can take $\omega_3(h) = (1 + M_a/\alpha)\omega_2(h) + \omega_1(h)/\alpha$. \square

In the following theorem we shall see that the assumptions of Theorem 1 are also, in a sense, necessary for the uniform convergence of T_Φ^h to T_Φ in $\mathcal{L}(H_\Phi, \Phi)$.

THEOREM 2. *Assume that the sequence T_Φ^h is bounded in $\mathcal{L}(\Phi', \Phi)$, and converges uniformly to T_Φ in $\mathcal{L}(H_\Phi, \Phi)$ (see (50)). Then, the ellipticity in the kernel property (DEK) holds true. Moreover, both the strong approximability of Φ_0^H and the weak approximability of Ξ_0^H are satisfied.*

PROOF. The (DEK) property can be obtained applying Proposition 3. Let ψ be an element of Φ_0^H . Then by definition of Φ_0^H there is $f \in H_\Phi$ such that $\psi = T_\Phi f$. Define $\psi^I := T_\Phi^h f$. Uniform convergence implies the *strong approximability* of Φ_0^H .

In a similar way, let χ be an element of Ξ_0^H . Then by definition of Ξ_0^H , $\chi = C_{\Xi}^* \circ S \circ C_\Phi f$ for some $f \in H_\Phi$. There might be more than one such f . We choose \bar{f} such that $\|\bar{f}\|_{H_\Phi} \leq \frac{3}{2} \inf\{\|f\|_{H_\Phi} : C_{\Xi}^* \circ S \circ C_\Phi f = \chi\} = \frac{3}{2} \|\chi\|_{\Xi_0^H}$. Let $\psi := T_\Phi \bar{f}$. Correspondingly let $\psi_h := T_\Phi^h \bar{f}$ and let χ_h be such that (ψ_h, χ_h) is a solution of (13) with the same right-hand side (such χ_h might not be unique). Then we obtain

$$\begin{aligned} \sup_{\varphi_h \in \mathbb{K}_h} \frac{b(\varphi_h, \chi)}{\|\varphi_h\|_\Phi} &= \sup_{\varphi_h \in \mathbb{K}_h} \frac{b(\varphi_h, \chi - \chi_h)}{\|\varphi_h\|_\Phi} = \sup_{\varphi_h \in \mathbb{K}_h} \frac{a(\psi - \psi_h, \varphi_h)}{\|\varphi_h\|_\Phi} \\ &\leq M_a \|\psi - \psi_h\|_\Phi \leq M_a \omega_3(h) \|f\|_{H_\Phi} \leq \frac{3}{2} M_a \omega_3(h) \|\chi\|_{\Xi_0^H}, \end{aligned}$$

which gives (48) with $\omega_1(h) = \frac{3}{2} M_a \omega_3(h)$, that is the *weak approximability* of Ξ_0^H . \square

EXAMPLE 1. We go back to the Example 1 of the Introduction (Stokes problem). Now $\Phi = V = (H_0^1(\Omega))^2$ and $\Xi = Q = L^2(\Omega)/_{\mathbb{R}}$. It is easy to see that if Ω is, for instance, a convex polygon, Ξ_0^H is $H^1(\Omega)/_{\mathbb{R}}$ and Φ_0^H is the subspace of $(H^2(\Omega) \cap H_0^1(\Omega))^2$ made of free divergence functions (see [19]). In particular we can check that $\|\underline{u}\|_{\Phi_0^H} = \|\Delta \underline{u}\|_0 \sim \|\underline{u}\|_2$ and $\|p\|_{\Xi_0^H} = \|\nabla p\|_0$ (with standard notation, here and in the following, we denote by $\|\cdot\|_k$ the norm in $H^k(\Omega)$ for $k \in \mathbb{N}$). Let V_h and Q_h be finite dimensional subspaces of V and Q respectively. The weak approximability of Ξ_0^H will surely hold if

$$\inf_{q_h \in Q_h} \|p - q_h\|_0 \leq \omega_1(h) \|p\|_1 \quad \text{for all } p \in H^1(\Omega)/_{\mathbb{R}},$$

which is satisfied by all choices of finite element spaces that one may seriously think to use in practice.

The strong approximability of Φ_0^H , which now reads

$$(52) \quad \|\underline{u} - \underline{u}^I\|_1 \leq \omega_2(h) \|\underline{u}\|_2 \quad \text{for all } \underline{u} \in \Phi_0^H,$$

is more delicate, as \underline{u}^I has to be chosen in \mathbb{K}_h . If the pair (V_h, Q_h) satisfies the inf-sup condition (DIS) then the property trivially holds. Remark, however, that the typical way to proving the inf-sup condition is to show, following [15], that: *for every \underline{u} in V there exists \underline{u}^I in V_h such that $\|\underline{u}^I\|_V \leq C \|\underline{u}\|_V$ (C independent of \underline{u} and h) and $b(\underline{u} - \underline{u}^I, q_h) = 0 \forall q_h \in Q_h$, which is more difficult than proving (52) directly. Moreover there are choices of elements that fail to satisfy the inf-sup*

condition, for which (52) holds true. For instance, we may think to the so-called $Q_1 - P_0$ element, where

$$(53) \quad \begin{aligned} V_h &:= \{\underline{v}_h \in (C^0(\Omega))^2 : \underline{v}_h|_K \in (Q_1(K))^2 \ \forall K \in \mathcal{T}_h\}, \\ Q_h &:= \{q_h : q_h|_K \in P_0(K) \ \forall K \in \mathcal{T}_h\}, \end{aligned}$$

where, with standard finite element notation, for k integer ≥ 0 $P_k(D)$ denotes the space of polynomials of degree $\leq k$ on a domain D , and $Q_k(D)$ the space of polynomials of degree $\leq k$ separately in each variable. Hence, here $Q_1(K)$ is the set of bilinear polynomials on K , and $P_0(K)$ is the set of the constant functions on K . We may assume, for simplicity, that Ω is a square and that the decomposition \mathcal{T}_h is made by $2N \times 2N$ equal subsquares. It is known that this choice of elements does not satisfy the inf-sup condition: the operator B_h^f has a non trivial kernel (the checkerboard mode), and by discarding it we still have at best (DIS $_h$) with $\beta_h \sim h$ (see [22], [18], [6]). Nevertheless, for $u \in \Phi_0^H \subset \mathbb{K}$, we can construct u^I as follows: let \hat{u} be the vector in V_h which is bilinear in each square of the $N \times N$ (coarser) grid and agrees with u at the vertices of the coarser grid. Let now \tilde{u} be the vector in V_h with the following properties. It vanishes at the vertices of the coarser grid; its tangential component vanishes on the midpoints of the edges of the coarser grid; its normal component at the midpoints of each edge e of the coarser grid is chosen in such a way that $\int_e (u - \hat{u} - \tilde{u}) \cdot n = 0$, and finally the values at the center of each element K of the coarser grid are chosen to satisfy $\int_K (u - \hat{u} - \tilde{u}) q_h = 0$ for $q_h = \text{sign}(x - x_c)$ and $q_h = \text{sign}(y - y_c)$ (where (x_c, y_c) is the center of K). It is not difficult to check that $u^I = \hat{u} + \tilde{u}$ satisfies (52) with $\omega_2(h) = O(h)$. For a similar construction see [9], pages 241-242. We have here a first example in which the eigenvalues are approximated correctly even though the global matrix associated to (13) is singular.

EXAMPLE 2. *Dirichlet problem with Lagrange multipliers.* Here $\Phi = V = H^1(\Omega)$ and $\Xi = M = H^{-1/2}(\partial\Omega)$. It is well-known (see e.g. [17]), that if Ω is, for instance, a convex polygon, then $\Phi_0^H = H^2(\Omega) \cap H_0^1(\Omega)$ and $\Xi_0^H = H^{1/2}(\partial\Omega)$. Let now $\{\mathcal{T}_h^\Omega\}$ be a regular sequence of decompositions of Ω (see e.g. [12]), $\{\mathcal{T}_h^\Gamma\}$ be a regular sequence of decompositions of $\partial\Omega$, k_1 and k_2 be integers with $k_1 \geq 1$ and $k_2 \geq 0$. Set

$$(54) \quad \begin{aligned} V_h^{k_1} &:= \{\underline{v}_h \in V : \underline{v}_h|_K \in P_{k_1}(K) \ \forall K \in \mathcal{T}_h^\Omega\}, \\ M_h^{k_2} &:= \{\mu_h \in M : \mu_h|_e \in P_{k_2}(e) \ \forall e \in \mathcal{T}_h^\Gamma\}. \end{aligned}$$

It is trivial to check that (48) and (49) hold for every choice of $\{\mathcal{T}_h^\Omega\}$, $\{\mathcal{T}_h^\Gamma\}$, k_1 and k_2 . In particular (DEK), which is now a sort of Poincaré inequality, only requires that $M_h \subset M$ contains at least a $\bar{\mu}_h$ such that $\langle \bar{\mu}_h, 1 \rangle \neq 0$.

Note that to have (DIS) one must ask rather strict compatibility conditions on $\{\mathcal{T}_h^\Omega\}$, $\{\mathcal{T}_h^\Gamma\}$, k_1 and k_2 , see [2]. Therefore, for a general choice, solvability of (13) might fail. Nevertheless, as we have seen, convergence of the eigenvalues is assured under weaker assumptions.

4. – Problems of the type $\begin{pmatrix} 0 \\ g \end{pmatrix}$

In this section, together with (1)-(4), we assume that, for every given $g \in \Xi'$ and $f = 0$, problem (6) has a unique solution (ψ, χ) and that there exists a constant C (independent of g) such that

$$(55) \quad \|\psi\|_{\Phi} + \|\chi\|_{\Xi} \leq C\|g\|_{\Xi'}.$$

It is easy to see that this implies (IS) but not (EK) (see Example 4 of the Introduction). Moreover we assume that we are given a Hilbert space H_{Ξ} (that we shall identify with its dual space H'_{Ξ}) such that

$$(56) \quad \Xi \subseteq H_{\Xi} \subseteq \Xi'$$

with continuous and dense embeddings. For simplicity, we assume that for every $\xi \in \Xi$, we have $\|\xi\|_{H_{\Xi}} \leq \|\xi\|_{\Xi}$ (with constant equal to 1).

We consider the eigenvalue problem

$$(57) \quad \begin{aligned} & \text{find } (\lambda, \chi) \text{ in } \mathbb{R} \times \Xi, \text{ with } \chi \neq 0, \text{ such that} \\ & \text{there exists } \psi \in \Phi \text{ satisfying} \\ & \begin{cases} a(\psi, \varphi) + b(\varphi, \chi) = 0 & \forall \varphi \in \Phi \\ b(\psi, \xi) = -\lambda(\chi, \xi)_{H_{\Xi}} & \forall \xi \in \Xi, \end{cases} \end{aligned}$$

which in the formalism of Section 2 can be written

$$(58) \quad \lambda T_{\Xi} \chi = -\chi.$$

As we shall see, problems of the type $\begin{pmatrix} 0 \\ g \end{pmatrix}$ are more closely related to the abstract theory of [14] than problems of the previous type $\begin{pmatrix} f \\ 0 \end{pmatrix}$.

From now on we assume that the operator T_{Ξ} is compact from H_{Ξ} into Ξ .

We introduce two finite dimensional subspaces Φ_h and Ξ_h of Φ and Ξ , respectively. Then the approximation of (57) reads

$$(59) \quad \begin{aligned} & \text{find } (\lambda_h, \chi_h) \text{ in } \mathbb{R} \times \Xi_h, \text{ with } \chi_h \neq 0, \text{ such that} \\ & \text{there exists } \psi_h \in \Phi_h \text{ satisfying} \\ & \begin{cases} a(\psi_h, \varphi_h) + b(\varphi_h, \chi_h) = 0 & \forall \varphi_h \in \Phi_h \\ b(\psi_h, \xi_h) = -\lambda_h(\chi_h, \xi_h)_{H_{\Xi}} & \forall \xi_h \in \Xi_h, \end{cases} \end{aligned}$$

that is

$$(60) \quad \lambda_h T_{\Xi}^h \chi_h = -\chi_h.$$

We are now looking for necessary and sufficient conditions that ensure the uniform convergence of T_{Ξ}^h to T_{Ξ} in $\mathcal{L}(H_{\Xi}, \Xi)$, which implies the convergence of eigenvalues and eigenvectors (see (23)).

To start with, we look for sufficient conditions.

We introduce some notation. Let Φ_H^0 and Ξ_H^0 be the subspaces of Φ and Ξ respectively, containing all the solutions $\psi \in \Phi$ and $\chi \in \Xi$, respectively, of problem (6) when $f = 0$; that is, with the formalism of Section 2,

$$(61) \quad \begin{aligned} \Phi_H^0 &= C_\Phi^* \circ S \circ C_\Xi(H_\Xi) \\ \Xi_H^0 &= C_\Xi^* \circ S \circ C_\Xi(H_\Xi) = T_\Xi(H_\Xi). \end{aligned}$$

It will also be useful to define the space $\Phi_{\Xi'}^0$ as the image of $C_\Phi^* \circ S \circ C_\Xi$ (from Ξ' to Φ).

As before, the spaces Φ_H^0 , Ξ_H^0 and $\Phi_{\Xi'}^0$ will be endowed with their natural norms (see for instance (47)).

DEFINITION 3. We say that the *weak approximability* of Ξ_H^0 with respect to $a(\cdot, \cdot)$ is verified if there exists $\omega_4(h)$, tending to zero as h goes to zero, such that for every $\chi \in \Xi_H^0$ and for every $\varphi_h \in \mathbb{K}_h$

$$(62) \quad b(\varphi_h, \chi) \leq \omega_4(h) \|\chi\|_{\Xi_H^0} \|\varphi_h\|_a.$$

Notice that (62) is indeed an approximation property, as we already pointed out for its counterpart (48).

DEFINITION 4. We say that the *strong approximability* of Ξ_H^0 is verified if there exists $\omega_5(h)$, tending to zero as h goes to zero, such that for every $\chi \in \Xi_H^0$ there exists $\chi^I \in \Xi_h$ such that

$$(63) \quad \|\chi - \chi^I\|_\Xi \leq \omega_5(h) \|\chi\|_{\Xi_H^0}.$$

Notice that (62) and (63) are (much) weaker forms of assumption H7 of [14].

DEFINITION 5. An operator Π_h from Φ (or from a subspace of it) into Φ_h is called a *Fortin operator* with respect to the bilinear form b and the subspace $\Xi_h \subset \Xi$ if it verifies, for all φ in its domain,

$$(64) \quad b(\varphi - \Pi_h \varphi, \xi_h) = 0 \quad \forall \xi_h \in \Xi_h.$$

The following assumptions for a Fortin operator will be useful.

– There exists a constant C_Π , independent of h such that:

$$(65) \quad \|\Pi_h\|_{\mathcal{L}(\Phi_{\Xi'}^0, \Phi)} \leq C_\Pi.$$

– There exists $\omega_6(h)$, tending to zero as h goes to zero, such that for every $\varphi \in \Phi_H^0$ it holds

$$(66) \quad \|\varphi - \Pi_h \varphi\|_a \leq \omega_6(h) \|\varphi\|_{\Phi_H^0}.$$

Notice that (66) is strongly related to assumption H5 of [14]. However, being interested in convergence, we have to assume that $\omega_6(h)$ goes to 0, while [14] only assumes it to be bounded and puts it in the right-hand side of a priori estimates. On the other hand, as we shall see, (66) is actually *necessary* for having convergence of eigenvalues. This was not pointed out in [14] for the very good reason that, first, their interest was in a priori bounds (and not on necessity) and, second, they were dealing with direct problems (and not with eigenvalues). In particular (66) is not necessary for having pointwise convergence of T_{Ξ}^h to T_{Ξ} where (DEK) and (DIS) are sufficient. Notice that (as it is also pointed out in Proposition 1 of [14]) (IS) and (DIS) imply (65), but, as we shall see later in this paper, (EK), (DEK), (IS) and (DIS) (all together) imply pointwise convergence but not (66).

THEOREM 3. *Let us assume that there exists a Fortin operator (see (64)) $\Pi_h : \Phi_{\Xi'}^0 \rightarrow \Phi_h$ satisfying (65) and (66). Assume moreover that the strong approximability of Ξ_H^0 is verified (see (63)) as well as the weak approximability of Ξ_H^0 with respect to a (see (62)). Then the sequence T_{Ξ}^h converges to T_{Ξ} uniformly from H_{Ξ} into Ξ , that is there exists $\omega_7(h)$, tending to zero as h goes to zero, such that*

$$(67) \quad \|T_{\Xi}g - T_{\Xi}^hg\|_{\Xi} \leq \omega_7(h)\|g\|_{H_{\Xi}}, \quad \text{for all } g \in H_{\Xi}.$$

PROOF. We remark first that, as it is well-known, (7) and (64)-(65) imply (DIS) (see [15] or [9]). Thanks to Proposition 4, T_{Ξ}^h is then well defined.

Let $g \in H_{\Xi}$ and let $(\psi, \chi) \in \Phi_H^0 \times \Xi_H^0$ be the solution of (6) with $f = 0$. Recall that $\chi = T_{\Xi}g$. Let $\chi_h := T_{\Xi}^hg$ and let ψ_h be such that (ψ_h, χ_h) is a solution of (13) (such ψ_h might be not unique). In order to prove the uniform convergence of T_{Ξ}^h to T_{Ξ} we have to find a priori estimates for the error $\|\chi - \chi_h\|_{\Xi}$. Let $\tilde{g} \in \Xi'$ be such that $\langle \tilde{g}, \chi - \chi_h \rangle = \|\chi - \chi_h\|_{\Xi}$ and $\|\tilde{g}\|_{\Xi'} = 1$. Take $\tilde{\varphi} := C_{\Phi}^* \circ S \circ C_{\Xi} \tilde{g}$, hence $\|\tilde{\varphi}\|_{\Phi_{\Xi'}^0} \leq \|\tilde{g}\|_{\Xi'} = 1$ (see (47)). Then we have

$$(68) \quad \begin{aligned} \|\chi - \chi_h\|_{\Xi} &= \langle \tilde{g}, \chi - \chi_h \rangle = b(\tilde{\varphi}, \chi - \chi_h) \\ &= b(\tilde{\varphi} - \Pi_h \tilde{\varphi}, \chi - \chi_h) + b(\Pi_h \tilde{\varphi}, \chi - \chi_h) \\ &= b(\tilde{\varphi} - \Pi_h \tilde{\varphi}, \chi - \chi^I) - a(\psi - \psi_h, \Pi_h \tilde{\varphi}). \end{aligned}$$

Let us estimate separately the two terms in the right-hand side:

$$(69) \quad \begin{aligned} b(\tilde{\varphi} - \Pi_h \tilde{\varphi}, \chi - \chi^I) &\leq M_b \|\tilde{\varphi} - \Pi_h \tilde{\varphi}\|_{\Phi} \|\chi - \chi^I\|_{\Xi} \\ &\leq M_b (\|\tilde{\varphi}\|_{\Phi} + \|\Pi_h \tilde{\varphi}\|_{\Phi}) \|\chi - \chi^I\|_{\Xi}; \\ a(\psi - \psi_h, \Pi_h \tilde{\varphi}) &\leq \|\Pi_h \tilde{\varphi}\|_a \|\psi - \psi_h\|_a. \end{aligned}$$

Using (65) we obtain the following estimate for $\Pi_h \tilde{\varphi}$

$$(70) \quad \|\Pi_h \tilde{\varphi}\|_{\Phi} \leq C_{\Pi} \|\tilde{\varphi}\|_{\Phi_{\Xi'}^0} \leq C_{\Pi}.$$

Putting together (68), (69) and (70) and using (63) we obtain

$$(71) \quad \begin{aligned} \|\chi - \chi_h\|_{\Xi} &\leq M_b(1 + C_{\Pi})\|\chi - \chi^I\|_{\Xi} + C_{\Pi}\|\psi - \psi_h\|_a \\ &\leq M_b(1 + C_{\Pi})\omega_5(h)\|\chi\|_{\Xi_H^0} + C_{\Pi}\|\psi - \psi_h\|_a. \end{aligned}$$

To conclude the proof it remains to estimate $\|\psi - \psi_h\|_a$. Thanks to the triangular inequality and to (66) we bound only $\|\Pi_h\psi - \psi_h\|_a$ using also (62) and (64). Notice that $\Pi_h\psi - \psi_h$ belongs to \mathbb{K}_h .

$$(72) \quad \begin{aligned} \|\Pi_h\psi - \psi_h\|_a^2 &= a(\Pi_h\psi - \psi, \Pi_h\psi - \psi_h) + a(\psi - \psi_h, \Pi_h\psi - \psi_h) \\ &\leq \|\psi - \Pi_h\psi\|_a\|\Pi_h\psi - \psi_h\|_a - b(\Pi_h\psi - \psi_h, \chi - \chi_h) \\ &= \|\psi - \Pi_h\psi\|_a\|\Pi_h\psi - \psi_h\|_a - b(\Pi_h\psi - \psi_h, \chi) \\ &\leq \|\Pi_h\psi - \psi_h\|_a \left(\|\psi - \Pi_h\psi\|_a + \omega_4(h)\|\chi\|_{\Xi_H^0} \right), \end{aligned}$$

which, due to (66), gives

$$(73) \quad \begin{aligned} \|\psi - \psi_h\|_a &\leq 2\|\psi - \Pi_h\psi\|_a + \omega_4(h)\|\chi\|_{\Xi_H^0} \\ &\leq 2\omega_6(h)\|\psi\|_{\Phi_H^0} + \omega_4(h)\|\chi\|_{\Xi_H^0} \end{aligned}$$

and (67) holds with $\omega_7(h) = M_b(1 + C_{\Pi})\omega_5(h) + 2C_{\Pi}\omega_6(h) + C_{\Pi}\omega_4(h)$. \square

REMARK 2. In Theorem 3 we have proved the uniform convergence of T_{Ξ}^h to T_{Ξ} in $\mathcal{L}(H_{\Xi}, \Xi)$. However in Section 2 we have seen that the convergence of the spectrum is equivalent to the uniform convergence of T_{Ξ}^h to T_{Ξ} in $\mathcal{L}(H_{\Xi})$. Indeed the latter holds under the weaker assumption that there exists a Fortin operator satisfying only (66) as we shall see in the following theorem. \square

THEOREM 4. *Let us assume that there exists a Fortin operator (see (64)) $\Pi_h : \Phi_{\Xi'}^0 \rightarrow \Phi_h$ satisfying (66). Assume moreover that both the strong approximability of Ξ_H^0 (see (63)) and the weak approximability of Ξ_H^0 with respect to a (see (62)) are verified. Then the sequence T_{Ξ}^h converges uniformly to T_{Ξ} in H_{Ξ} .*

PROOF. We observe that (7) and (64) imply the weak discrete inf-sup condition (DIS_h). Thanks to Proposition 4, T_{Ξ}^h is then well defined.

Let $g \in H_{\Xi}$ and let $(\psi, \chi) \in \Phi_H^0 \times \Xi_H^0$ be the solution of (6) with $f = 0$. Recall that $\chi = T_{\Xi}g$. Let $\chi_h := T_{\Xi}^hg$ and let ψ_h be such that (ψ_h, χ_h) is a solution of (13) with right-hand side $(0, g)$ (such ψ_h might be not unique). We estimate $\|\chi - \chi_h\|_{H_{\Xi}}$. Using a duality argument, let $(\tilde{\psi}, \tilde{\chi}) \in \Phi \times \Xi$ be defined by $(\tilde{\psi}, \tilde{\chi}) := S(0, \chi - \chi_h)$. Due to the definition (61), $\tilde{\psi}$ belongs to Φ_H^0 with

the following estimate $\|\tilde{\psi}\|_{\Phi_H^0} \leq \|\chi - \chi_h\|_{H_\Xi}$ (see (47)). Then

$$\begin{aligned}
\|\chi - \chi_h\|_{H_\Xi}^2 &= (\chi - \chi_h, \chi - \chi_h) = b(\tilde{\psi}, \chi - \chi_h) \\
&= b(\tilde{\psi} - \Pi_h \tilde{\psi}, \chi) + b(\Pi_h \tilde{\psi}, \chi - \chi_h) \\
&= -a(\psi, \tilde{\psi} - \Pi_h \tilde{\psi}) - a(\psi - \psi_h, \Pi_h \tilde{\psi}) \\
&\leq \|\psi\|_a \|\tilde{\psi} - \Pi_h \tilde{\psi}\|_a + \|\psi - \psi_h\|_a \|\Pi_h \tilde{\psi}\|_a \\
&\leq \|\psi\|_a \omega_6(h) \|\tilde{\psi}\|_{\Phi_H^0} + 2\|\psi - \psi_h\|_a \|\tilde{\psi}\|_{\Phi_H^0} \\
&\leq (\omega_6(h) \|\psi\|_a + 2\|\psi - \psi_h\|_a) \|\chi - \chi_h\|_{H_\Xi},
\end{aligned}$$

having assumed $\omega_6(h) \leq 1$. Hence

$$\|\chi - \chi_h\|_{H_\Xi} \leq \omega_6(h) \|\psi\|_a + 2\|\psi - \psi_h\|_a.$$

The rest of the proof follows the same lines as the one of Theorem 3, using (62) and (66) (see (72) and (73)). \square

The remaining part of this section is devoted to see what one can deduce from the uniform convergence of T_Ξ^h to T_Ξ .

THEOREM 5. *Assume that the sequence T_Ξ^h is bounded in $\mathcal{L}(\Xi', \Xi)$. Then there exists a Fortin operator (see (64)) $\Pi_h : \Phi_{\Xi'}^0 \rightarrow \Phi_h$ such that*

$$(74) \quad \|\psi - \Pi_h \psi\|_a \leq C \|\psi\|_{\Phi_{\Xi'}^0}.$$

PROOF. Let ψ belong to $\Phi_{\Xi'}^0$. Then by definition $\psi = C_\Phi^* \circ S \circ C_{\Xi} g$ for some $g \in \Xi'$. There is only one g in this condition, and therefore, by definition, $\|\psi\|_{\Phi_{\Xi'}^0} = \|g\|_{\Xi'}$ (see (47)). Let $\chi \in \Xi$ be such that $(\psi, \chi) = S(0, g)$. Let $\chi_h := T_\Xi^h g$; notice that, by assumption, $\|\chi_h\|_\Xi \leq C \|g\|_{\Xi'}$. By Propositions 5 and 4, there exists at least one ψ_h such that $(\psi_h, \chi_h) \in \Phi_h \times \Xi_h$ is a corresponding discrete solution of (13). If such ψ_h is unique, we define $\Pi_h \psi := \psi_h$. Otherwise we still define $\Pi_h \psi$ as the ψ_h having minimum norm in Φ . By construction we have (64) and

$$(75) \quad \|\Pi_h \psi\|_a^2 = \langle g, \chi_h \rangle \leq \|g\|_{\Xi'} \|T_\Xi^h g\|_\Xi \leq C \|g\|_{\Xi'}^2 = C \|\psi\|_{\Phi_{\Xi'}^0}^2.$$

Let us bound $\|\psi - \Pi_h \psi\|_a$:

$$\begin{aligned}
\|\psi - \Pi_h \psi\|_a^2 &= a(\psi - \Pi_h \psi, \psi - \Pi_h \psi) \\
(76) \quad &= a(\psi, \psi - \Pi_h \psi) - a(\Pi_h \psi, \psi - \Pi_h \psi) \\
&= -b(\psi - \Pi_h \psi, \chi) - a(\psi - \Pi_h \psi, \Pi_h \psi).
\end{aligned}$$

The first term in the right-hand side can be handled as follows:

$$\begin{aligned}
 (77) \quad b(\psi - \Pi_h \psi, \chi) &= b(\psi - \Pi_h \psi, \chi - \chi_h) \\
 &= \langle g, \chi - \chi_h \rangle - b(\Pi_h \psi, \chi - \chi_h) \\
 &= \langle g, \chi - \chi_h \rangle + a(\psi - \Pi_h \psi, \Pi_h \psi).
 \end{aligned}$$

Inserting (77) in (76), we obtain

$$\begin{aligned}
 (78) \quad \|\psi - \Pi_h \psi\|_a^2 &= -\langle g, \chi - \chi_h \rangle - 2a(\psi - \Pi_h \psi, \Pi_h \psi) \\
 &\leq \|g\|_{\Xi'} \|\chi - \chi_h\|_{\Xi} + 2\|\psi - \Pi_h \psi\|_a \|\Pi_h \psi\|_a \\
 &\leq \|g\|_{\Xi'} (\|\chi\|_{\Xi} + \|\chi_h\|_{\Xi}) + 2\|\psi - \Pi_h \psi\|_a \|\Pi_h \psi\|_a
 \end{aligned}$$

then the boundedness of T_{Ξ}^h and (75) imply (74). \square

THEOREM 6. *Assume that the sequence T_{Ξ}^h converges to T_{Ξ} uniformly from H_{Ξ} to Ξ , then for all $\chi \in \Xi_H^0$ there is $\chi^I \in \Xi_h$ such that (63) holds true.*

PROOF. Let χ belong to Ξ_H^0 , then $\chi = T_{\Xi} g$ for a suitable g in H_{Ξ} . Let $\chi_h := T_{\Xi}^h g$ be the corresponding discrete solution, then we define $\chi^I := \chi_h$ and the inequality (63) is an easy consequence of the uniform convergence of $T_{\Xi}^h g$ to $T_{\Xi} g$ in Ξ . \square

THEOREM 7. *Let us assume that the sequence T_{Ξ}^h is bounded in $\mathcal{L}(\Xi', \Xi)$ and converges uniformly to T_{Ξ} in $\mathcal{L}(H_{\Xi}, \Xi)$. In addition we assume that the following bound holds for the solutions of (13) with $f = 0$*

$$(79) \quad \|\psi_h\|_{\Phi} \leq C \|g\|_{\Xi'}.$$

Then there exists a Fortin operator $\Pi_h : \Phi_{\Xi'}^0 \rightarrow \Phi_h$ satisfying (65) and (66). Moreover we have (DIS) (see (15)), and the weak approximability of Ξ_H^0 with respect to a (see (62)) holds.

PROOF. From Proposition 5 we have that $C_{\Phi}^* \circ S \circ C_{\Xi}$ is also well defined and (DIS) holds. Let us check (65). For $\psi \in \Phi_{\Xi'}^0$, there exists $g \in \Xi'$ and $\chi \in \Xi$ such that $(\psi, \chi) = S(0, g)$. We set $\Pi_h \psi := C_{\Phi}^* \circ S_h \circ C_{\Xi} g$. As we have seen, (64) holds trivially, and now (65) also holds in virtue of (79), with $C_{\Pi} := C$.

Now let us check (66). Let ψ belong to Φ_H^0 ; by definition $\psi = C_{\Phi}^* \circ S \circ C_{\Xi} g$ for some $g \in H_{\Xi}$. As in the proof of Theorem 5, g is unique, and $\|\psi\|_{\Phi_H^0} = \|g\|_{H_{\Xi}}$. Let $\chi := T_{\Xi} g$; clearly $\chi \in \Xi_H^0$. Let $\chi_h := T_{\Xi}^h g$. By construction $(\Pi_h \psi, \chi_h)$ solves (13) with the right-hand side $(0, g)$. Moreover by the same computations as above, we arrive (see the first line in (78)) at

$$\|\psi - \Pi_h \psi\|_a^2 = -\langle g, \chi - \chi_h \rangle - 2a(\psi - \Pi_h \psi, \Pi_h \psi).$$

From this we have

$$\begin{aligned}
\|\psi - \Pi_h \psi\|_a^2 &= -(g, \chi - \chi_h) - 2b(\Pi_h \psi, \chi - \chi_h) \\
&\leq (\|g\|_{\Xi'} + 2M_b \|\Pi_h \psi\|_{\Phi}) \|\chi - \chi_h\|_{\Xi} \\
(80) \quad &\leq (1 + 2M_b C) \|g\|_{\Xi'} \omega_7(h) \|g\|_{H_{\Xi}} \\
&\leq (1 + 2M_b C) \omega_7(h) \|g\|_{H_{\Xi}}^2 \\
&= (1 + 2M_b C) \omega_7(h) \|\psi\|_{\Phi_H^0}^2,
\end{aligned}$$

where we used (79) and the uniform convergence of T_{Ξ}^h to T_{Ξ} in $\mathcal{L}(H_{\Xi}, \Xi)$ (see (67)). The bound (80) gives (66) with $\omega_6(h) = ((1 + 2M_b C) \omega_7(h))^{1/2}$.

Now let us check (62). If $\chi \in \Xi_H^0$, then $\chi = T_{\Xi} g$ for a suitable g in H_{Ξ} , let ψ be such that $(\psi, \chi) = S(0, g)$. Next we set $\chi_h := T_{\Xi}^h g$ and $\psi_h := \Pi_h \psi$. Then we get for every $\varphi_h \in \mathbb{K}_h$

$$(81) \quad b(\varphi_h, \chi) = b(\varphi_h, \chi - \chi_h) = a(\Pi_h \psi - \psi, \varphi_h) \leq M_a \|\Pi_h \psi - \psi\|_a \|\varphi_h\|_a$$

and (66) (already proved) ends the proof, since $\|\psi\|_{\Phi_H^0} = \|g\|_{H_{\Xi}} = \|\chi\|_{\Xi_H^0}$, by definition. \square

EXAMPLE 3. Let us consider the mixed formulation of second order linear elliptic problems. Recall that $\Phi = \Sigma = H(\text{div}; \Omega)$ and $\Xi = V = L^2(\Omega)$. As usual we identify $L^2(\Omega)$ with its dual space, so that in our notation we have $\Xi = H_{\Xi} = \Xi' = L^2(\Omega)$. It is easy to see (using e.g. [17]) that if Ω is, for instance, a convex polygon, then Ξ_H^0 is $H^2(\Omega) \cap H_0^1(\Omega)$ and $\Phi_H^0 \equiv \Phi_{\Xi'}^0 = \nabla(\Phi_H^0) \subseteq (H^1(\Omega))^2$.

Let Σ_h and V_h be finite dimensional subspaces of Σ and V respectively. We consider, first, classical approximations of $H(\text{div}; \Omega)$; for instance we can choose as Σ_h the spaces of the elements (RT) introduced in [24], or the elements (BDM) and (BDFM) introduced in [11], [10], respectively. For a unified presentation we refer to [9]. Correspondingly V_h will be the space $\text{div } \Sigma_h$. For convenience of the reader we recall for instance the definition of BDM spaces. For k integer ≥ 1 we set

$$\begin{aligned}
\Sigma_h^k &:= \{\underline{\tau}_h \in \Sigma : \underline{\tau}_h|_K \in (P_k(K))^2 \ \forall K \in \mathcal{T}_h\} \\
(82) \quad V_h^k &:= \{v_h \in V : v_h|_K \in P_{k-1}(K) \ \forall K \in \mathcal{T}_h\},
\end{aligned}$$

where, as usual, $\{\mathcal{T}_h\}$ is a regular sequence of triangulations of Ω . A Fortin operator satisfying (65) for all these choices of finite element spaces can be constructed using suitable degrees of freedom. Moreover it is well-known that (66) holds true, see e.g. [9], pag. 132.

Since $V_h = \text{div } \Sigma_h$ then $\mathbb{K}_h \subseteq \mathbb{K}$, hence (DEK) and (62) trivially hold. It remains only to verify the strong approximability of Ξ_H^0 , that is

$$(83) \quad \|v - v^I\|_0 \leq \omega_5(h) \|v\|_1 \quad \text{for all } v \in H^1(\Omega),$$

which also holds thanks to standard approximation properties of piecewise polynomial spaces.

For various reasons, see for instance [4], [26], one might want to approximate Σ_h by continuous functions, using therefore finite element spaces that are not especially fit for mixed formulations. In constructing these new spaces, one might believe that (DEK) and (DIS) should be sufficient in order to well approximate eigenvalues and eigenvectors, once V_h satisfies the *strong approximability in Ξ_H^0* assumption. However, while conditions (62), (64) and (65) can be deduced from (DEK) and (DIS) the bound (66) does not, as it is shown by the following choice of the so-called $P_1 - \text{div}(P_1)$ element on a criss-cross mesh. Let us assume that Ω is a square, which is divided into $2N \times 2N$ subsquares, each of them partitioned into four triangles K by its diagonals. Then we set

$$(84) \quad \begin{aligned} \Sigma_h &:= \{\underline{\tau}_h \in (C^0(\Omega))^2 : \underline{\tau}_h|_K \in (P_1(K))^2 \ \forall K \in \mathcal{T}_h\}, \\ V_h &:= \text{div}(\Sigma_h) \subset \{v_h : v_h|_K \in P_0(K) \ \forall K \in \mathcal{T}_h\}. \end{aligned}$$

In a recent paper [5], we proved that the pair (Σ_h, V_h) defined in (84) satisfies (DEK) and (DIS) but the sequence T_{Ξ}^h does not converge uniformly to T_{Ξ} in $L^2(\Omega)$. This fact produces in the numerical computations spurious eigenvalues which converge to points belonging to the resolvent set of T_{Ξ} .

Hence (66), which we have seen to be necessary, has to be checked independently of (DEK) and (DIS). On the other hand, (DEK) is not necessary, and we can obtain convergence of eigenvalues with finite element spaces that fail to satisfy it. For instance, on a quasi-uniform triangulation, one might take $\Sigma_h = \Sigma_h^3$ (see (82)) and $V_h = V_h^2$. Notice that the pair (Σ_h^3, V_h^3) , as we have seen, works. Now, however, having chosen a smaller V_h , we obtain a bigger \mathbb{K}_h (not anymore contained in \mathbb{K}). This will not jeopardize property (66) (the Π_h operator working for the pair (Σ_h^3, V_h^3) will also work for the pair (Σ_h^3, V_h^2)) but (62) is now at risk. However, by inverse inequality (see e.g. [12])

$$(85) \quad b(\underline{\tau}_h, v) = b(\underline{\tau}_h, v - v^I) \leq C \|\underline{\tau}_h\|_1 \|v - v^I\|_0 \leq Ch^{-1} \|\underline{\tau}_h\|_a h^2 \|v\|_2,$$

for $\underline{\tau}_h \in \mathbb{K}_h$ and $v^I = L^2$ -projection of v onto V_h^2 . Notice that this argument will work for any pair (Σ_h^k, V_h^r) provided $k \geq r > 1$ (in other words, (Σ_h^2, V_h^1) will not work).

EXAMPLE 4. Biharmonic problem. In this case we have $\Phi = Z = H^1(\Omega)$ and $\Xi = V = H_0^1(\Omega)$. We take $H_{\Xi} = L^2(\Omega)$. We always assume that Ω is a convex polygon. In that case we obtain $\Phi_{\Xi'}^0 = \{z \in H^1(\Omega) : \exists v \in H_0^2(\Omega) \text{ with } z = \Delta v\} = \{z \in H^1(\Omega) : (z, \mu) = 0 \ \forall \mu \in L^2(\Omega) \text{ with } \Delta \mu = 0\}$ and, with obvious notation, $\Xi_{\Xi'}^0 = H^3(\Omega) \cap H_0^2(\Omega)$. For any given polygon, Φ_H^0 and Ξ_H^0 will be slightly more regular, according to the maximum angle (see e.g. [17]).

For every given regular sequence $\{\mathcal{T}_h\}$ of triangulations of Ω and for every integer $k \geq 2$ we can take as in [16], [20], [13]:

$$(86) \quad \begin{aligned} Z_h^k &:= \{z_h \in Z : z_h|_K \in P_k(K) \ \forall K \in \mathcal{T}_h\} \\ V_h^k &:= \{v_h \in V : v_h|_K \in P_k(K) \ \forall K \in \mathcal{T}_h\}. \end{aligned}$$

Notice that $V_h^k = Z_h^k \cap H_0^1(\Omega)$. We can now define $\Pi_h w$ in Z_h as the solution of: $(\nabla \Pi_h w, \nabla z_h) = (\nabla w, \nabla z_h)$ for all $z_h \in Z_h^k$. Clearly (64), (65), (66) hold. Similarly (63) holds by taking χ^I (here u^I) as the usual interpolant. On the other hand, to check (62) we have to assume quasi-uniformity of the decomposition and then proceed as in (85): for $z_h \in \mathbb{K}_h$ and $v \in H^3(\Omega) \cap H_0^2(\Omega)$

$$(\nabla z_h, \nabla v) = (\nabla z_h, \nabla v - \nabla v^I) \leq Ch^{-1} \|z_h\|_a Ch^2 \|v\|_3.$$

This shows the utility of the requirement $k \geq 2$. However, a more sophisticated proof, following the arguments of Scholz [25], shows that (62) also holds for $k = 1$.

REFERENCES

- [1] I. BABUŠKA, *Error-Bounds for Finite Element Method*, Numer. Math. **16** (1971), 322-333.
- [2] I. BABUŠKA, *On the finite element method with Lagrangian multipliers*, Numer. Math. **20** (1973), 179-192.
- [3] I. BABUŠKA – J. E. OSBORN, “Handbook of Numerical Analysis”, vol. II, ch. Eigenvalue Problems, North-Holland, 1991, pp. 641-788.
- [4] K. J. BATHE – C. NITIKITPAIBOON – X. WANG, *A mixed displacement-based finite element formulation for acoustic fluid-structure interaction*, Computers & Structures **56** (1995), 225-237.
- [5] D. BOFFI – F. BREZZI – L. GASTALDI, *On the problem of spurious eigenvalues in the approximation of linear elliptic problems in mixed form*, submitted to Math. Comp., 1997.
- [6] J. M. BOLAND – R. NICOLAIDES, *On the stability of bilinear-constant velocity-pressure finite elements*, Numer. Math. **44** (1984), 219-222.
- [7] J. H. BRAMBLE – J. E. OSBORN, *Rate of convergence for nonselfadjoint eigenvalue approximations*, Math. Comp. **27** (1973), 525-549.
- [8] F. BREZZI, *On the existence, uniqueness and approximation of saddle-point problems arising from Lagrangian multipliers*, R.A.I.R.O. Anal. Numer. **8** (1974), 129-151.
- [9] F. BREZZI – M. FORTIN, “Mixed and Hybrid Finite Element Methods”, Springer-Verlag, New York, 1991.
- [10] F. BREZZI – J. DOUGLAS JR. – M. FORTIN – L. D. MARINI, *Efficient rectangular mixed finite elements in two and three space variables*, R.A.I.R.O. Model. Math. Anal. Numer. **21** (1987), 237-250.
- [11] F. BREZZI – J. DOUGLAS JR. – L. D. MARINI, *Two families of mixed finite elements for second order elliptic problems*, Numer. Math. **47** (1985), 217-235.
- [12] P. G. CIARLET, “The Finite Element Method for Elliptic Problems”, North-Holland, Amsterdam, 1978.
- [13] P. G. CIARLET – P.-A. RAVIART, *A mixed finite element method for the biharmonic equation*, in “Mathematical Aspects of Finite Element in Partial Differential Equations”, C. de Boor (ed.), Academic Press, New York, 1974, 125-143.
- [14] J. FALK – J. E. OSBORN, *Error estimates for mixed methods*, R.A.I.R.O. Anal. Numer. **4** (1980), 249-277.

- [15] M. FORTIN, *An analysis of the convergence of mixed finite element methods*, R.A.I.R.O. Anal. Numer. **11** (1977), 341-354.
- [16] R. GLOWINSKI, *Approximations externes par éléments finis de Lagrange d'ordre un et deux, du problème de Dirichlet pour l'opérateur biharmonique. Méthodes itératives de résolution des problèmes approchés*, "Topics in Numerical Analysis", J. Miller (ed.), Academic Press, New York, 1973, 123-171.
- [17] P. GRISVARD, "Elliptic Problems in Non-Smooth Domains", Pitman, Marshfields, Mass., 1985.
- [18] C. JOHNSON – J. PITKÄRANTA, *Analysis of some mixed finite element methods related to reduced integration*, Math. Comp. **38** (1982), 375-400.
- [19] R. B. KELLOGG – J. E. OSBORN, *A regularity result for the Stokes problem*, J. Funct. Anal. **21** (1976), 397-431.
- [20] B. MERCIER, *Numerical solution of the biharmonic problem by mixed finite elements of class C^0* , Boll. U.M.I. **10** (1974), 133-149.
- [21] B. MERCIER – J. E. OSBORN – J. RAPPAZ – P.-A. RAVIART, *Eigenvalue approximation by mixed and hybrid methods*, Math. Comp. **36** (1981), 427-453.
- [22] J. T. ODEN – O. JACQUOTTE, *Stability of some mixed finite element methods for Stokesian flows*, Comp. Methods Appl. Mech. Eng. **43** (1984), 231-247.
- [23] J. E. OSBORN, *Eigenvalue approximations by mixed methods*, in "Advances in Computer Methods for Partial Differential Equations III", R. Vichnevetsky and R. Stepleman (eds.), New Brunswick, 1979, 158-161.
- [24] P.-A. RAVIART – J. M. THOMAS, *A mixed finite element method for second order elliptic problems*, in "Mathematical Aspects of the Finite Element Method", I. Galligani and E. Magenes (eds.), Lecture Notes in Math., Springer-Verlag, New York, 1977, 292-315.
- [25] R. SCHOLZ, *A mixed method for fourth order problems using linear finite elements*, R.A.I.R.O. Anal. Numer. **12** (1978), 85-90.
- [26] X. WANG – K. J. BATHE, *On mixed elements for acoustic fluid-structure interaction*, M³AS, **7** (1997), 329-344.

Dipartimento di Matematica
 Università di Pavia
 Via Abbiategrasso, 215
 27100 Pavia, Italy

Dipartimento di Matematica
 Università di Pavia
 and
 I.A.N.-C.N.R.
 Via Abbiategrasso, 215
 27100 Pavia, Italy

Dipartimento di Matematica
 Università di Roma "La Sapienza"
 P.le A. Moro, 2
 00185 Roma, Italy