

ON THE CONVERGENCE OF NONLINEAR SEMI-GROUPS

ISAO MIYADERA

(Received June 14, 1968)

1. Introduction. Let X be a Banach space and let $\{T(\xi); \xi \geq 0\}$ be a family of (nonlinear) operators from X into itself satisfying the following conditions:

(i) $T(0) = I$ (the identity) and $T(\xi + \eta) = T(\xi)T(\eta)$ for $\xi, \eta \geq 0$.

(ii) For each $x \in X$, $T(\xi)x$ is strongly continuous in $\xi \geq 0$.

We call such a family $\{T(\xi); \xi \geq 0\}$ simply a *nonlinear semi-group*. If there is a non-negative constant c such that

(iii) $\|T(\xi)x - T(\xi)y\| \leq e^{c\xi} \|x - y\|$ for $x, y \in X$ and $\xi \geq 0$,

then a nonlinear semi-group $\{T(\xi); \xi \geq 0\}$ is said to be of *local type*. (In particular, if $c = 0$, it is called a *nonlinear contraction semi-group*.) We define the *infinitesimal generator* A_0 of a nonlinear semi-group $\{T(\xi); \xi \geq 0\}$ by

$$(1.1) \quad A_0x = \lim_{\delta \rightarrow 0+} \delta^{-1}(T(\delta) - I)x$$

and the *weak infinitesimal generator* A' by

$$(1.2) \quad A'x = \text{w-lim}_{\delta \rightarrow 0+} \delta^{-1}(T(\delta) - I)x,$$

if the right sides exist. (The notation “lim” (“w-lim”) means the strong limit (the weak limit) in X .)

REMARK. In case of *linear* semi-groups, it is well known that the weak infinitesimal generator coincides with the infinitesimal generator.

H. F. Trotter [9] proved the following convergence theorem of *linear* semi-groups.

THEOREM. Let $\{T_n(\xi); \xi \geq 0\}_{n=1,2,3,\dots}$ be a sequence of semi-groups (of linear operators) of class (C_0) satisfying the stability condition

$$\|T_n(\xi)\| \leq Me^{\omega\xi} \text{ for } \xi \geq 0, n = 1, 2, 3, \dots,$$

where M and ω are independent of n and ξ . Let A_n be the infinitesimal generator of $\{T_n(\xi); \xi \geq 0\}$ and define $Ax = \lim_n A_nx$.

Suppose that

(a) $D(A)$ (the domain of A) is dense in X ,

(b) for some $\lambda > \omega$, $R(\lambda - A) = X$ (or $\overline{R(\lambda - A)} = X$).

Then A (or the closure of A) generates a semi-group $\{T(\xi); \xi \geq 0\}$ of class (C_0) ; and for each $x \in X$

$$\lim_n T_n(\xi) x = T(\xi) x$$

for $\xi \geq 0$ and the convergence is uniform with respect to ξ in every finite interval.

In this paper we shall study the convergence of nonlinear semi-groups $\{T_n(\xi); \xi \geq 0\}$ ($n = 1, 2, 3, \dots$) of local type with the stability condition

$$(1.3) \quad \|T_n(\xi) x - T_n(\xi) y\| \leq e^{\omega \xi} \|x - y\|;$$

and we can prove the following (see Theorem 2.1):

"Let A_n be the infinitesimal generator of $\{T_n(\xi); \xi \geq 0\}$, and let A' be the weak infinitesimal generator of a semi-group $\{T(\xi); \xi \geq 0\}$ of local type. If there exists a dense set D_0 such that for each $x \in D_0$, $\lim_n A_n x = A'x$ and $\lim_n A_n T(\xi) x = A'T(\xi) x$ for a.a. ξ (with additional conditions $T_n(\xi) x \in D(A_n)$ for a.a. ξ), then for each $x \in X$,

$$T(\xi)x = \lim_n T_n(\xi)x$$

uniformly on every finite interval."

(We note here that we may take $\bigcup_{x \in D_0} \{T(\xi)x; \lim_n A_n T(\xi)x = A'T(\xi)x\}$ as a set D in Theorem 2.1.) In particular if X^* (the adjoint space of X) is uniformly convex, then the Trotter theorem holds good for our nonlinear case (see Theorem 2.3).

For linear semi-group $\{T(\xi); \xi \geq 0\}$ of class (C_0) , it is well known that

$$T(\xi)x = \lim_{\delta \rightarrow 0+} T_\delta(\xi)x \quad \text{for } x \in X, \xi \geq 0,$$

where $A_\delta = \delta^{-1}(T(\delta) - I)$ and $\{T_\delta(\xi); \xi \geq 0\}$ is the semi-group generated by A_δ . And, in this case, $T_\delta(\xi) (= \exp(\xi A_\delta))$ is continuous in $\xi \geq 0$ with respect to the uniform operator topology (see [3]). In §4 we shall give similar results for nonlinear semi-groups of local type.

2. Theorems. The main theorems are as follows.

THEOREM 2.1. Let $\{T_n(\xi); \xi \geq 0\}_{n=1,2,3,\dots}$ be a sequence of nonlinear semi-groups of local type satisfying the stability condition

$$(2.1) \quad \|T_n(\xi)x - T_n(\xi)y\| \leq e^{\omega\xi} \|x - y\|$$

for $\xi \geq 0$, $n = 1, 2, 3, \dots$ and $x, y \in X$, where ω is a non-negative constant independent of n, x, y , and ξ . Let A_n be the infinitesimal generator of $\{T_n(\xi); \xi \geq 0\}$ and let $\lim_n A_n x = Ax$ on a set $D \subset \bigcap_{n=1}^{\infty} D(A_n)$.

Suppose that

(a) A (defined on D) is a restriction of the weak infinitesimal generator of some nonlinear semi-group $\{T(\xi); \xi \geq 0\}$ such that for any $\beta > 0$, $\{T(\xi); 0 \leq \xi \leq \beta\}$ is equi-Lipschitz continuous on every bounded set,

(b) there exists a set $D_0 \subset D$ such that for each $x \in D_0$

(b₁) for each n , $T_n(\xi)x \in D(A_n)$ for a.a. $\xi \geq 0$,

(b₂) $T(\xi)x \in D$ for a.a. $\xi \geq 0$.

Then for each $x \in \overline{D_0}$ (the strong closure of D_0) we have

$$(2.2) \quad T(\xi)x = \lim_n T_n(\xi)x \quad \text{for each } \xi \geq 0,$$

and the convergence is uniform with respect to ξ in every finite interval.

REMARKS 1. If for any bounded set B there is a positive constant M_B such that $\|T(\xi)x - T(\xi)y\| \leq M_B \|x - y\|$ for $\xi \in [0, \beta]$ and $x, y \in B$, then the family $\{T(\xi); 0 \leq \xi \leq \beta\}$ is said to be *equi-Lipschitz continuous* on every bounded set.

2. The above theorem remains true even if the conditions " $D \subset \bigcap_{n=1}^{\infty} D(A_n)$ "

and (b₁) are replaced by " $D \subset \bigcup_{k=1}^{\infty} \bigcap_{n=k}^{\infty} D(A_n)$ " and the following (b'₁), respectively.

(b'₁) For sufficiently large n , $T_n(\xi)x \in D(A_n)$ for a. a. $\xi \geq 0$.

The proof is given in §3.

In the above theorem, if X is a reflexive Banach space and $D \supset D(A_0)$, where A_0 is the infinitesimal generator of $\{T(\xi); \xi \geq 0\}$ in the assumption (a), then the assumption (b) is automatically satisfied by taking $D_0 = D$. In fact, if $x \in D$, then $x \in D(A')$ and $x \in D(A_n)$, where A' is the weak infinitesimal generator of $\{T(\xi); \xi \geq 0\}$; and hence $T(\xi)x$ and $T_n(\xi)x$ are strongly absolutely continuous on every finite interval (see the proof of Lemma 3.3). Thus the reflexivity of X shows that $T(\xi)x$ and $T_n(\xi)x$ are

strongly differentiable at a.a. $\xi \geq 0$ (for example, see Y. Kōmura [5]), so that the semi-group property (i) (in §1) implies

$$T_n(\xi)x \in D(A_n) \quad \text{for a. a. } \xi \geq 0$$

and

$$T(\xi)x \in D(A_0) \subset D \quad \text{for a. a. } \xi \geq 0.$$

Thus we have the following

THEOREM 2.2. *Let $\{T_n(\xi); \xi \geq 0\}_{n=1,2,3,\dots}$ be a sequence of nonlinear semi-groups in Theorem 2.1 defined on a reflexive Banach space X , and let A_n be the infinitesimal generator of $\{T_n(\xi); \xi \geq 0\}$ and assume $\lim_n A_n x = Ax$ on a set D .*

If the condition (a) in Theorem 2.1 is satisfied and $D \supset D(A_0)$ (i.e., $A' \supset A \supset A_0$), then for each $x \in \overline{D}^{(1)}$ we have

$$T(\xi)x = \lim_n T_n(\xi)x \quad \text{for all } \xi \geq 0,$$

and the convergence is uniform with respect to ξ in every finite interval.

T.Kato proved a generation theorem of nonlinear contraction semi-groups defined on a Banach space such that the adjoint space is uniformly convex (see T.Kato [4] and F.E. Browder [1]), and his result has been extended to some class of nonlinear semi-groups (which contains semi-groups of local type) by S.Oharu [8]. Using Oharu's result, we can prove the following

THEOREM 2.3. *Let the adjoint space X^* of X be a uniformly convex Banach space. Let $\{T_n(\xi); \xi \geq 0\}_{n=1,2,3,\dots}$ be a sequence of nonlinear semi-groups in Theorem 2.1, and let A_n be the infinitesimal generator of $\{T_n(\xi); \xi \geq 0\}$ and define $Ax = \lim_n A_n x$.*

Suppose that

(a') $D(A)$ (the domain of A) is dense in X ,

(b') for some $h_0 \in (0, 1/\omega)$, $R(1 - h_0 A) = X$.

Then A is the weak infinitesimal generator of a nonlinear semi-group $\{T(\xi); \xi \geq 0\}$ of local type and for each $x \in X$

$$(*) \quad T(\xi)x = \lim_n T_n(\xi)x \quad \text{for all } \xi \geq 0,$$

and the convergence is uniform with respect to ξ in every finite interval.

1) It is easy to see that $\overline{D} = \overline{D(A')} = \overline{D(A_0)}$.

REMARK. If we omit the condition (a'), then A is the weak infinitesimal generator of a nonlinear semi-group $\{T(\xi); \xi \geq 0\}$ of local type defined on $D(A)$ and the convergence (*) holds on $\overline{D(A)}$.

PROOF. If we can prove the following

$$(2.3) \quad \left\{ \begin{array}{l} \text{the limit operator } A \text{ is the weak infinitesimal generator of a} \\ \text{nonlinear semi-group } \{T(\xi); \xi \geq 0\} \text{ such that for any } \beta > 0, \{T(\xi); \\ 0 \leq \xi \leq \beta\} \text{ is equi-Lipschitz continuous on every bounded set,} \end{array} \right.$$

then the convergence (*) follows from Theorem 2.2 by taking $D=D(A)$ because X is reflexive with X^* , and the convergence implies

$$\|T(\xi)x - T(\xi)y\| \leq e^{\omega\xi} \|x - y\|$$

for $\xi \geq 0, x, y \in X$.

We shall now prove (2.3). Let x and y be elements in $D(A)$. By Lemma 3.1, for each n , we have

$$\operatorname{Re}(A_n x - A_n y, f) \leq \omega \|x - y\|^2$$

for $f = F(x - y)$, where F denotes the duality map from X into X^* . Letting $n \rightarrow \infty$

$$(2.4) \quad \operatorname{Re}(Ax - Ay, f) \leq \omega \|x - y\|^2$$

This means that $B = A - \omega$ is a dissipative (i.e., $\operatorname{Re}(Bx - By, f) \leq 0$). And the assumption (b') implies

$$R(1 - h_0(1 - h_0\omega)^{-1}B) = X,$$

so that $R(1 - \varepsilon B) = X$ for all $\varepsilon > 0$ (see S. Oharu [7], Y. Kōmura [5], T. Kato [4]). This leads

$$(2.5) \quad R(1 - hA) = X \quad \text{for all } h \in (0, 1/\omega).$$

Let $h \in (0, 1/\omega)$. Since $\|x - y - h(Ax - Ay)\| \|x - y\| \geq \operatorname{Re}(x - y - h(Ax - Ay), f) = \|x - y\|^2 - h \operatorname{Re}(Ax - Ay, f) \geq (1 - h\omega) \|x - y\|^2$ ($x, y \in D(A), f = F(x - y)$) by (2.4), we obtain

$$\|x - y - h(Ax - Ay)\| \geq (1 - h\omega) \|x - y\|$$

for each $x, y \in D(A)$. Consequently

$$(2.6) \quad \text{for each } h \in (0, 1/\omega), (1 - hA)^{-1} \text{ exists on } X.$$

Now (2.3) follows from Oharu's results ([8; Theorems 4.1 and 4.2]).²⁾ Q.E.D.

3. Proof of Theorem 2.1. We start from the following

LEMMA 3.1 *If $\{T(\xi); \xi \geq 0\}$ is a nonlinear semi-group of local type with $\|T(\xi)x - T(\xi)y\| \leq e^{\omega\xi}\|x - y\|$ ($\xi \geq 0, x, y \in X$), and if A' is its weak infinitesimal generator, then for each $x, y \in D(A')$ we have*

$$\operatorname{Re}(A'x - A'y, f) \leq \omega\|x - y\|^2$$

for any $f \in F(x - y)$, where F is the duality map from X into X^* .

PROOF. Let $x, y \in D(A')$, and let $f \in F(x - y)$.

$$\begin{aligned} & \operatorname{Re}(\xi^{-1}[T(\xi)x - x] - \xi^{-1}[T(\xi)y - y], f) \\ &= \xi^{-1}\operatorname{Re}(T(\xi)x - T(\xi)y, f) - \xi^{-1}\operatorname{Re}(x - y, f) \\ &\leq \xi^{-1}\|T(\xi)x - T(\xi)y\|\|x - y\| - \xi^{-1}\|x - y\|^2 \\ &\leq \xi^{-1}(e^{\omega\xi} - 1)\|x - y\|^2. \end{aligned}$$

Letting $\xi \rightarrow 0 +$, we get

$$\operatorname{Re}(A'x - A'y, f) \leq \omega\|x - y\|^2.$$

Q. E. D.

LEMMA 3.2 (T.Kato [4]). *Let $x(\xi)$ be an X -valued function on an interval of real numbers. Suppose $x(\xi)$ has a weak derivative $x'(\eta) \in X$ at $\xi = \eta$ and $\|x(\xi)\|$ is differentiable at $\xi = \eta$. Then*

$$\|x(\eta)\| \left[\frac{d}{d\xi} \|x(\xi)\| \right]_{\xi=\eta} = \operatorname{Re}(x'(\eta), f)$$

for any $f \in F(x(\eta))$.

LEMMA 3.3. *Let $\{T(\xi); \xi \geq 0\}$ be a nonlinear semi-group with the*

2) We note that (2.4) implies the condition (S) in his theorem.

weak infinitesimal generator A' , and let for any $\beta > 0$ the family $\{T(\xi); 0 \leq \xi \leq \beta\}$ be equi-Lipschitz continuous on every bounded set. If $x \in D(A')$ and $T(\xi)x \in D(A')$ for a.a. $\xi \geq 0$, then $A'T(\xi)x$ is strongly measurable and essentially bounded (and hence, Bochner integrable) on every finite interval, and

$$T(\xi)x - x = \int_0^\xi A'T(\eta)x d\eta \quad \text{for all } \xi \geq 0.$$

Consequently $T(\xi)x$ is strongly differentiable at a.a. ξ and

$$(d/d\xi)T(\xi)x = A'T(\xi)x \quad \text{for a. a. } \xi \geq 0.$$

PROOF. Let $\beta > 0$ be an arbitrary given. If we put

$$B = \{T(\xi)x; 0 \leq \xi \leq 1\} \quad \text{and} \quad K = \sup_{0 < \delta \leq 1} \delta^{-1} \|T(\delta)x - x\|,$$

then B is a bounded set and K is finite. Since the family $\{T(\xi); 0 \leq \xi \leq \beta\}$ is equi-Lipschitz continuous on B , there exists a constant M_B such that

$$\|T(\xi)y - T(\xi)z\| \leq M_B \|y - z\|$$

for all $y, z \in B$ and $\xi \in [0, \beta]$. Therefore, for $0 \leq \xi \leq \beta$ and $0 \leq \delta \leq 1$, we have

$$(3.1) \quad \|T(\xi + \delta)x - T(\xi)x\| \leq M_B \|T(\delta)x - x\| \leq M_B K\delta.$$

This shows that $T(\xi)x$ is strongly absolutely continuous on $[0, \beta]$. Since $T(\xi)x \in D(A')$ for a. a. $\xi \geq 0$,

$$(3.2) \quad \begin{cases} \int A'T(\xi)x = \text{w-lim}_{\delta \rightarrow 0+} \delta^{-1}(T(\delta) - I)T(\xi)x \\ \quad = \text{w-lim}_{\delta \rightarrow 0+} \delta^{-1}(T(\xi + \delta)x - T(\xi)x) \end{cases}$$

for a.a. $\xi \geq 0$; hence $A'T(\xi)x$ is strongly measurable (for example, see [3, Theorem 3.5.4]). By (3.1) and (3.2)

$$\|A'T(\xi)x\| \leq M_B K \quad \text{for a. a. } \xi \in [0, \beta],$$

so that $A'T(\xi)x$ is essentially bounded on $[0, \beta]$. Consequently $A'T(\xi)x$ is Bochner integrable on $[0, \beta]$.

Let $f \in X^*$. Since $(T(\xi)x, f) (= f(T(\xi)x))$ is absolutely continuous on $[0, \beta]$, $(T(\xi)x, f)$ is differentiable at a.a. $\xi \in [0, \beta]$ and

$$(T(\xi)x, f) - (x, f) = \int_0^\xi \frac{d}{d\eta}(T(\eta)x, f) d\eta$$

for any $\xi \in [0, \beta]$. Moreover it follows from (3.2) that

$$\frac{d}{d\xi}(T(\xi)x, f) = (A'T(\xi)x, f)$$

for a. a. $\xi \in [0, \beta]$. Thus the above equalities and the Bochner integrability of $A'T(\xi)x$ on $[0, \beta]$ show that

$$\begin{aligned} (T(\xi)x, f) - (x, f) &= \int_0^\xi (A'T(\eta)x, f) d\eta \\ &= \left(\int_0^\xi A'T(\eta)x d\eta, f \right) \end{aligned}$$

for all $\xi \in [0, \beta]$. Hence we get

$$T(\xi)x - x = \int_0^\xi A'T(\eta)x d\eta \quad \text{for all } \xi \in [0, \beta]$$

and $(d/d\xi)T(\xi)x = A'T(\xi)x$ for a. a. $\xi \in [0, \beta]$.

Q. E. D.

LEMMA 3.4. *Under the assumptions of Theorem 2.1, for each $x \in D_0$ we have the following :*

(3.3) $\left\{ \begin{array}{l} AT(\xi)x \text{ is strongly measurable and essentially bounded on every} \\ \text{finite interval.} \end{array} \right.$

$$(3.4) \quad T(\xi)x - x = \int_0^\xi AT(\eta)x d\eta \quad \text{for all } \xi \geq 0$$

and $(d/d\xi)T(\xi)x = AT(\xi)x$ for a. a. $\xi \geq 0$.

$$(3.5) \quad T_n(\xi)x - x = \int_0^\xi A_n T_n(\eta)x d\eta \quad \text{for all } \xi \geq 0$$

and $(d/d\xi)T_n(\xi)x = A_n T_n(\xi)x$ for a. a. $\xi \geq 0$.

PROOF. If we denote the weak infinitesimal generator of $\{T(\xi); \xi \geq 0\}$ by A' , then the condition (a) is as follows ;

$$(3.6) \quad D \subset D(A') \text{ and } Ax = A'x \quad \text{for } x \in D.$$

Let $x \in D_0$. By (3.6) and (b_2)

$$x \in D(A'), \quad T(\xi)x \in D(A') \text{ and } A'T(\xi)x = AT(\xi)x$$

for a.a. $\xi \geq 0$. Therefore it follows from Lemma 3.3 that $AT(\xi)x$ ($=A'T(\xi)x$ a.a.) is strongly measurable and essentially bounded on every finite interval, and

$$T(\xi)x - x = \int_0^\xi AT(\eta)x \, d\eta \quad \text{for all } \xi \geq 0,$$

$$(d/d\xi)T(\xi)x = AT(\xi)x \quad \text{for a.a. } \xi \geq 0.$$

We remark that for any $\beta > 0$, $\{T_n(\xi); 0 \leq \xi \leq \beta\}$ is equi-Lipschitz continuous on X , because it is of local type. Since $x \in D(A_n)$ and $T_n(\xi)x \in D(A_n)$ for a.a. $\xi \geq 0$ (see $M(b_1)$), (3.5) also follows from Lemma 3.3.

Q. E. D.

PROOF OF THEORE 2.1. Let $x \in D_0$ and put

$$(3.7) \quad z_n(\xi) = T_n(\xi)x - T(\xi)x.$$

By Lemma 3.4

$$z_n(\xi) = \int_0^\xi (A_n T_n(\eta)x - AT(\eta)x) \, d\eta,$$

and each $z_n(\xi)$ has the strong derivative

$$z'_n(\xi) = A_n T_n(\xi)x - AT(\xi)x \quad \text{for a.a. } \xi \geq 0;$$

moreover each $\|z_n(\xi)\|$ is differentiable at a.a. $\xi \geq 0$ since $\|z_n(\xi)\|$ is absolutely continuous in $\xi \geq 0$. Therefore it follows from Lemma 3.2 that for a.a. $\xi \geq 0$

$$(3.8) \quad \begin{cases} \|z_n(\xi)\| [(d/d\xi)\|z_n(\xi)\|] = \operatorname{Re}(z'_n(\xi), f_\xi) \\ \quad = \operatorname{Re}(A_n T_n(\xi)x - AT(\xi)x, f_\xi) \end{cases}$$

for every $f_\xi \in F(z_n(\xi))$. And

$$(3.9) \quad \|z_n(\xi)\|^2 = \int_0^\xi (d/d\eta) \|z_n(\eta)\|^2 d\eta = 2 \int_0^\xi \|z_n(\eta)\| [(d/d\eta) \|z_n(\eta)\|] d\eta$$

for all $\xi \geq 0$.

Let $\beta > 0$ be arbitrarily given. We shall show that the sequence $\{\|z_n(\xi)\| [(d/d\xi) \|z_n(\xi)\|]\}$ is uniformly (essentially) bounded on $[0, \beta]$. Put

$$K_1 = \text{ess sup}_{0 \leq \xi \leq \beta} \|AT(\xi)x\| (< \infty)$$

(see (3.3)). Since $\|A_n T_n(\xi)x\| = \lim_{\delta \rightarrow 0+} \|\delta^{-1}(T_n(\xi + \delta)x - T_n(\xi)x)\| \leq e^{\omega\xi} \lim_{\delta \rightarrow 0+} \delta^{-1} \|T_n(\delta)x - x\| = e^{\omega\xi} \|A_n x\|$ (a.a. ξ) and since $\lim_n A_n x = Ax$, there is a constant K_2 independent of n such that

$$\text{ess sup}_{0 \leq \xi \leq \beta} \|A_n T_n(\xi)x\| \leq K_2.$$

Consequently, for all n , we get

$$\text{ess sup}_{0 \leq \xi \leq \beta} \|z'_n(\xi)\| = \text{ess sup}_{0 \leq \xi \leq \beta} \|A_n T_n(\xi)x - AT(\xi)x\| \leq K_1 + K_2$$

and

$$(3.10) \quad \|z_n(\xi)\| \leq \int_0^\xi \|A_n T_n(\eta)x - AT(\eta)x\| d\eta \leq (K_1 + K_2)\beta$$

for every $\xi \in [0, \beta]$. Hence by (3.8)

$$\begin{aligned} |\|z_n(\xi)\| [(d/d\xi) \|z_n(\xi)\|]| &\leq \|z'_n(\xi)\| \|f_\xi\| = \|z'_n(\xi)\| \|z_n(\xi)\| \\ &\leq (K_1 + K_2)^2 \beta \end{aligned}$$

for a.a. $\xi \in [0, \beta]$; so that $\{\|z_n(\xi)\| [(d/d\xi) \|z_n(\xi)\|]\}$ is uniformly (essentially) bounded on $[0, \beta]$. Thus by the Lebesgue convergence theorem

$$(3.11) \quad \left\{ \begin{aligned} \limsup_{n \rightarrow \infty} \|z_n(\xi)\|^2 &= \limsup_{n \rightarrow \infty} 2 \int_0^\xi \|z_n(\eta)\| [(d/d\eta) \|z_n(\eta)\|] d\eta \\ &\leq 2 \int_0^\xi \limsup_{n \rightarrow \infty} \|z_n(\eta)\| [(d/d\eta) \|z_n(\eta)\|] d\eta \end{aligned} \right.$$

for all $\xi \in [0, \beta]$.

Since $T(\xi)x \in D \subset D(A_n)$ and $T_n(\xi)x \in D(A_n)$ for a.a. ξ , it follows from

Lemma 3.1 that for a.a. $\xi \geq 0$

$$(3.12) \quad \operatorname{Re}(A_n T_n(\xi) x - A_n T(\xi) x, f_\xi) \leq \omega \|z_n(\xi)\|^2$$

for every $f_\xi \in F(z_n(\xi))$. Combining this with (3.8), for a.a. $\xi \in [0, \beta]$

$$\begin{aligned} \|z_n(\xi)\| [(d/d\xi)\|z_n(\xi)\|] &\leq \operatorname{Re}(A_n T(\xi) x - AT(\xi) x, f_\xi) + \omega \|z_n(\xi)\|^2 \\ &\leq \|A_n T(\xi) x - AT(\xi) x\| \|z_n(\xi)\| + \omega \|z_n(\xi)\|^2 \\ &\leq (K_1 + K_2)\beta \|A_n T(\xi) x - AT(\xi) x\| + \omega \|z_n(\xi)\|^2 \quad (\text{see (3.10)}); \end{aligned}$$

and hence

$$(3.13) \quad \limsup_{n \rightarrow \infty} \|z_n(\xi)\| [(d/d\xi)\|z_n(\xi)\|] \leq \omega \limsup_{n \rightarrow \infty} \|z_n(\xi)\|^2$$

for a.a. $\xi \in [0, \beta]$. If we put

$$g(\xi) = \limsup_{n \rightarrow \infty} \|z_n(\xi)\|^2 \quad \text{for } \xi \in [0, \beta],$$

then $0 \leq g(\xi) \leq (K_1 + K_2)^2 \beta^2$ on $[0, \beta]$ (see (3.10)), and from (3.11) and (3.13) we obtain

$$0 \leq g(\xi) \leq 2\omega \int_0^\xi g(\eta) d\eta$$

for every $\xi \in [0, \beta]$. It is easy to see that the above inequality implies $g(\xi) = 0$ for $\xi \in [0, \beta]$. Thus we get

$$\lim_n \|T_n(\xi) x - T(\xi) x\| (= \lim_n \|z_n(\xi)\|) = 0$$

for all $\xi \in [0, \beta]$. We shall show that the above convergence is uniform. Since

$$\|z_n(\xi)\|^2 \leq 2 \int_0^\xi \|z'_n(\eta)\| \|z_n(\eta)\| d\eta$$

(see (3.8) and (3.9)),

$$\sup_{0 \leq \xi \leq \beta} \|z_n(\xi)\|^2 \leq 2 \int_0^\beta \|z'_n(\eta)\| \|z_n(\eta)\| d\eta \rightarrow 0$$

as $n \rightarrow \infty$, because the integrand converges boundedly to zero. Thus the theorem holds for $x \in D_0$.

Finally let $x \in D_0$. There is a sequence $\{x_k\}$ ($x_k \in D_0$) such that $\lim_k x_k = x$. Now

$$\begin{aligned} \|T_n(\xi)x - T(\xi)x\| &\leq \|T_n(\xi)x - T_n(\xi)x_k\| \\ &\quad + \|T_n(\xi)x_k - T(\xi)x_k\| + \|T(\xi)x_k - T(\xi)x\| \\ &\leq e^{\omega\xi}\|x - x_k\| + \|T_n(\xi)x_k - T(\xi)x_k\| + M_B\|x_k - x\| \end{aligned}$$

for $\xi \in [0, \beta]$. (Note there is a constant M_B such that $\|T(\xi)x_k - T(\xi)x\| \leq M_B\|x_k - x\|$ for $\xi \in [0, \beta]$ and k , since the set $B = \{x, x_1, x_2, \dots\}$ is bounded and the family $\{T(\xi); 0 \leq \xi \leq \beta\}$ is equi-Lipschitz continuous on bounded set.) Hence we get

$$\lim_n \|T_n(\xi)x - T(\xi)x\| = 0$$

uniformly on $[0, \beta]$.

Q. E. D.

4. Approximation of semi-groups. Let $\{T(\xi); \xi \geq 0\}$ be a nonlinear semi-group of local type with $\|T(\xi)x - T(\xi)y\| \leq e^{\omega\xi}\|x - y\|$, and let A_0 be its infinitesimal generator, and put

$$A_\delta = \delta^{-1}(T(\delta) - I) \quad \text{for } \delta > 0.$$

THEOREM 4.1. I. *Each A_δ is the infinitesimal generator of a semi-group $\{T_\delta(\xi); \xi \geq 0\}$ of local type satisfying the following conditions:*

(a) *For each $x \in X$, $T_\delta(\xi)x \in C^1([0, \infty); X)^{3)}$ and*

$$(d/d\xi)T_\delta(\xi)x = A_\delta T_\delta(\xi)x \quad \text{for all } \xi \geq 0.$$

(b) *For each $\xi \geq 0$*

$$\sup_{x \neq y} \|T_\delta(\xi+h)x - T_\delta(\xi+h)y - (T_\delta(\xi)x - T_\delta(\xi)y)\| / \|x - y\| \rightarrow 0 \text{ as } h \rightarrow 0.$$

II. Suppose that

$$(4.1) \quad \begin{cases} \text{there exists a set } D_0 \text{ such that } D_0 \subset D(A_0) \text{ and for any } x \in D_0, \\ T(\xi)x \in D(A_0) \text{ for a.a. } \xi \geq 0. \end{cases}$$

3) $C^1([0, \infty); X)$ denotes the set of all strongly continuously differentiable X -valued functions defined on $[0, \infty)$.

Then for each $x \in \overline{D_0}$ we have

$$(4.2) \quad T(\xi)x = \lim_{\delta \rightarrow 0+} T_\delta(\xi)x \quad \text{for all } \xi \geq 0,$$

and the convergence is uniform with respect to ξ in every finite interval.

REMARK. In case of nonlinear contraction semi-groups, the theorem has been proved by the author [6] (see also J. R. Dorroh [2]).

If X is a reflexive Banach space, then the assumption (4.1) is satisfied by taking $D_0 = D(A_0)$. (For, if $x \in D(A_0)$, then $T(\xi)x$ is strongly absolutely continuous on every finite interval. It follows from the reflexivity of X that $T(\xi)x$ is strongly differentiable at a.a. ξ and a fortiori $T(\xi)x \in D(A_0)$ for a.a. $\xi \geq 0$.) Thus we have the following

COROLLARY 4.2. *If $\{T(\xi); \xi \geq 0\}$ is a nonlinear semi-group of local type defined on a reflexive Banach space, then for each $x \in \overline{D(A_0)}$*

$$T(\xi)x = \lim_{\delta \rightarrow 0+} T_\delta(\xi)x \quad \text{for all } \xi \geq 0,$$

and the convergence is uniform with respect to ξ in every finite interval.

We shall now prove Theorem 4.1.

PROOF. I. Fix $\delta > 0$. Since the map $x \rightarrow A_\delta x$ is Lipschitz continuous, uniformly in $x \in X$ (in fact, $\|A_\delta x - A_\delta y\| \leq \delta^{-1}(e^{\omega\delta} + 1)\|x - y\|$ for $x, y \in X$), the equation

$$\begin{cases} (d/d\xi)u(\xi; x) = A_\delta u(\xi; x) & \text{for } \xi \geq 0 \\ u(0; x) = x \end{cases}$$

has a unique solution $u(\xi; x) \in C^1([0, \infty); X)$ for any $x \in X$. If we define $T_\delta(\xi)$ by

$$T_\delta(\xi)x = u(\xi; x) \quad \text{for } \xi \geq 0, x \in X,$$

then $\{T_\delta(\xi); \xi \geq 0\}$ is a nonlinear semi-group satisfying the condition (a) and its infinitesimal generator is A_δ .

We shall now prove that $\{T_\delta(\xi); \xi \geq 0\}$ is of local type. Fix $x, y \in X$ and put

$$z(\xi) = T_\delta(\xi)x - T_\delta(\xi)y.$$

Clearly $z(\xi) \in C^1([0, \infty); X)$ and

$$\begin{cases} (d/d\xi) z(\xi) = A_\delta T_\delta(\xi) x - A_\delta T_\delta(\xi) y \\ z(0) = x - y. \end{cases}$$

Since $\|z(\xi)\|$ is absolutely continuous, $\|z(\xi)\|$ is differentiable at a.a. $\xi \geq 0$. By Lemma 3.2, for a.a. $\xi \geq 0$

$$\begin{aligned} \|z(\xi)\| [(d/d\xi) \|z(\xi)\|] &= \operatorname{Re}(z'(\xi), f_\xi) \\ &= \operatorname{Re}(A_\delta T_\delta(\xi)x - A_\delta T_\delta(\xi)y, f_\xi) \end{aligned}$$

for every $f_\xi \in F(z(\xi))$. Note that for each $u, v \in X$

$$\operatorname{Re}(A_\delta u - A_\delta v, f) \leq \delta^{-1}(e^{\omega\delta} - 1) \|u - v\|^2$$

for all $f \in F(u - v)$. Hence

$$\|z(\xi)\| [(d/d\xi) \|z(\xi)\|] \leq c_\delta \|z(\xi)\|^2 \quad \text{for a.a. } \xi \geq 0,$$

where $c_\delta = \delta^{-1}(e^{\omega\delta} - 1)$; and

$$\begin{aligned} \|z(\xi)\|^2 &= \|z(0)\|^2 + \int_0^\xi [(d/d\eta) \|z(\eta)\|^2] d\eta \\ &= \|z(0)\|^2 + 2 \int_0^\xi \|z(\eta)\| [(d/d\eta) \|z(\eta)\|] d\eta \\ &\leq \|z(0)\|^2 + 2c_\delta \int_0^\xi \|z(\eta)\|^2 d\eta \end{aligned}$$

for any $\xi \geq 0$. This leads the following inequality

$$\|z(\xi)\|^2 \leq \|z(0)\|^2 \sum_{k=0}^n (2c_\delta \xi)^k / k! + [(2c_\delta)^{n+1} / n!] \int_0^\xi (\xi - \eta)^n \|z(\eta)\|^2 d\eta$$

for all n and $\xi \geq 0$. Letting $n \rightarrow \infty$, we get $\|z(\xi)\|^2 \leq e^{2c_\delta \xi} \|z(0)\|^2$, i.e.,

$$(4.3) \quad \|T_\delta(\xi)x - T_\delta(\xi)y\| \leq e^{c_\delta \xi} \|x - y\| \quad \text{for all } \xi \geq 0,$$

so that $\{T_\delta(\xi); \xi \geq 0\}$ is of local type.

We shall show (b). Since $\|A_\delta x - A_\delta y\| \leq \delta^{-1}(e^{\omega\delta} + 1) \|x - y\|$ for all $x, y \in X$,

$$\begin{aligned} & \|T_\delta(\xi + h)x - T_\delta(\xi + h)y - (T_\delta(\xi)x - T_\delta(\xi)y)\| \\ &= \left\| \int_{\xi}^{\xi+h} (A_\delta T_\delta(\eta)x - A_\delta T_\delta(\eta)y) d\eta \right\| \\ &\leq \delta^{-1}(e^{\omega\delta} + 1) \left| \int_{\xi}^{\xi+h} \|T_\delta(\eta)x - T_\delta(\eta)y\| d\eta \right| \\ &\leq \delta^{-1}(e^{\omega\delta} + 1) e^{c_\delta(\xi+|h|)} \|x - y\| |h|. \end{aligned}$$

Hence we obtain (b).

II. Since $c_\delta = \delta^{-1}(e^{\omega\delta} - 1) \rightarrow \omega$ as $\delta \rightarrow 0+$, there is a constant $c > 0$ such that $c_\delta \leq c$ for $0 < \delta \leq 1$. Hence by (4.3) we obtain

$$(4.4) \quad \|T_\delta(\xi)x - T_\delta(\xi)y\| \leq e^{c\xi} \|x - y\|$$

for every $x, y \in X$, $\xi \geq 0$ and $\delta \in (0, 1]$.

Let $\{\delta_n\}$ be a sequence such that $\delta_n \rightarrow 0+$. Put

$$T_n(\xi) = T_{\delta_n}(\xi) \text{ and } A_n = A_{\delta_n} (= \delta_n^{-1}(T(\delta_n) - I)).$$

Since $\lim_n A_{\delta_n}x = A_0x$ on $D(A_0)$ and $D(A_n) = X$, the assumptions in Theorem 2.1 are satisfied by taking $A = A_0$ and $D = D(A_0)$. Therefore for each $x \in \overline{D_0}$ we have

$$T(\xi)x = \lim_n T_{\delta_n}(\xi)x \quad \text{for each } \xi \geq 0,$$

and the convergence is uniform with respect to ξ in every finite interval.

Q. E. D.

REFERENCES

- [1] F. E. BROWDER, Nonlinear equations of evolutions and nonlinear accretive operators in Banach spaces, Bull. Amer. Math. Soc., 73(1967), 867-874.
- [2] J. R. DORROH, Semigroups of nonlinear transformations, Notices Amer. Math. Soc., 15(1968), 128.
- [3] E. HILLE AND R. S. PHILLIPS, Functional analysis and semi-groups, Amer. Math. Soc. Colloq. Publ., 1957.

- [4] T. KATO, Nonlinear semi-groups and evolution equations, J. Math. Soc. Japan, 19 (1967), 508-520.
- [5] Y. KŌMURA, Nonlinear semi-groups in Hilbert space, J. Math. Soc. Japan, 19(1967), 493-507.
- [6] I. MIYADERA, Note on nonlinear contraction semi-groups, Proc. Amer. Math. Soc., 21 (1969), 219-225.
- [7] S. OHARU, Note on the representation of semi-groups of nonlinear operators, Proc. Japan Acad., 42(1966), 1149-1154.
- [8] S. OHARU, Nonlinear semi-groups in Banach spaces, to appear.
- [9] H. F. TROTTER, Approximation of semi-groups of operators, Pacific J. Math., 8(1958), 887-919.

DEPARTMENT OF MATHEMATICS
WASEDA UNIVERSITY
TOKYO, JAPAN

AND

DEPARTMENT OF MATHEMATICS
GEORGETOWN UNIVERSITY
WASHINGTON, D. C., U. S. A.