

On the equivalence of linear sets

Bence Csajbók* Corrado Zanella*

July 3, 2018

Abstract

Let L be a linear set of pseudoregulus type in a line ℓ in $\Sigma^* = \text{PG}(t-1, q^t)$, $t = 5$ or $t > 6$. We provide examples of q -order canonical subgeometries $\Sigma_1, \Sigma_2 \subset \Sigma^*$ such that there is a $(t-3)$ -subspace $\Gamma \subset \Sigma^* \setminus (\Sigma_1 \cup \Sigma_2 \cup \ell)$ with the property that for $i = 1, 2$, L is the projection of Σ_i from center Γ and there exists no collineation ϕ of Σ^* such that $\Gamma^\phi = \Gamma$ and $\Sigma_1^\phi = \Sigma_2$.

Condition (ii) given in Theorem 3 in Lavrauw and Van de Vonder (Des. Codes Cryptogr. 56:89–104, 2010) states the existence of a collineation between the projecting configurations (each of them consisting of a center and a subgeometry), which give rise by means of projections to two linear sets. It follows from our examples that this condition is not necessary for the equivalence of two linear sets as stated there. We characterize the linear sets for which the condition above is actually necessary.

Keywords: linear set, subgeometry, finite field, finite projective space, collineation.

MSC: 51E20

1 Introduction

If V is a vector space over the finite field \mathbb{F}_{q^t} , then $\text{PG}_{q^t}(V)$ denotes the projective space whose points are the one-dimensional \mathbb{F}_{q^t} -subspaces of V . If V has dimension n over \mathbb{F}_{q^t} , then $\text{PG}_{q^t}(V) = \text{PG}(n-1, q^t)$. For a point set $T \subset \text{PG}(n-1, q^t)$ we denote by $\langle T \rangle$ the projective subspace of $\text{PG}(n-1, q^t)$ spanned by the points in T . For $m \mid t$ and a set of elements $S \subset V$ we

*This research was supported by the Italian Ministry of Education, University and Research (PRIN 2012 project “Strutture geometriche, combinatoria e loro applicazioni”).

denote by $\langle S \rangle_{q^m}$ the \mathbb{F}_{q^m} -vector subspace of V spanned by the vectors in S . For the rest of the paper we assume that $q = p^e$ is a power of the prime p .

Let $R = \mathbb{F}_{q^t}^r$. The *field reduction map* [8] $\mathcal{F}_{r,t,q}$ is a map from the points of $\text{PG}_{q^t}(R) = \text{PG}(r-1, q^t)$ to the $(t-1)$ -subspaces of $\text{PG}_q(R) = \text{PG}(rt-1, q)$. Let $P = \text{PG}_{q^t}(T)$ be a point of $\text{PG}(r-1, q^t)$, where T is a one-dimensional \mathbb{F}_{q^t} -subspace of R . Then $\mathcal{F}_{r,t,q}(P) := \text{PG}_q(T)$. As $\dim_{\mathbb{F}_q}(T) = t$, it follows that $\mathcal{F}_{r,t,q}(P)$ is a $(t-1)$ -dimensional subspace of $\text{PG}(rt-1, q)$. Denote the point set of $\text{PG}(r-1, q^t)$ by \mathcal{P} . Then $\mathcal{F}_{r,t,q}(\mathcal{P})$ is a Desarguesian $(t-1)$ -spread of $\text{PG}(rt-1, q)$ [13]. Let S be a subspace of $\text{PG}(rt-1, q) = \text{PG}_q(R)$, then

$$\mathcal{B}(S) := \{P \in \text{PG}(r-1, q^t) : \mathcal{F}_{r,t,q}(P) \cap S \neq \emptyset\}.$$

When S is an \mathbb{F}_q -vector space of R , then we define $\mathcal{B}(S)$ analogously. A point set $L \subseteq \text{PG}(r-1, q^t)$ is said to be \mathbb{F}_q -*linear* (or just *linear*) of rank n if $L = \mathcal{B}(S)$ for some $(n-1)$ -subspace $S \subseteq \text{PG}(rt-1, q)$. If L has size $(q^n - 1)/(q - 1)$ (which is the maximum size for a linear set of rank n), then L is a *scattered* linear set. For further information on linear sets see [8, 17].

Definition 1.1. Let $\text{PG}_{q^t}(V) = \text{PG}(n-1, q^t)$ and let W be an \mathbb{F}_q -vector subspace of V . If $\dim_{\mathbb{F}_q}(W) = \dim_{\mathbb{F}_{q^t}}(V) = n$ and $\langle W \rangle_{q^t} = V$, then $\Sigma' = \{\langle w \rangle_{q^t} : w \in W^*\}$ is a canonical subgeometry of $\text{PG}_{q^t}(V)$.

Let $\Sigma' \cong \text{PG}(n-1, q)$ be a canonical subgeometry of $\Sigma^* = \text{PG}(n-1, q^t)$. Let $\Gamma \subset \Sigma^* \setminus \Sigma'$ be an $(n-1-r)$ -subspace and let $\Lambda \subset \Sigma^* \setminus \Gamma$ be an $(r-1)$ -subspace of Σ^* . The projection of Σ' from *center* Γ to *axis* Λ is the point set

$$L = p_{\Gamma, \Lambda}(\Sigma') := \{\langle \Gamma, P \rangle \cap \Lambda : P \in \Sigma'\}. \quad (1)$$

In [15] Lunardon and Polverino characterized linear sets as projections of canonical subgeometries. They proved the following.

Theorem 1.2 ([15, Theorems 1 and 2]). Let $\Sigma^*, \Sigma', \Lambda, \Gamma$ and $L = p_{\Gamma, \Lambda}(\Sigma')$ be defined as above. Then L is an \mathbb{F}_q -linear set of rank n and $\langle L \rangle = \Lambda$. Conversely, if L is an \mathbb{F}_q -linear set of rank n of $\Lambda = \text{PG}(r-1, q^t) \subset \Sigma^*$ and $\langle L \rangle = \Lambda$, then there is an $(n-1-r)$ -subspace Γ disjoint from Λ and a canonical subgeometry $\Sigma' \cong \text{PG}(n-1, q)$ disjoint from Γ such that $L = p_{\Gamma, \Lambda}(\Sigma')$.

Note that when $r = n$ in Theorem 1.2, then $L = p_{\Gamma, \Lambda}(\Sigma') = \Sigma'$, hence L is a canonical subgeometry. We rephrase here a theorem by Lavrauw and Van de Voorde.

Theorem 1.3 ([6, Theorem 3]). *For $i = 1, 2$, let $\Sigma_i \cong \text{PG}(n-1, q)$ be two canonical subgeometries of $\Sigma^* = \text{PG}(n-1, q^t)$ and let Γ_i be two $(n-1-r)$ -subspaces of Σ^* , such that $\Gamma_i \cap \Sigma_i = \emptyset$. Let $\Lambda_i \subset \Sigma^* \setminus \Gamma_i$ be two $(r-1)$ -subspaces and let $L_i = p_{\Gamma_i, \Lambda_i}(\Sigma_i)$. Suppose that L_i is not an \mathbb{F}_q -linear set of rank $k < n$. Then the following statements are equivalent.*

- (i) *There exists a collineation $\alpha: \Lambda_1 \rightarrow \Lambda_2$, such that $L_1^\alpha = L_2$,*
- (ii) *there exists a collineation β of Σ^* such that $\Sigma_1^\beta = \Sigma_2$ and $\Gamma_1^\beta = \Gamma_2$,*
- (iii) *for all canonical subgeometries $\Sigma \cong \text{PG}(n-1, q)$ in Σ^* , there exist δ, ϕ, ψ collineations of Σ^* , such that¹ $\Sigma^\delta = \Sigma$, $\Gamma_1^{\phi\delta} = \Gamma_2^\psi$, $\Sigma_1^\phi = \Sigma$ and $\Sigma_2^\psi = \Sigma$.*

We are going to prove that the implication (i) \Rightarrow (ii) does not hold. In Sections 3 and 4, we will show this in the case when $L_1 = L_2$ is a linear set of pseudoregulus type in $\text{PG}(1, q^n)$, for $n = 5$ or $n > 6$. In Section 5 a characterization will be given of the linear sets for which (i) \Rightarrow (ii) holds.

In this paper, the results which lead to counterexamples are presented in a general setting. In the remainder of this introduction some hints are given to a better understanding of the key facts.

1.1

The idea of a counterexample arose from the investigation of the linear sets of pseudoregulus type [14, 3]. A minimal counterexample to Theorem 1.3 (i) \Rightarrow (ii) can be obtained as follows. In $\text{PG}(4, q^5)$, with coordinates X_1, X_2, \dots, X_5 , let Γ be the plane of equations $X_1 = X_2 = 0$, and let ℓ be the line $X_3 = X_4 = X_5 = 0$. The sets

$$\Sigma_1 = \{ \langle (\lambda, \lambda^q, \lambda^{q^2}, \lambda^{q^3}, \lambda^{q^4}) \rangle_{q^5} : \lambda \in \mathbb{F}_{q^5}^* \}$$

and

$$\Sigma_2 = \{ \langle (\lambda, \lambda^{q^2}, \lambda^q, \lambda^{q^3}, \lambda^{q^4}) \rangle_{q^5} : \lambda \in \mathbb{F}_{q^5}^* \}$$

are canonical subgeometries. Defining $\text{PG}(4, q) = \{ \langle x \rangle_q : x \in \mathbb{F}_{q^5}^* \}$, the map $\varphi_1 : \text{PG}(4, q) \rightarrow \Sigma_1$ defined by $\langle x \rangle_q^{\varphi_1} = \langle x, x^q, x^{q^2}, x^{q^3}, x^{q^4} \rangle_{q^5}$ is a collineation, as well as the map $\varphi_2 : \text{PG}(4, q) \rightarrow \Sigma_2$ similarly defined. By [14, 3], the projections $p_{\Gamma, \ell}(\Sigma_1)$ and $p_{\Gamma, \ell}(\Sigma_2)$ coincide, and are a linear set of pseudoregulus type, say $\mathbb{L} = \{ \langle (\lambda, \lambda^q, 0, 0, 0) \rangle_{q^5} : \lambda \in \mathbb{F}_{q^5}^* \}$.

¹Compositions are executed from the left to the right.

Now assume that as stated in Theorem 1.3 (ii) a collineation β of $\text{PG}(4, q^5)$ exists such that $\Gamma^\beta = \Gamma$, and $\Sigma_1^\beta = \Sigma_2$. It is shown in Theorem 4.5 that a projectivity exists having the same two properties; call it ϕ . By the arguments in Lemma 4.4 in Section 4, the collineation $\varphi_2\phi^{-1}\varphi_1^{-1}$ of $\text{PG}(4, q)$ is the composition of a projectivity and either the map $\langle x \rangle_q \mapsto \langle x^{\theta_1} \rangle_q$ (cfr. ((6)), or the map

$$\langle x \rangle_q \mapsto \langle x^{-\theta_1} \rangle_q = \langle x^{\theta_4 - \theta_1} \rangle_q = \langle x^{q^2\theta_2} \rangle_q,$$

which in turn is the composition of a projectivity and $\langle x \rangle_q \mapsto \langle x^{\theta_2} \rangle_q$. However, by Corollary 3.6 in Section 3, no map of type $\langle x \rangle_q \mapsto \langle x^{\theta_s} \rangle_q$ with $0 < s < t = 5$ is a collineation, and this is a contradiction.

1.2

In section 5 the same counterexamples are found with essentially distinct arguments. Theorem 5.5 states that a counterexample to Theorem 1.3 (i) \Rightarrow (ii) can always be constructed in case the following condition does not hold: (A) For any two $(n-1)$ -subspaces $U, U' \subset \text{PG}(rt-1, q)$, such that $\mathcal{B}(U) = L = \mathcal{B}(U')$, a collineation $\gamma \in \text{P}\Gamma\text{L}(rt, q)$ exists, such that $U^\gamma = U'$, γ preserves the Desarguesian spread $\mathcal{F}_{r,t,q}(\mathcal{P})$, and the induced map on $\text{PG}(r-1, q^t)$ is a collineation.

If $L = \mathcal{B}(\mathcal{Q}_{t-1,q})$, where $\mathcal{Q}_{t-1,q} \subset \text{PG}(2t-1, q)$ is the nonsingular hypersurface of degree t studied in [5], then L is a linear set of pseudoregulus type in $\text{PG}(1, q^t)$.

In [5] all linear subspaces contained in $\mathcal{Q}_{t-1,q}$ are described for $q \geq t$. For $t \geq 4$ there exist two $(t-1)$ -subspaces, say U and U' , that are contained in $\mathcal{Q}_{t-1,q}$, and such that for any element γ of the stabilizer of $\mathcal{Q}_{t-1,q}$, preserving the standard Desarguesian spread, it holds $U^\gamma \neq U'$. For $t \geq 5$, $t \neq 6$, it is possible to choose U and U' with the additional property $\mathcal{B}(U) = L = \mathcal{B}(U')$. This leads to the counterexample of the previous subsection and its generalisations.

1.3

In the opinion of the authors of this paper, the proof of [6, Theorem 3] contains a wrong argument. The map δ defined at p. 93, line 21, is declared to be a projectivity, but is dealt with as a linear map e.g. in (1), as well as p_2 is. Such a δ is assumed to satisfy both conditions (i) δ maps an \mathbb{F}_q -vector space associated with Σ_1^χ onto an \mathbb{F}_q -vector space associated with Σ_2 , and

(ii) $p_2(a_i) = p_2(a_i^\delta)$, $i = 0, \dots, m$. However, in the case of the minimal example in subsection 1.1, no such δ exists.

2 Linear sets of pseudoregulus type in a projective line

A family of scattered \mathbb{F}_q -linear sets of rank tm of $\text{PG}(2m-1, q^t)$, called of *pseudoregulus type*, have been introduced in [16] for $m = 2$ and $t = 3$, further generalized in [7] for $m \geq 2$ and $t = 3$ and finally in [14] for $m \geq 1$ and $t \geq 2$ (for $t = 2$ they are the same as Baer subgeometries, see [14, Remark 3.4]). We will only consider linear sets of pseudoregulus type in $\text{PG}(1, q^t)$. It has been proved in [14, Section 4] and in [3, Remark 2.2] that all linear sets of pseudoregulus type in $\text{PG}(1, q^t)$ are $\text{PGL}(2, q^t)$ -equivalent and hence we can define them as follows.

Definition 2.1 ([14, 3]). *A point set L of $\text{PG}_{q^t}(\mathbb{F}_{q^t}^2) = \text{PG}(1, q^t)$, $t \geq 2$, is called a linear set of pseudoregulus type if L is projectively equivalent to*

$$\mathbb{L} := \{ \langle (\lambda, \lambda^q) \rangle_{q^t} : \lambda \in \mathbb{F}_{q^t}^* \}. \quad (2)$$

Definition 2.2. *Let $t \geq 2$ be an integer and let π be a permutation of $N_{t-1} := \{0, 1, 2, \dots, t-1\}$. We define Σ_π as the following set of points in $\text{PG}_{q^t}(\mathbb{F}_{q^t}^t) = \text{PG}(t-1, q^t)$.*

$$\Sigma_\pi := \{ \langle (\alpha^{q^{\pi(0)}}, \alpha^{q^{\pi(1)}}, \alpha^{q^{\pi(2)}}, \dots, \alpha^{q^{\pi(t-1)}}) \rangle_{q^t} : \alpha \in \mathbb{F}_{q^t}^* \}. \quad (3)$$

Proposition 2.3. *Consider \mathbb{F}_{q^t} as a vector space over \mathbb{F}_q and let $\Sigma = \text{PG}_q(\mathbb{F}_{q^t}) \cong \text{PG}(t-1, q)$. Let π be a permutation of N_{t-1} . Then the following statements hold.*

1. $S_\pi := \{ (\alpha^{q^{\pi(0)}}, \alpha^{q^{\pi(1)}}, \alpha^{q^{\pi(2)}}, \dots, \alpha^{q^{\pi(t-1)}}) : \alpha \in \mathbb{F}_{q^t} \}$ is an \mathbb{F}_q -subspace of $\mathbb{F}_{q^t}^t$.
2. $\phi'_\pi : \mathbb{F}_{q^t} \rightarrow S_\pi : \alpha \mapsto (\alpha^{q^{\pi(0)}}, \alpha^{q^{\pi(1)}}, \alpha^{q^{\pi(2)}}, \dots, \alpha^{q^{\pi(t-1)}})$ is a vector space isomorphism.
3. $\langle \Sigma_\pi \rangle = \text{PG}(t-1, q^t)$.
4. The map

$$\phi_\pi : \Sigma \rightarrow \Sigma_\pi : \langle \alpha \rangle_q \mapsto \langle (\alpha^{q^{\pi(0)}}, \alpha^{q^{\pi(1)}}, \alpha^{q^{\pi(2)}}, \dots, \alpha^{q^{\pi(t-1)}}) \rangle_{q^t} \quad (4)$$

is a collineation.

Proof. 1. and 2. are trivial, and 1., 2., 3. together imply 4., so it is enough to show 3. Let $\alpha_1, \alpha_2, \dots, \alpha_t \in \mathbb{F}_{q^t}$ and denote by M_π the $t \times t$ matrix over \mathbb{F}_{q^t} whose j -th row is $\phi'_\pi(\alpha_j)$ for $j = 1, 2, \dots, t$. According to [12, Lemma 3.51], $\det(M_{id}) \neq 0$ if and only if $\alpha_1, \alpha_2, \dots, \alpha_t$ are linearly independent over \mathbb{F}_q . As M_π can be obtained by permuting the columns of M_{id} , it follows that $\det(M_\pi) \neq 0$ if and only if $\det(M_{id}) \neq 0$. As $\dim_{\mathbb{F}_q}(\mathbb{F}_{q^t}) = t$, it follows that $\dim_{\mathbb{F}_{q^t}}(S_\pi) = t$ and hence 3. follows. \square \square

Let $\Sigma^* = \text{PG}(t-1, q^t)$, and denote the coordinates in Σ^* by X_1, X_2, \dots, X_t . Let $\Lambda \cong \text{PG}(1, q^t)$ be the line $X_3 = \dots = X_t = 0$ and let $\Gamma \cong \text{PG}(t-3, q^t)$ be the $(t-3)$ -subspace $X_1 = X_2 = 0$ in Σ^* . Let π be a permutation of N_{t-1} , it is easy to see that if $\gcd(\pi(1) - \pi(0), t) = 1$, then $p_{\Gamma, \Lambda}(\Sigma_\pi) = \mathbb{L}$, see for example [14, Remark 4.2]. If ϕ is a collineation of Σ^* such that $\Sigma_{id}^\phi = \Sigma_\pi$, then

$$\tilde{\phi} := \phi_\pi \phi^{-1} \phi_{id}^{-1} \quad (5)$$

(see (4)) is a collineation of Σ . To study the action of $\tilde{\phi}$ we need some results regarding the Θ_s map defined in the next section.

3 The Θ_s map

As before, let Σ denote the $\text{PG}(t-1, q)$ associated to \mathbb{F}_{q^t} .

Definition 3.1. For an integer $s \geq -1$ let θ_s denote the number of points of $\text{PG}(s, q)$, that is

$$\theta_s = \frac{q^{s+1} - 1}{q - 1}. \quad (6)$$

The map $\Theta_s: \Sigma \rightarrow \Sigma$ is defined as follows. For each $x \in \mathbb{F}_{q^t}^*$ let

$$\Theta_s(\langle x \rangle_q) = \langle x^{\theta_s} \rangle_q.$$

Remark 3.2. If $s, d \geq -1$ are two integers, then $\Theta_s = \Theta_d$ if and only if $s \equiv d \pmod{t}$. As we do not use this result, the proof is left to the reader.

In Section 4 we show the following. If ϕ is a collineation such that $\Sigma_{id}^\phi = \Sigma_\pi$ and $\Gamma^\phi = \Gamma$, then $\tilde{\phi} = \Theta_s \Omega$ (see (5)) for some collineation Ω and integer $0 \leq s < t$. Note that Θ_0 is the identity. In Corollary 3.6, we prove that Θ_s , with $0 < s < t$, is not a collineation of $\text{PG}(t-1, q)$. It will follow that in most cases $\tilde{\phi}$ is not a collineation and hence such ϕ cannot exist.

To determine whether Θ_s is a collineation or not we need some results regarding difference sets. A (v, k, λ) -difference set is a k -subset D of a group

G of order v such that the list of non-zero differences (if the group is written additively) contains each non-zero group element exactly λ times. If the group is cyclic etc., then we say that the difference set is cyclic etc. For our purposes here, it is enough to consider cyclic difference sets, so from now on we assume that G is cyclic. Let D be a (v, k, λ) -difference set of G and let j be an integer. By $D + j$ and jD we mean $\{d + j : d \in D\}$ modulo v and $\{jd : d \in D\}$ modulo v . Note that $D + j$ is also a (v, k, λ) -difference set of G . If $\gcd(j, v) \neq 1$, then let $t := v / \gcd(j, v)$ and let $d_1 - d_2 = t$, where $d_1, d_2 \in D$. As $v \mid tj$, it follows that $jd_1 \equiv jd_2 \pmod{v}$ and hence $|jD| < |D|$. On the other hand, if $\gcd(j, v) = 1$, then jD is also a (v, k, λ) -difference set of G . The integer m is a (numerical) multiplier of D if $mD = D + g$ for some integer g . The trace of $\beta \in \mathbb{F}_{q^t}$ is $\text{Tr}(\beta) = \text{Tr}_{\mathbb{F}_{q^t}/\mathbb{F}_q}(\beta) = \sum_{i=0}^{t-1} \beta^{q^i}$.

Theorem 3.3 ([18, Theorem 2.1.1]). *Let α be a generator of the multiplicative group of \mathbb{F}_{q^t} , $t \geq 3$. The set of integers $D := \{i : 0 \leq i < \theta_{t-1}, \text{Tr}(\alpha^i) = 0\}$ modulo θ_{t-1} forms a (cyclic) difference set in $\mathbb{Z}_{\theta_{t-1}}$ (written additively), with parameters $(\theta_{t-1}, \theta_{t-2}, \theta_{t-3})$. The set of points of $\text{PG}_q(\mathbb{F}_{q^t}) \cong \text{PG}(t-1, q)$ which corresponds to the set D , i. e. the point set $\mathcal{H}_D := \{\langle \alpha^i \rangle_q : i \in D\}$, is a hyperplane of $\text{PG}(t-1, q)$. Moreover, the hyperplanes of $\text{PG}(t-1, q)$ are the sets $\mathcal{H}_{D+j} := \{\langle \alpha^i \rangle_q : i \in D + j\}$, where $0 \leq j < \theta_{t-1}$.*

The difference sets constructed in Theorem 3.3 are called *classical Singer difference sets*. The next theorem describes the multipliers of these difference sets.

Theorem 3.4 ([18, Corollary 1.3.4 and Proposition 3.1.1]). *Let D be a classical Singer difference set with parameters as in Theorem 3.3. Then the multipliers of D are the powers of p modulo θ_{t-1} .*

For a subset $\mathcal{X} \subseteq \Sigma$ and an integer m , let $\mathcal{X}^m = \{\langle x^m \rangle_q : \langle x \rangle_q \in \mathcal{X}\}$.

Corollary 3.5. *Let $t \geq 3$ and m be two integers. The $\Sigma \rightarrow \Sigma$ map, $\langle x \rangle_q \mapsto \langle x^m \rangle_q$ is a collineation if and only if $m \equiv p^h \pmod{\theta_{t-1}}$, for some $h \in \mathbb{N}$.*

Proof. The map $\langle x \rangle_q \mapsto \langle x^m \rangle_q$ is a collineation if and only if it maps hyperplanes into hyperplanes. Let α , D and \mathcal{H}_{D+j} for $0 \leq j < \theta_{t-1}$ be defined as in Theorem 3.3. For any hyperplane \mathcal{H}_{D+j} of Σ the point set $\mathcal{H}_{D+j}^m = \{\langle \alpha^{mi} \rangle_q : i \in D + j\}$ is a hyperplane if and only if $\{mi : i \in D + j\} = mD + mj$ is a translate of D , i.e. when $mD + mj = D + k$ for some k and hence $mD = D + g$ (with $g = k - mj$), which is equivalent to saying that m is a multiplier of D . The assertion follows from Theorem 3.4. \square \square

Corollary 3.6. *Let $t \geq 3$. For $0 < s < t$ the map Θ_s is not a collineation.*

Proof. According to Corollary 3.5, it is enough to show that for any $0 < s < t$ there is no integer h such that $\theta_s \equiv p^h \pmod{\theta_{t-1}}$. Note that $q^t = (q-1)\theta_{t-1} + 1$, thus $p^0 = 1 \equiv q^t \pmod{\theta_{t-1}}$. It is therefore enough to show the assertion when $p^h < q^t$. Suppose, contrary to our claim, that $\theta_s \equiv p^h \pmod{\theta_{t-1}}$ for some $p^h < q^t$ and $0 < s < t$. Since $p^h \not\equiv 0 \pmod{\theta_{t-1}}$, we have $s \leq t-2$ and

$$\theta_s q^t / p^h \equiv q^t \equiv 1 \pmod{\theta_{t-1}}. \quad (7)$$

It is easy to see that $p^h \neq 1$. If $1 < p^h < \theta_{t-1}$, then $p^h = \theta_s$, which cannot be as p does not divide θ_s . So we may assume $\theta_{t-1} < p^h < q^t$. In this case $p^h < q^t(p^h - \theta_{t-1})$, thus $q^t \theta_{t-1} < p^h(q^t - 1)$ and hence $q^t/p^h < (q^t - 1)/\theta_{t-1} = q - 1$. On the other hand we have $1 < q^t/p^h$ and hence

$$1 < \theta_s q^t / p^h < \theta_{t-2}(q-1) = q^{t-1} - 1 < \theta_{t-1},$$

contrary to (7). \square \square

Proposition 3.7. *For $0 \leq s < t$, the map $\langle x \rangle_q \mapsto \langle x^{-\theta_s} \rangle_q$ is the composition of Θ_{t-s-2} and a projectivity of Σ .*

Proof. For each $x \in \mathbb{F}_{q^t}^*$ we have $x^{\theta_{t-1}} \in \mathbb{F}_q^*$, thus

$$\langle x^{-\theta_s} \rangle_q = \langle x^{\theta_{t-1} - \theta_s} \rangle_q = \Theta_{t-s-2} \langle x^{q^{s+1}} \rangle_q.$$

As $\langle x \rangle_q \mapsto \langle x^{q^h} \rangle_q$ is a projectivity for each h , the assertion follows. \square \square

4 Linear sets obtained via non-equivalent projections

Definition 4.1. *For an integer h let Φ_h denote the collineation of $\Sigma^* = \text{PG}(t-1, q^t)$, defined as*

$$\langle (\alpha_0, \alpha_1, \dots, \alpha_{t-1}) \rangle_{q^t}^{\Phi_h} = \langle (\alpha_0^{p^h}, \alpha_1^{p^h}, \dots, \alpha_{t-1}^{p^h}) \rangle_{q^t}.$$

Proposition 4.2. *For each integer h and permutation π of N_{t-1} , Φ_h fixes Σ_π .*

Proof. For each $\alpha \in \mathbb{F}_{q^t}^*$ we have

$$\left\langle \left(\alpha^{q^{\pi(0)}}, \alpha^{q^{\pi(1)}}, \dots, \alpha^{q^{\pi(t-1)}} \right) \right\rangle_{q^t}^{\Phi_h} = \left\langle \left(\beta^{q^{\pi(0)}}, \beta^{q^{\pi(1)}}, \dots, \beta^{q^{\pi(t-1)}} \right) \right\rangle_{q^t},$$

where $\beta = \alpha^{p^h}$. \square \square

Proposition 4.3. *Let π be a permutation of N_{t-1} and let ω be the permutation defined such that $\omega(i) \equiv \pi(i) - \pi(0) \pmod{t}$ for each $i \in N_{t-1}$. Then $\Sigma_\pi = \Sigma_\omega$.*

Proof. Let $h = -\pi(0)e$, where $q = p^e$, p prime. It is easy to see that $\Phi_h(\Sigma_\pi) = \Sigma_\omega$. As Φ_h fixes Σ_π the assertion follows. \square \square

Lemma 4.4. *Let $\Gamma \cong \text{PG}(t-3, q^t)$ be the $(t-3)$ -space $X_1 = X_2 = 0$ in $\Sigma^* = \text{PG}(t-1, q^t)$, $t \geq 3$. If ϕ is a projectivity of Σ^* and π is a permutation of N_{t-1} such that $\Sigma_{id}^\phi = \Sigma_\pi$ and $\Gamma^\phi = \Gamma$, then $\pi(1) - \pi(0) \equiv \pm 1 \pmod{t}$, or $\gcd(\pi(1) - \pi(0), t) > 1$.*

Proof. Suppose $\gcd(\pi(1) - \pi(0), t) = 1$. Let $M = (m_{ij})_{1 \leq i, j \leq t} \in \text{GL}(t, q^t)$ be a matrix associated with ϕ . As ϕ fixes Γ , we have $m_{ij} = 0$ when $i = 1, 2$ and $3 \leq j \leq t$. Denote the 2×2 matrix $(m_{ij})_{1 \leq i, j \leq 2}$ by A . As M is non-singular, the same holds for A . According to Proposition 4.3 we may assume $\pi(0) = 0$. Let $\mu = \pi(1)$, whence $\gcd(\mu, t) = 1$. As $\Sigma_{id}^\phi = \Sigma_\pi$, for each $\alpha \in \mathbb{F}_{q^t}^*$ there exist $\delta_\alpha, t_\alpha \in \mathbb{F}_{q^t}^*$ such that

$$\delta_\alpha t_\alpha = m_{11}\alpha + m_{12}\alpha^q, \quad (8)$$

$$\delta_\alpha t_\alpha^{q^\mu} = m_{21}\alpha + m_{22}\alpha^q. \quad (9)$$

Let N denote the norm function from \mathbb{F}_{q^t} to \mathbb{F}_q , that is $N(x) = x^{\theta_{t-1}}$. As $N(\delta_\alpha t_\alpha^{q^\mu}) = N(\delta_\alpha t_\alpha)N(t_\alpha^{q^\mu - 1}) = N(\delta_\alpha t_\alpha)$, it follows that

$$N(m_{11}\alpha + m_{12}\alpha^q) = N(m_{21}\alpha + m_{22}\alpha^q),$$

$$N(m_{11} + m_{12}\alpha^{q-1}) = N(m_{21} + m_{22}\alpha^{q-1}),$$

and hence for each $(q-1)$ -th power z of $\mathbb{F}_{q^t}^*$ we have

$$P(z) := N(m_{11} + m_{12}z) - N(m_{21} + m_{22}z) = 0,$$

where $P(z)$ is a polynomial of degree at most θ_{t-1} .

We claim $m_{11}m_{21} = 0$ and $m_{12}m_{22} = 0$. First suppose $m_{11}m_{21} \neq 0$. Then the coefficient of z^q in $P(z)$ is

$$m_{12}^q N(m_{11})/m_{11}^q - m_{22}^q N(m_{21})/m_{21}^q. \quad (10)$$

The coefficient of z^{q+1} is

$$m_{12}^{q+1} N(m_{11})/m_{11}^{q+1} - m_{22}^{q+1} N(m_{21})/m_{21}^{q+1}. \quad (11)$$

As $P(z)$ vanishes for each z with $z^{\theta t-1} = 1$, we have $P(z) = a(z^{\theta t-1} - 1)$ for some $a \in \mathbb{F}_{q^t}$. It follows that (10) and (11) are zero, thus if $m_{12}m_{22} \neq 0$, then

$$m_{12}/m_{11} = m_{22}/m_{21},$$

and hence $\det(A) = 0$, a contradiction. On the other hand if one of $\{m_{12}, m_{22}\}$ is zero, then both of them are zero, thus we obtain again $\det(A) = 0$. Now suppose $m_{12}m_{22} \neq 0$. Then the coefficient of $z^{\theta t-1-q}$ in $P(z)$ is

$$m_{11}^q N(m_{12})/m_{12}^q - m_{21}^q N(m_{22})/m_{22}^q. \quad (12)$$

As before, it follows that (12) is zero. Together with $m_{11}m_{21} = 0$, it means that m_{11} and m_{21} are both zero, and hence $\det(A) = 0$, a contradiction.

First we consider the case when A is diagonal, i.e. $m_{12} = m_{21} = 0$. We may assume $m_{11} = 1$. Then (8) and (9) yield

$$t_\alpha^{q^\mu-1} = m_{22}\alpha^{q-1}.$$

The last equation implies that a $\rho \in \mathbb{F}_{q^n}^*$ exists such that $\rho^{q-1} = m_{22}$. Then for any $x \in \mathbb{F}_{q^t}^*$ the following holds:

$$\begin{aligned} \left\langle \frac{x^{\theta_{\mu-1}}}{\rho} \right\rangle_q^{\phi_{id}\phi} &= \left\langle \left(\frac{x^{\theta_{\mu-1}}}{\rho}, \frac{x^{q\theta_{\mu-1}}}{\rho^q}, \dots, \frac{x^{q^{t-1}\theta_{\mu-1}}}{\rho^{q^{t-1}}} \right) \right\rangle_{q^t}^\phi = \\ &= \left\langle \left(\frac{x^{\theta_{\mu-1}}}{\rho}, \frac{x^{q\theta_{\mu-1}}}{\rho}, * \right) \right\rangle_{q^t} = \langle (1, x^{q^\mu-1}, *) \rangle_{q^t} \in \Sigma_\pi; \\ \langle x \rangle_q^{\phi_\pi} &= \langle (x, x^{q^\mu}, *) \rangle_{q^t} = \langle (1, x^{q^\mu-1}, *) \rangle_{q^t} \in \Sigma_\pi. \end{aligned}$$

Since μ and t are coprime, Σ_π contains exactly one element of the form $\langle (1, x^{q^\mu-1}, *) \rangle_{q^t}$. It follows that for any $x \in \mathbb{F}_{q^t}^*$

$$\langle x^{\theta_{\mu-1}}/\rho \rangle_q^{\phi_{id}\phi} = \langle x \rangle_q^{\phi_\pi},$$

and hence $\phi_\pi \phi^{-1} \phi_{id}^{-1}: \Sigma \rightarrow \Sigma$ which maps $\langle x \rangle_q$ to $\langle x^{\theta_{\mu-1}}/\rho \rangle_q$ is a collineation. Since the map $\langle y \rangle_q \mapsto \langle y/\rho \rangle_q$ is a collineation of Σ , Corollary 3.6 yields that $\phi_\pi \phi^{-1} \phi_{id}^{-1}$ is a collineation only if $\mu = 1$.

Now consider the case when $m_{11} = m_{22} = 0$. We may assume $m_{12} = 1$. Then (8) and (9) yield

$$t_\alpha^{q^\mu-1} = m_{21}\alpha^{1-q}.$$

This implies that a $\rho \in \mathbb{F}_{q^t}^*$ exists such that $\rho^{q-1} = m_{21}$. As before, it follows that

$$\langle x^{-\theta_{\mu-1}}\rho \rangle_q^{\phi_{id}\phi} = \langle x \rangle_q^{\phi_\pi},$$

for each $x \in \mathbb{F}_{q^t}^*$ and hence $\phi_\pi \phi^{-1} \phi_{id}^{-1} : \Sigma \rightarrow \Sigma$ which maps $\langle x \rangle_q$ to $\langle x^{-\theta_{\mu-1}} \rho \rangle_q$ is a collineation. Proposition 3.7 and Corollary 3.6 yield that this happens only if $\mu = t - 1$. \square \square

Theorem 4.5. *Let $\Gamma \cong \text{PG}(t - 3, q^t)$ be the $(t - 3)$ -space $X_1 = X_2 = 0$ in $\Sigma^* = \text{PG}(t - 1, q^t)$, $t \geq 3$. If ϕ is a collineation of Σ^* such that $\Sigma_{id}^\phi = \Sigma_\pi$ and $\Gamma^\phi = \Gamma$, then $\pi(1) - \pi(0) \equiv \pm 1 \pmod{t}$, or $\gcd(\pi(1) - \pi(0), t) > 1$.*

Proof. According to the Fundamental Theorem of Projective Geometry we may assume $\phi = \Phi_h \Omega$, where Ω is a projectivity and Φ_h is as in Definition 4.1. As Φ_h fixes Γ and Σ_{id} , it follows that Ω satisfies the conditions of Lemma 4.4 and hence the assertion follows. \square \square

Corollary 4.6. *If $t = 5$ or $t > 6$, then any linear set of pseudoregulus type in $\text{PG}(1, q^t)$ can be obtained as the projection of two different subgeometries, $\Sigma_1 \cong \Sigma_2 \cong \text{PG}(t - 1, q)$, from a center $\Gamma \cong \text{PG}(t - 3, q^t)$ to an axis $\Lambda \cong \text{PG}(1, q^t)$ in the ambient space $\Sigma^* = \text{PG}(t - 1, q^t)$, such that there exists no collineation ϕ of Σ^* with $\Gamma^\phi = \Gamma$ and $\Sigma_1^\phi = \Sigma_2$.*

Proof. Let Λ be the line $X_3 = \dots = X_t = 0$ and let Γ be the $(t - 3)$ -subspace $X_1 = X_2 = 0$ in Σ^* . As $t = 5$ or $t > 6$, we have $\varphi(t) \geq 3$, where φ is the Euler function. Thus we may choose a permutation π of N_{t-1} such that $\gcd(\pi(1) - \pi(0), t) = 1$ and $\pi(1) - \pi(0) \notin \{-1, 1\}$. Let $\Sigma_1 = \Sigma_{id}$ and $\Sigma_2 = \Sigma_\pi$. Then $p_{\Gamma, \Lambda}(\Sigma_1) = p_{\Gamma, \Lambda}(\Sigma_2) \cong \mathbb{L}$ and the assertion follows from Theorem 4.5. \square \square

Note that \mathbb{L} is not a linear set of rank $k < t$, thus (i) \Rightarrow (ii) in Theorem 1.3 cannot hold without further conditions.

5 Equivalence of linear sets

We say that the pair (L, n) , where L is a linear set of rank n in $\text{PG}(r - 1, q^t)$, satisfies condition (A), if the following holds.

- (A) For any two $(n - 1)$ -subspaces $U, U' \subset \text{PG}(rt - 1, q)$, such that $\mathcal{B}(U) = L = \mathcal{B}(U')$, a collineation $\gamma \in \text{P}\Gamma\text{L}(rt, q)$ exists, such that $U^\gamma = U'$, γ preserves the Desarguesian spread $\mathcal{F}_{r,t,q}(\mathcal{P})$, and the induced map on $\text{PG}(r - 1, q^t)$ is a collineation (which will again be denoted by γ).

In the next theorem we follow the proof of [2, Theorem 2.4] by Bonoli and Polverino.

Theorem 5.1. For $i = 1, 2$, let $\Sigma_i \cong \text{PG}(n-1, q)$ be two canonical subgeometries of $\Sigma^* = \text{PG}(n-1, q^t)$ and let Γ_i be two $(n-1-r)$ -subspaces of Σ^* , such that $\Gamma_i \cap \Sigma_i = \emptyset$. Let $\Lambda_i \subset \Sigma^* \setminus \Gamma_i$ be two $(r-1)$ -subspaces and let $L_i = p_{\Gamma_i, \Lambda_i}(\Sigma_i)$. Suppose that (L_2, n) satisfies condition (A). Then $(i) \Rightarrow (ii)$, where (i) and (ii) are the conditions stated in Theorem 1.3.

Proof. The collineation α in Theorem 1.3 can be extended to an element $\hat{\alpha} \in \text{PGL}(n, q^t)$ such that $\Gamma_1^{\hat{\alpha}} = \Gamma_2$. Then $L_2 = L_1^{\hat{\alpha}} = p_{\Gamma_2, \Lambda_2}(\Sigma_1^{\hat{\alpha}})$. Hence it is sufficient to prove the existence of a collineation $\phi \in \text{PGL}(n, q^t)$ such that $\Gamma_2^\phi = \Gamma_2$ and $\Sigma_1^{\hat{\alpha}\phi} = \Sigma_2$. If such ϕ exists, then $\beta := \hat{\alpha}\phi$.

So it is enough to prove the assertion when $\Gamma_1 = \Gamma_2 =: \Gamma$, $\Lambda_1 = \Lambda_2 =: \Lambda$ and $L_1 = L_2 =: L$. Let $\Sigma^* = \text{PG}_{q^t}(\mathbb{F}_{q^t}^n)$ and for $j = 1, 2$, let $\Sigma_j = \mathcal{B}(V_j)$, where V_j is an n -dimensional \mathbb{F}_q -subspace of $\mathbb{F}_{q^t}^n$. As Σ_j is a canonical subgeometry, we have $\langle V_j \rangle_{q^t} = \mathbb{F}_{q^t}^n$, for $j = 1, 2$. Let $V_1 = \langle v_1, v_2, \dots, v_n \rangle_q$. Also let $\Gamma = \text{PG}_{q^t}(Z)$ and $\Lambda = \text{PG}_{q^t}(R)$. Note that $V_j' := (V_j + Z) \cap R$ is an n -dimensional \mathbb{F}_q -vector space in R . As $p_{\Gamma, \Lambda}(\Sigma_1) = p_{\Gamma, \Lambda}(\Sigma_2) = L$, it follows that $\mathcal{B}(V_1') = \mathcal{B}(V_2')$. As (L, n) satisfies condition (A), it follows that there exists a non-singular \mathbb{F}_{q^t} -semilinear map $\gamma': R \rightarrow R$ such that $V_1'^{\gamma'} = V_2'$. The map γ' can be extended to a non-singular semilinear map $\hat{\gamma}: \mathbb{F}_{q^t}^n \rightarrow \mathbb{F}_{q^t}^n$ such that $Z^{\hat{\gamma}} = Z$. Then for each vector $v \in \mathbb{F}_{q^t}^n$ and $\mu \in \mathbb{F}_{q^t}$ we have $(\mu v)^{\hat{\gamma}} = \mu^{p^k} v^{\hat{\gamma}}$ for some integer k . Since all of $R + Z = \mathbb{F}_{q^t}^n$ and $V_j + Z$ ($j = 1, 2$) are direct sums, for any $v \in V_j \setminus \{0\}$ and $z \in Z$, the intersection $(\langle v \rangle_q + Z) \cap R$ is a one-dimensional \mathbb{F}_q -subspace, say $\langle v' \rangle_q$, with $v' \in V_j'$; this implies $hv' = v + z'$ for some $z' \in Z$ and $h \in \mathbb{F}_q$, whence $v + z = hv' + (z - z') \in V_j' + Z$. This implies $V_j + Z \subseteq V_j' + Z$ and therefore $V_j + Z = V_j' + Z$. Then we have

$$V_2 + Z = V_2' + Z = V_1'^{\hat{\gamma}} + Z^{\hat{\gamma}} = (V_1' + Z)^{\hat{\gamma}} = (V_1 + Z)^{\hat{\gamma}},$$

and hence for each $i = 1, 2, \dots, n$, $v_i^{\hat{\gamma}} = v_i' + z_i$ for some $v_i' \in V_2$ and $z_i \in Z$. We claim that v_1', v_2', \dots, v_n' is an \mathbb{F}_q -basis of V_2 . To see this, suppose $\sum_{i=1}^n \lambda_i v_i' = 0$ with $\lambda_i \in \mathbb{F}_q$ for $i = 1, 2, \dots, n$. Then $\sum_{i=1}^n \lambda_i v_i^{\hat{\gamma}} = \sum_{i=1}^n \lambda_i z_i$. As on the left-hand side $v_1^{\hat{\gamma}}, v_2^{\hat{\gamma}}, \dots, v_n^{\hat{\gamma}}$ are \mathbb{F}_q -independent, it follows that either $\lambda_i = 0$ for $i = 1, 2, \dots, n$, or there is a non-zero vector $z \in V_1^{\hat{\gamma}} \cap Z$. The map $\hat{\gamma}$ fixes Z and hence in the latter case $z^{\hat{\gamma}^{-1}} \in V_1 \cap Z$, a contradiction because of $V_1 \cap Z = \{0\}$. As Σ_2 is a canonical subgeometry, it follows that v_1', v_2', \dots, v_n' are linearly independent over \mathbb{F}_q . Let f be the \mathbb{F}_{q^t} -semilinear map of $\mathbb{F}_{q^t}^n$ such that $v_i^f = v_i'$ for each $i = 1, 2, \dots, n$ and $(\mu v)^f = \mu^{p^k} v^f$ for each $\mu \in \mathbb{F}_{q^t}$ and $v \in \mathbb{F}_{q^t}^n$. If $P = \langle z \rangle_{q^t} \in \Gamma$, then we have $z = \sum_{i=1}^n a_i v_i$

for some $a_i \in \mathbb{F}_{q^t}$. Then $z^f = \sum_{i=1}^n a_i^{p^k} v'_i = \sum_{i=1}^n a_i^{p^k} (v_i^{\widehat{\gamma}} - z_i) = z^{\widehat{\gamma}} - \sum_{i=1}^n a_i^{p^k} z_i \in Z$, and hence the collineation induced by f fixes Γ and maps Σ_1 to Σ_2 . \square \square

Remark 5.2. *In Theorem 5.1, it is not assumed, contrary to Theorem 1.3, that the linear sets are not of rank less than n .*

Remark 5.3. *By [2, Proposition 2.3], condition (A) holds for any \mathbb{F}_q -linear blocking set of exponent e (where $p^e = q$) in $\text{PG}(2, q^t)$.*

Remark 5.4. *Up to the knowledge of the authors, no linear set is known which is not of pseudoregulus type in $\text{PG}(1, q^t)$ and does not satisfy condition (A).*

Theorem 5.5. *If L is a linear set of rank n in $\Lambda = \text{PG}(r-1, q^t)$, $\langle L \rangle = \text{PG}(r-1, q^t)$ and (L, n) does not satisfy condition (A), then in $\text{PG}(n-1, q^t) \supset \Lambda$ there are a subspace $\Gamma = \Gamma_1 = \Gamma_2$ disjoint from Λ , and two q -order canonical subgeometries $\Sigma_1, \Sigma_2 \subset \text{PG}(n-1, q^t) \setminus \Gamma$ such that $L = p_{\Gamma, \Lambda}(\Sigma_1) = p_{\Gamma, \Lambda}(\Sigma_2)$, and such that condition (ii) in Theorem 1.3 does not hold.*

Proof. Let U_1 and U_2 be two $(n-1)$ -subspaces of $\text{PG}_q(R)$, where R is the r -dimensional \mathbb{F}_{q^t} -subspace of $\mathbb{F}_{q^t}^n$ such that $\Lambda = \text{PG}_{q^t}(R)$. Assume that $\mathcal{B}(U_1) = L = \mathcal{B}(U_2)$. Let $W_i = \langle w_1^{(i)}, w_2^{(i)}, \dots, w_n^{(i)} \rangle_q$, $i = 1, 2$, be the n -dimensional vector \mathbb{F}_q -subspaces of R whose associated projective subspaces in $\text{PG}(rt-1, q)$ are U_1 and U_2 , respectively. From $\langle L \rangle = \text{PG}(r-1, q^t)$ we may assume that $w_1^{(1)}, w_2^{(1)}, \dots, w_r^{(1)}$ are \mathbb{F}_{q^t} -linearly independent, and also that $w_1^{(2)}, w_2^{(2)}, \dots, w_r^{(2)}$ are \mathbb{F}_{q^t} -linearly independent. Let Γ be an $(n-r-1)$ -subspace in $\text{PG}(n-1, q^t)$ disjoint from Λ , associated with the vector subspace $Z = \langle z_1, z_2, \dots, z_{n-r} \rangle_{q^t}$. For $i = 1, 2$, let Σ_i be the \mathbb{F}_q -linear set defined by the following \mathbb{F}_q -subspace of $\mathbb{F}_{q^t}^n$:

$$V_i := \langle w_1^{(i)}, w_2^{(i)}, \dots, w_r^{(i)}, w_{r+1}^{(i)} + z_1, w_{r+2}^{(i)} + z_2, \dots, w_n^{(i)} + z_{n-r} \rangle_q.$$

Since those n vectors of $\mathbb{F}_{q^t}^n$ are also \mathbb{F}_{q^t} -linearly independent, Σ_i is a canonical q -order subgeometry. Furthermore, $w_1^{(i)}, w_2^{(i)}, \dots, w_n^{(i)}, z_1, z_2, \dots, z_{n-r}$ are \mathbb{F}_q -linearly independent, and this implies $\Sigma_i \cap \Gamma = \emptyset$. We summarize here some properties of our construction, which we will use later. For $i = 1, 2$ we have the following.

1. For each $w \in W_i$, there exists a unique $z_w \in Z$ such that $w + z_w \in V_i$,

2. if w and w' are two \mathbb{F}_q -independent vectors in W_i , then $w + z_w$ and $w' + z_{w'}$ are \mathbb{F}_{q^t} -independent vectors in V_i .

If $w = \sum_{j=1}^n \alpha_j w_j^{(i)}$ with $\alpha_j \in \mathbb{F}_q$, $j = 1, 2, \dots, n$, then let $z_w = \sum_{j=r+1}^n \alpha_j z_{j-r}$. The unicity of z_w follows from $\Sigma_i \cap \Gamma = \emptyset$. To see 2., note that $\Psi_i: U_i \rightarrow \Sigma_i: \langle w \rangle_q \mapsto \langle w + z_w \rangle_{q^t}$ is a bijection, as $|U_i| = |\Sigma_i| = \theta_{n-1}$.

Now assume that condition (ii) in Theorem 1.3 holds. Let $\beta' : \mathbb{F}_{q^t}^n \rightarrow \mathbb{F}_{q^t}^n$ be the semilinear map associated with the collineation β ; so, an integer ν exists such that for any $a \in \mathbb{F}_{q^t}$ and $v \in \mathbb{F}_{q^t}^n$ there holds $(av)^{\beta'} = a^{p^\nu} v^{\beta'}$. Denote by π the canonical projection from $Z \oplus R$ to R . Fix an element $w \in W_1$. Then 1. and the condition $\Sigma_1^\beta = \Sigma_2$ imply the existence of $z_w \in Z$ and $\lambda \in \mathbb{F}_{q^t}^*$ depending on z_w such that $\xi_\lambda := \lambda(w + z_w)^{\beta'} \in V_2$. Let $T_\lambda = \{v \in W_1 : \lambda v^{\beta'\pi} \in W_2\}$. Note that T_λ is an \mathbb{F}_q -vector subspace of R . As $\lambda(w + z_w)^{\beta'} \in V_2$, we have $\lambda(w + z_w)^{\beta'\pi} = \lambda w^{\beta'\pi} \in V_2^\pi = W_2$, thus $w \in T_\lambda$. We are going to show that $W_1 = T_\lambda$. Suppose to the contrary that there exists $w' \in W_1 \setminus T_\lambda$. As T_λ is an \mathbb{F}_q -vector space, w and w' are \mathbb{F}_q -independent. The same argument as above yields the existence of $z_{w'} \in Z$ and $\mu \in \mathbb{F}_{q^t}^*$ such that $\xi_\mu := \mu(w' + z_{w'})^{\beta'} \in V_2$ and hence $\mu w'^{\beta'\pi} \in W_2$. Then 2. yields that $\langle w + z_w \rangle_{q^t}$ and $\langle w' + z_{w'} \rangle_{q^t}$ are distinct points of Σ_1 , and hence $\langle \xi_\lambda \rangle_{q^t}$ and $\langle \xi_\mu \rangle_{q^t}$ are distinct points of Σ_2 . First we prove

$$\langle \xi_\lambda, \xi_\mu \rangle_{q^t} \cap V_2 = \langle \xi_\lambda, \xi_\mu \rangle_q. \quad (13)$$

Denote by X de left hand side of (13). The inclusion \supseteq trivially holds. Suppose to the contrary $\dim_{\mathbb{F}_q}(X) \geq 3$, then there exist three \mathbb{F}_q -independent vectors, $v_1, v_2, v_3 \in X$. We can extend the set $\{v_1, v_2, v_3\}$ to a basis of V_2 . As $\dim_{\mathbb{F}_{q^t}}(\langle v_1, v_2, v_3 \rangle_{q^t}) = 2$, it follows that $\langle V_2 \rangle_{q^t} \neq \mathbb{F}_{q^t}^n$, a contradiction since Σ_2 is a canonical subgeometry.

Let $P_1 = \langle w + z_w \rangle_{q^t}$ and $P_2 = \langle w' + z_{w'} \rangle_{q^t}$. Denote by ℓ the q -order subline $\langle P_1, P_2 \rangle \cap \Sigma_1$. Then $\langle (w + z_w)^{\beta'} + (w' + z_{w'})^{\beta'} \rangle_{q^t} = \langle \xi_\lambda + \lambda\mu^{-1}\xi_\mu \rangle_{q^t} \in \ell^\beta \cap \Sigma_2$. It follows that there exists some $\delta \in \mathbb{F}_{q^t}^*$ such that $\delta(\xi_\lambda + \lambda\mu^{-1}\xi_\mu) \in V_2$. It follows from (13) that $\delta \in \mathbb{F}_q^*$ and $\delta\lambda\mu^{-1} \in \mathbb{F}_q^*$. Hence $\lambda\mu^{-1} \in \mathbb{F}_q^*$. This is a contradiction, as in this case $(\lambda\mu^{-1})\mu w'^{\beta'\pi} = \lambda w'^{\beta'\pi} \in W_2$, contradicting the choice of w' . Hence $T_\lambda = W_1$, that is, $\lambda w^{\beta'\pi} \in W_2$ for any $w \in W_1$.

Now define the map $\varphi' : R \rightarrow R: v \mapsto \lambda v^{\beta'\pi}$. As β' is semilinear, π is linear and $v \mapsto \lambda v$ is also linear, it follows that φ' is \mathbb{F}_{q^t} -semilinear. Also, φ' is non-singular. Let φ be the associated collineation of Λ . Then

$$U_1^\varphi = \{\langle \lambda w^{\beta'\pi} \rangle_q : w \in W_1^*\} = \text{PG}_q(W_2) = U_2,$$

because of $T_\lambda = W_1$.

We have constructed an example for which, if condition (ii) in Theorem 1.3 is satisfied, then also condition (A) must be satisfied: this concludes the proof. \square \square

Scattered linear sets of pseudoregulus type in a line are, under the field reduction map, embeddings of minimum dimension of $\text{PG}(t-1, q) \times \text{PG}(t-1, q)$, which have been studied in [5]. These embeddings are projections of Segre varieties. A similar representation of the clubs as projections of Segre varieties can be found in [10] (see also [9]). By [5, Theorem 16], (\mathbb{L}, t) does not fulfill condition (A), for $t = 5$ or $t > 6$, where \mathbb{L} is the scattered linear set of pseudoregulus type in $\text{PG}(1, q^t)$. By Theorem 5.5 this yields the counterexamples as shown in the previous sections. Many results in [6] are on linear sets in $\text{PG}(1, q^3)$; hence, the following result can be of interest:

Proposition 5.6. *Condition (A) is fulfilled by linear sets of rank three in $\text{PG}(1, q^3)$, $q > 2$, for $n = 3$.*

Proof. A linear set L of rank three in $\text{PG}(1, q^3)$ is either a point, or a club, or a scattered linear set.

If L is a point, then from

$$\mathcal{B}(U) = \mathcal{B}(U'), \quad \dim(U) = \dim(U') = 2, \quad (14)$$

one obtains $U = U' = \mathcal{F}_{2,3,q}(L)$. This implies (A).

Assume now that L is a club. Denote by H its head, which is uniquely defined [4, 6]. Assume that the subspaces U and U' satisfy (14). Take a point $P = \langle (v_0, v_1) \rangle_{q^3} \in L \setminus H$ and let $A = \mathcal{F}_{2,3,q}(P) \cap U$ and $X = \mathcal{F}_{2,3,q}(P) \cap U'$. Then $A = \langle \alpha(v_0, v_1) \rangle_q$ and $X = \langle \beta(v_0, v_1) \rangle_q$ for some $\alpha, \beta \in \mathbb{F}_{q^3}^*$, and the map $\gamma: \langle (x_0, x_1) \rangle_q \mapsto \langle \beta\alpha^{-1}(x_0, x_1) \rangle_q$ is a projectivity of $\text{PG}(5, q)$ belonging to the elementwise stabilizer of the Desarguesian spread, and mapping A into X . Then $X \in U^\gamma \cap U' \setminus \mathcal{F}_{2,3,q}(H)$. The element of $\text{PGL}(2, q^3)$ associated with γ is the identity map. Denote by ℓ_1 and ℓ_2 the lines $U^\gamma \cap \mathcal{F}_{2,3,q}(H)$ and $U' \cap \mathcal{F}_{2,3,q}(H)$, respectively. If $\ell_1 = \ell_2$, then $U^\gamma = \langle \ell_1, X \rangle = U'$. Otherwise let ℓ' be a line in U' through X such that $\ell_1 \cap \ell_2 \notin \ell'$. The club L does not contain irregular sublines [6, Corollary 13], [4, Proposition 5]. Applying this result to the q -order subline $\mathcal{B}(\ell')$, we have that a line ℓ in U^γ exists such that $\mathcal{B}(\ell) = \mathcal{B}(\ell')$. The point X is on ℓ . The lines ℓ, ℓ' of $\text{PG}(5, q)$ are transversal lines to the regulus $\mathcal{F}_{2,3,q}(\mathcal{B}(\ell))$, having a common point, whence $\ell = \ell'$. This implies $U^\gamma = \langle \ell, \ell_1 \cap \ell_2 \rangle = U'$ and (A).

Up to projectivities there is a unique scattered linear set in $\text{PG}(1, q^3)$. To see this, fix a q -order canonical subgeometry Σ of $\Sigma^* = \text{PG}(2, q^3)$ and

denote by \mathcal{I} the set of points of Σ^* not contained in a line of Σ . Denote by G the stabilizer of Σ in the group of projectivities of Σ^* . Each scattered linear set of rank three of $\text{PG}(1, q^3)$ can be obtained as a projection of Σ from a point in \mathcal{I} . In Theorem 1.3 the implication $(ii) \Rightarrow (i)$ holds and hence the number of projectively non-equivalent scattered linear sets of rank three in $\text{PG}(1, q^3)$ is at most the number of orbits of G on the points of \mathcal{I} . It follows from [2, Proposition 3.1] that G is transitive on \mathcal{I} and hence there is a unique scattered linear set of rank three in $\text{PG}(1, q^3)$. An alternate proof can be obtained from [1], where the uniqueness of an exterior splash is proved, taking into account that the exterior splashes dealt with in that paper are precisely the scattered linear sets of rank three [11].

Assume that L is a scattered linear set of rank three, hence a linear set of pseudoregulus type, and that (14) holds. Then U and U' are planes contained in the hypersurface $\mathcal{Q}_{2,q}$ defined in [5, eqs. (7),(8)]. By [5, Corollary 10], $U = S_{h,k}$, $U' = S_{h',k'}$ for $h, h' \in \{1, 2\}$, $N(k) = N(k') = 1$. A collineation γ satisfying (A) can be obtained in the form $\varphi_{0,0}(a, b)$ if $h = h'$, and $\psi_{0,0}(a, b)$ if $h \neq h'$, for suitable $a, b \in \mathbb{F}_{q^3}$. \square \square

References

- [1] Barwick S.G., Jackson W.A., Exterior splashes and linear sets of rank 3, <<http://arxiv.org/abs/1404.1641v1>>
- [2] Bonoli G., Polverino O., \mathbb{F}_q -linear blocking sets in $\text{PG}(2, q^4)$, Innov. Incidence Geom. 2, 35–56 (2005)
- [3] Donati G., Durante N., Scattered linear sets generated by collineations between pencils of lines, J. Algebraic Combin. 40, 1121–1134 (2014)
- [4] Fancsali Sz.L., Sziklai P., Description of the clubs, Ann. Univ. Sci. Budapest. Eötvös Sect. Math. 51, 141–146 (2008)
- [5] Lavrauw M., Sheekey J., Zanella C., On embeddings of minimum dimension of $\text{PG}(n, q) \times \text{PG}(n, q)$, Des. Codes Cryptogr. 74, 427–440 (2015)
- [6] Lavrauw M., Van de Voorde G., On linear sets on a projective line, Des. Codes Cryptogr. 56, 89–104 (2010)
- [7] Lavrauw M., Van de Voorde G., Scattered linear sets and pseudoreguli, Electron. J. Combin. 20(1), #P15 (2013)

- [8] Lavrauw M., Van de Voorde G., Field reduction and linear sets on a projective line, *Topics in finite fields*, 271–293, *Contemp. Math.*, 632, Amer. Math. Soc., Providence, RI, 2015
- [9] Lavrauw M., Zanella C., Segre embeddings and finite semifields, *Finite Fields Appl.* 25, 8–18 (2014)
- [10] Lavrauw M., Zanella C., Subspaces intersecting each element of a regulus in one point, André-Bruck-Bose representation and clubs, <<http://arxiv.org/abs/1409.1081>>
- [11] Lavrauw M., Zanella C., Subgeometries and linear sets on a projective line, *Finite Fields Appl.* 34, 95–106 (2015)
- [12] Lidl R., Niederreiter H., *Finite Fields*, second edition, Cambridge University Press (1997)
- [13] Lunardon G., Normal spreads, *Geom. Dedicata* 75, 245–261 (1999)
- [14] Lunardon G., Marino G., Polverino O., Trombetti R., Maximum scattered linear sets of pseudoregulus type and the Segre Variety $\mathcal{S}_{n,n}$, *J. Algebraic Combin.* 39, 807–831 (2014)
- [15] Lunardon G., Polverino O., Translation ovoids of orthogonal polar spaces, *Forum Math.* 16, 663–669 (2004)
- [16] Marino G., Polverino O., Trombetti R., On \mathbb{F}_q -linear sets of $\text{PG}(3, q^3)$ and semifields, *J. Combin. Theory Ser. A* 114, 769–788 (2007)
- [17] Polverino O., Linear sets in finite projective spaces, *Discrete Math.* 310, 3096–3107 (2010)
- [18] Pott A., *Finite Geometry and Character Theory*, Springer-Verlag, Berlin Heidelberg (1995)

Bence Csajbók

Dipartimento di Tecnica e Gestione dei Sistemi Industriali,
 Università di Padova, Stradella S. Nicola, 3, I-36100 Vicenza, Italy
 e-mail: csajbok.bence@gmail.com

Corrado Zanella

Dipartimento di Tecnica e Gestione dei Sistemi Industriali,
 Università di Padova, Stradella S. Nicola, 3, I-36100 Vicenza, Italy
 e-mail: corrado.zanella@unipd.it