

ON THE EXISTENCE OF WEIERSTRASS POINTS WITH A CERTAIN SEMIGROUP GENERATED BY 4 ELEMENTS

By

Jiryo KOMEDA

Introduction

Let X be a smooth, proper 1-dimensional algebraic variety (of genus ≥ 2) over an algebraically closed field k of characteristic 0, and let P be a point of X . Then a positive integer ν is called a *gap* at P if $h^0(X, \mathcal{O}_X((\nu-1)P)) = h^0(X, \mathcal{O}_X(\nu P))$, and G_P denotes the set of gaps at P . If we denote by N and H_P respectively the additive semigroup of non-negative integers and the complement of G_P in N , then H_P is a semigroup. A subsemigroup H of N whose complement is finite is called a *numerical semigroup*. The following problem is fundamental and is a long-standing problem.

Is there a pair (X, P) with X a smooth, proper 1-dimensional algebraic variety over k and P its point, such that $H = H_P$?

Using the deformation theory on algebraic varieties with G_m -action, Pinkham [7] constructed a moduli space \mathcal{M}_H which classifies the set of isomorphic classes of pairs (X, P) consisting of a smooth, proper 1-dimensional algebraic variety X together with its point P such that $H_P = H$. But he did not claim that \mathcal{M}_H is non-empty. Using the Pinkham's construction of \mathcal{M}_H , some mathematicians showed that for some H , \mathcal{M}_H is non-empty. To state their results we prepare some notation. Let $M(H) = \{a_1, \dots, a_n\}$ be the minimal set of generators for the semigroup H , which is uniquely determined by H . I_H denotes the kernel of the k -algebra homomorphism $\varphi: k[X] = k[X_1, \dots, X_n] \rightarrow k[t]$ defined by $\varphi(X_i) = t^{a_i}$ where $k[X]$ and $k[t]$ are polynomial rings over k , and $\mu(H)$ denotes the least number of generators for the ideal I_H . When we set $C_H = \text{Spec } k[X]/I_H$, we denote by $T_{C_H}^1 = \bigoplus_{l \in \mathbb{Z}} T_{C_H}^1(l)$ the k -vector space of first order deformations of C_H with a natural graded structure. Moreover, $g(H)$ and $C(H)$ denote the cardinal number of the set $N - H$ and the least integer c with $c + N \subseteq H$, respectively. Then \mathcal{M}_H is non-empty in the following cases:

- 1) H is a complete intersection, i. e., $\mu(H) = n - 1$,
- 2) H is a special almost complete intersection (Waldi [10]),

3) H is negatively graded, i. e., $T_{C_H}^1(l)=0$ for $l>0$ (Pinkham [7], Rim-Vitulli [8]),

4) H is generated by 4 elements and is symmetric, i. e., $C(H)=2g(H)$ (Buchweitz [2], Waldi [9]).

In this paper we shall give some examples of numerical semigroups H generated by 4 elements with $\mathcal{M}_H \neq \emptyset$, because for any numerical semigroup H generated by 2 or 3 elements, 1) and 2) imply $\mathcal{M}_H \neq \emptyset$. Throughout the paper, we are devoted to a numerical semigroup H of torus embedding type (see Definition 1.1), roughly speaking, C_H is the fibre of a torus embedding. For such an H , we can prove that \mathcal{M}_H is non-empty. In Section 2 we show that numerical semigroups H generated by 2 or 3 elements are of torus embedding type. When H is a neat numerical semigroup (see Definition 3.1) generated by 4 elements, we construct a torus embedding, any irreducible component of whose fibre over the origin is isomorphic to C_H , in Section 4. Moreover, if H is 1-neat (see Definition 4.10), we can show that H is of torus embedding type. Using this we can show that symmetric or almost symmetric numerical semigroups H generated by 4 elements are of torus embedding type.

Notation

Throughout this paper we will use the following notation without further warning. We denote by k an algebraically closed field and by N the additive semigroup of non-negative integers. For elements a_1, \dots, a_n, m and l of N , $\langle a_1, \dots, a_n \rangle$ (resp. (a_1, \dots, a_n) , resp. $[l, m]$) denotes the subsemigroup of N generated by a_1, \dots, a_n (resp. the greatest common measure of a_1, \dots, a_n , resp. the set of integers which is larger than or equal to l , and which is smaller than or equal to m). For a weighted ring R and a homogeneous element f of R , $\partial(f)$ means the weight of f . Let H be a numerical semigroup, i. e., the subsemigroup of N whose complement in N is finite. Then \mathcal{M}_H denotes the moduli space, which is obtained by Pinkham, consisting of isomorphic classes of pairs (X, P) with a smooth, proper 1-dimensional algebraic variety X over k and with its point P whose gaps are $N-H$. Moreover, we denote by $g(H)$ the cardinal number of the set $N-H$, by $C(H)$ the least integer c with $c+N \subseteq H$ and by $M(H) = \{a_1, \dots, a_n\}$ the minimal set of generators for the semigroup H . We set

$$\alpha_i = \text{Min} \{ \alpha \in N - \{0\} \mid \alpha a_i \in \langle a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n \rangle \}$$

for all $i=1, \dots, n$. For any non-zero element h of H let

$$L_h(H) = \{0 = \omega_h(1) < \dots < \omega_h(h)\}$$

be the set of the least elements of H in respective congruence classes mod h . φ_H denotes the k -algebra homomorphism from $k[X_1, \dots, X_n]$ to $k[t]$ defined by sending X_i to t^{a_i} , hence assigning $\partial(X_i)=a_i$ for $1 \leq i \leq n$ and $\partial(c)=0$ for $c \in k^\times$, $k[X_1, \dots, X_n]$ is made into a weighted k -algebra. We denote by I_H the kernel of φ_H , by $\mu(H)$ the least number of generators for the ideal I_H and by C_H the affine curve $\text{Spec } k[X_1, \dots, X_n]/I_H$.

1. Numerical semigroups of torus embedding type.

In this paper we are concerned with the following numerical semigroups :

DEFINITION 1.1. A numerical semigroup H with $M(H)=\{a_1, \dots, a_n\}$ is of torus embedding type if there exist a positive integer $m \geq n$, homogeneous elements $g_i (1 \leq i \leq m)$ of $k[X]=k[X_1, \dots, X_n]$ of weight >0 , and a saturated subsemigroup S of \mathbb{Z}^{m+1-n} which is generated by b_1, \dots, b_m and which generates a subgroup of rank $m+1-n$ of \mathbb{Z}^{m+1-n} as a group, such that the kernel of the k -algebra homomorphism

$$\pi : k[Y]=k[Y_1, \dots, Y_m] \longrightarrow k[S]=k[T^s]_{s \in S}$$

defined by $\pi(Y_i)=T^{b_i}$, is generated by homogeneous elements $F_k (1 \leq k \leq u)$ with $I_H=(F_1(g_1, \dots, g_m), \dots, F_u(g_1, \dots, g_m))$ where the weight on $k[Y]$ is defined by $\partial(Y_i)=\partial(g_i)$ for $1 \leq i \leq m$ and $\partial(c)=0$ for $c \in k^\times$.

A sufficient condition that a numerical semigroup is of torus embedding type, which we will use, is the following :

LEMMA 1.2. Let H be a numerical semigroup with $M(H)=\{a_1, \dots, a_n\}$. Assume that there exist a positive integer $m \geq n$, non-constant monomials $g_i (1 \leq i \leq m)$ in $k[X]=k[X_1, \dots, X_n]$, and a saturated subsemigroup S of \mathbb{Z}^{m+1-n} which is generated by b_1, \dots, b_m and which generates a subgroup of rank $m+1-n$ of \mathbb{Z}^{m+1-n} as a group, such that if we let

$$\pi : k[Y]=k[Y_1, \dots, Y_m] \longrightarrow k[T^s]_{s \in S} \quad (\text{resp. } \eta : k[Y] \rightarrow k[X])$$

be the k -algebra homomorphism defined by $\pi(Y_i)=T^{b_i}$ (resp. $\eta(Y_i)=g_i$), then the ideal I_H is generated by the elements of $\eta(\text{Ker } \pi)$. Then H is of torus embedding type.

PROOF. When we define a weight on $k[Y]$ in virtue of $\partial(Y_i)=\partial(g_i)$ for $1 \leq i \leq m$ and $\partial(c)=0$ for $c \in k^\times$, it suffices to show that there exists a set $\{F_k\}_{1 \leq k \leq u}$ of homogeneous generators for the ideal $\text{Ker } \pi$, because the ideal I_H is generated

by $\eta(F_k)$ ($1 \leq k \leq u$). Now by [5] we may take generators F_k ($1 \leq k \leq u$) of the ideal $\text{Ker } \pi$ as follows:

$$F_k = \prod_{i=1}^m Y_i^{\nu_{ki}} - \prod_{i=1}^m Y_i^{\mu_{ki}}$$

where $\nu_{ki} \cdot \mu_{ki} = 0$ for all $1 \leq k \leq u$ and all $1 \leq i \leq m$. If we put $g_i = X_1^{\gamma_{i1}} \cdots X_n^{\gamma_{in}}$ for all $1 \leq i \leq m$, then we have

$$\begin{aligned} 0 &= \varphi_H(\eta(F_k)) = \varphi_H\left(\prod_{i=1}^m g_i^{\nu_{ki}} - \prod_{i=1}^m g_i^{\mu_{ki}}\right) \\ &= t^{\sum_{i=1}^m \nu_{ki} \sum_{j=1}^n \gamma_{ij} a_j} - t^{\sum_{i=1}^m \mu_{ki} \sum_{j=1}^n \gamma_{ij} a_j}, \end{aligned}$$

which implies $\sum_{i=1}^m \nu_{ki} \sum_{j=1}^n \gamma_{ij} a_j = \sum_{i=1}^m \mu_{ki} \sum_{j=1}^n \gamma_{ij} a_j$. Therefore F_k 's are homogeneous.

Q. E. D.

Here we give a few examples of numerical semigroups of torus embedding type.

EXAMPLE 1.3. (1) $H = \langle 3, 7 \rangle$ is of torus embedding type. In fact, let $a_1 = 3$ and $a_2 = 7$. If we set $n = m = 2$, $g_1 = X_1^7$, $g_2 = X_2^3$ and $b_1 = b_2 = 1$, then these satisfy the assumption of Lemma 1.2. In this case $\text{Ker } \pi$ contains a homogeneous element $F_1 = Y_1 - Y_2$. See Lemma 2.3 for a generalization.

(2) $H = \langle 4, 7, 13 \rangle$ is of torus embedding type. In fact, let $a_1 = 4$, $a_2 = 7$ and $a_3 = 13$. If we set $n = 3$, $m = 6$, $g_1 = X_1^2$, $g_2 = X_2$, $g_3 = X_3$, $g_4 = X_1^3$, $g_5 = X_2^2$, $g_6 = X_3$, $b_1 = (1, 0, 0, 0)$, $b_2 = (0, 1, 0, 0)$, $b_3 = (0, 0, 1, 0)$, $b_4 = (-1, 1, 1, 0)$, $b_5 = (0, 0, 0, 1)$ and $b_6 = (-1, 1, 0, 1)$, then these satisfy the assumption of Lemma 1.2. In this case we can see that $\text{Ker } \pi$ contains homogeneous elements F_k ($1 \leq k \leq 3$) as follows:

$$F_1 = Y_1 Y_4 - Y_2 Y_3, \quad F_2 = Y_2 Y_5 - Y_1 Y_6 \quad \text{and} \quad F_3 = Y_3 Y_6 - Y_4 Y_5.$$

See Proposition 2.5 for a generalization.

(3) $H = \langle 4, 9, 14, 15 \rangle$ is of torus embedding type. In fact, let $a_1 = 15$, $a_2 = 9$, $a_3 = 4$ and $a_4 = 14$. If we set $n = 4$, $m = 9$, $g_1 = X_1$, $g_2 = X_2$, $g_3 = X_3^4$, $g_4 = X_4$, $g_5 = X_1$, $g_6 = X_2$, $g_7 = X_3$, $g_8 = X_4$, $g_9 = X_3$, $b_i = e_i$ ($1 \leq i \leq 4$), $b_5 = (-1, 0, 1, 1, 0, 0)$, $b_6 = e_5$, $b_7 = e_6$, $b_8 = (0, 1, 0, 0, 1, -1)$ and $b_9 = (1, 1, -1, 0, 0, -1)$ where for any $i \in [1, 6]$ we denote by $e_i \in \mathbb{Z}^6$ the vector whose i -th component equals to 1 and whose j -th component equals to 0 if $j \neq i$, then these satisfy the assumption of Lemma 1.2. In this case we can see that $\text{Ker } \pi$ contains homogeneous elements F_k ($1 \leq k \leq 6$) as follows:

$$F_1 = Y_1 Y_5 - Y_3 Y_4, \quad F_2 = Y_2 Y_6 - Y_7 Y_8, \quad F_3 = Y_3 Y_7 Y_9 - Y_1 Y_2,$$

$$F_4=Y_4Y_8-Y_5Y_6Y_9, \quad F_5=Y_1Y_8-Y_3Y_6Y_9 \quad \text{and} \quad F_6=Y_2Y_4-Y_5Y_7Y_9.$$

See Theorem 4.11 for a generalization.

(4) $H=\langle 5, 8, 9, 11 \rangle$ is of torus embedding type. In fact, let $a_1=5, a_2=8, a_3=9$ and $a_4=11$. If we set $n=4, m=9, g_i=X_i (1 \leq i \leq 4), g_5=X_1^2, g_{4+i}=X_i (2 \leq i \leq 4), g_9=X_1, b_i=e_i (1 \leq i \leq 6), b_7=(0, 1, -1, 0, 1, 0), b_8=(-1, 1, 0, 0, 0, 1)$ and $b_9=(-1, 0, 1, 1, -1, 0)$ where e_i 's are as in (3), then these satisfy the assumption of Lemma 1.2. In this case, $\text{Ker } \pi$ contains homogeneous elements $F_k (1 \leq k \leq 5)$ as follows:

$$F_1=Y_1Y_5Y_9-Y_3Y_4, \quad F_2=Y_2Y_6-Y_1Y_8, \quad F_3=Y_3Y_7-Y_2Y_5, \\ F_4=Y_4Y_8-Y_6Y_7Y_9 \quad \text{and} \quad F_5=Y_1Y_7Y_9-Y_2Y_4.$$

See Theorem 4.11 for a generalization. Now we get $g(H)=7$ and $C(H)=13$, which imply $C(H)=2g(H)-1$, i. e., H is almost symmetric (see Theorem 6.4).

In the remains of this section we assume that k is of characteristic 0. If H is of torus embedding type, then we can show $\mathcal{M}_H \neq \emptyset$. For this purpose we show the following:

PROPOSITION 1.4. *Let a_1, \dots, a_n be positive integers and let $k[X]=k[X_1, \dots, X_n]$ be a polynomial ring on which the weight is defined by $\partial(X_i)=a_i$ for $1 \leq i \leq n$ and $\partial(c)=0$ for $c \in k^\times$. Let $k[Y]=k[Y_1, \dots, Y_m]$ and $k[Y, W]=k[Y_1, \dots, Y_m, W_1, \dots, W_l]$ be two polynomial rings. Let r be a non-negative integer with $n-l \geq r$, let J be an ideal in $k[Y]$ such that $R=k[Y]/J$ is a Cohen-Macaulay domain of dimension $m+l+r-n$ and that the singular locus of $\text{Spec } R$ has codimension larger than r , and let $R[X]=R[X_1, \dots, X_n]$. Assume that there exist homogeneous elements $g_i (1 \leq i \leq m)$ and $h_j (1 \leq j \leq l)$ of $k[X]$ of weight > 0 such that we have the fibre product:*

$$\begin{array}{ccc} \phi^{-1}(\text{the origin}) & \longrightarrow & \text{Spec } R[X] \\ \downarrow & & \downarrow \phi \\ \text{Spec } k & \longrightarrow & \text{Spec } k[Y, W] \end{array}$$

with $\dim \phi^{-1}(\text{the origin})=r$, where ϕ is the morphism which is induced by the k -algebra homomorphism $\phi^*: k[Y, W] \rightarrow R[X]$ defined by $\phi^*(Y_i)=g_i-Y_i \text{ mod } J$ and $\phi^*(W_j)=h_j$, and such that the ideal J is homogeneous where the weight on $k[Y]$ is defined by $\partial(Y_i)=\partial(g_i)$ for $1 \leq i \leq m$ and $\partial(c)=0$ for $c \in k^\times$. Then ϕ is flat and there exists a non-empty open subset V of $\text{Spec } k[Y, W]$ such that the

restriction $\phi^{-1}(V) \rightarrow V$ is smooth.

PROOF. We define a weight on $k[Y, W]$ as follows:

$$\partial(Y_i) = \partial(g_i), \quad \partial(W_j) = \partial(h_j) \quad \text{and} \quad \partial(c) = 0 \quad \text{for } c \in k^\times.$$

Since the ideal J in $k[Y]$ is homogeneous, ϕ is a G_m -equivariant morphism. For any $s \in \mathbb{Z}$, the closed subset

$$F_s = \{x \in \text{Spec } R[X] \mid \dim_x \phi^{-1}(\phi(x)) \geq s\}$$

contains the origin if $F_s \neq \emptyset$, because ϕ is G_m -equivariant and the weights of Y_i, X_k are positive. ϕ is dominating in virtue of

$$\dim \text{Spec } R[X] - \dim \text{Spec } k[Y, W] = m + l + r - (m + l) = r$$

and

$$\dim \phi^{-1}(\text{the origin}) = r,$$

which implies $\dim_x \phi^{-1}(\phi(x)) \geq r$ for all $x \in \text{Spec } R[X]$. Moreover, in virtue of $\partial(Y_i) > 0$ and $\partial(W_j) > 0$ the map ϕ send the origin in $\text{Spec } R[X]$ to the one in $\text{Spec } k[Y, W]$. Assume that $F_{r+1} \neq \emptyset$. Since the origin belongs to F_{r+1} , we get

$$r + 1 \leq \dim_{\substack{\phi^{-1}(\phi(\text{the origin})) \\ \text{the origin}} \phi^{-1}(\phi(\text{the origin})) = \dim_{\substack{\phi^{-1}(\phi(\text{the origin})) \\ \text{the origin}} \phi^{-1}(\phi(\text{the origin}))$$

$$\leq \dim \phi^{-1}(\text{the origin}) = r,$$

a contradiction, which implies $F_{r+1} = \emptyset$. Therefore we get $\dim_x \phi^{-1}(\phi(x)) = r$ for all $x \in \text{Spec } R[X]$, i. e., ϕ is equidimensional. Since R is a Cohen-Macaulay domain, ϕ is flat ([3]). Let $Z_i (i \in I)$ be the irreducible components in the singular locus $\text{Sing}(\text{Spec } R[X])$ of $\text{Spec } R[X]$ and let η be the generic point of $\text{Spec } k[Y, W]$. Assume that $\phi^{-1}(\eta) \cap \text{Sing}(\text{Spec } R[X]) \neq \emptyset$, i. e., there exists $i \in I$ such that $\phi^{-1}(\eta) \cap Z_i \neq \emptyset$. Since the restriction $Z_i \subset \text{Spec } R[X] \rightarrow \text{Spec } k[Y, W]$ is dominating, we have

$$0 \leq \dim Z_i - \dim \text{Spec } k[Y, W] \leq \dim \text{Sing}(\text{Spec } R[X]) - \dim \text{Spec } k[Y, W]$$

$$< \dim \text{Spec } R[X] - r - \dim \text{Spec } k[Y, W] = 0,$$

a contradiction. Hence we get $\phi^{-1}(\eta) \cap \text{Sing}(\text{Spec } R[X]) = \emptyset$, which implies that the set

$$\{y \in \text{Spec } k[Y, W] \mid \phi^{-1}(y) \cap \text{Sing}(\text{Spec } R[X]) = \emptyset\}$$

contains a non-empty open subset U . Then we have

$$\phi^{-1}(U) \subseteq \text{Spec } R[X] - \text{Sing}(\text{Spec } R[X])$$

Hence there is a non-empty open subset V in $\text{Spec } k[Y, W]$ such that the restric-

tion $\phi^{-1}(V) \rightarrow V$ is smooth, because the restriction $\phi^{-1}(U) \rightarrow \text{Spec } k[Y, W]$ is a morphism of varieties with smooth $\phi^{-1}(U)$ over the algebraically closed field k of characteristic 0 ([4]). Q. E. D.

Pinkham [7] showed the following:

REMARK 1.5. Let H be a numerical semigroup with $M(H) = \{a_1, \dots, a_n\}$. Then we have $\mathcal{M}_H \neq \emptyset$ if and only if there exists a flat homogeneous homomorphism $\phi^*: A = \bigoplus_{i \in \mathbb{Z}} A_i \rightarrow B = \bigoplus_{i \in \mathbb{Z}} B_i$ of affine graded k -algebras with $A_0 \cong k$ and $B_0 \cong k$ such that 1) C_H is the fibre of the morphism $\phi: \text{Spec } B \rightarrow \text{Spec } A$ associated to ϕ^* over a homogeneous k -rational point on $\text{Spec } A$, 2) A is a domain and the generic fibre of ϕ is smooth, and 3) $A_i = 0$ for all $i < 0$.

Combining Proposition 1.4 with Remark 1.5, we get the following:

COROLLARY 1.6. Let H be a numerical semigroup with $M(H) = \{a_1, \dots, a_n\}$ and let $k[X]$, $k[Y]$ and $k[Y, W]$ be polynomial rings as in Proposition 1.4. Let J be an ideal in $k[Y]$ such that $R = k[Y]/J$ is a normal Cohen-Macaulay domain of dimension $m+l+1-n$. Assume that there exist homogeneous elements $g_i (1 \leq i \leq m)$ and $h_j (1 \leq j \leq l)$ of $k[X]$ of weight > 0 such that we have the fibre product:

$$\begin{array}{ccc}
 C_H & \longrightarrow & \text{Spec } R[X] \\
 \downarrow & & \downarrow \phi \\
 \text{Spec } k & \longrightarrow & \text{Spec } k[Y, W] \\
 & & \downarrow \\
 [(0)] & \longleftarrow & \text{the origin}
 \end{array}$$

where ϕ is the morphism induced by the k -algebra homomorphism $\phi^*: k[Y, W] \rightarrow R[X]$ defined by $\phi^*(Y_i) = g_i - Y_i \pmod{J}$ and $\phi^*(W_j) = h_j$, and such that the ideal J is homogeneous where the weight on $k[Y]$ is defined by $\partial(Y_i) = \partial(g_i)$ for $1 \leq i \leq m$ and $\partial(c) = 0$ for $c \in k^*$. Then we have $\mathcal{M}_H \neq \emptyset$.

If we apply Corollary 1.6 to numerical semigroups of torus embedding type, we see:

THEOREM 1.7. For any numerical semigroup H of torus embedding type, we have $\mathcal{M}_H \neq \emptyset$.

PROOF. We use the notation in Definition 1.1. Since S is a saturated sub-

semigroup of \mathbb{Z}^{m+1-n} which is finitely generated and which generates a subgroup of rank $m+1-n$ of \mathbb{Z}^{m+1-n} as a group, by [6] $\text{Spec } k[T^s]_{s \in S}$ is a normal affine equivariant embedding of $(G_m)^{m+1-n}$ and is a Cohen-Macaulay scheme. Hence $R = k[Y]/\text{Ker } \pi$ is a normal Cohen-Macaulay domain of dimension $m+1-n$ and the ideal $J = \text{Ker } \pi$ is generated by homogeneous elements $F_k (1 \leq k \leq u)$. Since the ideal I_H is generated by the $F_k(g_1, \dots, g_m)$'s, we have a fibre product:

$$\begin{array}{ccc}
 C_H & \longrightarrow & \text{Spec } R[X] \\
 \downarrow & & \downarrow \phi \\
 \text{Spec } k & \longrightarrow & \text{Spec } k[Y] \\
 [(0)] & \longmapsto & \text{the origin}
 \end{array}$$

where ϕ is the morphism induced by the k -algebra homomorphism $\phi^*: k[Y] \rightarrow R[X]$ defined by $\phi^*(Y_i) = g_i - Y_i \text{ mod } J$. If we apply Corollary 1.6 to the case $l=0$, we obtain $\mathcal{M}_H \neq \emptyset$. Q. E. D.

2. Numerical semigroups generated by 2 or 3 elements.

In this section we will show that numerical semigroups generated by 2 or 3 elements are of torus embedding type. First we consider the following numerical semigroups:

DEFINITION 2.1. A numerical semigroup H with $M(H) = \{a_1, \dots, a_n\}$ is called a *strictly complete intersection* if renumbering a_1, \dots, a_n the least common multiple of (a_1, \dots, a_{i-1}) and a_i belongs to $\langle a_1, \dots, a_{i-1} \rangle$ for $2 \leq i \leq n$. In this case by [5] a set of generators for the ideal I_H is well-known.

REMARK 2.2. For a numerical semigroup H as in Definition 2.1 we have $\alpha_i = (a_1, \dots, a_{i-1}) / (a_1, \dots, a_i)$ for $2 \leq i \leq n$. If we set

$$\alpha_i a_i = \sum_{j=1}^{i-1} \alpha_{ij} a_j \quad \text{with } \alpha_{ij} \in \mathbb{N}$$

for $2 \leq i \leq n$, then the ideal I_H is generated by f_2, \dots, f_n where we set $f_i = X_i^{\alpha_i} - X_1^{\alpha_{i1}} \dots X_{i-1}^{\alpha_{i,i-1}}$.

LEMMA 2.3. A numerical semigroup H which is a strictly complete intersection, is of torus embedding type.

PROOF. We use the notation in Remark 2.2. The set

$$U = \{(i, j) \in \mathbb{N}^2 \mid 2 \leq i \leq n \text{ and } 1 \leq j \leq i-1\}$$

is a totally ordered set, where we define $(i, j) \leq (i', j')$ if $i < i'$ or if $i = i'$ and $j \leq j'$. If we set $P = \{(i, j) \in U \mid \alpha_{ij} \neq 0\}$ and $l = \#P$, then we have the isomorphism $\xi: P \rightarrow [1, l]$ of ordered sets. Let

$$\pi: k[Y_{ij}((i, j) \in P); Z_k(2 \leq k \leq n)] \longrightarrow k[t_1, \dots, t_l]$$

be the k -algebra homomorphism of polynomial rings, defined by $\pi(Y_{ij}) = t_{\xi(i, j)}$ and $\pi(Z_k) = \prod_{j \in P(k)} t_{\xi(k, j)}$ where $P(k) = \{j \in [1, k-1] \mid (k, j) \in P\}$. We set

$$g_{\xi(i, j)} = X_j^{\alpha_{ij}} \text{ for } (i, j) \in P \text{ and } g_{l+k-1} = X_k^{\alpha_k} \text{ for } 2 \leq k \leq n.$$

Let $\eta: k[Y_{ij}; Z_k] \rightarrow k[X] = k[X_1, \dots, X_n]$ (resp. $\zeta: k[t_1, \dots, t_l] \rightarrow k[t]$) be the k -algebra homomorphism defined by $\eta(Y_{ij}) = g_{\xi(i, j)}$ and $\eta(Z_k) = g_{l+k-1}$ (resp. $\zeta(t_{\xi(i, j)}) = t^{\alpha_{ij}}$). In virtue of $\varphi_H \circ \eta = \zeta \circ \pi$, we get $\eta(\text{Ker } \pi) \subseteq \text{Ker } \varphi_H = I_H$. If we set $F_k = Z_k - \prod_{j \in P(k)} Y_{kj}$ for $2 \leq k \leq n$, then $F_k \in \text{Ker } \pi$ and $\eta(F_k) = f_k$. Therefore by Remark 2.2 the ideal I_H is generated by the elements of $\eta(\text{Ker } \pi)$. By Lemma 1.2 H is of torus embedding type. Q. E. D.

COROLLARY 2.4. 1) Numerical semigroups with $M(H) = \{a_1, a_2\}$ are of torus embedding type.

2) Symmetric numerical semigroups, i.e., $C(H) = 2g(H)$, with $M(H) = \{a_1, a_2, a_3\}$ are of torus embedding type.

PROOF. It is trivial that numerical semigroups with $M(H) = \{a_1, a_2\}$ are strictly complete intersections. Herzog [5] proved that numerical semigroups H with $M(H) = \{a_1, a_2, a_3\}$ are strictly complete intersections if and only if they are symmetric. Q. E. D.

In the non-symmetric case H with $M(H) = \{a_1, a_2, a_3\}$, H is also of torus embedding type in the following way: by [5] there exist positive integers $\alpha_{ij} < \alpha_j$ such that

$$\alpha_1 a_1 = \alpha_{12} a_2 + \alpha_{13} a_3, \quad \alpha_2 a_2 = \alpha_{21} a_1 + \alpha_{23} a_3 \quad \text{and} \quad \alpha_3 a_3 = \alpha_{31} a_1 + \alpha_{32} a_2,$$

in this case

$$\alpha_1 = \alpha_{21} + \alpha_{31}, \quad \alpha_2 = \alpha_{12} + \alpha_{32} \quad \text{and} \quad \alpha_3 = \alpha_{13} + \alpha_{23}.$$

Moreover, Herzog showed that the ideal I_H is generated by

$$f_1 = X_1^{\alpha_1} - X_2^{\alpha_{12}} X_3^{\alpha_{13}}, \quad f_2 = X_2^{\alpha_2} - X_1^{\alpha_{21}} X_3^{\alpha_{23}} \quad \text{and} \quad f_3 = X_3^{\alpha_3} - X_1^{\alpha_{31}} X_2^{\alpha_{32}}.$$

Let S be the subsemigroup of \mathbb{Z}^4 generated by

$$b_{21}=(1, 0, 0, 0), \quad b_{12}=(0, 1, 0, 0), \quad b_{13}=(0, 0, 1, 0), \quad b_{31}=(-1, 1, 1, 0), \\ b_{32}=(0, 0, 0, 1) \quad \text{and} \quad b_{23}=(-1, 1, 0, 1).$$

Then it can be easily seen that $S = \sum \mathbf{R}_+ b_{ij} \cap \mathbf{Z}^4$ where \mathbf{R}_+ is the set of non-negative real numbers. Hence S is saturated. When we let

$$\pi : k[Y_{ij}]_{1 \leq i \neq j \leq 3} \longrightarrow k[T^s]_{s \in S} \quad (\text{resp. } \eta : k[Y_{ij}] \rightarrow k[X_1, X_2, X_3])$$

be the k -algebra homomorphism defined by $\pi(Y_{ij}) = T^{b_{ij}}$ (resp. $\eta(Y_{ij}) = X_j^{i_j}$), there exists a k -algebra homomorphism $\zeta : k[T^s]_{s \in S} \rightarrow k[t]$ such that $\varphi_H \circ \eta = \zeta \circ \pi$, which implies $\eta(\text{Ker } \pi) \subseteq I_H$. Since

$$F_1 = Y_{21}Y_{31} - Y_{12}Y_{13}, \quad F_2 = Y_{12}Y_{32} - Y_{21}Y_{23} \quad \text{and} \quad F_3 = Y_{13}Y_{23} - Y_{31}Y_{32}$$

belong to $\text{Ker } \pi$ and we have $\eta(F_i) = f_i$ for $1 \leq i \leq 3$, the ideal I_H is generated by the elements of $\eta(\text{Ker } \pi)$, hence H is of torus embedding type. Therefore combining this with Corollary 2.4 2), we obtain the following:

PROPOSITION 2.5. *Numerical semigroups with $M(H) = \{a_1, a_2, a_3\}$ are of torus embedding type.*

3. Neat numerical semigroups.

Hereafter we are concerned with the following numerical semigroups:

DEFINITION 3.1. For a numerical semigroup H with $M(H) = \{a_1, \dots, a_n\}$,

$$\mathcal{R} : \begin{cases} \alpha_i a_i = \sum_{j \neq i} \alpha_{ij} a_j & \text{with } 0 \leq \alpha_{ij} < \alpha_j, \quad \text{for } 1 \leq i \leq n, \\ \sum_{i \neq j} \alpha_{ij} = \alpha_j & \text{for } 1 \leq j \leq n \end{cases}$$

is called a *neat system of relations with respect to H and $\{a_1, \dots, a_n\}$* . When H has a neat system of relations, it is called to be *neat*.

EXAMPLE 3.2. (1) $H = \langle 4, 7, 13 \rangle$ is neat. In fact, let $a_1 = 4$, $a_2 = 7$ and $a_3 = 13$. Then

$$\mathcal{R} : 5a_1 = a_2 + a_3, \quad 3a_2 = 2a_1 + a_3, \quad 2a_3 = 3a_1 + 2a_2$$

is a neat system of relations.

(2) $H = \langle 4, 9, 14, 15 \rangle$ is neat. In fact, let $a_1 = 15$, $a_2 = 9$, $a_3 = 4$ and $a_4 = 14$. Then

$$\mathcal{R} : 2a_1 = 4a_3 + a_4, \quad 2a_2 = a_3 + a_4, \quad 6a_3 = a_1 + a_2, \quad 2a_4 = a_1 + a_2 + a_3$$

is a neat system of relations.

(3) $H = \langle 10, 11, 13, 14 \rangle$ is neat. In fact, let $a_1 = 10, a_2 = 11, a_3 = 14$ and $a_4 = 13$. Then

$$\mathcal{R} : 4a_1 = a_3 + 2a_4, \quad 3a_2 = 2a_1 + a_4, \quad 3a_3 = 2a_1 + 2a_2, \quad 3a_4 = a_2 + 2a_3$$

is a neat system of relations.

(4) $H = \langle 5, 7, 9, 11, 13 \rangle$ is neat. In fact, let $a_1 = 5, a_2 = 7, a_3 = 9, a_4 = 11$ and $a_5 = 13$. Then

$$\mathcal{R} : 4a_1 = a_2 + a_5, \quad 2a_2 = a_1 + a_3, \quad 2a_3 = a_2 + a_4, \quad 2a_4 = a_3 + a_5, \quad 2a_5 = 3a_1 + a_4$$

is a neat system of relations.

In this section, let H be a neat numerical semigroup with $M(H) = \{a_1, \dots, a_n\}$, and let \mathcal{R} be a neat system of relations with respect to H and $\{a_1, \dots, a_n\}$. We can see easily:

REMARK 3.3. We put

$$P = P_{\mathcal{R}} = \{(i, j) \in [1, n]^2 \mid i \neq j, \alpha_{i,j} \neq 0\}, \quad P^i = \{j \in [1, n] \mid (i, j) \in P\}$$

$$\text{for } 1 \leq i \leq n \text{ and } P_j = \{i \in [1, n] \mid (i, j) \in P\} \quad \text{for } 1 \leq j \leq n.$$

Then $*P^i \geq 2$ and $*P_j \geq 2$. Hence we have $*P \geq 2n$, for

$$P = \bigcup_{1 \leq i \leq n} \{(i, j) \mid j \in P^i\} = \bigcup_{1 \leq j \leq n} \{(i, j) \mid i \in P_j\}.$$

Moreover, we make P into a totally ordered set by defining an order on it as follows: for a fixed $j \in [1, n]$ and any $1 \leq k \leq *P_j$, we define inductively

$$i_j(k) = \text{Min} \{i \in [1, n] \mid i \in P_j - \{i_j(1), \dots, i_j(k-1)\}\}.$$

For any (i, j) and $(i', j') \in P$ with $i = i_j(k)$ and $i' = i_{j'}(k')$, we define $(i, j) \leq (i', j')$ if $k < k'$ or if $k = k'$ and $j \leq j'$.

DEFINITION 3.4. An element (i, j) of P has a *v-relation* (resp. an *h-relation*) if we have

$$i = \text{Max} \{i' \in [1, n] \mid i' \in P_j\} \quad \text{and} \quad P^j(i, j) = \emptyset$$

$$\text{where } P^j(i, j) = \{j' \in P^j \mid (j, j') > (i, j)\}$$

$$\text{(resp. } (i, j) = \text{Max} \{(i, j') \mid j' \in P^i\} \quad \text{and} \quad P_i(i, j) = \emptyset$$

$$\text{where } P_i(i, j) = \{i' \in P_i \mid (i', i) > (i, j)\}.$$

v-relations and *h-relations* have the following properties:

LEMMA 3.5. 1) $(i_0, j_0) = \text{Max } P$ has a *v-relation* and an *h-relation*.

2) For any $1 \leq l \leq n$, there exists $i \in [1, n]$ such that (i, l) has a *v-relation* or

$j \in [1, n]$ such that (l, j) has an h -relation.

3) We have $*Q \leq n-1$ where

$$Q = \{(i, j) \in P \mid (i, j) \text{ has either a } v\text{-relation or an } h\text{-relation}\}.$$

PROOF. 1) is trivial. We set

$$i = \text{Max } P_l \quad \text{and} \quad (l, j) = \text{Max}\{(l, j') \mid j' \in P^l\}.$$

Assume that (i, l) does not have a v -relation and that (l, j) does not have an h -relation. Then there exist $j' \in P^l(i, l)$ and $i' \in P_l(l, j)$, which imply

$$(i, l) \geq (i', l) > (l, j) \geq (l, j') > (i, l),$$

a contradiction. This proves 2). Let $l \in [1, n]$. If (i, l) has a v -relation, then we define $\zeta(l) = (i, l)$. If (l, j) has an h -relation, then we define $\zeta(l) = (l, j)$. Then the map $\zeta: [1, n] \rightarrow Q$ is well-defined. In fact, if (i, l) (resp. (i', l)) has a v -relation, then $i = \text{Max } P_l = i'$. If (l, j) (resp. (l, j')) has an h -relation, then $(l, j) = \text{Max}\{(l, k) \mid k \in P^l\} = (l, j')$, hence $j = j'$. If (i, l) (resp. (l, j)) has a v -relation (resp. an h -relation), then we have $(i, l) \leq (l, j) \leq (i, l)$, hence $l = j$, a contradiction. To prove 3) it suffices to show that ζ is surjective, because we have $\zeta(i_0) = (i_0, j_0) = \zeta(j_0)$. If $(i, j) \in Q$ has a v -relation (resp. an h -relation), then $\zeta(j) = (i, j)$ (resp., $\zeta(i) = (i, j)$). Hence ζ is surjective. Q. E. D.

Finally we define the subset P_H of $\mathcal{S}_n = \{(i, j) \in [1, n]^2 \mid i \neq j\}$ associated to a neat numerical semigroup H with $M(H) = \{a_1, \dots, a_n\}$ as follows:

DEFINITION 3.6. We define an order on the set of subsets of \mathcal{S}_n in the following way:

1) for any (i, j) and $(i', j') \in \mathcal{S}_n$, we define $(i, j) \leq (i', j')$ if $i < i'$ or if $i = i'$ and $j \leq j'$,

2) for two subsets P and P' of \mathcal{S}_n with $*P = *P' = *\mathcal{S}_n - r$, we define $P \leq P'$ if there exists $0 \leq q \leq r$ such that

$$(i_1, j_1) = (i'_1, j'_1), \dots, (i_q, j_q) = (i'_q, j'_q) \quad \text{and} \quad (i_{q+1}, j_{q+1}) < (i'_{q+1}, j'_{q+1})$$

where

$$\mathcal{S}_n - P = \{(i_1, j_1) < \dots < (i_r, j_r)\} \quad \text{and} \quad \mathcal{S}_n - P' = \{(i'_1, j'_1) < \dots < (i'_r, j'_r)\},$$

3) for two subsets P and P' of \mathcal{S}_n we define $P \leq P'$ if $*P < *P'$ or if $*P = *P'$ and $P \leq P'$.

Then the set of subsets of \mathcal{S}_n becomes a totally ordered set. Using this order, we define the subset P_H of \mathcal{S}_n :

$$P_H = \text{Min}\{P_{H, \{a_{\sigma(1)}, \dots, a_{\sigma(n)}\}} \mid \sigma \text{ runs over the set of permutations of } [1, n]\}$$

where

$$P_{H, \{a_1, \dots, a_n\}} = \text{Min}\{P_{\mathcal{R}} \mid \mathcal{R} \text{ runs over the set of neat systems of relations with respect to } H \text{ and } \{a_1, \dots, a_n\}\}.$$

4. Neat numerical semigroups generated by 4 elements.

In this section, we are devoted to neat numerical semigroups H with $M(H) = \{a_1, a_2, a_3, a_4\}$. In the case $*M(H) = 4$ we can explain v -relations and h -relations in detail.

LEMMA 4.1. *Let \mathcal{R} be a neat system of relations with respect to H and $\{a_1, a_2, a_3, a_4\}$. Then*

- 1) $(i, j) \in P_{\mathcal{R}}$ has a v -relation and an h -relation if and only if $(i, j) = \text{Max } P_{\mathcal{R}}$,
- 2) we have $*Q = 3$ where

$$Q = \{(i, j) \in P_{\mathcal{R}} \mid (i, j) \text{ has either a } v\text{-relation or an } h\text{-relation}\}.$$

PROOF. To check 1), by Lemma 3.5 1) it suffices to show the “only if” part. For brevity, we put $P = P_{\mathcal{R}}$. Let us take $(i, j) \in P$ which has a v -relation and an h -relation. Then for any $k \in [1, 4]$ the following hold:

- a) if $(i, k) \in P$, then $(i, k) \leq (i, j)$, b) if $(j, k) \in P$, then $(j, k) < (i, j)$, c) if $(k, i) \in P$, then $(k, i) < (i, j)$, d) if $(k, j) \in P$, then $(k, j) \leq (i, j)$.

From now on we will see that for $(k, l) \in P$ with $k, l \in [1, 4] - \{i, j\}$, $(k, l) < (i, j)$. The case $i = 1$ does not occur, because (i, j) has a v -relation. Moreover, since for $k = 1$ we have $(k, l) < (i, j)$, we may assume $j = 1$ or $l = 1$.

(A) $j = 1$. Then $i = 3$ or 4 , because $i \geq i_1(2) \geq 3$.

- 1) $i = 3$. Then $(i_3(2), 3) < (3, 1) = (i_1(2), 1)$, a contradiction.
- 2) $i = 4$. Then $(k, l) = (2, 3)$ or $(3, 2)$. If $(k, l) = (2, 3)$, then

$$(k, l) \leq (i_3(2), 3) < (i_4(2), 4) < (4, 1) = (i, j).$$

If $(k, l) = (3, 2)$, then

$$(k, l) \leq (i_2(2), 2) < (i_4(2), 4) < (4, 1) = (i, j).$$

(B) $l = 1$. Then $k = 2$ or 3 or 4 .

- 1) $k = 2$. Then $(k, l) = (i_1(1), 1) < (i, j)$.
- 2) $k = 3$. Then $(k, l) \leq (i_1(2), 1) < (i_j(2), j) \leq (i, j)$.
- 3) $k = 4$. Then $(i, j) = (2, 3)$ or $(3, 2)$. If $i = i_j(3)$, then

$$(k, l) = (4, 1) \leq (i_1(3), 1) < (i_j(3), j) = (i, j).$$

Assume $i=i_j(2)$. Then

$$(i, j)=(i_j(2), j)<(i_4(2), 4)<(i, j),$$

because $i_4(2)=2$ or 3 . This is a contradiction. Hence we have $(i, j)=\text{Max } P$.

By the proof of Lemma 3.5 3), we can define a surjective map $\zeta: [1, 4] \rightarrow Q$ by sending l to (i_l, l) (resp. (l, i_l)) if (i_l, l) has a ν -relation (resp. if (l, i_l) has an h -relation). Let l and l' be two distinct elements of $[1, 4]$ such that $\zeta(l)=\zeta(l')$. Then $\zeta(l)=\zeta(l')$ has a ν -relation and an h -relation. Hence if we set $(i, j)=\text{Max } P$, by 1) we get $\{l, l'\}=\{i, j\}$. So $\zeta(k), \zeta(k')$ and $\zeta(i)$ are distinct where we set $[1, 4]=\{i, j, k, k'\}$. Therefore we obtain $*Q=3$, because ζ is surjective.

Q. E. D.

From now on, we will construct a torus embedding $T_H \times A_k^4$, any irreducible component of whose fibre over the origin of $\text{Spec } k[Y_{ij}]_{(i,j) \in P_H}$ is isomorphic to C_H . First let \mathcal{R} be a neat system of relations with respect to H and $\{a_1, a_2, a_3, a_4\}$, i. e., $\alpha_i a_i = \sum_{j \neq i} \alpha_{ij} a_j$ for $1 \leq i \leq 4$ and $\alpha_j = \sum_{i \neq j} \alpha_{ij}$ for $1 \leq j \leq 4$, with $0 \leq \alpha_{ij} < \alpha_j$, and let $Y_{ij}, (i, j) \in P_{\mathcal{R}}$, (resp. t_1, \dots, t_{m-3}) be independent variables over k where we put $m = *P_{\mathcal{R}}$. Q denotes the set of $(i, j) \in P_{\mathcal{R}}$ which has either a ν -relation or an h -relation. For brevity, we put $P = P_{\mathcal{R}}$, and let the order on Q (resp. $P-Q$) be induced by that on P defined in Definition 3.3. Then by Lemma 4.1 2) the set Q consists of three elements

$$(i', j') < (i'', j'') < (i_0, j_0),$$

and there exists a unique isomorphism $\xi: P-Q \rightarrow [1, m-3]$ of ordered sets. Now we will define a k -algebra homomorphism

$$\pi: k[Y_{ij}]_{(i,j) \in P} \longrightarrow k[t_1^{\pm 1}, \dots, t_{m-3}^{\pm 1}]$$

inductively as follows:

- 1) $\pi_1: k[Y_{ij}]_{(i,j) \in P < (i', j')} \rightarrow k[t_1^{\pm 1}, \dots, t_{m-3}^{\pm 1}]$ is defined by

$$\pi_1(Y_{ij}) = t_{\xi(ij)} \quad \text{if } (i, j) < (i', j'),$$

$$\pi_1(Y_{i'j'}) = \begin{cases} \prod_{i \in P_{j'-(i')}} t_{\xi(ij')}^{-1} \prod_{j \in P_{j'}} t_{\xi(i'j')} & \text{if } (i', j') \text{ has a } \nu\text{-relation,} \\ \prod_{j \in P_{i'-(j')}} t_{\xi(i'j')}^{-1} \prod_{i \in P_{i'}} t_{\xi(i'j')} & \text{if } (i', j') \text{ has an } h\text{-relation,} \end{cases}$$

and

$$\pi_1(Y_{ij}) = t_{\xi(ij)} \quad \text{if } (i', j') < (i, j) < (i'', j''),$$

- 2) $\pi_2: k[Y_{ij}]_{(i,j) \in P < (i_0, j_0)} \rightarrow k[t_1^{\pm 1}, \dots, t_{m-3}^{\pm 1}]$ is defined by

$$\pi_2(Y_{ij}) = \pi_1(Y_{ij}) \quad \text{if } (i, j) < (i'', j''),$$

$$\pi_2(Y_{i''j''}) = \begin{cases} \prod_{i \in P_{j''-(i')}} \pi_1(Y_{ij''})^{-1} \prod_{j \in P_{j''}} \pi_1(Y_{j''j}) & \text{if } (i'', j'') \text{ has a } v\text{-relation,} \\ \prod_{j \in P_{i''-(j')}} \pi_1(Y_{i''j})^{-1} \prod_{i \in P_{i''}} \pi_1(Y_{ii'}) & \text{if } (i'', j'') \text{ has an } h\text{-relation,} \end{cases}$$

and

$$\pi_2(Y_{ij}) = t_{\xi(ij)} \quad \text{if } (i'', j'') < (i, j) < (i_0, j_0),$$

3) $\pi : k[Y_{ij}]_{(i,j) \in P} \rightarrow k[t_1^{\pm 1}, \dots, t_{m-3}^{\pm 1}]$ is defined by

$$\pi(Y_{ij}) = \pi_2(Y_{ij}) \quad \text{if } (i, j) < (i_0, j_0)$$

and

$$\pi(Y_{i_0j_0}) = \prod_{i \in P_{j_0-(i_0)}} \pi_2(Y_{ij_0})^{-1} \prod_{j \in P_{j_0}} \pi_2(Y_{j_0j}).$$

We note that

$$\prod_{i \in P_{j_0-(i_0)}} \pi_2(Y_{ij_0})^{-1} \prod_{j \in P_{j_0}} \pi_2(Y_{j_0j}) = \prod_{j \in P_{i_0-(j_0)}} \pi_2(Y_{i_0j})^{-1} \prod_{i \in P_{i_0}} \pi_2(Y_{ii_0}).$$

DEFINITION 4.2. If we canonically identify $k[t_1^{\pm 1}, \dots, t_{m-3}^{\pm 1}]$ with the semigroup k -algebra $k[T^b]_{b \in \mathbb{Z}^{m-3}}$, in the above situation for any $(i, j) \in P$ there exists a unique $b_{ij} \in \mathbb{Z}^{m-3}$ such that $\pi(Y_{ij}) = T^{b_{ij}}$. Then the subsemigroup S of \mathbb{Z}^{m-3} generated by $b_{ij} ((i, j) \in P)$ is called the *semigroup associated to P* and the surjective k -algebra homomorphism $\pi : k[Y_{ij}]_{(i,j) \in P} \rightarrow k[T^s]_{s \in S}$ is called the *homomorphism associated to P* .

LEMMA 4.3. Let $\eta : k[Y_{ij}]_{(i,j) \in P} \rightarrow k[X] = k[X_1, X_2, X_3, X_4]$ be the k -algebra homomorphism defined by sending Y_{ij} to $X_1^{a_i} X_2^{a_j}$. Then we have $I_H \cong \eta(\text{Ker } \pi)$.

PROOF. The k -algebra homomorphism $\zeta' : k[T^{b_{ij}}]_{(i,j) \in P} \rightarrow k[t^h]_{h \in H}$ defined by $\zeta'(T^{b_{ij}}) = t^{\alpha_i \alpha_j}$ extends uniquely to the k -algebra homomorphism $\zeta : k[T^s]_{s \in S} \rightarrow k[t^h]_{h \in H}$. Moreover,

$$\varphi_H \circ \eta(Y_{ij}) = \varphi_H(X_1^{a_i} X_2^{a_j}) = t^{\alpha_i \alpha_j}$$

and

$$\zeta \circ \pi(Y_{ij}) = \zeta(T^{b_{ij}}) = t^{\alpha_i \alpha_j},$$

hence $\varphi_H \circ \eta = \zeta \circ \pi$, which implies $I_H = \text{Ker } \varphi_H \cong \eta(\text{Ker } \pi)$.

Q. E. D.

Let us recall the definition of P_H in Definition 3.6 which is determined by a neat numerical semigroup H . In our case $M(H) = \{a_1, a_2, a_3, a_4\}$, elementary computations show the following:

PROPOSITION 4.4. P_H is one of the following:

- (1) the case $*P_H = 12$, then $P_H = S_4 = \{(i, j) \in [1, 4]^2 \mid i \neq j\}$,
- (2) the case $*P_H = 11$, then $P_H = S_4 - \{(1, 2)\}$,

(3) the case $*P_H=10$, then $P_H=S_4-\{(1, 2)\} \cup G$ where G is one of the following:

a) $\{(2, 1)\}$, b) $\{(2, 3)\}$, c) $\{(3, 4)\}$,

(4) the case $*P_H=9$, then $P_H=S_4-\{(1, 2)\} \cup G$ where G is one of the following:

a) $\{(2, 1), (3, 4)\}$, b) $\{(2, 3), (3, 1)\}$, c) $\{(2, 3), (3, 4)\}$,

(5) the case $*P_H=8$, then $P_H=S_4-\{(1, 2)\} \cup G$ where G is one of the following:

a) $\{(2, 1), (3, 4), (4, 3)\}$ and b) $\{(2, 3), (3, 4), (4, 1)\}$.

DEFINITION-PROPOSITION 4.5. Let S_H be the semigroup associated to P_H . Then the subsemigroup S_H of \mathbf{Z}^{m-3} is saturated and generates \mathbf{Z}^{m-3} as a group. Therefore $T_H = \text{Spec } k[Y_{ij}]_{(i,j) \in P_H} / \text{Ker } \pi$, which is isomorphic to $\text{Spec } k[T^s]_{s \in S_H}$, is called the *torus embedding associated to the neat numerical semigroup H* with $M(H) = \{a_1, a_2, a_3, a_4\}$.

PROOF. By the construction of S_H , S_H generates \mathbf{Z}^{m-3} as a group. For any $i \in [1, m-3]$ we denote by $e_i \in \mathbf{Z}^{m-3}$ the vector whose i -th component equals to 1 and whose j -th component equals to 0 if $j \neq i$. Let

$$\sigma : [1, m] \longrightarrow P_H = \{(i, j) \in [1, 4]^2 \mid i \neq j, \alpha_{ij} \neq 0\}$$

be the isomorphism of ordered sets, and for brevity we set $b_i = b_{\sigma(i)}$ for all $i \in [1, m]$. Let the situation be as in Proposition 4.4. Then

(1) $b_i = e_i$ ($1 \leq i \leq 8$), $b_9 = (-1, 1, 1, 1, -1, 0, 0, 0, 0)$, $b_{10} = (1, -1, 0, 0, 0, -1, 1, 1, 0)$, $b_{11} = e_9$, $b_{12} = (0, 0, 1, 0, -1, -1, 1, 0, 1)$,

(2) $b_i = e_i$ ($1 \leq i \leq 7$), $b_8 = (-1, 1, 0, 0, 0, 1, -1, 0)$, $b_9 = (-1, 0, 1, 1, -1, 0, 0, 0)$, $b_{10} = e_8$, $b_{11} = (0, -1, 1, 0, -1, 0, 1, 1)$,

(3) a) $b_i = e_i$ ($1 \leq i \leq 4$), $b_5 = (-1, 0, 1, 1, 0, 0, 0)$, $b_6 = e_5$, $b_7 = e_6$, $b_8 = (0, 1, 0, 0, 1, -1, 0)$, $b_9 = e_7$, $b_{10} = (-1, -1, 1, 0, 0, 1, 1)$,

b) $b_i = e_i$ ($1 \leq i \leq 7$), $b_8 = (-1, 1, 0, 0, 0, 1, 0)$, $b_9 = (-1, 0, 1, 1, -1, 0, 0)$, $b_{10} = (0, -1, 1, 0, -1, 0, 1)$,

c) $b_i = e_i$ ($1 \leq i \leq 7$), $b_8 = (-1, 1, 0, 0, 0, 1, -1)$, $b_9 = (-1, 0, 1, 1, -1, 0, 0)$, $b_{10} = (0, 1, -1, 0, 1, 0, -1)$,

(4) a) $b_i = e_i$ ($1 \leq i \leq 4$), $b_5 = (-1, 0, 1, 1, 0, 0)$, $b_6 = e_5$, $b_7 = e_6$, $b_8 = (0, 1, 0, 0, 1, -1)$, $b_9 = (1, 1, -1, 0, 0, -1)$,

b) $b_i = e_i$ ($1 \leq i \leq 4$), $b_5 = (-1, 0, 1, 1, 0, 0)$, $b_6 = e_5$, $b_7 = e_6$, $b_8 = (-1, 1, 0, 0, 1, 0)$, $b_9 = (0, -1, 1, 0, 0, 1)$,

c) $b_i = e_i$ ($1 \leq i \leq 6$), $b_7 = (0, 1, -1, 0, 1, 0)$, $b_8 = (-1, 1, 0, 0, 0, 1)$, $b_9 = (-1, 0, 1, 1, -1, 0)$,

(5) a) $b_i = e_i$ ($1 \leq i \leq 4$), $b_5 = (-1, 0, 1, 1, 0)$, $b_6 = e_5$, $b_7 = (1, 1, -1, 0, 0)$,
 $b_8 = (-1, 0, 1, 0, 1)$,

b) $b_i = e_i$ ($1 \leq i \leq 4$), $b_5 = (-1, 0, 1, 1, 0)$, $b_6 = e_5$, $b_7 = (-1, 1, 0, 1, 0)$,
 $b_8 = (-1, 1, 0, 0, 1)$.

By computation the subsemigroups S_H of \mathbf{Z}^{m-3} generated by b_1, \dots, b_m are saturated. For example, we check the case (4) c). It suffices to show that $\sum_{i=1}^9 \mathbf{R}_+ b_i \cap \mathbf{Z}^6 \subseteq S_H$ where \mathbf{R}_+ is the set of non-negative real numbers. Let us take $z = \sum_{i=1}^9 \lambda_i b_i \in \mathbf{Z}^6$ with $\lambda_i \in \mathbf{R}_+$, and set $\lambda_i = m_i + \beta_i$ with $m_i \in \mathbf{N}$ and $0 \leq \beta_i < 1$ for $1 \leq i \leq 9$. Hence it suffices to show that $y = \sum_{i=1}^9 \beta_i b_i \in S_H$. Now we get

$$y = (\beta_1 - \beta_8 - \beta_9, \beta_2 + \beta_7 + \beta_8, \beta_3 - \beta_7 + \beta_9, \beta_4 + \beta_9, \beta_5 + \beta_7 - \beta_9, \beta_6 + \beta_8) \in \mathbf{Z}^6,$$

hence

$$\beta_1 - \beta_8 - \beta_9 = -1 \text{ or } 0, \beta_2 + \beta_7 + \beta_8 = 0 \text{ or } 1 \text{ or } 2, \beta_3 - \beta_7 + \beta_9 = 0 \text{ or } 1,$$

$$\beta_4 + \beta_9 = 0 \text{ or } 1, \beta_5 + \beta_7 - \beta_9 = 0 \text{ or } 1, \text{ and } \beta_6 + \beta_8 = 0 \text{ or } 1.$$

First assume $\beta_1 - \beta_8 - \beta_9 = 0$. Since $e_i \in S_H$ for all $1 \leq i \leq 6$, we get $y \in S_H$. Secondly assume $\beta_1 - \beta_8 - \beta_9 = -1$. Then we have $\beta_8 > 0$ and $\beta_9 > 0$, which imply $\beta_2 + \beta_7 + \beta_8 = 1$ or 2 , $\beta_4 + \beta_9 = 1$ and $\beta_6 + \beta_8 = 1$. Then $y \in S_H$, because $(-1, 1, 0, 1, 0, 1) = b_4 + b_8 \in S_H$. Therefore S_H is saturated. The other cases work similarly.

Q. E. D.

For our purposes it is necessary to investigate generators of the ideal I_H . When H is a neat numerical semigroup with $M(H) = \{a_1, a_2, a_3, a_4\}$, the following Lemma gives us a set of generators for I_H .

LEMMA 4.6. *Let H be a numerical semigroup with $M(H) = \{a_1, a_2, a_3, a_4\}$, such that for any $1 \leq i \leq 4$*

$$\alpha_i a_i = \alpha_{ij} a_j + \alpha_{ik} a_k + \alpha_{il} a_l \text{ with } \alpha_{ij} > 0, \alpha_{ik} > 0 \text{ and } \alpha_{il} \geq 0$$

where i, j, k and l are distinct. For any $1 \leq i \leq 4$ we denote $X_i^{\alpha_i} - X_j^{\alpha_{ij}} X_k^{\alpha_{ik}} X_l^{\alpha_{il}}$ by f_i . Set

$$A_1 = \{f_1, f_2, f_3, f_4\}, \quad A_2 = \{X_1^{\beta_1} X_2^{\beta_2} - X_3^{\beta_3} X_4^{\beta_4} \in I_H \mid 0 < \beta_i < \alpha_i\},$$

$$A_3 = \{X_1^{\beta_1} X_3^{\beta_3} - X_2^{\beta_2} X_4^{\beta_4} \in I_H \mid 0 < \beta_i < \alpha_i\}, \quad A_4 = \{X_1^{\beta_1} X_4^{\beta_4} - X_2^{\beta_2} X_3^{\beta_3} \in I_H \mid 0 < \beta_i < \alpha_i\}.$$

Moreover, for any $2 \leq i \leq 4$ we put

$$A_i^* = \{X_1^{\beta_1} X_i^{\beta_i} - X_j^{\beta_j} X_k^{\beta_k} \in A_i \mid \text{for any } X_1^{\gamma_1} X_i^{\gamma_i} - X_j^{\gamma_j} X_k^{\gamma_k} \in A_i, \text{ different from } X_1^{\beta_1} X_i^{\beta_i} - X_j^{\beta_j} X_k^{\beta_k}, \gamma_1 \leq \beta_1 \text{ and } \gamma_i \leq \beta_i \text{ do not hold}\}.$$

Then 1) the ideal I_H is generated by the elements of the set $A_1 \cup A_2^* \cup A_3^* \cup A_4^*$,
 2) if $\alpha_i a_i \neq \alpha_j a_j$ for $i \neq j$, then $\mu(H)$ is equal to $4 + {}^*A_2^* + {}^*A_3^* + {}^*A_4^*$.

PROOF. 1) Let (A') (resp. (A) , resp. (A^*)) be the ideal generated by the set
 $A' = A_1 \cup \{X_i^{\gamma_i} X_j^{\gamma_j} - X_k^{\gamma_k} X_l^{\gamma_l} \in I_H \mid \gamma_i, \gamma_j, \gamma_k, \gamma_l > 0 \text{ and } (i, j, k, l)$
 is a permutation of $[1, 4]\}$

(resp. the set $A = A_1 \cup A_2 \cup A_3 \cup A_4$, resp. the set $A^* = A_1 \cup A_2^* \cup A_3^* \cup A_4^*$).

First we show: $I_H = (A')$, that is, $g = X_i^{\lambda_i} - X_j^{\nu_j} X_k^{\nu_k} X_l^{\nu_l} \in I_H$, with $\lambda_i \geq \alpha_i$ and a permutation (i, j, k, l) of $[1, 4]$, belongs to (A') , i.e., $g = f + \left(\prod_{s=1}^4 X_s^{\gamma_s}\right)h$ with $f \in (A')$ and $\partial h < \partial g$ if $h \neq 0$. If we set $\lambda_i = \alpha_i q + r$ with $q > 0$ and $0 \leq r < \alpha_i$, then

$$G = g - X_i^r (X_i^{\alpha_i q} - X_j^{\alpha_j q} X_k^{\alpha_k q} X_l^{\alpha_l q}) = X_i^r X_j^{\alpha_j q} X_k^{\alpha_k q} X_l^{\alpha_l q} - X_j^{\nu_j} X_k^{\nu_k} X_l^{\nu_l}.$$

Then we can write $G = f + \left(\prod_{s=1}^4 X_s^{\gamma_s}\right)h$ with $f \in (A')$ and $\partial h < \partial g$ if $h \neq 0$.

Secondly we see: $I_H = (A)$, that is, $g = X_i^{\gamma_i} X_j^{\gamma_j} - X_k^{\gamma_k} X_l^{\gamma_l} \in I_H$, with $\gamma_i, \gamma_j, \gamma_k, \gamma_l > 0$ and a permutation (i, j, k, l) of $[1, 4]$, belongs to (A) . We may assume that $\gamma_i = \alpha_i q + r$ with $q > 0$ and $0 \leq r < \alpha_i$. Hence we have

$$G = g - X_i^r X_j^{\gamma_j} (X_i^{\alpha_i q} - X_j^{\alpha_j q} X_k^{\alpha_k q} X_l^{\alpha_l q}) = X_i^r X_j^{\gamma_j + \alpha_j q} X_k^{\alpha_k q} X_l^{\alpha_l q} - X_i^r X_j^{\gamma_j} X_k^{\alpha_k q} X_l^{\alpha_l q}.$$

Then we can write $G = \left(\prod_{s=1}^4 X_s^{\gamma_s}\right)h$ with $\partial h < \partial g$ if $h \neq 0$.

Lastly we check: $I_H = (A^*)$. Let us take $g = X_1^{\gamma_1} X_i^{\gamma_i} - X_j^{\gamma_j} X_k^{\gamma_k} \in A_i$ such that there exists $g_i = X_1^{\beta_1} X_i^{\beta_i} - X_j^{\beta_j} X_k^{\beta_k} \in A_i^*$ with $\gamma_1 \geq \beta_1$, $\gamma_i \geq \beta_i$ and $(\gamma_1, \gamma_i) \neq (\beta_1, \beta_i)$. Then

$$G = g - X_1^{\gamma_1 - \beta_1} X_i^{\gamma_i - \beta_i} g_i = X_1^{\gamma_1 - \beta_1} X_i^{\gamma_i - \beta_i} X_j^{\beta_j} X_k^{\beta_k} - X_j^{\gamma_j} X_k^{\gamma_k} = X_j^{\gamma_j} X_k^{\gamma_k} \cdot h$$

with $\partial h < \partial g$.

2) It suffices to show that the images of elements of $A_1 \cup A_2^* \cup A_3^* \cup A_4^*$ in $I_H / (X_1, X_2, X_3, X_4)I_H$ are linearly independent over k . By the assumptions $\alpha_i a_i \neq \alpha_j a_j$ and the minimality of α_i , the weights of elements of $A_1 \cup A_2 \cup A_3 \cup A_4$ are distinct. For brevity, the ideal (X_1, X_2, X_3, X_4) (resp. $X_1^{\beta_1} X_i^{\beta_i} - X_j^{\beta_j} X_k^{\beta_k} \in A_i^*$) is denoted by (X) (resp. $g_{\beta_1^i \beta_i}$). Let

$$\sum_{i=1}^4 c_i f_i + \sum c_{\beta_1^2 \beta_2}^{(2)} g_{\beta_1^2 \beta_2}^{(2)} + \sum c_{\beta_1^3 \beta_3}^{(3)} g_{\beta_1^3 \beta_3}^{(3)} + \sum c_{\beta_1^4 \beta_4}^{(4)} g_{\beta_1^4 \beta_4}^{(4)} \in (X)I_H,$$

with $c_i, c_{\beta_1^2 \beta_2}^{(2)}, c_{\beta_1^3 \beta_3}^{(3)}, c_{\beta_1^4 \beta_4}^{(4)} \in k$. First assume that $c_i \neq 0$. Since the ideal $(X)I_H$ is homogeneous, we get $c_i f_i \in (X)I_H$, which has an expression:

$$c_i f_i = \sum_{m=1}^4 h_m f_m + \sum h_{\beta_1^2 \beta_2}^{(2)} g_{\beta_1^2 \beta_2}^{(2)} + \sum h_{\beta_1^3 \beta_3}^{(3)} g_{\beta_1^3 \beta_3}^{(3)} + \sum h_{\beta_1^4 \beta_4}^{(4)} g_{\beta_1^4 \beta_4}^{(4)}$$

with $h_m, h_{\beta_1\beta_2}^{(2)}, h_{\beta_1\beta_3}^{(3)}, h_{\beta_1\beta_4}^{(4)} \in (X)$. If we substitute 0 for X_j , all j different from i , then we get $c_i X_i^{\alpha_i} = c X_i^{\beta_i + \alpha_i}$ with $c \in k$ and $\beta_i > 0$, a contradiction. Hence $c_i = 0$ for all $i = 1, \dots, 4$. Secondly assume that $c_{\beta_1\beta_i}^{(i)} \neq 0$. Then $c_{\beta_1\beta_i}^{(i)} g_{\beta_1\beta_i}^{(i)} \in (X)I_H$, which has an expression :

$$c_{\beta_1\beta_i}^{(i)} g_{\beta_1\beta_i}^{(i)} = \sum h_{\beta_1\beta_2}^{(2)} g_{\beta_1\beta_2}^{(2)} + \sum h_{\beta_1\beta_3}^{(3)} g_{\beta_1\beta_3}^{(3)} + \sum h_{\beta_1\beta_4}^{(4)} g_{\beta_1\beta_4}^{(4)}$$

because of $g_{\beta_1\beta_i}^{(i)} \in A_i$ and the minimality of α_j . Substituting 0 for X_j and X_k , where $(1, i, j, k)$ is a permutation of $[1, 4]$, we obtain

$$c_{\beta_1\beta_i}^{(i)} X_1^{\beta_1} X_i^{\beta_i} = \sum_{(\gamma_1, \gamma_i) \neq (\beta_1, \beta_i)} h_{\gamma_1\gamma_i}^{(i)}(X_1, 0, X_i, 0) X_1^{\gamma_1} X_i^{\gamma_i},$$

hence there exists $(\lambda_1, \lambda_i) \in \mathbb{N}^2, \neq (0, 0)$ such that

$$\beta_1 a_1 + \beta_i a_i = (\gamma_1 + \lambda_1) a_1 + (\gamma_i + \lambda_i) a_i.$$

If $\beta_1 \geq \gamma_1 + \lambda_1$, in virtue of $\alpha_1 > \beta_1$ we have $\beta_1 = \gamma_1 + \lambda_1$ and $\beta_i = \gamma_i + \lambda_i$, which contradict $g_{\beta_1\beta_i}^{(i)} \in A_i^*$. If $\beta_1 < \gamma_1 + \lambda_1$, we have

$$(\beta_i - \gamma_i - \lambda_i) a_i = (\gamma_1 + \lambda_1 - \beta_1) a_1,$$

which contradicts the minimality of α_i . Hence we get $c_{\beta_1\beta_i}^{(i)} = 0$. *Q. E. D.*

For a neat system $\mathcal{R} : \alpha_i a_i = \sum \alpha_{ij} a_j$ for $1 \leq i \leq 4$ and $\alpha_j = \sum \alpha_{ij}$ for $1 \leq j \leq 4$, of relations with respect to H with $M(H) = \{a_1, a_2, a_3, a_4\}$, the following holds:

LEMMA 4.7. *We have*

$$D = \begin{vmatrix} \alpha_1 & -\alpha_{12} & -\alpha_{13} \\ -\alpha_{21} & \alpha_2 & -\alpha_{23} \\ -\alpha_{31} & -\alpha_{32} & \alpha_3 \end{vmatrix} > 0.$$

PROOF. Since we have $\alpha_j = \sum_{i \neq j} \alpha_{ij}$ for $1 \leq j \leq 4$, we obtain

$$\begin{aligned} D &= \begin{vmatrix} \alpha_1 & -\alpha_{12} & -\alpha_{13} \\ -\alpha_{21} & \alpha_2 & -\alpha_{23} \\ \alpha_{41} & \alpha_{42} & \alpha_{43} \end{vmatrix} = \alpha_{41} \begin{vmatrix} -\alpha_{12} & -\alpha_{13} \\ \alpha_2 & -\alpha_{23} \end{vmatrix} - \alpha_{42} \begin{vmatrix} \alpha_1 & -\alpha_{13} \\ -\alpha_{21} & -\alpha_{23} \end{vmatrix} \\ &\quad + \alpha_{43} \begin{vmatrix} \alpha_1 & -\alpha_{12} \\ -\alpha_{21} & \alpha_2 \end{vmatrix} \\ &= \alpha_{41}(\alpha_{12}\alpha_{23} + \alpha_2\alpha_{13}) + \alpha_{42}(\alpha_1\alpha_{23} + \alpha_{21}\alpha_{13}) + \alpha_{43}\{\alpha_1(\alpha_{32} + \alpha_{42}) + (\alpha_{31} + \alpha_{41})\alpha_{12}\} \end{aligned}$$

If $\alpha_{43} > 0$, then $D > 0$ because of $\alpha_{43}\alpha_1(\alpha_{32} + \alpha_{42}) > 0$. If $\alpha_{43} = 0$, then $\alpha_{41} > 0$ and $\alpha_{13} > 0$, hence we get $D > 0$. *Q. E. D.*

Hereafter we are in the following situation, which is similar to that in Corollary 1.6: let $P=P_H$ be as in Definition 3.6 and let $T_H=\text{Spec } k[Y_{ij}]_{(i,j)\in P}/\text{Ker } \pi$ be the torus embedding associated to the neat numerical semigroup H with $M(H)=\{a_1, a_2, a_3, a_4\}$. Let us consider the fibre product:

$$\begin{array}{ccc} \phi^{-1}(O) & \longrightarrow & T_H \times A_k^4 \cong \text{Spec}(k[Y_{ij}]/J)[X_1, X_2, X_3, X_4] \\ \downarrow & & \downarrow \phi \\ \text{Spec } k & \longrightarrow & \text{Spec } k[Y_{ij}]_{(i,j)\in P} \end{array}$$

where O and J are respectively the origin of $\text{Spec } k[Y_{ij}]$ and the ideal $\text{Ker } \pi$, and ϕ is the morphism corresponding to the k -algebra homomorphism $\phi^*: k[Y_{ij}] \rightarrow (k[Y_{ij}]/J)[X_1, X_2, X_3, X_4]$ by sending Y_{ij} to $X_j^{a_{ij}} - Y_{ij} \pmod J$. If J_0 is the ideal in $k[X]=k[X_1, X_2, X_3, X_4]$ generated by the set $\eta(J)$ where $\eta: k[Y_{ij}] \rightarrow k[X]$ is the k -algebra homomorphism defined by $\eta(Y_{ij})=X_j^{a_{ij}}$, then $\phi^{-1}(O)$ is isomorphic to $\text{Spec } k[X]/J_0$.

PROPOSITION 4.8. C_H is an irreducible component in $\phi^{-1}(O)=\text{Spec } k[X]/J_0$.

PROOF. We use the notation in Lemma 4.6. Since

$$F_i = \prod_{j \in P^i} Y_{ji} - \prod_{j \in P^i} Y_{ij} \in J$$

for all i implies $(f_1, f_2, f_3, f_4) \subseteq J_0$ and by Lemma 4.3 we have $I_H \supseteq J_0$, we will check that the ideal I_H is minimal prime over (f_1, f_2, f_3, f_4) . Let \mathfrak{p} be any prime ideal in $k[X]$ with $(f_1, f_2, f_3, f_4) \subseteq \mathfrak{p} \subseteq I_H$. Let us take

$$g = X_1^{\beta_1} X_2^{\beta_2} - X_3^{\beta_3} X_4^{\beta_4} \in A_2, \quad \text{hence } \beta_1 a_1 + \beta_2 a_2 - \beta_3 a_3 = \beta_4 a_4.$$

By Lemma 4.7, there exists a positive integer μ such that

$$\mu(\beta_1, \beta_2, -\beta_3) = \nu_1(\alpha_1, -\alpha_{12}, -\alpha_{13}) + \nu_2(-\alpha_{21}, \alpha_2, -\alpha_{23}) + \nu_3(-\alpha_{31}, -\alpha_{32}, \alpha_3)$$

with $\nu_i \in \mathbb{Z}$, which implies $\mu\beta_4 = \nu_1\alpha_{14} + \nu_2\alpha_{24} + \nu_3\alpha_{34}$. Since $\beta_i > 0$ for $1 \leq i \leq 4$, this case is divided into the following:

- 1) $\nu_1 > 0, \nu_2 > 0, \nu_3 \geq 0,$ 2) $\nu_1 > 0, \nu_2 > 0, \nu_3 < 0,$
- 3) $\nu_1 > 0, \nu_2 < 0, \nu_3 < 0,$ 4) $\nu_1 \leq 0, \nu_2 > 0, \nu_3 < 0.$

If $\nu_1 > 0, \nu_2 > 0$ and $\nu_3 \geq 0$, then

$$\begin{aligned} & X_1^{\nu_2\alpha_{21} + \nu_3\alpha_{31}} X_2^{\nu_1\alpha_{12} + \nu_3\alpha_{32}} X_3^{\nu_3\alpha_3} (X_1^{\mu\beta_1} X_2^{\mu\beta_2} - X_3^{\mu\beta_3} X_4^{\mu\beta_4}) \\ &= X_2^{\nu_2\alpha_2} X_3^{\nu_3\alpha_3} (X_1^{\nu_1\alpha_1} - X_2^{\nu_1\alpha_{12}} X_3^{\nu_1\alpha_{13}} X_4^{\nu_1\alpha_{14}}) \\ &\quad + X_2^{\nu_1\alpha_{12}} X_3^{\nu_1\alpha_{13} + \nu_3\alpha_3} X_4^{\nu_1\alpha_{14}} (X_2^{\nu_2\alpha_2} - X_1^{\nu_2\alpha_{21}} X_3^{\nu_2\alpha_{23}} X_4^{\nu_2\alpha_{24}}) \\ &\quad + X_1^{\nu_2\alpha_{21}} X_2^{\nu_1\alpha_{12}} X_3^{\nu_1\alpha_{13} + \nu_2\alpha_{23}} X_4^{\nu_1\alpha_{14} + \nu_2\alpha_{24}} (X_3^{\nu_3\alpha_3} - X_1^{\nu_3\alpha_{31}} X_2^{\nu_3\alpha_{32}} X_4^{\nu_3\alpha_{34}}) \\ &\in (f_1, f_2, f_3) \subseteq \mathfrak{p} \subseteq I_H. \end{aligned}$$

Since

$$X_1^{\nu_2\alpha_{21}+\nu_3\alpha_{31}}X_2^{\nu_1\alpha_{12}+\nu_3\alpha_{32}}X_3^{\nu_3\alpha_{33}}(X_1^{(\mu-1)\beta_1}X_2^{(\mu-1)\beta_2}+\dots+X_3^{(\mu-1)\beta_3}X_4^{(\mu-1)\beta_4})\in I_H,$$

we get $g=X_1^{\beta_1}X_2^{\beta_2}-X_3^{\beta_3}X_4^{\beta_4}\in\mathfrak{p}$. The other cases work similarly. For $g\in A_3\cup A_4$, the proof of $g\in\mathfrak{p}$ is similar. By Lemma 4.6 \mathfrak{p} coincides with I_H , hence we get our desired result. Q. E. D.

If $\phi^{-1}(O)$ and C_H are respectively regarded as the algebraic subsets $V(J_0)$ and $V(I_H)$ of the affine space A_k^4 , we see:

PROPOSITION 4.9. 1) For any $x=(x_1, x_2, x_3, x_4)\in\phi^{-1}(O)$, different from the origin, we have $x_i\neq 0$ for any $1\leq i\leq 4$.

2) For any $x=(x_1, x_2, x_3, x_4)\in\phi^{-1}(O)$, different from the origin, we have $x^{-1}=(x_1^{-1}, x_2^{-1}, x_3^{-1}, x_4^{-1})\in\phi^{-1}(O)$.

3) Any irreducible component in $\phi^{-1}(O)$ is isomorphic to C_H .

PROOF. In the proof we use the notation in Lemma 4.6.

1) If $x_i=0$ for some i , x must be the origin of A_k^4 , because J_0 contains the ideal (f_1, f_2, f_3, f_4) .

2) We may take generators $F_k(1\leq k\leq u)$ of the ideal J as follows:

$$F_k = \prod_{(i,j)\in P} Y_{ij}^{\nu_{ij}} - \prod_{(i,j)\in P} Y_{ij}^{\mu_{ij}}$$

with $\nu_{ij}\mu_{ij}=0$. In virtue of $x\in\phi^{-1}(O)=V(J_0)=V(\eta(J))$, we have

$$\prod x_j^{\nu_{ij}\alpha_{ij}} - \prod x_j^{\mu_{ij}\alpha_{ij}} = 0,$$

which implies

$$\prod (x_j^{-1})^{\nu_{ij}\alpha_{ij}} - \prod (x_j^{-1})^{\mu_{ij}\alpha_{ij}} = 0.$$

This means $x^{-1}\in\phi^{-1}(O)$.

3) For any $x=(x_1, x_2, x_3, x_4)\in\phi^{-1}(O)$, different from the origin, let $\varphi_x:k[X]\rightarrow k[X]/J_0$ be the k -algebra homomorphism defined by $\varphi_x(X_i)=x_iX_i+J_0$. Then $\text{Ker } \varphi_x$ contains the ideal J_0 , because

$$\begin{aligned} \varphi_x(\eta(F_k)) &= \prod (x_jX_j)^{\alpha_{ij}\nu_{ij}} - \prod (x_jX_j)^{\alpha_{ij}\mu_{ij}} + J_0 \\ &= \prod x_j^{\alpha_{ij}\nu_{ij}} (\prod (X_j)^{\alpha_{ij}\nu_{ij}} - \prod (X_j)^{\alpha_{ij}\mu_{ij}}) + J_0 \\ &= \prod x_j^{\alpha_{ij}\nu_{ij}} \eta(F_k) + J_0 = J_0. \end{aligned}$$

Therefore φ_x induces the homomorphism $\bar{\varphi}_x:k[X]/J_0\rightarrow k[X]/J_0$, which is an isomorphism by 2). Since J_0 is homogeneous, $\phi^{-1}(O)$ has a natural G_m -action. Then we see that for any $x\in\phi^{-1}(O)$, different from the origin, we have

$$\phi_{x^{-1}}(\text{the closure of } G_m \cdot x) = C_H$$

where $\phi_{x^{-1}}$ is the automorphism of $\phi^{-1}(O)$ corresponding to $\varphi_{x^{-1}}$. Using Proposition 4.8 any irreducible component in $\phi^{-1}(O)$ is isomorphic to C_H . *Q.E.D.*

Lastly, for our purpose we classify neat numerical semigroups H with $M(H) = \{a_1, a_2, a_3, a_4\}$ as follows:

DEFINITION 4.10. In virtue of $(a_1, a_2, a_3, a_4) = 1$ and Lemma 4.7, there exists a unique positive integer ν such that

$$\nu a_4 = \begin{vmatrix} \alpha_1 & -\alpha_{12} & -\alpha_{13} \\ -\alpha_{21} & \alpha_2 & -\alpha_{23} \\ -\alpha_{31} & -\alpha_{32} & \alpha_3 \end{vmatrix} = D.$$

Then the numerical semigroup H is called to be ν -neat.

Our main result in this section is the following:

THEOREM 4.11. *1-neat numerical semigroups H are of torus embedding type, hence if the characteristic of k is 0, then we get $\mathcal{M}_H \neq \emptyset$.*

PROOF. Let the situation be as in Proposition 4.4. Since $a_4 = D$, by computation we get:

(1) $L_{a_3}(H) = \{\beta_1 a_1 + \beta_2 a_2 + \beta_4 a_4 \mid \beta_i \in \mathbb{N} \text{ and } (\beta_1, \beta_2, \beta_4) \text{ satisfies one of the following: 1) } \beta_1 < \alpha_{21} + \alpha_{31}, \beta_2 < \alpha_{32}, \beta_4 < \alpha_4, \text{ 2) } \beta_1 < \alpha_{31}, \alpha_{32} \leq \beta_2 < \alpha_2, \beta_4 < \alpha_{14} + \alpha_{34}, \text{ 3) } \alpha_{21} + \alpha_{31} \leq \beta_1 < \alpha_1, \beta_2 < \alpha_{32}, \beta_4 < \alpha_{24} + \alpha_{34}, \text{ 4) } \alpha_{31} \leq \beta_1 < \alpha_1, \alpha_{32} \leq \beta_2 < \alpha_{32} + \alpha_{42}, \beta_4 < \alpha_{34}, \text{ 5) } \beta_1 < \alpha_{31}, \alpha_{32} \leq \beta_2 < \alpha_{12} + \alpha_{32}, \alpha_{14} + \alpha_{34} \leq \beta_4 < \alpha_4, \text{ 6) } \alpha_{31} \leq \beta_1 < \alpha_{31} + \alpha_{41}, \alpha_{32} + \alpha_{42} \leq \beta_2 < \alpha_2, \beta_4 < \alpha_{34}\}$,

(2) $L_{a_3}(H) = \{\beta_1 a_1 + \beta_2 a_2 + \beta_4 a_4 \mid \beta_i \in \mathbb{N} \text{ and } (\beta_1, \beta_2, \beta_4) \text{ satisfies one of the following: 1) } \beta_1 < \alpha_{21} + \alpha_{31}, \beta_2 < \alpha_{32}, \beta_4 < \alpha_4, \text{ 2) } \beta_1 < \alpha_{31}, \alpha_{32} \leq \beta_2 < \alpha_2, \beta_4 < \alpha_{14} + \alpha_{34}, \text{ 3) } \alpha_{21} + \alpha_{31} \leq \beta_1 < \alpha_1, \beta_2 < \alpha_{32}, \beta_4 < \alpha_{24} + \alpha_{34}, \text{ 4) } \alpha_{31} \leq \beta_1 < \alpha_1, \alpha_{32} \leq \beta_2 < \alpha_2, \beta_4 < \alpha_{34}\}$,

(3) a) $L_{a_3}(H) = \{\beta_1 a_1 + \beta_2 a_2 + \beta_4 a_4 \mid \beta_i \in \mathbb{N} \text{ and } (\beta_1, \beta_2, \beta_4) \text{ satisfies one of the following: 1) } \beta_1 < \alpha_1, \beta_2 < \alpha_2, \beta_4 < \alpha_{34}, \text{ 2) } \beta_1 < \alpha_{31}, \beta_2 < \alpha_{32}, \alpha_{34} \leq \beta_4 < \alpha_4, \text{ 3) } \alpha_{31} \leq \beta_1 < \alpha_1, \beta_2 < \alpha_{32}, \alpha_{34} \leq \beta_4 < \alpha_{24} + \alpha_{34}, \text{ 4) } \beta_1 < \alpha_{31}, \alpha_{32} \leq \beta_2 < \alpha_2, \alpha_{34} \leq \beta_4 < \alpha_{14} + \alpha_{34}\}$,

b) $L_{a_1}(H) = \{\beta_2 a_2 + \beta_3 a_3 + \beta_4 a_4 \mid \beta_i \in \mathbb{N} \text{ and } (\beta_2, \beta_3, \beta_4) \text{ satisfies one of the following: 1) } \beta_2 < \alpha_2, \beta_3 < \alpha_3, \beta_4 < \alpha_{14}, \text{ 2) } \beta_2 < \alpha_2, \beta_3 < \alpha_{13}, \alpha_{14} \leq \beta_4 < \alpha_{14} + \alpha_{34}, \text{ 3) } \beta_2 < \alpha_{32}, \beta_3 < \alpha_{13}, \alpha_{14} + \alpha_{34} \leq \beta_4 < \alpha_4\}$,

c) $L_{a_3}(H) = \{\beta_1 a_1 + \beta_2 a_2 + \beta_4 a_4 \mid \beta_i \in \mathbb{N} \text{ and } (\beta_1, \beta_2, \beta_4) \text{ satisfies one of the following: 1) } \beta_1 < \alpha_{21} + \alpha_{31}, \beta_2 < \alpha_{32}, \beta_4 < \alpha_4, \text{ 2) } \beta_1 < \alpha_{31}, \alpha_{32} \leq \beta_2 < \alpha_2, \beta_4 < \alpha_{14}, \text{ 3) } \alpha_{21} + \alpha_{31} \leq \beta_1 < \alpha_1, \beta_2 < \alpha_{32}, \beta_4 < \alpha_{24}\}$,

(4) a) $L_{\alpha_3}(H) = \{\beta_1 a_1 + \beta_2 a_2 + \beta_4 a_4 \mid \beta_i \in N \text{ and } (\beta_1, \beta_2, \beta_4) \text{ satisfies one of the following: 1) } \beta_1 < \alpha_{31}, \beta_2 < \alpha_{32}, \beta_4 < \alpha_4, \text{ 2) } \beta_1 < \alpha_{31}, \alpha_{32} \leq \beta_2 < \alpha_2, \beta_4 < \alpha_{14}, \text{ 3) } \alpha_{31} \leq \beta_1 < \alpha_1, \beta_2 < \alpha_{32}, \beta_4 < \alpha_{24}\}$,

b) $L_{\alpha_4}(H) = \{\beta_1 a_1 + \beta_2 a_2 + \beta_3 a_3 \mid \beta_i \in N \text{ and } (\beta_1, \beta_2, \beta_3) \text{ satisfies one of the following: 1) } \beta_1 < \alpha_1, \beta_2 < \alpha_2, \beta_3 < \alpha_{43}, \text{ 2) } \beta_1 < \alpha_1, \beta_2 < \alpha_{42}, \alpha_{43} \leq \beta_3 < \alpha_3, \text{ 3) } \beta_1 < \alpha_{41}, \alpha_{42} \leq \beta_2 < \alpha_2, \alpha_{43} \leq \beta_3 < \alpha_3\}$,

c) $L_{\alpha_1}(H) = \{\beta_2 a_2 + \beta_3 a_3 + \beta_4 a_4 \mid \beta_i \in N \text{ and } (\beta_2, \beta_3, \beta_4) \text{ satisfies one of the following: 1) } \beta_2 < \alpha_2, \beta_3 < \alpha_3, \beta_4 < \alpha_{14}, \text{ 2) } \beta_2 < \alpha_{32}, \beta_3 < \alpha_{13}, \alpha_{14} \leq \beta_4 < \alpha_4\}$,

(5) a) $L_{\alpha_1}(H) = \{\beta_2 a_2 + \beta_3 a_3 + \beta_4 a_4 \mid \beta_i \in N \text{ and } (\beta_2, \beta_3, \beta_4) \text{ satisfies one of the following: 1) } \beta_2 < \alpha_2, \beta_3 < \alpha_{13}, \beta_4 < \alpha_{14}, \text{ 2) } \beta_2 < \alpha_{42}, \alpha_{13} \leq \beta_3 < \alpha_3, \beta_4 < \alpha_{14}, \text{ 3) } \beta_2 < \alpha_{32}, \beta_3 < \alpha_{13}, \alpha_{14} \leq \beta_4 < \alpha_4\}$,

b) $L_{\alpha_1}(H) = \{\beta_2 a_2 + \beta_3 a_3 + \beta_4 a_4 \mid \beta_i \in N \text{ and } (\beta_2, \beta_3, \beta_4) \text{ satisfies one of the following: 1) } \beta_2 < \alpha_2, \beta_3 < \alpha_3, \beta_4 < \alpha_{14}, \text{ 2) } \beta_2 < \alpha_{32}, \beta_3 < \alpha_{13}, \alpha_{14} \leq \beta_4 < \alpha_4\}$.

Using the above and Lemma 4.6 we get $J_0 = I_H$. For example, in the case (4) c) we will show that $J_0 = I_H$. It suffices to show that $J_0 \supseteq I_H$. We use the notation in Lemma 4.6. Assume that $A_2 \neq \emptyset$, i. e., take

$$X_1^{\beta_1} X_2^{\beta_2} - X_3^{\beta_3} X_4^{\beta_4} \in A_2, \quad \text{hence } \beta_1 a_1 + \beta_2 a_2 = \beta_3 a_3 + \beta_4 a_4.$$

Then 1) implies $\beta_4 \geq \alpha_{14}$, hence by 2) we get $\beta_3 \geq \alpha_{13}$. Therefore we have

$$\begin{aligned} \beta_1 a_1 + \beta_2 a_2 &= (\beta_3 - \alpha_{13}) a_3 + (\beta_4 - \alpha_{14}) a_4 + \alpha_{13} a_3 + \alpha_{14} a_4 \\ &= (\beta_3 - \alpha_{13}) a_3 + (\beta_4 - \alpha_{14}) a_4 + \alpha_1 a_1, \end{aligned}$$

which implies

$$\beta_2 a_2 = (\alpha_1 - \beta_1) a_1 + (\beta_3 - \alpha_{13}) a_3 + (\beta_4 - \alpha_{14}) a_4.$$

Since $0 < \beta_2 < \alpha_2$, this contradicts the minimality of α_2 , hence $A_2 = \emptyset$, which implies $A_2^* = \emptyset$. Now we have

$$g_3 = X_1^{\alpha_{21} + \alpha_{41}} X_3^{\alpha_{43}} - X_2^{\alpha_{32}} X_4^{\alpha_{14}} \in A_3.$$

Take $X_1^{\beta_1} X_3^{\beta_3} - X_2^{\beta_2} X_4^{\beta_4} \in A_3$, different from g_3 . Then 1) implies $\beta_4 \geq \alpha_{14}$, hence by 2) we get $\beta_2 \geq \alpha_{32}$. Therefore we get

$$A_3^* = \{g_3 = X_1^{\alpha_{21} + \alpha_{41}} X_3^{\alpha_{43}} - X_2^{\alpha_{32}} X_4^{\alpha_{14}}\}.$$

Lastly 1) implies $A_4 = \emptyset$. Hence by Lemma 4.6 the ideal I_H is generated by f_1, f_2, f_3, f_4 and g_3 . Since we have

$$\pi(Y_{21} Y_{41} Y_{43} - Y_{32} Y_{14}) = t_1 t_1^{-1} t_5^{-1} t_3 t_4 t_3^{-1} t_5 t_2 - t_2 t_4 = 0$$

and

$$\eta(Y_{21} Y_{41} Y_{43} - Y_{32} Y_{14}) = X_1^{\alpha_{21} + \alpha_{41}} X_3^{\alpha_{43}} - X_2^{\alpha_{32}} X_4^{\alpha_{14}} = g_3,$$

we get $I_H \subseteq J_0$. The other cases work similarly. Using Lemma 1.2, H is of torus embedding type. Q. E. D.

REMARK 4.12. 1) By calculation, any neat numerical semigroup with $M(H) = \{a_1, a_2, a_3, a_4\}$ and $g(H) \leq 8$ is 1-neat.

2) For a ν -neat numerical semigroup H with $\nu \geq 2$, $\phi^{-1}(O) = \text{Spec } k[X]/J_0$ does not necessarily coincide with $C_H = \text{Spec } k[X]/I_H$. For example, let H be the numerical semigroup with $M(H) = \{10, 11, 14, 13\}$. Then $g(H) = 16$ and H is 2-neat. Using Lemma 4.6, I_H is generated by

$$\begin{aligned} f_1 &= X_1^4 - X_3 X_4^2, & f_2 &= X_2^3 - X_1^2 X_4, & f_3 &= X_3^3 - X_1^2 X_2^2, & f_4 &= X_4^3 - X_2 X_3^2, \\ f_5 &= X_1^3 X_2 - X_3^2 X_4, & f_6 &= X_1 X_3 - X_2 X_4 & \text{and} & & f_7 &= X_1 X_4^2 - X_2^2 X_3, \end{aligned}$$

hence $\mu(H) = 7$. But J_0 is generated by f_1, f_2, f_3, f_4 and $X_1^2 X_3^2 - X_2^2 X_4^2$. More explicitly, as an algebraic subset of A_k^4 we have $V(J_0) \cong V(I_H)$, because $(-1, 1, 1, 1) \in V(J_0) - V(I_H)$.

5. Symmetric numerical semigroups generated by 4 elements.

In this section, we always assume that H is a numerical semigroup with $M(H) = \{a_1, a_2, a_3, a_4\}$. Then using Bresinsky's result [1] we will show that any symmetric H is of torus embedding type, in this case if H is not a complete intersection then it is 1-neat. In the symmetric case, a set of generators for the ideal I_H is given by the following, which is due to Bresinsky:

REMARK 5.1. Let H be symmetric, i. e., $2g(H) = C(H)$.

(1) When H is a complete intersection, renumbering a_1, a_2, a_3, a_4 we may assume that $X_1^{\alpha_1} - X_2^{\alpha_2} \in I_H$.

a) The case $X_3^{\alpha_3} - X_4^{\alpha_4} \in I_H$. Then $(a_1, a_2)(a_3, a_4) \in \langle a_1, a_2 \rangle \cap \langle a_3, a_4 \rangle$, hence we put

$$(a_1, a_2)(a_3, a_4) = \beta_1 a_1 + \beta_2 a_2 = \beta_3 a_3 + \beta_4 a_4.$$

In this case,

$$I_H = (f_1 = X_1^{\alpha_1} - X_2^{\alpha_2}, f_2 = X_3^{\alpha_3} - X_4^{\alpha_4}, f_3 = X_1^{\beta_1} X_2^{\beta_2} - X_3^{\beta_3} X_4^{\beta_4}).$$

b) The case $X_3^{\alpha_3} - X_4^{\alpha_4} \in I_H$. Then H is a strictly complete intersection.

(2) If H is not a complete intersection, renumbering a_1, a_2, a_3, a_4 we have

$$\begin{aligned} I_H &= (f_1 = X_1^{\alpha_1} - X_3^{\alpha_3} X_4^{\alpha_4}, f_2 = X_2^{\alpha_2} - X_1^{\alpha_2} X_4^{\alpha_4}, f_3 = X_3^{\alpha_3} - X_1^{\alpha_3} X_2^{\alpha_2}, \\ & f_4 = X_4^{\alpha_4} - X_2^{\alpha_4} X_3^{\alpha_3}, f_5 = X_1^{\alpha_2} X_3^{\alpha_3} - X_2^{\alpha_2} X_4^{\alpha_4}) \end{aligned}$$

where

$$0 < \alpha_{ij} < \alpha_j, \quad \alpha_1 = \alpha_{21} + \alpha_{31}, \quad \alpha_2 = \alpha_{32} + \alpha_{42}, \quad \alpha_3 = \alpha_{13} + \alpha_{43}, \quad \alpha_4 = \alpha_{14} + \alpha_{24}.$$

In this case,

$$a_1 = \alpha_2 \alpha_3 \alpha_{14} + \alpha_{32} \alpha_{13} \alpha_{24}, \quad a_2 = \alpha_{21} \alpha_3 \alpha_4 + \alpha_{31} \alpha_{43} \alpha_{24}, \quad a_3 = \alpha_1 \alpha_{32} \alpha_4 + \alpha_{31} \alpha_{42} \alpha_{14}$$

and

$$a_4 = \alpha_1 \alpha_2 \alpha_{43} + \alpha_{21} \alpha_{42} \alpha_{13},$$

hence H is 1-neat.

PROPOSITION 5.2. *Any symmetric H is of torus embedding type.*

PROOF. In virtue of Lemma 2.3 and Theorem 4.11, it suffices to show that in the case of Remark 5.1 (1) a) H is of torus embedding type. Renumbering a_1 and a_2 (resp. a_3 and a_4), we may assume that $\beta_1 \neq 0$ and $\beta_3 \neq 0$, hence the following four cases occur:

- 1) $\beta_2 \neq 0$ and $\beta_4 \neq 0$, 2) $\beta_2 \neq 0$ and $\beta_4 = 0$, 3) $\beta_2 = 0$ and $\beta_4 \neq 0$

and

- 4) $\beta_2 = 0$ and $\beta_4 = 0$.

For the case 1), let

$$\pi : k[Z, Y] = k[Z_1, \dots, Z_4, Y_1, \dots, Y_4] \longrightarrow k[t_1^{\pm 1}, \dots, t_5^{\pm 1}]$$

$$(\text{resp. } \eta : k[Z, Y] \longrightarrow k[X] = k[X_1, \dots, X_4])$$

be the k -algebra homomorphism defined by $\pi(Z_i) = t_1$ for $i = 1, 2$, $\pi(Z_j) = t_2$ for $j = 3, 4$, $\pi(Y_k) = t_{2+k}$ for $k = 1, 2, 3$ and $\pi(Y_4) = t_3 t_4 t_5^{-1}$ (resp. $\eta(Z_i) = X_i^{\alpha_i}$ and $\eta(Y_i) = X_i^{\beta_i}$ for $1 \leq i \leq 4$). Then we see easily that $I_H \supseteq \eta(\text{Ker } \pi)$. Moreover, since $F_1 = Z_1 - Z_2$, $F_2 = Z_3 - Z_4$ and $F_3 = Y_1 Y_2 - Y_3 Y_4 \in \text{Ker } \pi$, we have $I_H = (\eta(F_1), \eta(F_2), \eta(F_3))$, which is generated by the set $\eta(\text{Ker } \pi)$. Using Lemma 1.2, H is of torus embedding type. The other cases 2), 3), 4) work similarly. Q. E. D.

6. Almost symmetric numerical semigroups generated by 4 elements.

In the last section we will give another examples of 1-neat numerical semigroups, which are called to be *almost symmetric*, i. e., $C(H) = 2g(H) - 1$. In this section we are devoted to proving that any almost symmetric numerical semigroup H with $M(H) = \{a_1, a_2, a_3, a_4\}$ is 1-neat. First we investigate the properties of almost symmetric H with $M(H) = \{a_1, \dots, a_n\}$.

LEMMA 6.1. *Let H be a numerical semigroup with $M(H) = \{a_1, \dots, a_n\}$ and h be its element.*

0) For any $1 \leq i \leq h$ there exists a unique $1 \leq h_i \leq h$ such that $\omega_h(h) - \omega_h(i) \equiv \omega_h(h_i) \pmod{h}$.

1) H is almost symmetric if and only if there exists a unique $i_0 \in [2, h-1]$ such that $2\omega_h(i_0) = \omega_h(h) + h$ and that $\omega_h(i) + \omega_h(h_i) = \omega_h(h)$ for all $i \neq i_0$.

PROOF. The definition of $L_h(H) = \{\omega_h(1) < \dots < \omega_h(h)\}$ means 0). We see easily :

$$g(H) = \sum_{i=1}^h [\omega_h(i)/h] \quad \text{and} \quad C(H) - g(H) = \sum_{i=1}^h [(\omega_h(h) - \omega_h(i))/h]$$

where $[\]$ is the Gauss symbol. For any $1 \leq i \leq h$ there exists a unique $n_i \in \mathcal{N}$ such that $\omega_h(h) - \omega_h(i) = \omega_h(h_i) - n_i h$. Hence H is almost symmetric if and only if $\sum_{i=1}^h n_i = 1$. This implies 1). Q. E. D.

PROPOSITION 6.2. Let H be an almost symmetric numerical semigroup with $M(H) = \{a_1, \dots, a_n\}$, and let j, k be two distinct element of $[1, n]$ such that $\alpha_j a_j = \sum_{l \neq j} \alpha_{jl} a_l$ with $\alpha_{jk} \geq 1$.

1) If $\alpha_{jk} \geq 2$, then $\omega_{a_k}(a_k) - (\alpha_j - 1)a_j \in L_{a_k}(H)$.

2) We have

$$\omega_{a_k}(a_k) = \begin{cases} \sum_{l \in [1, n] - \{k, j\}} \beta_l a_l + (\alpha_j - 1)a_j & \text{if } \omega_{a_k}(a_k) - (\alpha_j - 1)a_j \in L_{a_k}(H). \\ \sum_{l \in [1, n] - \{k, j\}} \alpha_{jl} a_l + (\alpha_j - 2)a_j & \text{otherwise.} \end{cases}$$

PROOF. 1) Since $(\alpha_j - 1)a_j \in L_{a_k}(H)$, by Lemma 6.1 it suffices to show that

$$(\alpha_j - 1)a_j \neq \omega_{a_k}(i_0) \quad \text{where} \quad 2\omega_{a_k}(i_0) = \omega_{a_k}(a_k) + a_k.$$

Assume $(\alpha_j - 1)a_j = \omega_{a_k}(i_0)$. Then

$$\omega_{a_k}(a_k) + a_k = 2(\alpha_j - 1)a_j = (\alpha_j - 2)a_j + \sum_{l \neq j} \alpha_{jl} a_l.$$

Hence we have

$$\omega_{a_k}(a_k) - a_k = (\alpha_j - 2)a_j + (\alpha_{jk} - 2)a_k + \sum_{l \in [1, n] - \{j, k\}} \alpha_{jl} a_l.$$

This contradicts $\omega_{a_k}(a_k) - a_k \notin H$.

2) In view of $\alpha_{jk} \geq 1$, if $\omega_{a_k}(a_k) - (\alpha_j - 1)a_j \in L_{a_k}(H)$, then

$$\omega_{a_k}(a_k) = \sum_{l \in [1, n] - \{k, j\}} \beta_l a_l + (\alpha_j - 1)a_j.$$

If $\omega_{a_k}(a_k) - (\alpha_j - 1)a_j \notin L_{a_k}(H)$, we have

$$2(\alpha_j - 1)a_j = 2\omega_{a_k}(i_0) = \omega_{a_k}(a_k) + a_k,$$

hence

$$\begin{aligned} \omega_{a_k}(a_k) &= \alpha_j a_j + (\alpha_j - 2)a_j - a_k = \sum_{l \in [1, n] - \{j, k\}} \alpha_{jl} a_l + (\alpha_j - 2)a_j + (\alpha_{jk} - 1)a_k \\ &= \sum_{l \in [1, n] - \{j, k\}} \alpha_{jl} a_l + (\alpha_j - 2)a_j. \end{aligned} \quad Q. E. D.$$

For the remainder of this section we assume that H is a numerical semi-group with $M(H) = \{a_1, a_2, a_3, a_4\}$.

PROPOSITION 6.3. *Let H be almost symmetric and let $k \in [1, 4]$ such that for any $j \in [1, 4]$, different from k , we have $\alpha_j a_j = \sum_{l \neq j} \alpha_{jl} a_l$ with $\alpha_{jk} \geq 1$.*

1) *For any $j \in [1, 4]$, different from k , the following are equivalent:*

a) $\omega_{a_k}(a_k) = \sum_{l \in [1, 4] - \{k, j\}} \beta_l a_l + (\alpha_j - 2)a_j,$

b) $\omega_{a_k}(a_k) - (\alpha_j - 1)a_j \in L_{a_k}(H).$

In this case, $\alpha_{jk} = 1$ and $\beta_l = \alpha_{jl}$ for $l \in [1, 4] - \{k, j\}$.

2) *We have*

$$\omega_{a_k}(a_k) = (\alpha_i - 1)a_i + (\alpha_l - 1)a_l + (\alpha_j - 2)a_j$$

and

$$L_{a_k}(H) = \{\beta_i a_i + \beta_l a_l + \beta_j a_j \mid 0 \leq \beta_i < \alpha_i, 0 \leq \beta_l < \alpha_l, 0 \leq \beta_j < \alpha_j - 1\} \cup \{(\alpha_j - 1)a_j\}$$

for some permutation (k, i, l, j) of $[1, 4]$.

PROOF. 1) Proposition 6.2 2) implies b) \Rightarrow a). By the assumption we have $\beta_l < \alpha_l$ for $l \in [1, 4] - \{k, j\}$, which induces $\beta_l = \alpha_{jl}$. Assume that $\omega_{a_k}(a_k) - (\alpha_j - 1)a_j \in L_{a_k}(H)$. Then we have

$$\sum_{l \in [1, 4] - \{k, j\}} \beta_l a_l + (\alpha_j - 2)a_j = \sum_{l \in [1, 4] - \{k, j\}} \beta_l a_l + (\alpha_j - 1)a_j.$$

This is a contradiction.

2) Renumbering a_1, \dots, a_4 , we may assume $k = 1$. Now assume $\omega_{a_1}(a_1) - (\alpha_j - 1)a_j \in L_{a_1}(H)$ for all $j \in [2, 4]$. Then by Proposition 6.2 and the assumption, we get

$$\omega_{a_1}(a_1) = (\alpha_2 - 1)a_2 + (\alpha_3 - 1)a_3 + (\alpha_4 - 1)a_4,$$

which implies

$$L_{a_1}(H) = \{\beta_2 a_2 + \beta_3 a_3 + \beta_4 a_4 \mid 0 \leq \beta_i < \alpha_i\}.$$

This contradicts Lemma 6.1 1). Hence there exists a unique $j \in [2, 4]$ such that $2(\alpha_j - 1)a_j = \omega_{a_1}(a_1) + a_1$, which implies

$$\omega_{a_1}(a_1) = \sum_{l \in [2, 4] - \{j\}} \beta_l a_l + (\alpha_j - 2)a_j.$$

Therefore we get

$$\omega_{a_1}(a_1) = (\alpha_i - 1)a_i + (\alpha_l - 1)a_l + (\alpha_j - 2)a_j$$

for some permutation (i, l, j) or $(2, 3, 4)$. Hence we have

$$L_{a_1}(H) \cong \{\beta_i a_i + \beta_l a_l + \beta_j a_j \mid 0 \leq \beta_i < \alpha_i, 0 \leq \beta_l < \alpha_l, 0 \leq \beta_j < \alpha_j - 1\} \cup \{(\alpha_j - 1)a_j\}.$$

Assume $z = \gamma_i a_i + \gamma_l a_l + (\alpha_j - 1)a_j \in L_{a_1}(H)$ with $(\gamma_i, \gamma_l) \neq (0, 0)$. Since $\omega_{a_1}(a_1) - z \in L_{a_1}(H)$, we put

$$\omega_{a_1}(a_1) - z = \nu_i a_i + \nu_l a_l + \nu_j a_j$$

where $\nu_i < \alpha_i, \nu_l < \alpha_l$ and $\nu_j < \alpha_j$, hence

$$(\alpha_i - 1 - \gamma_i)a_i + (\alpha_l - 1 - \gamma_l)a_l - a_j = \nu_i a_i + \nu_l a_l + \nu_j a_j,$$

which implies $\nu_j + 1 = 0$, a contradiction.

Q. E. D.

By tedious computations using Proposition 6.3 we can give generators of the ideal I_H in the case of almost symmetric H .

THEOREM 6.4. *Let H be almost symmetric. Then renumbering a_1, a_2, a_3, a_4 the ideal I_H is generated by*

$$f_1 = X_1^{\alpha_1} - X_3^{\alpha_{13}} X_4^{\alpha_{14}}, \quad f_2 = X_2^{\alpha_2} - X_1^{\alpha_{21}} X_4^{\alpha_{24}}, \quad f_3 = X_3^{\alpha_3} - X_1^{\alpha_{31}} X_2^{\alpha_{32}},$$

$$f_4 = X_4^{\alpha_4} - X_1^{\alpha_{41}} X_2^{\alpha_{42}} X_3^{\alpha_{43}} \quad \text{and} \quad g = X_1^{\alpha_{21} + \alpha_{41}} X_3^{\alpha_{43}} - X_2^{\alpha_{32}} X_4^{\alpha_{14}}$$

where $0 < \alpha_{ij} < \alpha_j, \alpha_1 = \alpha_{21} + \alpha_{31} + \alpha_{41}, \alpha_2 = \alpha_{32} + \alpha_{42}, \alpha_3 = \alpha_{13} + \alpha_{43}$ and $\alpha_4 = \alpha_{14} + \alpha_{24}$, which imply $\mu(H) = 5$. More explicitly we obtain $\alpha_{13} = 1, \alpha_{14} = \alpha_4 - 1, \alpha_{24} = 1, \alpha_{31} = \alpha_1 - \alpha_{21} - 1, \alpha_{32} = 1, \alpha_{41} = 1, \alpha_{42} = \alpha_2 - 1$ and $\alpha_{43} = \alpha_3 - 1$. Hence using Proposition 6.3 2). We can show that H is 1-neat.

PROOF. For any $i \in [1, 4]$, let $f_i \in I_H$ be a polynomial of the type $X_i^{\alpha_i} - \prod_{j \in [1, 4] - \{i\}} X_j^{\alpha_j}$. First, assume that there exist two distinct $i, j \in [1, 4]$ with $X_i^{\alpha_i} - X_j^{\alpha_j} \in I_H$. Then renumbering a_1, \dots, a_4 we may assume $i = 1$ and $j = 2$. They are divided into the four cases:

- 1) $X_i^{\alpha_i} - X_j^{\alpha_j} \in I_H$ for all $\{i, j\} \neq \{1, 2\}$,
- 2) $X_1^{\alpha_1} - X_3^{\alpha_3} \in I_H$ and $X_1^{\alpha_1} - X_4^{\alpha_4} \in I_H$,
- 3) $X_3^{\alpha_3} - X_4^{\alpha_4} \in I_H$ and $X_1^{\alpha_1} - X_3^{\alpha_3} \in I_H$,
- 4) $X_1^{\alpha_1} - X_3^{\alpha_3} \in I_H$ and $X_1^{\alpha_1} - X_4^{\alpha_4} \in I_H$.

The case 1). Then $f_3 = X_3^{\alpha_3} - X_1^{\alpha_{31}} X_2^{\alpha_{32}} X_4^{\alpha_{34}}$ and $f_4 = X_4^{\alpha_4} - X_1^{\alpha_{41}} X_2^{\alpha_{42}} X_3^{\alpha_{43}}$. These are divided into the following:

- a) $\alpha_{31} > 0, \alpha_{32} > 0, \alpha_{41} > 0, \alpha_{42} > 0,$ b) $\alpha_{31} > 0, \alpha_{32} > 0, \alpha_{41} > 0, \alpha_{42} = 0,$
 c) $\alpha_{31} > 0, \alpha_{32} = 0, \alpha_{41} > 0, \alpha_{42} = 0,$ d) $\alpha_{31} > 0, \alpha_{32} = 0, \alpha_{41} = 0, \alpha_{42} > 0.$

a) Then we have

$$\begin{aligned} \omega_{a_1}(a_1) &= (\alpha_2 - 1)a_2 + (\alpha_3 - 1)a_3 + (\alpha_4 - 2)a_4 = \alpha_{42}a_2 + \alpha_{43}a_3 + (\alpha_4 - 2)a_4, \\ \omega_{a_2}(a_2) &= (\alpha_1 - 1)a_1 + (\alpha_3 - 1)a_3 + (\alpha_4 - 2)a_4 = \alpha_{41}a_1 + \alpha_{43}a_3 + (\alpha_4 - 2)a_4, \end{aligned}$$

which imply $\alpha_1 = \alpha_2 = 2$, hence $a_1 = a_2$, a contradiction.

b) Similarly, we get $a_1 = a_2$, a contradiction.

c) We have

$$\omega_{a_1}(a_1) = (\alpha_2 - 1)a_2 + (\alpha_i - 1)a_i + (\alpha_j - 2)a_j$$

with $\{i, j\} = \{3, 4\}$. This is a contradiction.

d) We get

$$\omega_{a_1}(a_1) = (\alpha_2 - 1)a_2 + (\alpha_3 - 1)a_3 + \beta_4 a_4, \quad \omega_{a_2}(a_2) = (\alpha_1 - 1)a_1 + (\alpha_4 - 1)a_4 + \beta_3 a_3,$$

which implies $\beta_4 = \alpha_4 - 1$. Hence we have

$$L_{a_1}(H) = \{\beta_2 a_2 + \beta_3 a_3 + \beta_4 a_4 \mid 0 \leq \beta_i < \alpha_i\},$$

which implies $C(H) = 2g(H)$, a contradiction.

The case 2). Then $f_4 = X_4^{\alpha_4} - X_1^{\alpha_{41}} X_2^{\alpha_{42}} X_3^{\alpha_{43}}$, where we may assume $\alpha_{41} > 0$. In the similar manner to 1) a), we get $a_1 = a_2$, a contradiction.

The case 3). We have $\omega_{a_3}(a_3) = \gamma_1 a_1 + \gamma_2 a_2 + (\alpha_4 - 1)a_4$. Set $d = (a_3, a_4)$ and $H' = \langle d, a_1, a_2 \rangle$. Then $L_d(H') \subseteq L_{a_3}(H)$. If $\nu_1 a_1 + \nu_2 a_2 + \nu_4 a_4 = \mu_1 a_1 + \mu_2 a_2 + \mu_4 a_4$ with $\nu_4 < \alpha_4$ and $\mu_4 < \alpha_4$, then $\nu_4 = \mu_4$. Using this, for any $\omega' \in \langle a_1, a_2 \rangle$ with $\omega_{a_3}(a_3) - \omega' \in L_{a_3}(H)$ we have

$$\omega_{a_3}(a_3) - \omega' = \mu_1 a_1 + \mu_2 a_2 + (\alpha_4 - 1)a_4$$

with $\mu_1, \mu_2 \in \mathbb{N}$. Hence if $\omega' \in L_d(H')$ with $\omega_{a_3}(a_3) - \omega' \in L_{a_3}(H)$, then for any $\nu_4 \in [0, \alpha_4 - 1]$ we get $\omega' + \nu_4 a_4 \in L_{a_3}(H)$. Therefore we can see:

$$L_{a_3}(H) = \{\omega' + \nu_4 a_4 \mid \omega' \in L_d(H'), 0 \leq \nu_4 < \alpha_4\} \quad \text{and} \quad \omega_{a_3}(a_3) = \omega_d(d) + (\alpha_4 - 1)a_4.$$

Since we have $\omega_d(d) - \omega' \in L_d(H')$ for any $\omega' \in L_d(H')$, we get $\omega_{a_3}(a_3) - \omega \in L_{a_3}(H)$ for any $\omega \in L_{a_3}(H)$, i. e., $C(H) = 2g(H)$, a contradiction.

The case 4). Then H is a complete intersection ([1]), which implies $C(H) = 2g(H)$, a contradiction.

Secondly, assume: each f_i contains at least three variables and there exists $j \in [1, 4]$ such that the variable X_j appears only in the f_j . Then we may assume that

$$f_1 = X_1^{\alpha_1} - X_2^{\alpha_{12}} X_3^{\alpha_{13}} X_4^{\alpha_{14}}, \quad f_2 = X_2^{\alpha_2} - X_3^{\alpha_{23}} X_4^{\alpha_{24}}, \quad f_3 = X_3^{\alpha_3} - X_2^{\alpha_{32}} X_4^{\alpha_{34}},$$

and

$$f_4 = X_4^{\alpha_4} - X_2^{\alpha_{42}} X_3^{\alpha_{43}}$$

with $\alpha_{13} > 0$, $\alpha_{14} > 0$. Hence we get

$$\omega_{a_3}(a_3) = (\alpha_2 - 1)a_2 + (\alpha_4 - 1)a_4 + (\alpha_1 - 2)a_1,$$

$$\omega_{a_4}(a_4) = (\alpha_2 - 1)a_2 + (\alpha_3 - 1)a_3 + (\alpha_1 - 2)a_1,$$

which imply $\alpha_4 a_4 = \alpha_3 a_3 = \alpha_{32} a_2 + \alpha_{34} a_4$, a contradiction.

Thirdly, assume: each f_i contains at least three variables and there exists $j \in [1, 4]$ such that the variable X_j appears twice in the f_i 's. Then we may assume that

$$f_1 = X_1^{\alpha_1} - X_2^{\alpha_{12}} X_3^{\alpha_{13}} X_4^{\alpha_{14}}, \quad f_2 = X_2^{\alpha_2} - X_1^{\alpha_{21}} X_3^{\alpha_{23}} X_4^{\alpha_{24}}, \quad f_3 = X_3^{\alpha_3} - X_2^{\alpha_{32}} X_4^{\alpha_{34}}$$

and

$$f_4 = X_4^{\alpha_4} - X_2^{\alpha_{42}} X_3^{\alpha_{43}}.$$

The case $\alpha_{12} > 0$. Then we have

$$\omega_{a_2}(a_2) = (\alpha_3 - 1)a_3 + (\alpha_4 - 1)a_4 + (\alpha_1 - 2)a_1,$$

$$\omega_{a_3}(a_3) = (\alpha_4 - 1)a_4 + (\alpha_i - 1)a_i + (\alpha_j - 2)a_j,$$

$$\omega_{a_4}(a_4) = (\alpha_3 - 1)a_3 + (\alpha_j - 1)a_j + (\alpha_i - 2)a_i,$$

with $\{i, j\} = \{1, 2\}$. If $j=1$ (resp. 2), then $(\alpha_4 - \alpha_{34})a_4 = a_1 + (\alpha_{32} - 1)a_2$ (resp. $(\alpha_3 - \alpha_{43})a_3 = a_1 + (\alpha_{42} - 1)a_2$), a contradiction. The case $\alpha_{12} = 0$. We have

$$\omega_{a_3}(a_3) = (\alpha_1 - 1)a_1 + (\alpha_4 - 1)a_4 + (\alpha_2 - 2)a_2,$$

$$\omega_{a_4}(a_4) = (\alpha_1 - 1)a_1 + (\alpha_3 - 1)a_3 + (\alpha_2 - 2)a_2,$$

which implies $\alpha_4 a_4 = \alpha_3 a_3$, a contradiction.

Lastly, assume: each f_i contains at least three variables and all the variables X_j appear at least three times in the f_i 's. Renumbering a_1, \dots, a_4 , these are divided into the 10 cases in Proposition 4.4.

The case (1). Then we may assume:

$$\omega_{a_1}(a_1) = (\alpha_2 - 1)a_2 + (\alpha_3 - 1)a_3 + (\alpha_4 - 2)a_4,$$

$$\omega_{a_4}(a_4) = (\alpha_i - 1)a_i + (\alpha_j - 1)a_j + (\alpha_k - 2)a_k.$$

Using $\omega_{a_1}(a_1) - a_1 = \omega_{a_4}(a_4) - a_4$, this is a contradiction.

The case (2). We have

$$\omega_{a_3}(a_3) = (\alpha_1 - 1)a_1 + (\alpha_i - 1)a_i + (\alpha_j - 2)a_j,$$

$$\omega_{a_4}(a_4) = (\alpha_k - 1)a_k + (\alpha_l - 1)a_l + (\alpha_m - 2)a_m.$$

This is a contradiction.

The case (3) a). We have

$$\omega_{a_3}(a_3) = (\alpha_1 - 1)a_1 + (\alpha_2 - 1)a_2 + (\alpha_4 - 2)a_4,$$

$$\omega_{a_4}(a_4) = (\alpha_1 - 1)a_1 + (\alpha_2 - 1)a_2 + (\alpha_3 - 2)a_3.$$

Then $(\alpha_4 - 1)a_4 = (\alpha_3 - 1)a_3$, a contradiction.

The case (3) b). We have

$$\omega_{a_4}(a_4) = (\alpha_1 - 1)a_1 + (\alpha_2 - 1)a_2 + (\alpha_3 - 2)a_3,$$

$$\omega_{a_1}(a_1) = (\alpha_2 - 1)a_2 + (\alpha_3 - 1)a_3 + (\alpha_4 - 2)a_4.$$

Moreover,

$$\omega_{a_3}(a_3) = (\alpha_1 - 1)a_1 + \gamma_2 a_2 + \gamma_4 a_4 \quad \text{or} \quad (\alpha_1 - 2)a_1 + \alpha_{14} a_4.$$

Using $\omega_{a_4}(a_4) - a_4 = \omega_{a_3}(a_3) - a_3 = \omega_{a_1}(a_1) - a_1$, this is a contradiction.

The case (3) c). We have

$$\omega_{a_1}(a_1) = (\alpha_3 - 1)a_3 + (\alpha_i - 1)a_i + (\alpha_j - 2)a_j,$$

$$\omega_{a_3}(a_3) = (\alpha_1 - 1)a_1 + (\alpha_j - 1)a_j + (\alpha_i - 2)a_i$$

This is a contradiction.

The case (4) a). We have

$$\omega_{a_3}(a_3) = (\alpha_1 - 1)a_1 + (\alpha_2 - 1)a_2 + (\alpha_4 - 2)a_4$$

and

$$\omega_{a_1}(a_1) = (\alpha_3 - 1)a_3 + \gamma_2 a_2 + \gamma_4 a_4 \quad \text{or} \quad (\alpha_3 - 2)a_3 + \alpha_{32} a_2.$$

This is a contradiction.

The case (4) b). $\omega_{a_4}(a_4) = (\alpha_i - 1)a_i + (\alpha_j - 1)a_j + (\alpha_l - 2)a_l$, a contradiction.

The case (5) a). We have

$$\omega_{a_1}(a_1) = (\alpha_3 - 1)a_3 + \gamma_2 a_2 + \gamma_4 a_4 \quad \text{or} \quad (\alpha_3 - 2)a_3 + \alpha_{32} a_2$$

Moreover,

$$\omega_{a_1}(a_1) = (\alpha_4 - 1)a_4 + \beta_2 a_2 + \beta_3 a_3 \quad \text{or} \quad (\alpha_4 - 2)a_4 + \alpha_{42} a_2$$

This is a contradiction.

The case (5) b). We have

$$\omega_{a_1}(a_1) = (\alpha_2 - 1)a_2 + (\alpha_3 - 1)a_3 + \gamma_4 a_4 \quad \text{or} \quad (\alpha_2 - 1)a_2 + (\alpha_3 - 2)a_3,$$

$$\omega_{a_2}(a_2) = (\alpha_3 - 1)a_3 + (\alpha_4 - 1)a_4 + \gamma_1 a_1 \quad \text{or} \quad (\alpha_3 - 1)a_3 + (\alpha_4 - 2)a_4,$$

$$\omega_{a_3}(a_3) = (\alpha_4 - 1)a_4 + (\alpha_1 - 1)a_1 + \gamma_2 a_2 \quad \text{or} \quad (\alpha_4 - 1)a_4 + (\alpha_1 - 2)a_1$$

and

$$\omega_{a_4}(a_4) = (\alpha_1 - 1)a_1 + (\alpha_2 - 1)a_2 + \gamma_3 a_3 \quad \text{or} \quad (\alpha_1 - 1)a_1 + (\alpha_2 - 2)a_2.$$

If we renumber a_1, \dots, a_4 , each latter case is reduced to the case (4) c). For example, let $\omega_{a_1}(a_1) = (\alpha_2 - 1)a_2 + (\alpha_3 - 2)a_3$. If $\omega_{a_2}(a_2) = (\alpha_3 - 1)a_3 + (\alpha_4 - 1)a_4 + \gamma_1 a_1$, then $\alpha_2 a_2 = (\gamma_1 + 1)a_1 + a_3 + (\alpha_4 - 1)a_4$, whose case is reduced to (4) c). If $\omega_{a_2}(a_2) = (\alpha_3 - 1)a_3 + (\alpha_4 - 2)a_4$, then $\alpha_2 a_2 = a_1 + a_3 + (\alpha_4 - 2)a_4$. If $\alpha_4 = 2$, we replace f_2 by $X_2^2 - X_1 X_3$, which is reduced to the third case, a contradiction. Hence we have $\alpha_4 \geq 3$, whose case is reduced to (4) c). Therefore for any $i \in [1, 4]$, $\omega_{a_i}(a_i)$ is equal to the former case. Then we see:

$$\alpha_{21} + \alpha_{31} = \alpha_1, \quad \alpha_{32} + \alpha_{42} = \alpha_2, \quad \alpha_{13} + \alpha_{43} = \alpha_3 \quad \text{and} \quad \alpha_{14} + \alpha_{24} = \alpha_4.$$

Using $\omega_{a_1}(a_1) - a_1 = \omega_{a_4}(a_4) - a_4$ we obtain

$$\begin{aligned} \omega_{a_1}(a_1) &= (\alpha_2 - 1)a_2 + (\alpha_3 - 1)a_3 + (\alpha_{14} - 1)a_4 \\ &= (\alpha_{32} - 1)a_2 + (\alpha_{13} - 1)a_3 + (\alpha_4 + \alpha_{14} - 1)a_4, \end{aligned}$$

which implies

$L_{a_1}(H) \supseteq \{\beta_2 a_2 + \beta_3 a_3 + \beta_4 a_4 \mid \beta_i \in N \text{ and } (\beta_2, \beta_3, \beta_4) \text{ satisfies one of the following:}$

$$1) \beta_2 < \alpha_2, \beta_3 < \alpha_3, \beta_4 < \alpha_{14}, \quad 2) \beta_2 < \alpha_{32}, \beta_3 < \alpha_{13}, \alpha_{14} \leq \beta_4 < \alpha_4\}.$$

Since there exists a positive integer ν such that

$$a_1 = \nu^{-1}(\alpha_2 \alpha_3 \alpha_{14} + \alpha_{32} \alpha_{13} \alpha_{24}),$$

the above inclusion becomes the equality, hence for any $\omega \in L_{a_1}(H)$ we have $\omega_{a_1}(a_1) - \omega \in L_{a_1}(H)$, i. e., $C(H) = 2g(H)$, a contradiction.

Therefore, if H is almost symmetric, renumbering a_1, \dots, a_4 it is reduced to the case (4) c), i. e.,

$$f_1 = X_1^{\alpha_1} - X_3^{\alpha_{13}} X_4^{\alpha_{14}}, \quad f_2 = X_2^{\alpha_2} - X_1^{\alpha_{21}} X_4^{\alpha_{24}}, \quad f_3 = X_3^{\alpha_3} - X_1^{\alpha_{31}} X_2^{\alpha_{32}}$$

and

$$f_4 = X_4^{\alpha_4} - X_1^{\alpha_{41}} X_2^{\alpha_{42}} X_3^{\alpha_{43}}.$$

Moreover,

$$\omega_{a_1}(a_1) = (\alpha_2 - 1)a_2 + (\alpha_3 - 1)a_3 + (\alpha_4 - 2)a_4 = \alpha_{42}a_2 + \alpha_{43}a_3 + (\alpha_4 - 2)a_4,$$

which implies $\alpha_{41} = 1$, $\alpha_{42} = \alpha_2 - 1$ and $\alpha_{43} = \alpha_3 - 1$. Now

$$\omega_{a_1}(a_1) = (\alpha_2 - 1)a_2 + (\alpha_3 - 1 - \alpha_{13})a_3 + (\alpha_4 - 2 - \alpha_{14})a_4 + \alpha_1 a_1,$$

which implies $\alpha_{14} = \alpha_4 - 1$. Moreover, we get

$$\begin{aligned} \omega_{a_4}(a_4) &= (\alpha_1 - 1)a_1 + (\alpha_2 - 1)a_2 + (\alpha_3 - 1 - \alpha_{13})a_3 \\ &= (\alpha_1 - 1 - \alpha_{21} - \alpha_{31} - \alpha_{41})a_1 + (\alpha_2 - 1 - \alpha_{42} + \alpha_2 - \alpha_{32})a_2 \\ &\quad + (\alpha_3 - \alpha_{43})a_3 + (\alpha_4 - \alpha_{24})a_4. \end{aligned}$$

If $\alpha_2 \geq \alpha_{32}$, then $\alpha_1 \leq \alpha_{21} + \alpha_{31} + \alpha_{41}$. If $\alpha_2 < \alpha_{32}$, then we have

$$\begin{aligned} \omega_{a_4}(a_4) &= (\alpha_1 - 1)a_1 + (\alpha_2 - 1)a_2 + (\alpha_3 - 2)a_3 \\ &= (\alpha_{21} + \alpha_{31})a_1 + (\alpha_{32} - \alpha_2)a_2 + (\alpha_3 - 2)a_3, \end{aligned}$$

which implies $\alpha_{21} + \alpha_{31} = \alpha_1 - 1$, hence $\alpha_1 \leq \alpha_{21} + \alpha_{31} + \alpha_{41}$. Since

$$\begin{aligned} \alpha_1 a_1 + \alpha_2 a_2 + \alpha_3 a_3 + \alpha_4 a_4 &= (\alpha_{21} + \alpha_{31} + \alpha_{41})a_1 + (\alpha_{32} + \alpha_{42})a_2 \\ &\quad + (\alpha_{13} + \alpha_{43})a_3 + (\alpha_{14} + \alpha_{34})a_4, \end{aligned}$$

we have

$$\alpha_1 = \alpha_{21} + \alpha_{31} + \alpha_{41}, \quad \alpha_2 = \alpha_{32} + \alpha_{42}, \quad \alpha_3 = \alpha_{13} + \alpha_{43} \quad \text{and} \quad \alpha_4 = \alpha_{14} + \alpha_{34},$$

which imply

$$\begin{aligned} \alpha_{41} &= 1, \quad \alpha_{31} = \alpha_1 - \alpha_{21} - 1, \quad \alpha_{32} = 1, \quad \alpha_{42} = \alpha_2 - 1, \\ \alpha_{13} &= 1, \quad \alpha_{43} = \alpha_3 - 1, \quad \alpha_{14} = \alpha_4 - 1, \quad \alpha_{34} = 1. \end{aligned}$$

Since we have

$$L_{a_1}(H) = \{\beta_2 a_2 + \beta_3 a_3 + \beta_4 a_4 \mid 0 \leq \beta_2 < \alpha_2, 0 \leq \beta_3 < \alpha_3, 0 \leq \beta_4 < \alpha_4 - 1\} \cup \{(\alpha_4 - 1)a_4\},$$

H is 1-neat.

Q. E. D.

Conversely, by simple calculations we get the following:

THEOREM 6.5. *Let $\alpha_i > 1$ for $1 \leq i \leq 4$ and let $0 < \alpha_{21} < \alpha_1 - 1$. If $a_1 = \alpha_2 \alpha_3 (\alpha_4 - 1) + 1$, $a_2 = \alpha_{21} \alpha_3 \alpha_4 + (\alpha_1 - \alpha_{21} - 1)(\alpha_3 - 1) + \alpha_3$, $a_3 = \alpha_1 \alpha_4 + (\alpha_1 - \alpha_{21} - 1)(\alpha_2 - 1)(\alpha_4 - 1) - \alpha_4 + 1$, $a_4 = \alpha_1 \alpha_2 (\alpha_3 - 1) + \alpha_{21}(\alpha_2 - 1) + \alpha_2$ and $(a_1, a_2, a_3, a_4) = 1$, then $H = \langle a_1, a_2, a_3, a_4 \rangle$ is an almost symmetric numerical semigroup with $M(H) = \{a_1, a_2, a_3, a_4\}$ and the ideal I_H is generated by*

$$\begin{aligned} f_1 &= X_1^{\alpha_1} - X_3 X_4^{\alpha_4 - 1}, \quad f_2 = X_2^{\alpha_2} - X_1^{\alpha_{21}} X_4, \quad f_3 = X_3^{\alpha_3} - X_1^{\alpha_1 - \alpha_{21} - 1} X_2, \\ f_4 &= X_4^{\alpha_4} - X_1 X_2^{\alpha_2 - 1} X_3^{\alpha_3 - 1} \quad \text{and} \quad g = X_1^{\alpha_{21} + 1} X_3^{\alpha_3 - 1} - X_2 X_4^{\alpha_4 - 1}. \end{aligned}$$

PROOF. By the assumption, we have

$$\alpha_1 a_1 = a_3 + (\alpha_4 - 1)a_4, \quad \alpha_2 a_2 = \alpha_{21} a_1 + a_4 \quad \text{and} \quad \alpha_3 a_3 = (\alpha_1 - \alpha_{21} - 1)a_1 + a_2,$$

which imply $\alpha_4 a_4 = a_1 + (\alpha_2 - 1)a_2 + (\alpha_3 - 1)a_3$. Using the relations, we get

$$L_{a_1}(H) = \{\beta_2 a_2 + \beta_3 a_3 + \beta_4 a_4 \mid 0 \leq \beta_2 < \alpha_2, 0 \leq \beta_3 < \alpha_3, 0 \leq \beta_4 < \alpha_4 - 1\} \cup \{(\alpha_4 - 1)a_4\}$$

and

$$\omega_{a_1}(a_1) = (\alpha_2 - 1)a_2 + (\alpha_3 - 1)a_3 + (\alpha_4 - 2)a_4,$$

which show that H is almost symmetric. Moreover, since we have

$$\begin{aligned}
L_{a_4}(H) = & \{ \beta_1 a_1 + \beta_2 a_2 + \beta_3 a_3 \mid 0 \leq \beta_1 < \alpha_1, 0 \leq \beta_2 < \alpha_2, 0 \leq \beta_3 < \alpha_3 - 1 \} \\
& \cup \{ \beta_1 a_1 + \beta_2 a_2 + (\alpha_3 - 1) a_3 \mid 0 \leq \beta_1 \leq \alpha_{21}, 0 \leq \beta_2 < \alpha_2 - 1 \} \\
& \cup \{ (\alpha_2 - 1) a_2 + (\alpha_3 - 1) a_3 \},
\end{aligned}$$

we get $a_1 \in \langle a_2, a_3, a_4 \rangle$, $a_2 \in \langle a_1, a_3, a_4 \rangle$, $a_3 \in \langle a_1, a_2, a_4 \rangle$, $a_4 \in \langle a_1, a_2, a_3 \rangle$. Using the above relations, we get

$$\begin{aligned}
L_{a_2}(H) = & \{ \beta_1 a_1 + \beta_3 a_3 + \beta_4 a_4 \mid 0 \leq \beta_1 < \alpha_{21}, 0 \leq \beta_3 < \alpha_3, 0 \leq \beta_4 < \alpha_4 \} \\
& \cup \{ \beta_1 a_1 + \beta_3 a_3 \mid \alpha_{21} \leq \beta_1 < \alpha_1, 0 \leq \beta_3 < \alpha_3 - 1 \} \cup \{ \alpha_{21} a_1 + (\alpha_3 - 1) a_3 \}.
\end{aligned}$$

The complete descriptions of $L_{a_1}(H)$, $L_{a_2}(H)$ and $L_{a_4}(H)$ show that the above relations are minimal. Q. E. D.

References

- [1] Bresinsky, H., Symmetric semigroups of integers generated by 4 elements, *manuscripta math.* 17 (1975) 205-219.
- [2] Buchweitz, R.O., On deformations of monomial curves, *Séminaire sur les singularités des surfaces*, Palaiseau, France 1976-1977, Springer, Lecture Notes 777, 1980, 205-220.
- [3] Grothendieck, A. (with Dieudonné, J.), *Éléments de géométrie algébrique IV 3^{ème} partie*, Publ. Math. I. H. E. S., no. 28, 1966.
- [4] Hartshorne, R., *Algebraic geometry*, Springer, GTM 52, p. 272, 1977.
- [5] Herzog, J., Generators and relations of abelian semigroups and semigroup rings, *manuscripta math.* 3 (1970) 175-193.
- [6] Kempf, G., Knudsen, F., Mumford, D. and Saint-Donat, B., *Toroidal embeddings I*, Springer, Lecture Notes 339, 1973.
- [7] Pinkham, H.C., Deformations of algebraic varieties with G_m action, *Astérisque*, 20 (1974) 1-131.
- [8] Rim, D.S. and Vitulli, M.A., Weierstrass points and monomial curves, *J. Algebra*, 48 (1977) 454-476.
- [9] Waldi, R., Deformation von Gorenstein-Singularitäten der Kodimension 3, *Math. Ann.* 242 (1979) 201-208.
- [10] Waldi, R., Zur Konstruktion von Weierstrasspunkten mit vorgegebener Halbgruppe, *manuscripta math.* 30 (1980) 257-278.

Institute of Mathematics
University of Tsukuba
Ibaraki, Japan