

AD-A177 077

ON THE INTENSITY OF CROSSINGS BY A SHOT NOISE PROCESS

1/1

(U) NORTH CAROLINA UNIV AT CHAPEL HILL CENTER FOR  
STOCHASTIC PROCESSES T HSING JUL 86 TR-141


UNCLASSIFIED

AFOSR-TR-87-0122 F49620-82-C-0009

F/G 12/1

NL





UNCLASSIFIED

(2)

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

AD-A177 077

1a. REPORT SECURITY CLASSIFICATION <b>UNCLASSIFIED</b>		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY <b>NA</b>		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE <b>NA</b>			
4. PERFORMING ORGANIZATION REPORT NUMBER(S)  Technical Report No. 141		5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR- 87-0122</b>	
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6c. ADDRESS (City, State and ZIP Code) Center for Stochastic Processes, Statistics Department, Phillips Hall 039-A, Chapel Hill, NC 27514		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER <del>F49620 82 C 0000</del> F49620 82 C 0009	
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC		10. SOURCE OF FUNDING NOS.	
11. TITLE (Include Security Classification) "On the intensity of crossings by a shot noise process"		PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304
12. PERSONAL AUTHOR(S) Hsing, T		TASK NO. AS	WORK UP NO.
13a. TYPE OF REPORT technical	13b. TIME COVERED FROM 9/84 TO 8/85	14. DATE OF REPORT (Yr., Mo., Day) July 1986	15. PAGE COUNT 6
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	Keywords: Level crossing; shot noise.
XXXXX	XXXXXXXXXX	XXXXX	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)  The crossing intensity of a level by a shot noise process with a monotone impulse response is studied. It is shown that the intensity can be naturally expressed in terms of a marginal probability. Also some examples are given to illustrate how the marginal probability can be obtained.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Peggy Revitch Maj Crawley		22b. TELEPHONE NUMBER (Include Area Code) 202 767 5033 <del>910 662 2207</del>	22c. OFFICE SYMBOL AFOSR/NM


DTIC FILE COPY

DTIC ELECTED FEB 25 1987

# CENTER FOR STOCHASTIC PROCESSES

Department of Statistics  
University of North Carolina  
Chapel Hill, North Carolina

AFOSR-TR- 87 - 0122



ON THE INTENSITY OF CROSSINGS BY A  
SHOT NOISE PROCESS

by

T. Hsing

Technical Report No. 141

July 1986

Approved for public release;  
distribution unlimited.

87 2 20 185

ON THE INTENSITY OF CROSSINGS BY A  
SHOT NOISE PROCESS

Tailen Hsing  
The University of Texas at Arlington

Summary. The crossing intensity of a level by a shot noise process with a monotone impulse response is studied. It is shown that the intensity can be naturally expressed in terms of a marginal probability. Also some examples are given to illustrate how the marginal probability can be obtained.

AMS 1980 Subject Classification: 60K99.

Key Words and Phrases: Level crossing, shot noise.

Research partially supported by the Air Force Office of Scientific Research  
Grant No. AFOSR F49620 82 C 0009.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



## 1. Introduction.

Consider the shot noise process

$$X(t) = \sum_{\tau \leq t} h(t - \tau), \quad t \in R,$$

where the  $\tau$ 's are the points of a stationary Poisson process on  $R$  with mean rate  $\lambda > 0$ , and  $h$ , the impulse response, is a non-negative function on  $[0, \infty)$  such that

- i)  $h$  is non-increasing,
- ii)  $h$  is finite except possibly at zero, and
- iii)  $\int_u^\infty h(x) dx < \infty$  for some large  $u$ .

By Daley (1971), Theorem 1, the conditions (ii) and (iii) ensure that  $X(t) < \infty$  a.s. for each  $t$ .

Observe that the sample function of  $X$  increases only at the points of  $\eta$ . Thus it is unambiguous to define that  $X$  upcrosses the level  $u$  at  $t$ , where  $u \geq 0$ , if  $X(t-) \leq u$  and  $X(t) > u$ . For  $u \geq 0$ , write  $N_u$  for the point process (cf. Kallenberg (1976)) that consists of the points at which upcrossings of level  $u$  by  $X$  occur. Thus for each Borel set  $B$ ,  $N_u(B)$  denotes the number of upcrossings of  $u$  by  $X$  in  $B$ .  $N_u$  is a stationary point process, which may be viewed as a thinned process of  $\eta$ . The purpose of this paper is to derive the following result.

Theorem 1. For each  $u \geq 1$ ,  $EN_u[0,1] = \lambda P[u - h(0) < X(0) \leq u]$ .

Note that the "downcrossing" intensity of a level by  $X$  is also given by Theorem 1.

It is worth mentioning that similar problems were treated by Rice (1944), and Bar-David and Nemirovsky (1972) in other settings. A result in the latter paper can be reduced to one which is similar to Theorem 1. However, our assumptions on  $h$  are considerably simpler.

We prove Theorem 1 in Section 2 using an approach which appears to be most natural for the present purpose. In Section 3, we illustrate the manner in which Theorem 1 can be made useful for a number of situations.

## 2. Derivation.

It is convenient to enumerate the points of  $\eta$  in  $(-\infty, 0)$  by letting  $\rho_i$  be the  $i$ th largest point of  $\eta$  to the left of zero for  $i = 1, 2, \dots$ . The  $\rho_i$  are well-defined with probability one (w.p.1), and  $-\rho_1, \rho_1 - \rho_2, \rho_2 - \rho_3, \dots$  are independent and identically distributed (i.i.d.) exponential random variables. The following result is useful.

Lemma 2. For each  $i = 1, 2, \dots$ ,  $P[X(\rho_i^-) = \sum_{j \geq i+1} h(\rho_i - \rho_j)] = 1$  where  $X(\rho_i^-)$  denotes the left-hand limit of  $X$  at  $\rho_i$ . From this, it follows immediately that  $X(\rho_i^-)$  is independent of  $\rho_i$ , and  $X(\rho_i^-)$  has the same distribution as  $X(0)$ .

Proof. Let  $i \geq 1$  be fixed. Since  $h$  is monotone, it is almost everywhere continuous. Using the continuity of  $\rho_i - \rho_j$ ,  $j \geq i + 1$ , we obtain

$$\lim_{\epsilon \downarrow 0} h(\rho_i - \rho_j - \epsilon) = h(\rho_i - \rho_j) \text{ w.p.1 for } j \geq i + 1.$$

Also by the monotonicity of  $h$ ,  $h(\rho_i - \rho_j - \epsilon) \leq h(\rho_{i+1} - \rho_j)$  for

$0 < \varepsilon < \rho_i - \rho_{i+1}$ ,  $j \geq i + 2$ , where  $\sum_{j \geq i+2} h(\rho_{i+1} - \rho_j)$  is equal in distribution to  $X(0)$  which is finite w.p.1. Thus it follows from dominated convergence that w.p.1,

$$\lim_{\varepsilon \rightarrow 0} X(\rho_i - \varepsilon) = \lim_{\varepsilon \rightarrow 0} \sum_{j \geq i+1} h(\rho_i - \rho_j - \varepsilon) = \sum_{j \geq i+1} h(\rho_i - \rho_j). \quad \square$$

Proof of Theorem 1. By stationarity, it apparently suffices to show that  $EN_u(B)$  equals  $\lambda m(B)P[u - h(0) < X(0) \leq u]$  for each Borel set  $B$  in  $(-\infty, 0)$ , where  $m(B)$  denotes the Lebesgue measure of  $B$ . Since  $X(\rho_i) = h(0) + \sum_{j \geq i+1} h(\rho_i - \rho_j)$ , it follows from Lemma 2 that w.p.1,

$$N_u(B) = \sum_{i \geq 1} 1(u - h(0) < X(\rho_i^-) \leq u, \rho_i \in B),$$

where  $1(\cdot)$  is the indicator function. Applying the facts that  $X(\rho_i^-)$  is independent of  $\rho_i$  and  $X(\rho_i^-)$  is equal in distribution to  $X(0)$ , we get

$$\begin{aligned} EN_u(B) &= \sum_{i \geq 1} E1(u - h(0) < X(\rho_i^-) \leq u)E1(\rho_i \in B) \\ &= P[u - h(0) < X(0) \leq u]\lambda m(B). \quad \square \end{aligned}$$

### 3. Marginal Distribution.

The usefulness of Theorem 1 obviously depends on the availability of the marginal probability  $P[u - h(0) < X(0) \leq u]$ . The Laplace transform of  $X(0)$  is (cf. Gilbert and Pollak (1960))

$$(3.1) \quad L(s) = Ee^{-sX(0)} = \exp\{-\lambda \int_0^\infty (1 - e^{-sh(x)})dx\}, \quad s \geq 0.$$



For some impulse responses  $h$ , the distribution of  $X(0)$  can be expressed analytically, while for a class of others, a recursive method due to Gilbert and Pollak (1960) is applicable. If it is of interest to study the asymptotic crossing intensity for increasingly high levels, certain Tauberian theorems (cf. Embrechts et. al. (1985)) are useful. We consider three examples.

(a) Suppose  $h(x) = \begin{cases} \infty, & x = 0 \\ -\log x, & 0 < x < 1 \\ 0, & x \geq 1 \end{cases}$  . Then

$$L(s) = \exp\{-\lambda \int_0^\infty (1 - e^{-sx})e^{-x} dx\}, \quad s \geq 0,$$

which is the Laplace transform of the Bessel density (cf. Feller (1971)):

$$f(x) = e^{-(x+\lambda)} \sqrt{\frac{\lambda}{x}} I_1(2\sqrt{\lambda x}), \quad x > 0.$$

(b) For  $h(x) = e^{-x}$ ,  $x \geq 0$ , Gilbert and Pollak (1960) showed that the density  $f$  of  $X(0)$  can be obtained recursively as follows:

$$f(x) = \begin{cases} \frac{e^{-\lambda\gamma}}{\Gamma(\lambda)} x^{\lambda-1}, & 0 < x < 1, \\ x^{\lambda-1} \left[ \frac{e^{-\lambda\gamma}}{\Gamma(\lambda)} - \lambda \int_1^x f(y-1)y^{-\lambda} dy \right], & x \geq 1, \end{cases}$$

where  $\gamma$  is Euler's constant.

(c) Assume that  $h$  is boundedly supported, say, on  $[0,1]$ . Then by a change-of-variable, (3.1) becomes

$$L(s) = \exp\{-\lambda + \lambda \int_{[0, \infty)} e^{-sy} \mu(dy)\}$$

where  $\mu$  is a probability measure on  $[0, \infty)$  such that

$$\mu(B) = \text{Lebesgue measure of } \{0 \leq x \leq 1 : h(x) \in B\}$$

for each Borel set  $B$  in  $[0, \infty)$ . Thus  $X(0)$  has a compound Poisson distribution. For  $h$  satisfying certain regularity conditions, Embrecht et. al. (1985) showed that

$$P[X(0) > x] \sim \frac{\exp\{-\lambda[1 - \psi(t)] - e^{-\lambda} - t(x - 1)\}}{t\sqrt{2\pi\lambda\psi''(t)}} \text{ as } x \rightarrow \infty$$

where  $\psi(s) = \int e^{-su} \mu(du)$ , and  $t$  satisfies  $\lambda\psi'(t) = x$ .

## REFERENCES

- Bar-David, I. and Nemirovsky, A. (1972). Level crossings of nondifferentiable shot process. IEEE Trans. Inform. Theory, Vol. IT-18, No. 1, 27-34.
- Daley, D.J. (1971). The definition of a multi-dimensional generalization of shot noise. J. Appl. Prob. 8, 128-135.
- Embrechts, P., Jensen, J. L., Maejima, M., and Teugels, J. L. (1985). Approximation for compound Poisson and Pólya process. Adv. Appl. Prob. 17, 623-637.
- Feller, W. (1971). An Introduction to Probability Theory and Its Applications, Vol. 2, Second Ed. New York: Wiley.
- Gilbert, E. N. and Pollak, H. O. (1960). Amplitude distribution of shot noise. Bell System Tech. J. 30, 333-350.
- Kallenberg, O. (1976). Random Measures. Berlin: Akademie-Verlag, London-New York: Academic Press.
- Rice, S. O. (1944). Mathematical analysis of random noise. Bell System Tech. J. 24, 46-156.

END

3-87

DTIC