

## ON THE INTERPOLATION CONSTANT FOR ORLICZ SPACES

ALEXEI YU. KARLOVICH AND LECH MALIGRANDA

(Communicated by Jonathan M. Borwein)

ABSTRACT. In this paper we deal with the interpolation from Lebesgue spaces  $L^p$  and  $L^q$ , into an Orlicz space  $L^\varphi$ , where  $1 \leq p < q \leq \infty$  and  $\varphi^{-1}(t) = t^{1/p} \rho(t^{1/q-1/p})$  for some concave function  $\rho$ , with special attention to the interpolation constant  $C$ . For a bounded linear operator  $T$  in  $L^p$  and  $L^q$ , we prove modular inequalities, which allow us to get the estimate for both the Orlicz norm and the Luxemburg norm,

$$\|T\|_{L^\varphi \rightarrow L^\varphi} \leq C \max \left\{ \|T\|_{L^p \rightarrow L^p}, \|T\|_{L^q \rightarrow L^q} \right\},$$

where the interpolation constant  $C$  depends only on  $p$  and  $q$ . We give estimates for  $C$ , which imply  $C < 4$ . Moreover, if either  $1 < p < q \leq 2$  or  $2 \leq p < q < \infty$ , then  $C < 2$ . If  $q = \infty$ , then  $C \leq 2^{1-1/p}$ , and, in particular, for the case  $p = 1$  this gives the classical Orlicz interpolation theorem with the constant  $C = 1$ .

### 1. INTRODUCTION

The classical Riesz-Thorin interpolation theorem says that  $(L^p, L^q)$  are interpolation spaces for linear operators between  $(L^{p_0}, L^{q_0})$  and  $(L^{p_1}, L^{q_1})$ , where

$$(1.1) \quad \frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}, \quad \frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1},$$

with the estimation of the norm

$$\|T\|_{L^p \rightarrow L^q} \leq C (\|T\|_{L^{p_0} \rightarrow L^{q_0}})^{1-\theta} (\|T\|_{L^{p_1} \rightarrow L^{q_1}})^\theta$$

where  $C \leq \sqrt{2}$ . The constant  $C$  is 1 when either the spaces are complex or the spaces are real and  $p_i \leq q_i, i = 0, 1$  (see, e.g., [5, Section 1.7]).

After the Riesz-Thorin interpolation theorem several results have been proved about the interpolation of Orlicz spaces. The problem was the following: *if  $T$  is any bounded linear operator from  $L^{\varphi_i}$  into  $L^{\psi_i}, i = 0, 1$ , then under what conditions on  $\varphi$  and  $\psi$  is it true that  $T$  is also bounded from  $L^\varphi$  to  $L^\psi$ ?*

The assumption corresponding to (1.1) which appears naturally here is

$$(1.2) \quad \varphi^{-1} = \varphi_0^{-1} \rho \left( \frac{\varphi_1^{-1}}{\varphi_0^{-1}} \right), \quad \psi^{-1} = \psi_0^{-1} \rho \left( \frac{\psi_1^{-1}}{\psi_0^{-1}} \right),$$

for some concave function  $\rho$ .

---

Received by the editors January 24, 2000.

1991 *Mathematics Subject Classification*. Primary 46B70, 46E30; Secondary 26D07.

*Key words and phrases*. Orlicz spaces, interpolation constant, interpolation of operators,  $K$ -functional, convex function, concave function.

The first result on interpolation of Orlicz spaces appeared in the case when  $\rho(t) = t^\theta$ ,  $0 \leq \theta \leq 1$ , and we mention here Ya. B. Rutickii (1963), A. P. Calderón (1964), M. M. Rao (1966) as the precursors of such results. The constants found by Ya. B. Rutickii were 4 in the complex case and 8 in the real case. The corresponding constants in two other papers are 2 and 4, respectively. Theorems with general concave  $\rho$  (and the interpolation theorems for, in fact, the Calderón-Lozanovskii construction) were done by several authors; among others, we mention V. I. Ovchinnikov (1976, 1984), J. Gustavsson-J. Peetre (1977), E. I. Bereznoi (1980), V. A. Shestakov (1981), E. I. Pustylnik (1983), P. Nilsson (1985) and L. Maligranda (1985, 1989). For the precise references and the proofs, see [16] and also [5, 8, 11, 17, 18].

Only some of these results on interpolation of Orlicz (or Calderón-Lozanovskii) spaces take care about the estimate of the operator norm. One known result is the following:

**Theorem 1.1.** *If (1.2) holds for some concave function  $\rho$ , then  $(L^\varphi, L^\psi)$  are interpolation spaces for linear operators between  $(L^{\varphi_0}, L^{\psi_0})$  and  $(L^{\varphi_1}, L^{\psi_1})$ , and*

$$\|T\|_{L^\varphi \rightarrow L^\psi} \leq C \max\{M_0, M_1\}, \quad M_i := \|T\|_{L^{\varphi_i} \rightarrow L^{\psi_i}}, \quad i = 0, 1,$$

where  $C \leq 26$  and all norms in Orlicz spaces are the Luxemburg norms.

The proof of this statement can be found in [16, Theorem 14.2]. The careful analysis of the proof shows that, in fact,

$$(1.3) \quad C \leq 2(3 + 2\sqrt{2})C_\psi < 12C_\psi, \quad \text{where } C_\psi = \sup_{t>0} \frac{\psi^{-1}(2t)}{\psi^{-1}(t)} \leq 2.$$

In the general case we have only the fairly rough estimate of the norm  $C < 24$ , but for the “diagonal case” and Lebesgue spaces, i.e., when  $L^{\varphi_0} = L^{\psi_0} = L^p$  and  $L^{\varphi_1} = L^{\psi_1} = L^q$ , one can obtain more precise estimates. Moreover, for most of the operators, we have information about boundedness between  $L^p$  spaces and we would like to get estimates in more general spaces, for example, in Orlicz spaces. Our problem here starts with the bounded linear operators  $T$  from  $L^p$  into  $L^p$  which are also bounded from  $L^q$  into  $L^q$  (with  $1 \leq p < q \leq \infty$ ) and we want to have boundedness of  $T$  from the Orlicz space  $L^\varphi$  into itself with the best possible estimate of the norm

$$(1.4) \quad \|T\|_{L^\varphi \rightarrow L^\varphi} \leq C \max\{M_0, M_1\}, \quad M_0 := \|T\|_{L^p \rightarrow L^p}, \quad M_1 := \|T\|_{L^q \rightarrow L^q},$$

where  $\varphi^{-1}(u) = u^{1/p} \rho(u^{1/q-1/p})$  for some concave function  $\rho$ .

Besides the above-mentioned authors working in the general case, there were also others working either with weak type operators or the diagonal case and Lebesgue spaces. I. B. Simonenko [21], among others, showed in 1964 that if

$$(1.5) \quad 1 \leq p < a_\varphi := \inf_{u>0} \frac{u\varphi'(u)}{\varphi(u)} \leq b_\varphi := \sup_{u>0} \frac{u\varphi'(u)}{\varphi(u)} < q < \infty,$$

then the Orlicz space  $L^\varphi$  is an interpolation space between  $L^p$  and  $L^q$  but the constant  $C$  in the estimation of the norm can be large:  $C \approx \left((a_\varphi - p)(q - b_\varphi)\right)^{-1}$ . B. W. Boyd [4] extended in 1967 this theorem to rearrangement-invariant spaces and his constant in the estimation of the norm increases to infinity when one of the Boyd indices of the space is going either to  $p$  or to  $q$ .

Let us also mention that interpolation theorems of Marcinkiewicz type (operators are of weak type, i.e., maps  $L^p$  into weak  $L^p$  and  $L^q$  into weak  $L^q$ ) in Orlicz spaces were done by A. Zygmund (1956), A. Torchinsky (1976), A. Cianchi (1998) (see [6] and the references given there) and in symmetric spaces by E. M. Semenov (1968), D. W. Boyd (1969), M. Zippin (1971), S. G. Krein and E. M. Semenov (1973) (see [10, 14] and references given there). Observe that the constants in these theorems are still large.

The paper is organized as follows. Section 2, called preliminaries, contains necessary definitions and some auxiliary results from the theory of Orlicz spaces and interpolation theory. In Section 3 we consider useful properties of concave and convex functions on  $[0, \infty)$ . In particular, the equivalence between representations  $\varphi^{-1}(u) = u^{1/p}\rho(u^{1/q-1/p})$  and  $\varphi(u) = u^p\rho(u^{q-p})$  is proved. In Section 4 we prove modular estimates of bounded linear operators in  $L^p$  and  $L^q$  spaces which are the keys for the estimates in the norms, both the Luxemburg one and the Orlicz one. In the case  $q < \infty$ , the idea of the proof goes back to J. Peetre [20]. In the proof we use essentially the exact estimation of the Sparr functional  $K_{p,q}^*$  [22]. In the case  $q = \infty$  our proof is based on the Krée formula, the Hardy-Littlewood-Pólya majorization theorem and the convexity of  $\varphi(u^{1/p})$ . Finally, in Section 5, we put all our pieces of results together and prove our main Theorem 5.1, which shows that the interpolation constant  $C$  in the estimate (1.4) is always less than 4. Moreover, if either  $1 < p < q \leq 2$  or  $2 \leq p < q < \infty$ , then  $C < 2$ . If  $q = \infty$ , then  $C \leq 2^{1-1/p}$ , and, in particular, for the case  $p = 1$  this gives the classical Orlicz interpolation theorem with the constant  $C = 1$ .

## 2. PRELIMINARIES

Let  $(\Omega, \mu)$  be a  $\sigma$ -finite measure space. Let  $\varphi : [0, \infty) \rightarrow [0, \infty]$  be a convex nondecreasing function such that  $\varphi(0) = 0$  and  $\lim_{u \rightarrow 0^+} \varphi(u) = 0$  but not identically zero or infinity on  $(0, \infty)$ . For a measurable real or complex-valued function  $x$ , define a functional (*modular*)

$$(2.1) \quad I_\varphi(x) := \int_\Omega \varphi(|x(s)|) d\mu(s) = \int_0^\infty \varphi(x^*(t)) dt,$$

where  $x^*$  is the nonincreasing rearrangement of  $x$  (see [1, 10, 12]). The *Orlicz space*  $L^\varphi = L^\varphi(\Omega, \mu)$  is the space of all equivalence classes of measurable functions on  $\Omega$  such that  $I_\varphi(\lambda x) < \infty$  for some  $\lambda = \lambda(x) > 0$ . This space is a Banach space with two norms: the *Luxemburg norm*

$$\|x\|_\varphi := \inf \{ \lambda > 0 : I_\varphi(x/\lambda) \leq 1 \}$$

and the *Orlicz norm* (in the Amemiya form)

$$\|x\|_\varphi^0 := \inf_{k>0} \frac{1}{k} \left( 1 + I_\varphi(kx) \right).$$

It is well known that  $\|x\|_\varphi \leq \|x\|_\varphi^0 \leq 2\|x\|_\varphi$ , and  $\|x\|_\varphi \leq 1$  if and only if  $I_\varphi(x) \leq 1$  (cf. [9]). The Orlicz space  $L^\varphi$  with each of the above two norms is a rearrangement-invariant space (= symmetric space with the Fatou property) (see [1, 10]). If  $\varphi$  satisfies the  $\Delta_2$ -condition, then the dual of the Orlicz space  $L^\varphi$  is an Orlicz space  $L^{\varphi^*}$  generated by the conjugate function  $\varphi^*$ , defined by

$$\varphi^*(u) := \sup_{v>0} \left( uv - \varphi(v) \right).$$

Moreover,  $(\|\cdot\|_\varphi)^* = \|\cdot\|_{\varphi^*}^0$  and  $(\|\cdot\|_\varphi^0)^* = \|\cdot\|_{\varphi^*}$ .

Let  $(X_0, X_1)$  be a couple of Banach spaces and let  $X$  be an intermediate Banach space between  $X_0$  and  $X_1$ , that is,  $X_0 \cap X_1 \subset X \subset X_0 + X_1$  with continuous imbeddings ( $X_0 \cap X_1, X_0 + X_1$  are standard spaces; see [1, 3, 5, 10, 12]). By  $\mathcal{A}(X_0, X_1)$  we denote the class of all admissible operators, i.e., linear operators  $T : X_0 + X_1 \rightarrow X_0 + X_1$  which restriction to  $X_i$  is bounded from  $X_i$  into itself for  $i = 0, 1$ . We denote

$$M := \max \{M_0, M_1\}, \quad \text{where } M_i := \|T|_{X_i}\|_{X_i \rightarrow X_i}, \quad i = 0, 1.$$

The space  $X$  is said to be an *interpolation space* between  $X_0$  and  $X_1$  if every admissible operator  $T \in \mathcal{A}(X_0, X_1)$  maps  $X$  into itself and

$$\|T\|_{X \rightarrow X} \leq C \max \{M_0, M_1\}$$

for some  $C > 0$ .

For  $0 < p, q < \infty, t > 0$  and  $x \in X_0 + X_1$ , we define a functional  $K_{p,q}$  known as the Peetre  $L$ -functional (Peetre [20]; see also [3, Th. 5.2.2], [5, Definition 3.1.22]):

$$K_{p,q}(t, x; X_0, X_1) := \inf \left\{ \|x_0\|_{X_0}^p + t \|x_1\|_{X_1}^q : x = x_0 + x_1, x_0 \in X_0, x_1 \in X_1 \right\}.$$

In the case  $p = q = 1$  this is the classical Peetre  $K$ -functional, which we shortly denote by  $K(t, x; X_0, X_1)$ .

**Proposition 2.1.** *If  $T \in \mathcal{A}(X_0, X_1)$ , then*

$$K_{p,q} \left( t, \frac{Tx}{M}; X_0, X_1 \right) \leq K_{p,q}(t, x; X_0, X_1) \quad \text{for all } t > 0.$$

The proof is standard.

Following [22], we consider the functional  $K_{p,q}^*$  on the couple of Lebesgue spaces defined by

$$K_{p,q}^*(t, x; L^p, L^q) := \int_{\Omega} \min \left( |x(s)|^p, t|x(s)|^q \right) d\mu(s).$$

**Lemma 2.2** (G. Sparr [22, Lemma 5.1, Example 5.3]). *Suppose  $1 \leq p < q < \infty$ . If  $x, y \in L^p + L^q$  and*

$$K_{p,q}(t, x; L^p, L^q) \leq K_{p,q}(t, y; L^p, L^q) \quad \text{for all } t > 0,$$

*then*

$$K_{p,q}^*(t, x; L^p, L^q) \leq \gamma_{p,q} K_{p,q}^*(t, y; L^p, L^q) \quad \text{for all } t > 0,$$

*where*

$$\gamma_{p,q} := \inf \left\{ \gamma > 0 : \inf_{\substack{x+y=\gamma, \\ x,y \geq 0}} \left( x^p + y^q \right) = 1 \right\}$$

*is the Sparr constant, which cannot be replaced by any smaller constant, and which satisfies the inequalities  $1 < \gamma_{p,q} < 2$ .*

3. SOME PROPERTIES OF CONCAVE AND CONVEX FUNCTIONS

We need some properties of concave and convex functions. More information about concave and convex functions and their properties can be found in [5, Ch. 3], [9], [10, Ch. 2, Section 1].

We denote by  $\mathcal{P}$  the set of all quasi-concave functions  $\rho : [0, \infty) \rightarrow [0, \infty)$  which are continuous, positive on  $\mathbf{R}_+ := (0, \infty)$  and such that  $\rho(st) \leq \max\{1, s\}\rho(t)$  for all  $s, t > 0$ . Let  $\tilde{\mathcal{P}}$  denote the subset of all concave functions in  $\mathcal{P}$ . Note that if  $\rho \in \mathcal{P}$ , then  $\tilde{\rho}$  defined by

$$\tilde{\rho}(t) := \inf_{s>0} \left(1 + \frac{t}{s}\right) \rho(s)$$

belongs to  $\tilde{\mathcal{P}}$  and

$$(3.1) \quad \rho(t) \leq \tilde{\rho}(t) \leq 2\rho(t) \quad \text{for all } t > 0.$$

Later on  $p'$  will always denote the conjugate number to  $p$ ,  $1 \leq p \leq \infty$ , that is,  $1/p + 1/p' = 1$  ( $1/\infty$  means 0).

**Lemma 3.1.** *Let  $1 < p < \infty$  and  $\varphi$  be a convex function on  $\mathbf{R}_+$ . The function  $u^{-p}\varphi(u)$  is increasing (decreasing) on  $\mathbf{R}_+$  if and only if the function  $u^{-p'}\varphi^*(u)$  is decreasing (increasing) on  $\mathbf{R}_+$ .*

*Proof* (cf. [7, Lemma 6.1.4]). Let  $u^{-p}\varphi(u)$  be increasing. If  $0 < u_1 \leq u_2$ , then

$$\begin{aligned} \varphi^*(u_2) &= \sup_{v>0} (vu_2 - \varphi(v)) = \sup_{v>0} \left( \left(\frac{u_2}{u_1}\right)^{p'-1} vu_2 - \varphi\left(\left(\frac{u_2}{u_1}\right)^{p'-1} v\right) \right) \\ &\leq \sup_{v>0} \left( \left(\frac{u_2}{u_1}\right)^{p'} vu_1 - \left(\frac{u_2}{u_1}\right)^{(p'-1)p} \varphi(v) \right) \\ &= \left(\frac{u_2}{u_1}\right)^{p'} \sup_{v>0} (vu_1 - \varphi(v)) = \left(\frac{u_2}{u_1}\right)^{p'} \varphi^*(u_1), \end{aligned}$$

which means that  $u^{-p'}\varphi^*(u)$  is decreasing. The remaining implications can easily be proved by also using the fact that for convex function  $\varphi$  we have  $\varphi^{**} = \varphi$ .  $\square$

Now we derive relations between some representations of  $\varphi$  and  $\varphi^{-1}$ .

**Lemma 3.2.** *Suppose that  $1 \leq p < q \leq \infty$  and, for some  $\rho \in \tilde{\mathcal{P}}$ ,*

$$\varphi^{-1}(u) = u^{1/p}\rho(u^{1/q-1/p}) \quad \text{for all } u > 0.$$

(a) *Then  $\varphi$  is convex.*

(b) *If  $q < \infty$ , then there exists a function  $h \in \mathcal{P}$  such that*

$$\varphi(u) = u^q h(u^{p-q}) \quad \text{for all } u > 0.$$

(c) *If  $1 < p < q < \infty$ , then there exists a function  $k \in \mathcal{P}$  such that*

$$\varphi^*(u) = u^{p'} k(u^{q'-p'}) \quad \text{for all } u > 0.$$

(d) *If  $q = \infty$  and  $\rho_*(t) := t\rho(1/t)$  satisfies  $\rho_*(\mathbf{R}_+) = \mathbf{R}_+$ , then  $\varphi(u) = \rho_*^{-1}(u)^p$  and  $\psi(u) = \varphi(u^{1/p})$  is a convex function.*

*Proof.* Proposition (a) follows from [16, Lemma 14.2].

(b) Since  $\rho(t)$  is increasing and  $\rho(t)/t$  is decreasing, for all  $s, t > 0$  we have

$$\min \{s^{1/p}, s^{1/q}\} \varphi^{-1}(t) \leq \varphi^{-1}(st) \leq \max \{s^{1/p}, s^{1/q}\} \varphi^{-1}(t).$$

Hence,  $\varphi^{-1}$  is strictly increasing on  $\mathbf{R}_+$  and  $\varphi^{-1}(\mathbf{R}_+) = \mathbf{R}_+$ . It is easy to see that the following statements are equivalent:

- (i)  $\varphi^{-1}(st) \leq \max \{s^{1/p}, s^{1/q}\} \varphi^{-1}(t)$  for all  $s, t > 0$ ;
- (ii)  $\varphi(st) \leq \max \{s^p, s^q\} \varphi(t)$  for all  $s, t > 0$ ;
- (iii)  $u^{-p} \varphi(u)$  is increasing and  $u^{-q} \varphi(u)$  is decreasing;
- (iv)  $h \in \mathcal{P}$ , where  $h$  is given by  $\varphi(u) = u^q h(u^{p-q})$ .

Now we prove (c). If  $1 < p < q < \infty$ , then, by Lemma 3.1, (iii) is equivalent to each of the following properties:

- (v)  $u^{-p'} \varphi^*(u)$  is decreasing and  $u^{-q'} \varphi^*(u)$  is increasing;
- (vi)  $\varphi^*(st) \leq \max \{s^{p'}, s^{q'}\} \varphi^*(t)$  for all  $s, t > 0$ ;
- (vii)  $k \in \mathcal{P}$ , where  $k$  is given by  $\varphi^*(u) = u^{p'} k(u^{q'-p'})$ .

Let us prove (d). First observe that  $\rho_*$  is concave (see, e.g., [16, Lemma 8.7]). By the assumption  $\rho_*(\mathbf{R}_+) = \mathbf{R}_+$ , we can see that  $\varphi^{-1}(\mathbf{R}_+) = \mathbf{R}_+$  and  $\varphi^{-1}$  is concave. Hence,  $\varphi$  is a finite valued convex function vanishing only at zero,  $\varphi(u) = \rho_*^{-1}(u)^p$ . We show that even  $\psi(u) = \varphi(u^{1/p})$  is convex. Since  $\rho_*$  is concave,  $\varphi_0 := \rho_*^{-1}$  is convex. Hence,

$$\varphi_0(u) = \varphi_0^{**}(u) = \sup_{v \geq 0} (uv - \varphi_0^*(v)).$$

To prove that

$$\varphi(u^{1/p}) = \varphi_0(u^{1/p})^p = \sup_{v > 0} (u^{1/p}v - \varphi_0^*(v))^p$$

is convex, it is enough to show that  $f(u) := (u^{1/p}v - \varphi_0^*(v))^p$  is a convex function. Since

$$f'(u) = (u^{1/p}v - \varphi_0^*(v))^{p-1} u^{1/p-1}v = \left( \frac{u^{1/p}v - \varphi_0^*(v)}{u^{1/p}} \right)^{p-1} v =: g(u)^{p-1}v$$

and  $g(u)$  is increasing, it follows that  $f$  is convex. □

Note that the previous lemma guarantees only that  $h, k \in \mathcal{P}$  (and  $h$  and  $k$  need not necessarily to be concave).

**Example 3.3.** If  $1 \leq p < q < \infty$  and  $\varphi^{-1}(u) = u^{1/p} \rho(u^{1/q-1/p})$  with  $\rho(t) = \min \{1, t\}$ , then  $\varphi(u) = u^q h(u^{p-q})$  with  $h(t) = \max \{1, t\}$ . Obviously,  $\rho \in \tilde{\mathcal{P}}$ , but  $h \in \mathcal{P} \setminus \tilde{\mathcal{P}}$ . In particular, if  $p = 2$  and  $q = 3$ , then  $\varphi^*(u) = u^{3/2} k(u^{2-3/2})$  with

$$k(t) = \begin{cases} t/4, & 0 \leq t \leq \sqrt{2}, \\ t^{-1} - t^{-3}, & \sqrt{2} \leq t \leq \sqrt{3}, \\ 2 \cdot 3^{-3/2}, & t \geq \sqrt{3}. \end{cases}$$

One can prove that  $k \in \tilde{\mathcal{P}}$ .

4. MODULAR ESTIMATES

For the proof of the first modular estimate, we need the following representation of concave functions, which goes back to J. Peetre [19] (see also [3, Lemma 5.4.3]).

**Lemma 4.1.** *Every function  $h \in \tilde{\mathcal{P}}$  can be represented in the following form:*

$$(4.1) \quad h(u) = a_h + b_h u + \int_0^\infty \min(u, t) dm(t), \quad \text{for all } u > 0,$$

where

$$a_h := \lim_{u \rightarrow 0^+} h(u), \quad b_h := \lim_{u \rightarrow \infty} \frac{h(u)}{u},$$

and  $m : \mathbf{R}_+ \rightarrow \mathbf{R}_+$  is a nondecreasing function (in fact,  $m(t) = -h'(t)$ ).

Now we are ready to prove some modular inequalities, which are the keys for the above-mentioned estimations of norms.

**Theorem 4.2.** *Let  $1 \leq p < q \leq \infty$  and  $T \in \mathcal{A}(L^p, L^q)$ .*

(a) *If  $q < \infty$  and  $\varphi(u) = u^q h(u^{p-q})$  for some  $h \in \tilde{\mathcal{P}}$  ( $\varphi$  not necessarily convex), then*

$$I_\varphi \left( \frac{Tx}{M} \right) \leq \gamma_{p,q} I_\varphi(x) \quad \text{for all } x \in L^p \cap L^q.$$

(b) *If  $q = \infty$  and  $\psi(u) = \varphi(u^{1/p})$  is convex, then*

$$I_\varphi \left( \frac{Tx}{2^{1-1/p} M} \right) \leq I_\varphi(x) \quad \text{for all } x \in L^p + L^\infty.$$

*Proof.* (a) Due to Lemma 4.1,  $h$  can be represented in the form (4.1). Hence,

$$(4.2) \quad \varphi(u) = u^q h(u^{p-q}) = a_h u^q + b_h u^p + \int_0^\infty \min(u^p, tu^q) dm(t), \quad u \in \mathbf{R}_+.$$

Consequently,

$$(4.3) \quad \begin{aligned} I_\varphi \left( \frac{Tx}{M} \right) &= \int_\Omega \varphi \left( \frac{|Tx(s)|}{M} \right) d\mu(s) = a_h \left\| \frac{Tx}{M} \right\|_q^q + b_h \left\| \frac{Tx}{M} \right\|_p^p \\ &+ \int_\Omega \left[ \int_0^\infty \min \left( \left( \frac{|Tx(s)|}{M} \right)^p, t \left( \frac{|Tx(s)|}{M} \right)^q \right) dm(t) \right] d\mu(s). \end{aligned}$$

Since the operator  $T$  is bounded in  $L^p$  and  $L^q$ , we get

$$(4.4) \quad \begin{aligned} a_h \left\| \frac{Tx}{M} \right\|_q^q + b_h \left\| \frac{Tx}{M} \right\|_p^p &\leq a_h \left( \frac{M_1}{M} \right)^q \|x\|_q^q + b_h \left( \frac{M_0}{M} \right)^p \|x\|_p^p \\ &\leq a_h \|x\|_q^q + b_h \|x\|_p^p \end{aligned}$$

and, according to Proposition 2.1,

$$K_{p,q} \left( t, \frac{Tx}{M}; L^p, L^q \right) \leq K_{p,q}(t, x; L^p, L^q) \quad \text{for all } t > 0.$$

By using Sparr's Lemma 2.2 we obtain

$$K_{p,q}^* \left( t, \frac{Tx}{M}; L^p, L^q \right) \leq \gamma_{p,q} K_{p,q}^*(t, x; L^p, L^q) \quad \text{for all } t > 0.$$

Hence, by the Fubini theorem and in view of the definition of  $K_{p,q}^*$ , we get

$$\begin{aligned}
 (4.5) \quad & \int_{\Omega} \left[ \int_0^{\infty} \min \left( \left( \frac{|Tx(s)|}{M} \right)^p, t \left( \frac{|Tx(s)|}{M} \right)^q \right) dm(t) \right] d\mu(s) \\
 &= \int_0^{\infty} K_{p,q}^* \left( t, \frac{Tx}{M}; L^p, L^q \right) dm(t) \\
 &\leq \gamma_{p,q} \int_0^{\infty} K_{p,q}^*(t, x; L^p, L^q) dm(t) \\
 &= \gamma_{p,q} \int_{\Omega} \left[ \int_0^{\infty} \min(|x(s)|^p, t|x(s)|^q) dm(t) \right] d\mu(s).
 \end{aligned}$$

Combining (4.4)–(4.5) and taking into account that  $\gamma_{p,q} > 1$  we obtain

$$\begin{aligned}
 I_{\varphi} \left( \frac{Tx}{M} \right) &\leq a_h \|x\|_q^q + b_h \|x\|_p^p + \gamma_{p,q} \int_{\Omega} \left[ \int_0^{\infty} \min(|x(s)|^p, t|x(s)|^q) dm(t) \right] d\mu(s) \\
 &\leq \gamma_{p,q} \left( a_h \|x\|_q^q + b_h \|x\|_p^p + \int_{\Omega} \left[ \int_0^{\infty} \min(|x(s)|^p, t|x(s)|^q) dm(t) \right] d\mu(s) \right) \\
 &= \gamma_{p,q} \int_{\Omega} \varphi(|x(s)|) d\mu(s) = \gamma_{p,q} I_{\varphi}(x).
 \end{aligned}$$

(b) For all  $x \in L^p + L^{\infty}$  and  $t > 0$ , according to the Krée formula (see [3, Theorem 5.2.1]),

$$(4.6) \quad \left( \int_0^t x^*(s)^p ds \right)^{1/p} \leq K(t^{1/p}, x; L^p, L^{\infty}) \leq 2^{1-1/p} \left( \int_0^t x^*(s)^p ds \right)^{1/p}.$$

The constant  $2^{1-1/p}$  cannot be improved (see [2]). Due to Proposition 2.1,

$$(4.7) \quad K \left( t, \frac{Tx}{M}; L^p, L^{\infty} \right) \leq K(t, x; L^p, L^{\infty}) \quad \text{for all } t > 0.$$

From (4.6) and (4.7) it follows that

$$\int_0^t \left( \frac{(Tx)^*(s)}{2^{1-1/p}M} \right)^p ds \leq \int_0^t x^*(s)^p ds \quad \text{for all } t > 0.$$

Since  $\psi(u) = \varphi(u^{1/p})$  is convex, by the Hardy-Littlewood-Pólya majorization theorem (see, e.g., [1, p. 88]),

$$\begin{aligned}
 & \int_0^{\infty} \varphi \left( \frac{(Tx)^*(s)}{2^{1-1/p}M} \right) ds = \int_0^{\infty} \psi \left( \left[ \frac{(Tx)^*(s)}{2^{1-1/p}M} \right]^p \right) ds \\
 & \leq \int_0^{\infty} \psi(x^*(s)^p) ds = \int_0^{\infty} \varphi(x^*(s)) ds.
 \end{aligned}$$

Since the modular  $I_{\varphi}$  is rearrangement-invariant (see (2.1)), the latter inequality gives (b).  $\square$

The method of the proof of part (a) is due to J. Peetre [20]. In this proof the estimation of the functional  $K_{p,q}^*$  (“ $K_{p,q}^*$ -monotonicity” property) is very essential. G. Sparr [22] proved that  $\gamma_{p,q}$  is the best possible constant in the estimation of the functional  $K_{p,q}^*$  (cf. Lemma 2.2). He also proved that  $1 < \gamma_{p,q} < 2$  for  $1 < p, q < \infty$ . Now we give more precise information about this constant.



**Proposition 4.3.** *Suppose  $1 \leq p, q < \infty$ .*

- (a) *Then  $\gamma_{p,q} = \gamma_{q,p}$  and  $\gamma_{1,1} = 1$ .*
- (b) *If  $q > 1$ , then*

$$(4.8) \quad \gamma_{p,q} = \inf \left\{ x + \left( \frac{p}{q} x^{p-1} \right)^{1/(q-1)} : x^p + \left( \frac{p}{q} x^{p-1} \right)^{q/(q-1)} = 1 \right\}.$$

*In particular,*

$$\gamma_{q,q} = 2^{1-1/q}, \quad \gamma_{1,q} = 1 + q^{1/(1-q)} - q^{q/(1-q)}.$$

- (c)  *$\gamma_{p,q}$  continuously increases in  $p$  and  $q$ .*
- (d) *If  $p \leq q$ , then  $2^{1-1/p} \leq \gamma_{p,q} \leq 2^{1-1/q}$ .*

*Proof.* Property (a) is obvious. Suppose  $q > 1$  and rewrite  $\gamma_{p,q}$  in the form

$$\gamma_{p,q} = \left\{ \gamma > 0 : \min_{0 \leq x \leq \gamma} F(\gamma, x, p, q) = 1 \right\},$$

where  $F(\gamma, x, p, q) = x^p + (\gamma - x)^q$  and  $\gamma \in (0, 2), x \in [0, \gamma], p \in [1, \infty), q \in (1, \infty)$ . Obviously, for  $x \in (0, \gamma)$ ,

$$(4.9) \quad \frac{\partial F}{\partial x} = px^{p-1} - q(\gamma - x)^{q-1}, \quad \frac{\partial^2 F}{\partial x^2} = p(p-1)x^{p-2} + q(q-1)(\gamma - x)^{q-2} > 0.$$

Hence,  $F(\gamma, x, p, q)$  is strictly convex in  $x$  and it has a unique minimum on  $[0, \gamma]$  at a point  $x(\gamma, p, q) \in (0, \gamma)$ , which is the solution of the equation

$$\frac{\partial F}{\partial x} = px^{p-1} - q(\gamma - x)^{q-1} = 0.$$

Clearly,

$$(4.10) \quad 0 < \gamma - 1 < x(\gamma, p, q) < 1,$$

and

$$\gamma = x(\gamma, p, q) + \left( \frac{p}{q} x(\gamma, p, q)^{p-1} \right)^{1/(q-1)},$$

$$\min_{0 \leq x \leq \gamma} F(\gamma, x, p, q) = x(\gamma, p, q)^p + \left( \frac{p}{q} x(\gamma, p, q)^{p-1} \right)^{q/(q-1)}.$$

So, (4.8) is proved. Using (4.8), one can easily calculate  $\gamma_{1,q}$  and  $\gamma_{q,q}$ . The proof of (b) is finished.

Let us prove (c). Due to (a), it is sufficient to prove that  $\gamma_{p,q}$  continuously increases in  $p$ . Consider  $\gamma = \gamma(p)$  such that

$$\min_{0 \leq x \leq \gamma} F(\gamma, x, p, q) = 1.$$

From conditions

$$(4.11) \quad \begin{cases} \min_{0 \leq x \leq \gamma} F(\gamma, x, p, q) - 1 = x^p + (\gamma - x)^q - 1 = 0, \\ \frac{\partial F}{\partial x}(\gamma, x, p, q) = px^{p-1} - q(\gamma - x)^{q-1} = 0, \end{cases}$$

taking into account that  $\frac{\partial^2 F}{\partial x^2} > 0$  (see (4.9)), one can find the derivative of the implicit function  $\gamma(p)$ :

$$\frac{d\gamma}{dp} = -\frac{x^p \log x}{q(\gamma - x)^{q-1}}.$$

Taking into account (4.10), we see that  $\frac{d\gamma}{dp} > 0$  for all  $\gamma(p)$  satisfying (4.11). Hence, such  $\gamma(p)$  continuously increases in  $p$ .

On the other hand,  $\gamma_{p,q}$  is the smallest  $\gamma(p)$  satisfying (4.11) (note that conditions (4.11) depend on  $p$ ). Thus,  $\gamma_{p,q}$  continuously increases in  $p$ . Property (c) is proved.

Property (d) follows from the monotonicity of  $\gamma_{p,q}$ :

$$2^{1-1/p} = \gamma_{p,p} \leq \gamma_{p,q} \leq \gamma_{q,q} = 2^{1-1/q}. \quad \square$$

## 5. THE MAIN INTERPOLATION THEOREM

Our main result reads:

**Theorem 5.1.** *Suppose  $1 \leq p < q \leq \infty$  and  $\rho \in \tilde{\mathcal{P}}$ . If  $q = \infty$  assume in addition that  $\rho_*(\mathbf{R}_+) = \mathbf{R}_+$ , where  $\rho_*(t) := t\rho(1/t)$ . If  $\varphi^{-1}(u) = u^{1/p}\rho(u^{1/q-1/p})$ , then the Orlicz space  $L^\varphi$  (with both the Luxemburg and the Orlicz norm) is an interpolation space for linear operators between  $L^p$  and  $L^q$ , and*

$$\|T\|_{L^\varphi \rightarrow L^\varphi} \leq C \max \{ \|T\|_{L^p \rightarrow L^p}, \|T\|_{L^q \rightarrow L^q} \},$$

where

(a)  $C \leq 2\gamma_{1,q} = 2(1 + q^{1/(q-1)} - q^{q/(q-1)}) \leq 2^{2-1/q} < 4$ , when  $1 = p < q < \infty$ .

(b)  $C \leq \min \left\{ (2\gamma_{p,q})^{1/p}, (2\gamma_{q',p'})^{1/q'} \right\} \leq 2^{1/(pq') + \min\{1/p, 1/q'\}} < 4$ , when  $1 < p < q < \infty$ .

(c)  $C \leq 2^{1-1/p} < 2$ , when  $1 \leq p < q = \infty$ .

In particular, if either  $1 < p < q \leq 2$  or  $2 \leq p < q \leq \infty$ , then  $C < 2$ .

*Proof.* First observe that the function  $\varphi$  is convex, due to Lemma 3.2(a). Hence,  $\varphi$  generates an Orlicz space  $L^\varphi$ .

Let  $1 \leq p < q < \infty$ . By Lemma 3.2(b), there is a function  $h \in \mathcal{P}$  such that  $\varphi(u) = u^q h(u^{p-q})$ . From (3.1) we see that  $\tilde{h} \in \tilde{\mathcal{P}}$  and

$$(5.1) \quad \varphi(u) \leq u^q \tilde{h}(u^{p-q}) \leq 2\varphi(u) \quad \text{for all } u > 0.$$

Applying Theorem 4.2(a) to the function  $\psi(u) = u^q \tilde{h}(u^{p-q})$  and taking into account (5.1), we obtain

$$(5.2) \quad I_\varphi \left( \frac{Tx}{M} \right) \leq I_\psi \left( \frac{Tx}{M} \right) \leq \gamma_{p,q} I_\psi(x) \leq 2\gamma_{p,q} I_\varphi(x) \quad \text{for all } x \in L^p \cap L^q.$$

Note that if  $A > 0$ , then

$$(5.3) \quad \begin{aligned} \varphi(Au) &= (Au)^q h((Au)^{p-q}) \leq A^q u^q \max \{ 1, A^{p-q} \} h(u^{p-q}) \\ &= \max \{ A^p, A^q \} \varphi(u). \end{aligned}$$

In particular,  $\varphi$  satisfies the  $\Delta_2$ -condition for all  $u \geq 0$ . If  $A = (2\gamma_{p,q})^{-1/p}$ , then from (5.2) and (5.3) we conclude

$$I_\varphi \left( \frac{Tx}{(2\gamma_{p,q})^{1/p} M} \right) \leq \frac{1}{2\gamma_{p,q}} I_\varphi \left( \frac{Tx}{M} \right) \leq I_\varphi(x) \quad \text{for all } x \in L^p \cap L^q,$$

which gives

$$(5.4) \quad \|Tx\|_\varphi \leq (2\gamma_{p,q})^{1/p} M \|x\|_\varphi, \quad \|Tx\|_\varphi^0 \leq (2\gamma_{p,q})^{1/p} M \|x\|_\varphi^0$$

for all  $x \in L^p \cap L^q$ . Since  $\varphi$  satisfies the  $\Delta_2$ -condition for all  $u \geq 0$ , it follows that  $L^p \cap L^q$  is dense in  $L^\varphi$  (see [9, Ch. 2] for the case of  $N$ -functions and a finite

measure, in the general case this result can be obtained analogously). Hence, (5.4) is true for all  $x \in L^\varphi$ . This fact and Proposition 4.3(d) show that

$$(5.5) \quad C \leq (2\gamma_{p,q})^{1/p} \leq 2^{(2-1/q)/p} < 4.$$

Now we will prove the second estimate in (b) by using duality arguments. Suppose  $1 < p < q < \infty$  and  $T \in \mathcal{A}(L^p, L^q)$ . Then  $T$  maps  $L^p + L^q$  into itself, but  $T$  maps also  $L^p \cap L^q$  into itself (see, e.g., [10, Ch. 1, Lemma 4.1]). Since  $L^p \cap L^q$  is dense in  $L^p$  and  $L^q$ , then  $(L^p \cap L^q)^* = L^{p'} + L^{q'}$  (see, e.g., [10, Ch. 1, Th. 3.1]). Therefore,  $T^* \in \mathcal{A}(L^{p'}, L^{q'})$ . Due to Lemma 3.2(c), there is a function  $k \in \mathcal{P}$  such that  $\varphi^*(u) = u^{p'}k(u^{q'-p'})$  for all  $u > 0$ . As above one can prove that

$$(5.6) \quad \|T^*x\|_{\varphi^*} \leq (2\gamma_{q',p'})^{1/q'} M\|x\|_{\varphi^*}, \quad \|T^*x\|_{\varphi^*}^0 \leq (2\gamma_{q',p'})^{1/q'} M\|x\|_{\varphi^*}^0$$

for all  $x \in L^{p'} \cap L^{q'}$ , and  $\varphi^*$  satisfies the  $\Delta_2$ -condition for all  $u \geq 0$ . Consequently, (5.6) holds for all  $x \in L^{\varphi^*}$ . Taking into account duality of the Orlicz and Luxemburg norms and Proposition 4.3, this gives

$$(5.7) \quad C \leq (2\gamma_{q',p'})^{1/q'} \leq 2^{(2-1/p')/q'} < 4.$$

Since

$$\min \left\{ 2^{(2-1/q)/p}, 2^{(2-1/p')/q'} \right\} = 2^{1/(pq') + \min\{1/p, 1/q'\}},$$

we obtain from (5.5) and (5.7) that  $C < 2$  in the cases  $1 < p < q \leq 2$  or  $2 \leq p < q < \infty$ . The proof of (b) is complete.

Let  $1 \leq p < q = \infty$ . In view of Lemma 3.2(d), the function  $\psi(u) = \varphi(u^{1/p})$  is convex. Hence, by Theorem 4.2(b), we obtain the modular estimate

$$I_\varphi \left( \frac{Tx}{2^{1-1/p}M} \right) \leq I_\varphi(x) \quad \text{for all } x \in L^\varphi,$$

which implies

$$\|Tx\|_\varphi \leq 2^{1-1/p}M\|x\|_\varphi, \quad \|Tx\|_\varphi^0 \leq 2^{1-1/p}M\|x\|_\varphi^0$$

for all  $x \in L^\varphi$ . This gives (c). □

*Remark 5.2.* Obviously Theorem 5.1 (a), (b) can, in fact, be reformulated in terms of Simonenko indices (1.5): if

$$1 \leq p \leq a_\varphi \leq b_\varphi \leq q < \infty,$$

then the Orlicz space  $L^\varphi$  is an interpolation space between  $L^p$  and  $L^q$  with the same interpolation constant as in Theorem 5.1 (a), (b).

*Remark 5.3.* If  $1 \leq p < q < \infty$  and  $\varphi(u) = u^q h(u^{p-q})$ , where  $h \in \tilde{\mathcal{P}}$ , then from the proof of the above theorem it follows that  $L^\varphi$  is an interpolation space between  $L^p$  and  $L^q$ , and we have a better estimate of the interpolation constant

$$C \leq (\gamma_{p,q})^{1/p} \leq 2^{1/(q'p)} < 2.$$

We illustrate Remark 5.3 with the following example:

**Example 5.4.** If  $r \geq (3 + \sqrt{5})/2$ , then  $\varphi(u) = u^r(1 + |\log u|)$  is convex. For every  $p$  and  $q$  such that  $1 \leq p < r < q < \infty$  and  $r - p = q - r$ , we have  $\varphi(u) = u^q h(u^{p-q})$  with  $h(u) = \sqrt{u}(1 + (q - p)^{-1}|\log u|)$ . One can prove that  $h \in \tilde{\mathcal{P}}$ .

*Remark 5.5.* In connection with Theorem 5.1(c) let us mention that for the case  $p = 1$  and  $q = \infty$  it coincides with the well-known Orlicz interpolation theorem. More precisely, Orlicz proved it in 1934 with certain constant  $C > 1$  but from the Calderón-Mitjagin interpolation theorem it follows with the constant 1 (see [10, Ch. 2, Th. 4.9]; cf. also [15] for the direct proof). Moreover, G. G. Lorentz and T. Shimogaki [13, Theorem 7] observed that for the function

$$\varphi(u) = \int_0^u (u-t)^p dm(t)$$

with increasing function  $m : \mathbf{R}_+ \rightarrow \mathbf{R}_+$ , the interpolation constant  $C$  is 1.

#### ACKNOWLEDGEMENTS

This paper was done during a visit of the first author at Luleå University of Technology in November-December 1999. The research was supported by a grant from the Royal Swedish Academy of Sciences for cooperation between Sweden and the former Soviet Union (project 35124). We are grateful to both institutions for support and hospitality.

#### REFERENCES

- [1] C. Bennett, R. Sharpley, *Interpolation of Operators*, Academic Press, London 1988. MR **89e**:46001
- [2] J. Bergh, *A generalization of Steffensen's inequality*, J. Math. Anal. Appl. **41** (1973), 187–191. MR **47**:3621
- [3] J. Bergh, J. Löfström, *Interpolation Spaces. An Introduction*, Springer-Verlag, Berlin 1976. MR **58**:2349
- [4] D. W. Boyd, *Spaces between a pair of reflexive Lebesgue spaces*, Proc. Amer. Math. Soc. **18** (1967), 215–219. MR **35**:3427
- [5] Yu. A. Brudnyi, N. Ya. Krugljak, *Interpolation Functors and Interpolation Spaces, I*, North-Holland, Amsterdam–New York–Tokyo 1991. MR **93b**:46141
- [6] A. Cianchi, *An optimal interpolation theorem of Marcinkiewicz type in Orlicz spaces*, J. Funct. Anal. **153** (1998), 357–381. MR **99d**:46039
- [7] I. Genebashvili, A. Gogatishvili, V. Kokilashvili, M. Krbeč, *Weight Theory for Integral Transforms on Spaces of Homogeneous Type*, Pitman Monographs and Surveys in Pure and Applied Math. **92**, Addison Wesley Longman 1998. CMP 2001:03
- [8] J. Gustavsson, J. Peetre, *Interpolation of Orlicz spaces*, Studia Math. **60** (1977), 33 – 59. MR **55**:11021
- [9] M. A. Krasnoselskii, Ya. B. Rutickii, *Convex Functions and Orlicz Spaces*, Noordhoff Ltd., Groningen 1961. MR **23**:A4016
- [10] S. G. Krein, Ju. I. Petunin, E. M. Semenov, *Interpolation of Linear Operators*, AMS Trans. of Math. Monographs **54**, Providence, R.I. 1982. MR **84j**:46103
- [11] N. Krugljak, L. Maligranda, L.-E. Persson, *A Carlson type inequality with blocks and interpolation*, Studia Math. **104** (1993), 161–180. MR **94k**:46153
- [12] J. Lindenstrauss, L. Tzafriri, *Classical Banach Spaces II. Function Spaces*, Springer Verlag, New York–Berlin 1979. MR **81c**:46001
- [13] G. G. Lorentz, T. Shimogaki, *Interpolation theorems for the pairs of spaces  $(L^p, L^\infty)$  and  $(L^1, L^q)$* , Trans. Amer. Math. Soc. **159** (1971), 207–221. MR **52**:1347
- [14] L. Maligranda, *Indices and interpolation*, Dissert. Math. **234** (1985), 1–49. MR **87k**:46059
- [15] L. Maligranda, *Some remarks on Orlicz's interpolation theorem*, Studia Math. **95** (1989), 43–58. MR **90k**:46160
- [16] L. Maligranda, *Orlicz Spaces and Interpolation*, Seminars in Mathematics **5**, Univ. Estadual de Campinas, Campinas SP, Brazil 1989.
- [17] V. I. Ovchinnikov, *Interpolation theorems resulting from an inequality of Grothendieck*, Funkts. Analiz i Prilozh., **10** (1976), 45–54 (in Russian); English transl.: Funct. Anal. Appl. **10** (1976), 287–294. MR **55**:3818

- [18] V. I. Ovchinnikov, *The Method of Orbits in Interpolation Theory*, Math. Rep. 1 (1984), No. 2, 349–515. MR **88d**:46136
- [19] J. Peetre, *On interpolation functions*, Acta Math. Sci. Szeged **27** (1966), 167–171. MR **34**:6523
- [20] J. Peetre, *A new approach in interpolation spaces*, Studia Math. **34** (1970), 23–42. MR **41**:8985; Correction MR **58**:23647
- [21] I. B. Simonenko, *Interpolation and extrapolation of linear operators in Orlicz spaces*, Mat. Sb. (N.S.) **63**(1964), 536–553 (in Russian). MR **33**:7839
- [22] G. Sparr, *Interpolation of weighted  $L_p$ -spaces*, Studia Math. **62** (1978), 229 – 271. MR **80d**:46055

DEPARTMENT OF MATHEMATICS AND PHYSICS, SOUTH UKRAINIAN STATE PEDAGOGICAL UNIVERSITY, STAROPORTOFRANKOVSKAYA 26, 65020 ODESSA, UKRAINE

*E-mail address*: `karlik@paco.net`

*Current address*: Departamento de Matemática, Instituto Superior Técnico, Av. Rovisco Pais 1, 1049-001, Lisbon, Portugal

*E-mail address*: `akarlov@math.ist.utl.pt`

DEPARTMENT OF MATHEMATICS, LULEÅ UNIVERSITY OF TECHNOLOGY, 971 87 LULEÅ, SWEDEN

*E-mail address*: `lech@sm.luth.se`