

ON THE PARTIALLY ORDERED SET OF PRIME IDEALS OF A DISTRIBUTIVE LATTICE

RAYMOND BALBES

1. Introduction. For a distributive lattice L , let $\mathcal{P}(L)$ denote the poset of all prime ideals of L together with \emptyset and L . This paper is concerned with the following type of problem. Given a class \mathcal{C} of distributive lattices, characterize all posets P for which $P \cong \mathcal{P}(L)$ for some $L \in \mathcal{C}$. Such a poset P will be called *representable over \mathcal{C}* . For example, if \mathcal{C} is the class of all relatively complemented distributive lattices, then P is representable over \mathcal{C} if and only if P is a totally unordered poset with $0, 1$ adjoined. One of our main results is a complete characterization of those posets P which are representable over the class of distributive lattices which are generated by their meet irreducible elements. The problem of determining which posets P are representable over the class of all distributive lattices appears to be very difficult. (See [2].) It will be shown that this problem is equivalent to the embeddability of P as the set of meet irreducible elements of a certain distributive algebraic lattice.

Results concerning the degree to which $\mathcal{P}(L)$ determines L are presented in §§ 4 and 5. It is shown that if $\mathcal{P}(L)$ is isomorphic with the power set of a non-empty set X , then L is a free distributive lattice.

2. Preliminaries. Let P be a poset and S a non-empty subset of P . Denote by $(S]_P$, or simply $(S]$, the set $\{x \in P \mid x \leq s \text{ for some } s \in S\}$; abbreviate $(\{s\}]$ by $(s]$; $[S)$ is defined dually. S is *hereditary* if $x \leq y$ and $y \in S$ imply $x \in S$. For a non-empty set X , 2^X will denote the poset of all subsets of X .

The class of all distributive lattices will be denoted by \mathcal{L} . As stated above, $\mathcal{P}(L)$ is the poset of prime ideals of L together with \emptyset and L (we avoid unnecessary technical complications by *not* excluding \emptyset and L from $\mathcal{P}(L)$). For each $x \in L$, let $x^* = \{I \in \mathcal{P}(L) \mid x \notin I\}$. Note that $\emptyset \in x^*$ and that $L \notin x^*$. It is well known (see, e.g., [4]) that the prime ideal theorem implies (i) $L \cong \{x^* \mid x \in L\}$, and (ii) if T_1, T_2 are non-empty subsets of L and

$$\cap \{x^* \mid x \in T_1\} \subseteq \cup \{y^* \mid y \in T_2\},$$

then there exist finite subsets $\emptyset \neq T_1' \subseteq T_1, \emptyset \neq T_2' \subseteq T_2$ such that

$$\cap \{x^* \mid x \in T_1'\} \subseteq \cup \{y^* \mid y \in T_2'\}.$$

Finally, recall that an element $x \in L$ is meet irreducible (*M.I.*) if $yz \leq x$ implies that $y \leq x$ or $z \leq x$. So $(x] \in \mathcal{P}(L)$ if and only if x is *M.I.* The class

Received April 30, 1971. This research was supported, in part, by NSF Grant GP11893.

of all distributive lattices which are generated by *M.I.* elements will be denoted by \mathcal{A} .

3. Posets representable over \mathcal{A} . We begin by giving a sufficient condition for a poset P to be representable over \mathcal{L} .

Definition 1. Let P be a poset with $0, 1$. A non-zero element $k \in P$ is *weakly compact* provided that if $\emptyset \neq D \subseteq P$, $\sum_P D$ exists, and $k \leq \sum_P D$, then there exists $\{d_1, \dots, d_n\} \subseteq D$ such that $[d_1] \cap \dots \cap [d_n] \subseteq [k]$.

Definition 2. A non-empty subset D of a poset P will be called *prime* provided that if $\{s, \dots, s_n, t_1, \dots, t_m\}$ are weakly compact in P , $\{t_1, \dots, t_m\} \subseteq D$ and $[t_1] \cap \dots \cap [t_m] \subseteq [s_1] \cup \dots \cup [s_n]$, then $s_i \in D$ for some $i \in \{1, \dots, n\}$.

Let P be a poset with $0 < 1$ and K the weakly compact members of P . Consider the following two conditions on P :

- (C1) If $p \not\leq q$ then there exists $k \in K$ such that $k \leq p$ and $k \not\leq q$.
- (C2) If D is a prime subset of P then $\sum_P D$ exists.

THEOREM 3. *If P is a poset with $0 < 1$ that satisfies (C1) and (C2) then P is representable over \mathcal{L} .*

Proof. Let R be the ring of sets generated by $\{[k]' \mid k \in K\}$ where $[k]' = P \sim [k]$. For each $p \in P$, let $\psi(p) = \{A \in R \mid p \notin A\}$. Then $\psi(p) \in \mathcal{P}(R)$, so $p \rightarrow \psi(p)$ defines a function from P into $\mathcal{P}(R)$. Now $[k]'$ is a hereditary subset of P for each $k \in K$, so R is a ring of hereditary sets. It follows that ψ preserves order. Condition (C1) implies that $[k]' \in \psi(p) \sim \psi(q)$, so $p \leq q$ if and only if $\psi(p) \leq \psi(q)$.

Next, observe that $\psi(0) = \emptyset$ and $\psi(1) = R$. Now let I be a prime ideal in R and set $D = \{k \in K \mid [k]' \in I\}$. Then $D \neq \emptyset$ since $I \neq \emptyset$. Also, if

$$\{s_1, \dots, s_n, t_1, \dots, t_m\} \subseteq K, \quad \{t_1, \dots, t_m\} \subseteq D,$$

and

$$[t_1] \cap \dots \cap [t_m] \subseteq [s_1] \cup \dots \cup [s_n],$$

then

$$[s_1]' \cap \dots \cap [s_n]' \subseteq [t_1]' \cup \dots \cup [t_m]';$$

but I is a prime and $\{[t_1]', \dots, [t_m]'\} \subseteq I$, so $s_i \in D$ for some $i \in \{1, \dots, n\}$. That is, D is a prime subset of P .

By (C2), $p = \sum_P D$ exists and the proof will be completed by showing that $\psi(p) = I$. Let

$$A = \bigcap_{i=1}^n \left(\bigcup_{j=1}^{n_i} [k_{ij}]' \right).$$

First, suppose that $A \in \psi(p)$; then $p \notin A$, so

$$p \in \bigcap_{j=1}^{n_{i_0}} [k_{i_0 j}]$$

for some $i_0 \in \{1, \dots, n\}$. Let $j \in \{1, \dots, n_{i_0}\}$. Then $k_{i_0j} \leq p = \sum_P D$, and since k_{i_0j} is weakly compact there exists $\{q_1, \dots, q_n\} \subseteq D$ such that $[q_1] \cap \dots \cap [q_n] \subseteq [k_{i_0j}]$. But D is prime, so $k_{i_0j} \in D$; hence $[k_{i_0j}]' \in I$. This means that $[k_{i_0j}]' \in I$ for $i = 1, \dots, n_{i_0}$, and so $A \in I$. To show the reverse inclusion, let $A \in I$. Say

$$\bigcup_{j=1}^{n_1} [k_{1j}]' \in I.$$

Then $\{k_{1j}, \dots, k_{1n_1}\} \subseteq D$ so $k_{1j} \leq \sum D = p$ for each j . Therefore,

$$p \notin \bigcup_{j=1}^{n_1} [k_{1j}]'$$

and $p \notin A$ which means that $A \in \psi(p)$.

Next, we determine the weakly compact elements in $\mathcal{P}(L)$ for $L \in \mathcal{A}$.

LEMMA 4. *If $L \in \mathcal{A}$ then the following are equivalent:*

- (i) *I is weakly compact in $\mathcal{P}(L)$;*
- (ii) *$I = [x]_L$ for some M.I. element $x \in L$.*

Proof. Let M be the M.I. elements of I and set $P = \mathcal{P}(L)$.

(i) \Rightarrow (ii). Let $D = \{(u)_L \mid u \in M \cap I\}$. $D \neq \emptyset$ since $I \neq \emptyset$. Now I is the ideal generated by $\cup D$ so, in fact, $I = \sum_P D$. Since I is weakly compact, there exist $\{(u_1)_L, \dots, (u_n)_L\} \subseteq D$ such that

$$[(u_1)_L]_P \cap \dots \cap [(u_n)_L]_P \subseteq [I]_P.$$

Now let $x = u_1 + \dots + u_n$. We will show $[x] = I$. Indeed, $u_i \in I$ for each $i \in \{1, \dots, n\}$ so $[x] \subseteq I$. If $I \not\subseteq [x]$ then there exists $y \in I$ such that $y \not\leq x$ and therefore a prime ideal J such that $x \in J$, $y \notin J$. But $x \in J$ implies $J \in [(u_1)_L]_P \cap \dots \cap [(u_n)_L]_P \subseteq [I]_P$ so $y \in I \subseteq J$, which is a contradiction. Thus, $I = [x]$ and since I is prime, x is M.I.

(ii) \Rightarrow (i) Firstly, $x \in M$ implies that $[x] \in P$. Suppose that $\emptyset \neq D \subseteq P$, $\sum_P D$ exists, and that $[x] \subseteq \sum_P D$. Let J be the ideal generated by $\cup D$ and suppose that $x \notin J$. Then there is a prime ideal J' such that $x \notin J'$, $J \subseteq J'$. So $K \subseteq J \subseteq J'$ for each $K \in D$ and hence $[x] \subseteq \sum_P D \subseteq J'$, which is a contradiction. Thus, since $x \in J$, there exist I_1, \dots, I_n in D and $x_i \in I_i$ such that $x \leq x_1 + \dots + x_n$. Finally,

$$[I_1] \cap \dots \cap [I_n] \subseteq [(x)_L]_P;$$

for if $K \in P$ and $K \in [I_1] \cap \dots \cap [I_n]$, then for each $i \in \{1, \dots, n\}$, $x_i \in I_i \subseteq K$, so $x \in K$. Hence $[x]_L \subseteq K$.

THEOREM 5. *A poset P is representable over \mathcal{A} if and only if P satisfies (C1) and (C2).*

Proof. Suppose first that P satisfies (C1) and (C2). Now the ring R constructed in Theorem 3 was generated by $\{[k]' \mid k \in K\}$ for some non-empty subset $K \subseteq P$. From this, it is easily verified that $[k]'$ is M.I. for each $k \in K$.

Conversely, suppose that $P = \mathcal{P}(L)$, where L is generated by the set M of $M.I.$ elements of L . For (C1), suppose that $\{I, J\} \subseteq P$ and that $I \not\subseteq J$. Then there is an element $x \in I \sim J$. But since x is a sum of products of members of M , there is an element $u \in M \cap (I \sim J)$. By Lemma 4, $[u]$ is weakly compact in P , and also $[u] \subseteq I$, $[u] \not\subseteq J$. For (C2), suppose that D is prime in P . Let J be the ideal in L generated by $\cup D$. Clearly, if $J \in P$, then $\sum D$ will exist and equal J . Now to prove that J is a prime ideal, it is sufficient to show that if $\{u_1, \dots, u_n\} \subseteq M$ and $u_1 \cdot \dots \cdot u_n \in J$, then $u_i \in J$ for some $i \in \{1, \dots, n\}$. But if $u_1 \cdot \dots \cdot u_n \in J$, then there are members I_1, \dots, I_m of D and elements $x_i \in I_i$ such that $u_1 \cdot \dots \cdot u_n \leq x_1 + \dots + x_m$. Now I_i is weakly compact since D is prime, so $I_i = [y_i]$ where $y_i \in M$ for each $i \in \{1, \dots, m\}$. Hence $u_1 \cdot \dots \cdot u_n \leq y_1 + \dots + y_m$ and so

$$[(y_1]_L)_P \cap \dots \cap [(y_m]_L)_P \subseteq [(u_1]_L)_P \cup \dots \cup [(u_n]_L)_P.$$

Invoking the primeness of D again, we find that $[u_i]_L \in D$ for some i , so $u_i \in J$.

To show how conditions (C1) and (C2) can be applied in specific cases we present the following corollary:

COROLLARY 6. *If P is a poset with $0 < 1$ and $[p]$ is finite for each $p \neq 0$, then it is representable over \mathcal{A} .*

Proof. Let D be a non-empty subset of $P \sim \{0\}$. For each finite, non-empty subset $T \subseteq D$, $\cap_{t \in T} [t]$ is finite and contains 1. Let n be the least number of elements in $\cap_{t \in T} [t]$ for any such $T \subseteq D$ and let T_0 be a finite non-empty subset of D such that $\cap_{t \in T_0} [t] = \{x_1, \dots, x_n\}$. Then the elements $\{x_1, \dots, x_n\}$ are all upper bounds of D , for clearly $t \leq x_i$ for all $t \in T_0$ and if $d \in D \sim T_0$, then

$$(\cap_{t \in T_0} [t]) \cap [d] \subseteq \cap_{t \in T_0} [t] = \{x_1, \dots, x_n\},$$

so by the minimality of n ,

$$(\cap_{t \in T_0} [t]) \cap [d] = \{x_1, \dots, x_n\}$$

and hence $d \leq x_i$.

We now proceed to verify (C1). Suppose that $p \neq 0$, $\sum_P D$ exists, and that $p \leq \sum_P D$. But then $u \in \cap_{t \in T_0} [t]$ implies that $x_i \leq u$ for some $i \in \{1, \dots, n\}$ and as x_i is an upper bound for D , $p \leq \sum_P D \leq x_i \leq u$. Thus, $\cap_{t \in T_0} [t] \subseteq [p]$. For (C2), suppose that D is prime in P . Let $\{u_1, \dots, u_m\}$ be the minimal elements of $\{x_1, \dots, x_n\}$ so that $\cap_{t \in T_0} [t] = [u_1] \cup \dots \cup [u_m]$. By the definition of prime, $u_{i_0} \in D$ for some $i_0 \in \{1, \dots, m\}$. But then

$$(\cap_{t \in T_0} [t]) \cap [u_{i_0}] \subseteq \{x_1, \dots, x_n\}$$

and again by the minimality of n ,

$$[u_{i_0}] = (\cap_{t \in T_0} [t]) \cap [u_{i_0}] = \{x_1, \dots, x_n\}.$$

It follows that $\sum_P D$ exists and equals u_{i_0} .

COROLLARY 7. *Every finite poset and every totally unordered poset, with 0 and 1 adjoined, is representable over \mathcal{A} .*

4. Uniqueness of posets representable over \mathcal{A} . Corollary 7 implies that posets representable over \mathcal{A} may also be representable by distributive lattices outside of \mathcal{A} . Indeed, choose a non-atomic Boolean algebra B . Then there is a lattice $L \in \mathcal{A}$ such that $\mathcal{P}(B) = \mathcal{P}(L)$. However, within the class \mathcal{A} , we do have uniqueness.

THEOREM 8. *If L and L' are members of \mathcal{A} and $\mathcal{P}(L) \cong \mathcal{P}(L')$, then $L \cong L'$.*

Proof. Let M and M' be the sets of *M.I.* elements of L and L' , respectively, and let $f: \mathcal{P}(L) \rightarrow \mathcal{P}(L')$ be an isomorphism. Since f induces an isomorphism between the set of weakly compact elements of $\mathcal{P}(L)$ and the set of weakly compact elements of $\mathcal{P}(L')$, Lemma 4 implies the existence of an isomorphism $g: M \rightarrow M'$ such that $f([x]) = [g(x)]$. To show that g can be extended to a homomorphism $G: L \rightarrow L'$, it is sufficient to prove that if S and T are finite non-empty subsets of M , and $\prod S \leq \sum T$, then $\prod g(S) \leq \sum g(T)$. Indeed, this condition implies that the function $G: L \rightarrow L'$ defined by

$$G(\prod S_1 + \dots + \prod S_n) = \prod g(S_n) + \dots + \prod g(S_n)$$

is well defined. It is easy then to verify that G is a homomorphism; the details can be found, for example, in [1, Lemma 1.7].

Now suppose that $\prod g(S) \not\leq \sum g(T)$. Then there exists $I \in P(L')$ such that $\sum g(T) \in I$ and $\prod g(S) \notin I$. For each $t \in T$, $g(t) \in I$, so $f([t]) = [g(t)] \subseteq I$. Hence, $t \in [t] \subseteq f^{-1}(I)$. But then $\sum T \in f^{-1}(I)$, so $s \in f^{-1}(I)$ for some $s \in S$ and, therefore, $[s] \subseteq f^{-1}(I)$. Finally, $g(s) \in [g(s)] = f([s]) \subseteq I$, which is a contradiction. Thus, there is a homomorphism $G: L \rightarrow L'$ such that $G|M = g$. Similarly, there is a homomorphism $G': L' \rightarrow L$ such that $G'|M' = g^{-1}$. It follows that G is an isomorphism.

The existence and uniqueness of representable chains can now be described completely.

THEOREM 9. *If C is a chain which is representable over \mathcal{L} , then C is complete and each interval $(a, b]$ contains an element with an immediate predecessor. Moreover, the representation of C over \mathcal{L} is unique. Conversely, if C is a complete chain in which each interval $(a, b]$ contains an element with an immediate predecessor then $C \cong \mathcal{P}(C_1)$ for some chain C_1 .*

Proof. If $C = \mathcal{P}(L)$ for some $L \in \mathcal{L}$, then C is closed under arbitrary unions, so C is complete. For $\{I, J\} \subseteq \mathcal{P}(L)$, if $I \subset J$, then there is an element $x \in J \sim I$, so $I \subset [x] \subseteq J$. Since C is a chain, so is L , and hence $[x] \in \mathcal{P}(L)$. The immediate predecessor of $[x]$ is $\{u \in L | u < x\}$. Next, if $\mathcal{P}(L) \cong C \cong \mathcal{P}(L')$, then L and L' are chains and hence in \mathcal{M} . By Theorem 8, $L \cong L'$. For the converse, it is sufficient to prove that if $c \in C$ has an immediate

predecessor c' , then c is weakly compact. Thus, suppose that $c \leq \sum_P D$. If $d < c$ for all $d \in D$, then $d \leq c'$, so $c \leq \sum_P D \leq c' < c$. Hence, $c \leq d_0$ for some $d_0 \in D$.

5. Free distributive lattices. In this section we show that $P = 2^X$ is representable only as the free distributive lattice on $|X|$ free generators. The fact that $\mathcal{P}(L) \cong 2^X$ when L is free is well known.

LEMMA 10. *Let L be a distributive lattice and suppose that $P = \mathcal{P}(L)$ is complete. If T is a finite non-empty set of M.I. elements of L , then*

$$(3) \quad \sum_P \{ \langle t \rangle \mid t \in T \} = \langle \sum T \rangle,$$

and $\sum T$ is M.I. in L .

Proof. For each $t \in T$, $t \in \langle t \rangle \subseteq \sum_P \{ \langle t \rangle \mid t \in T \}$, so

$$\langle \sum T \rangle \subseteq \sum_P \{ \langle t \rangle \mid t \in T \}.$$

Conversely, if $u \notin \langle \sum T \rangle$, then there is a prime ideal I such that $u \notin I$, $\sum T \in I$. But then $T \subseteq I$, so $\langle t \rangle \subseteq I$ for each $t \in T$. Hence $\sum_P \{ \langle t \rangle \mid t \in T \} \subseteq I$, and so $u \notin \sum_P \{ \langle t \rangle \mid t \in T \}$. Since $\langle \sum T \rangle \in P$, $\sum T$ is M.I.

LEMMA 11. *Let $L \in \mathcal{L}$ and let $P = \mathcal{P}(L) \cong 2^X$ for some $X \neq \emptyset$. Then I is an atom in P if and only if $I = \langle m \rangle$, where m is M.I. in L and is minimal in the set of all M.I. elements in L .*

Proof. Sufficiency. Suppose that I has no greatest element. Then for each $u \in I$, there exists $v_u \in I$ and $I(u, v_u) \in P$ such that $v_u \not\leq u$, $u \in I(u, v_u)$, and $v_u \notin I(u, v_u)$. Let $S = \{ I(u, v_u) \mid u \in I \}$. Since $v_u \in I \sim I(u, v_u)$, we have $I \not\subseteq J$ for all $J \in S$. But I is an atom in P which implies that $I \cdot J = 0_P$ for all $J \in S$. Now $I \subseteq \cup S \subseteq \sum_P S$, and since the Boolean algebra P is $(2, \infty)$ -distributive,

$$\begin{aligned} I &= I \cdot \sum_P S \\ &= \sum_P \{ I \cdot J \mid J \in S \} \\ &= 0_P, \end{aligned}$$

contradicting the definition of an atom.

So I has a greatest element m . It follows that $I = \langle m \rangle$, m is M.I., and is, in fact, minimal in the set of all M.I. elements in L .

Necessity. Under the conditions of the hypothesis, $\langle m \rangle \in P$ and $\langle m \rangle \neq \emptyset = 0_P$. So there is an atom $J \in P$ such that $J \subseteq \langle m \rangle$. But from the converse, $J = \langle n \rangle$ where n is M.I. Since m is minimal in the set of M.I. elements and $n \leq m$, we have $n = m$, so $\langle m \rangle = J$ is an atom in P .

THEOREM 12. *If L is a distributive lattice and $\mathcal{P}(L) \cong 2^X$ for some non-empty set X , then L is the free distributive lattice on $|X|$ free generators.*

Proof. Let $P = \mathcal{P}(L)$ and let S be the minimal elements in the set of $M.I.$ elements of L . $|S| = |X| > 0$ by Lemma 11. We prove first that S is an independent set.

Let T_1 and T_2 be finite non-empty subsets of S such that $\prod T_1 \leq \sum T_2$. By Lemma 10, $\sum T_2$ is $M.I.$, so there exists $t_1 \in T_1$ such that $t_1 \leq \sum T_2$.
Now

$$(t_1] \subseteq (\sum T_2] = \sum_P \{(t] | t \in T_2\},$$

and since $t_1 \in S$, $(t_1]$ is an atom in P ; so there exists $t_2 \in T_2$ such that $(t_1] \subseteq (t_2]$. The minimality of t_2 and $t_1 \leq t_2$ imply $t_1 = t_2$.

Since independent sets generate free distributive lattices, it suffices to prove that S generates L . For this purpose, let $S^2 = \{\sum T | T \text{ is a finite non-empty subset of } S\}$. Recall from Lemma 10 that the members of S^2 are $M.I.$ in L . We prove that if $I \in P$ and $I \neq \emptyset$, then $I \cap S^2 \neq \emptyset$ and

$$\sum_P \{(t] | t \in I \cap S^2\} = \cup \{(t] | t \in I \cap S^2\}.$$

Since $I \neq \emptyset$, it is a sum of atoms in P . By Lemma 11, there is a member $y \in S$ such that $(y] \subseteq I$, so $y \in I \cap S \subseteq I \cap S^2$. For the second part of the assertion it is easily verified that

$$\cup \{(t] | t \in I \cap S^2\} \in P.$$

We will now show that each $x \in L$ is a finite product of members of S^2 . The work is divided into two cases.

Firstly, assume that $S^2 \cap [x] \neq \emptyset$. Then

$$(4) \quad x^* = \cap \{y^* | y \in S^2 \cap [x]\}.$$

To see this, let $I \in x^*$. Then $x \notin I$, so $y \in S^2 \cap [x]$ implies that $y \notin I$ and, therefore, that $I \in y^*$. Conversely, suppose that

$$I \in \cap \{y^* | y \in S^2 \cap [x]\} \sim x^*.$$

Now

$$\begin{aligned} I &= \sum_P \{(t] | t \in S^2 \cap I\} \\ &= \cup \{(t] | t \in S^2 \cap I\}, \end{aligned}$$

and as $x \in I$, $x \in (t]$ for some $t \in S^2 \cap I$. So $t \in S^2 \cap [x]$ and, therefore, $I \in t^*$, which is a contradiction. But by (2), (4) implies that $x^* = y_1^* \cap \dots \cap y_n^*$ for some $y_i \in S^2$ and hence that $x = y_1 \cdot \dots \cdot y_n$, which completes the proof for this case.

Finally, suppose that $S^2 \cap [x] = \emptyset$. Thus, $x \not\leq t$ for all $t \in S^2$. Then

$$(5) \quad x^* = \cup \{s^* | s \in S\}.$$

Indeed, if $I \notin x^*$ then $x \in I = \cup \{(t] | t \in I \cap S^2\}$, so $x \leq t$ for some $t \in S^2$, which is a contradiction. If $I \in x^*$, then $x \notin I$, so either $I = \emptyset$ or there is an atom $(s]$, $s \in S$, such that $(s] \not\subseteq I$ and therefore, in either case, $I \in \cup \{s^* | s \in S\}$.

From (5), it follows that $x = s_1 + \dots + s_n \in S^2$.

6. Posets representable over \mathcal{L} . A characterization of those posets representable over L can be obtained immediately from (1) and (2).

THEOREM 13. *A poset P is representable over \mathcal{L} if and only if P has $0 < 1$ and there is a ring R of non-empty, proper, hereditary subsets of P satisfying:*

- (i) *If $p \not\leq q$ then there exists $A \in R$ such that $p \notin A$ and $q \in A$.*
- (ii) *If $\{A_i\}_{i \in I}$ and $\{B_j\}_{j \in J}$ are non-empty families in R and*

$$\cap \{A_i | i \in I\} \subseteq \{B_j | j \in J\},$$

then there exist finite non-empty subsets $I' \subseteq I$ and $J' \subseteq J$ such that

$$\cap \{A_i | i \in I'\} \subseteq \cup \{B_j | j \in J'\}.$$

Another characterization can be obtained by distinguishing the prime ideals in the class of all ideals of a distributive lattice.

THEOREM 14. *A poset P with $0 < 1$ is representable over \mathcal{L} if and only if P is the set of M.I. elements of a distributive algebraic lattice L in which the non-zero compact elements K form a sublattice of L .*

Proof. For the sufficiency of the condition, we show that $P \cong \mathcal{P}(K)$. For each $p \in P$, let $\psi(p) = \{q \in K | q \leq p\}$. The relation $p \mapsto \psi(p)$ establishes a function from P into $\mathcal{P}(K)$ which is order preserving in both directions. To show that ψ is onto, first note that $\psi(0) = \emptyset$ and that $\psi(1) = K$. Now let I be a prime ideal in K . Set $p = \sum_{L} I$. To show that p is M.I., suppose that $xy \leq p$ but $x \not\leq p$ and $y \not\leq p$ for some $\{x, y\} \subseteq P$. Since L is algebraic, there exists $\{s, t\} \subseteq K$ such that $s \leq x, s \not\leq p$ and $t \leq y, t \not\leq p$. But $st \leq xy \leq p = \sum_{L} I$, and since K is a sublattice of L , there exists $\{x_1, \dots, x_n\} \subseteq I$ such that $st \leq x_1 + \dots + x_n$. But I is prime, so $s \in I$ or $t \in I$. Thus, either $s \leq \sum I = p$ or $t \leq \sum I = p$, which is a contradiction. Hence, $p \in P$ and it follows that $\psi(p) = I$.

Conversely, suppose that L is a distributive lattice and that $P = \mathcal{P}(L)$. Let $\mathcal{I}(L)$ be the poset of all ideals in L together with \emptyset . $\mathcal{I}(L)$ is a complete lattice where $\prod S = \cap S$ and $\sum S$ is the ideal generated by $\cup S$. Since L is distributive, it follows that $\mathcal{I}(L)$ is also distributive. Since $I \in \mathcal{I}(L)$ can be represented by $I = \sum \{(x] | x \in I\}$, it is easily verified that $\mathcal{I}(L)$ is an algebraic lattice. It remains to show that $I \in \mathcal{P}(L)$ if and only if I is M.I. in $\mathcal{I}(L)$. Let $I \in \mathcal{P}(L)$ and let $J \cdot J_1 \subseteq I$. If $J \not\subseteq I$, then there is an element $u \in J \sim I$. But then $J_1 \subseteq I$; for, if $x \in J_1$, then $xu \in J \cap J_1 = J \cdot J_1 \subseteq I$, and so $x \in I$. On the other hand, if I is M.I. in $\mathcal{I}(L)$ and $xy \in I$, then $(x] \cdot (y] = (xy] \subseteq I$, so $(x] \subseteq I$ or $(y] \subseteq I$. Hence, $x \in I$ or $y \in I$, and $I \in \mathcal{P}(L)$.

Neither Theorem 13 nor Theorem 14 is an optimal solution, since neither really tells us much about P itself. We therefore ask for a characterization of posets representable over \mathcal{L} , which is analogous the solution for \mathcal{A} given in Theorem 5.

REFERENCES

1. R. Balbes, *Projective and injective distributive lattices*, Pacific J. Math. 21 (1967), 405–420.
2. G. Grätzer, *Lattice theory: First concepts and distributive lattices* (Freeman, San Francisco, 1971).
3. L. Nachbin, *Une propriété caractéristique des algèbres Booléennes*, Portugal. Math. 6 (1947), 115–118.
4. M. H. Stone, *Topological representations of distributive lattices and Brouwerian logics*, Časopis Pěst. Mat. 67 (1937), 1–25.

*University of Missouri,
St. Louis, Missouri*