

T. Kanzaki
Nagoya Math. J.
Vol. 49 (1973), 127-141

ON THE QUADRATIC EXTENSIONS AND THE EXTENDED WITT RING OF A COMMUTATIVE RING

TERUO KANZAKI

Let B be a ring and A a subring of B with the common identity element 1. If the residue A -module B/A is invertible as an A - A -bimodule, i.e. $B/A \otimes_A \text{Hom}_A(B/A, A) \approx \text{Hom}_A(B/A, A) \otimes_A B/A \approx A$, then B is called a quadratic extension of A . In the case where B and A are division rings, this definition coincides with in P. M. Cohn [2]. We can see easily that if B is a Galois extension of A with the Galois group G of order 2, in the sense of [3], and if $\text{Tr}_G(B) = \{\sum_{\sigma \in G} \sigma(b) : b \in B\} = A$, B is a quadratic extension of A . A generalized crossed product $\Delta(f, A, \Phi, G)$ of a ring A and a group G of order 2, in [4], is also a quadratic extension of A .

In this note, we study the case of commutative quadratic extensions, where A is a commutative ring and B is an A -algebra. Let A be a commutative ring with the identity element 1. We shall say that B is a quadratic extension of A if B is a ring extension of A with the common identity element and B is a finitely generated projective A -module of rank 2 so that B is a commutative ring. We denote by $Q(A)$ (resp. $Q_s(A)$) the set of all A -algebra isomorphism classes of quadratic (resp. separable quadratic) extensions of A . It is known that $Q_s(A)$ forms a group under a certain product, and in [1], [6] and [7], the group $Q_s(A)$ is investigated. In this note, in §1, we define a product in $Q(A)$, which coincides with the product defined in [1], [6] and [7] in the subset $Q_s(A)$. Then, $Q(A)$ forms an abelian semi-group containing the subsemi-group $Q_s(A)$ which is a group, and an element $[B]$ in $Q(A)$ is contained in $Q_s(A)$ if and only if $[B]^2 = [B][B]$ is the identity element of $Q(A)$. In §2, we give a generalization of a quadratic module and define A -isomorphisms between them. Then, we can consider a category consisting of these

Received April 24, 1972.
Revised September 4, 1972.

extended quadratic modules and A -isomorphisms. From this category we can construct a commutative ring $W^*(A)$. In §3, we shall show that $W^*(A)$ is a commutative ring with the identity element, and there exists a ring homomorphism of the Witt ring $W(A)$ to $W^*(A)$ for which the image is an ideal of $W^*(A)$. Especially, if 2 is invertible in A , then $W(A)$ and $W^*(A)$ are isomorphic. In §4, we shall give a group homomorphism of $Q_s(A)$ to the unit group $U(W^*(A))$ of $W^*(A)$.

1. Quadratic extension.

Let A be an arbitrary commutative ring with the identity element 1. A commutative extension ring B of A is called a quadratic extension of A if B is a finitely generated projective A -module of rank 2 and B has the same identity element 1. If B is a quadratic extension of A , then there exist a finitely generated projective A -module U of rank 1 and quadratic forms $q: U \rightarrow A$ and $q': U \rightarrow U^0$ such that $B = A \oplus U$ and $x^2 = q(x) + q'(x)$ for all x in U .

LEMMA 1. *Let U be a finitely generated projective A -module of rank 1, and $q': U \rightarrow U$ a quadratic form. Then there exists an A -homomorphism $f: U \rightarrow A$ such that $q'(x) = f(x)x$ for all x in U .*

Proof. For the quadratic form $q': U \rightarrow U$, there exists a bilinear form $B: U \times U \rightarrow U$ such that $q'(x) = B(x, x)$ for all x in U , (cf. (2.3) in [2]). We may consider that B is an element in $\text{Hom}_A(U \otimes_A U, U)$. Then by the following natural isomorphisms; $\text{Hom}_A(U \otimes_A U, U) \approx \text{Hom}_A(U \otimes_A U, A) \otimes_A U \approx \text{Hom}_A(U, A) \otimes_A \text{Hom}_A(U, A) \otimes_A U \approx \text{Hom}_A(U, A) \otimes_A A$, there exist f_i in $\text{Hom}_A(U, A)$ and a_i in A , $i = 1, 2, \dots, n$ such that $B(x, y) = \sum_{i=1}^n f_i(x)a_i y$ for all x and y in U . Put $f = \sum_{i=1}^n a_i f_i$ in $\text{Hom}_A(U, A)$, then we have $q'(x) = B(x, x) = f(x)x$ for all x in U .

LEMMA 2. *Let U be a finitely generated projective A -module of rank 1, and f and g elements in $\text{Hom}_A(U, A)$. If $f(x)x = g(x)x$ for all x in U , then $f = g$.*

Proof. If $f(x)x = g(x)x$ for all x in U , then we have also $f \otimes I(x)x = g \otimes I(x)x$ for all x in $U_m = U \otimes A_m$ and for every maximal ideal m of A . For the local ring A , this lemma is clear, therefore we get easily $f = g$.

¹⁾ cf. p. 490 in [5].

Thus, for a given quadratic extension B of A there exist a finitely generated projective A -module U of rank 1, an A -homomorphism $f: U \rightarrow A$ and a quadratic form $q: U \rightarrow A$ such that $B = A \oplus U$ and $x^2 = f(x)x + q(x)$ for all x in U . Conversely, if a finitely generated projective A -module U of rank 1, A -homomorphism $f: U \rightarrow A$ and a quadratic form $q: U \rightarrow A$ are given, then a quadratic extension $B = A \oplus U$ of A is constructed by $x^2 = f(x)x + q(x)$ for x in U . We denote such a quadratic extension of A by $B = (U, f, q)$.

In general, we can define as follows:

DEFINITION. Let P be a finitely generated projective and faithful A -module, $f: P \rightarrow A$ an A -homomorphism and $q: P \rightarrow A$ a quadratic form. Let $T(P) = A \oplus P \oplus P \otimes_A P \oplus \dots$ be the tensor algebra of P over A . We denote by (P, f, q) the residue ring $T(P)/(x \otimes x - f(x)x - q(x); x \in P)$ of $T(P)$ by the ideal generated from the set $\{x \otimes x - f(x)x - q(x); x \in P\}$.²⁾

PROPOSITION 1. Let (U, f, q) and (U', f', q') be quadratic extensions of A . Then (U, f, q) and (U', f', q') are A -algebra-isomorphic if and only if there exist an A -isomorphism $\sigma_1: U \rightarrow U'$ and an A -homomorphism $g: U \rightarrow A$ satisfying the following identities;

$$\begin{aligned} q' \circ \sigma_1 &= fg + q - g^2 \\ f' \circ \sigma_1 &= f - 2g, \end{aligned}$$

where fg, g^2 and $q' \circ \sigma_1$ are defined by $fg(x) = f(x)g(x)$, $g^2(x) = g(x)^2$ and $q' \circ \sigma_1(x) = q'(\sigma_1(x))$ for x in U .

Proof. Let $\sigma: (U, f, q) = A \oplus U \rightarrow (U', f', q') = A \oplus U'$ be an A -algebra-isomorphism. Then there exist an A -isomorphism $\sigma_1: U \rightarrow U'$ and an A -homomorphism $g: U \rightarrow A$ such that $\sigma(x) = g(x) + \sigma_1(x)$ for x in U . Since σ satisfies $\sigma(x^2) = \sigma(x)^2$ for x in U , we get the following identity

$$f(x)g(x) + q(x) + f(x)\sigma_1(x) = g(x)^2 + q'(\sigma_1(x)) + (f'(\sigma_1(x)) + 2g(x))\sigma_1(x)$$

for all x in U . Therefore we have

$$f(x)g(x) + q(x) = g(x)^2 + q'(\sigma_1(x)) \tag{1}$$

$$f(x)\sigma_1(x) = (f'(\sigma_1(x)) + 2g(x))\sigma_1(x) \tag{2}$$

²⁾ The composition of natural homomorphisms $A \oplus U \hookrightarrow T(U) \rightarrow T(U)/(x \otimes x - f(x)x - q(x); x \in U)$ is an A -isomorphism as A -modules. For any quadratic extension $C = A \oplus U$ satisfying $x^2 = f(x)x + q(x)$ for all $x \in U$, $C \approx T(U)/(x \otimes x - f(x)x - q(x); x \in U)$ as A -algebras.

for all x in U . From (2) we have $f(x)x = (f'(\sigma_1(x)) + 2g(x))x$ for x in U , and by Lemma 2, we get $f(x) = f'(\sigma_1(x)) + 2g(x)$ for x in U . Thus, we have the identities of this proposition. The converse is obvious.

LEMMA 3. *Let (U_i, f_i, q_i) and (U'_i, f'_i, q'_i) be A -algebra-isomorphic quadratic extensions of A , $i=1, 2$. Then $(U_1 \otimes_A U_2, f_1 \otimes f_2, f_1^2 \overline{\otimes} q_2 + q_1 \overline{\otimes} f_2^2 + 2q_1 \otimes q_2)$ and $(U'_1 \otimes_A U'_2, f'_1 \otimes f'_2, f_1'^2 \overline{\otimes} q'_2 + q'_1 \overline{\otimes} f_2'^2 + 2q'_1 \otimes q'_2)$ are also A -algebra-isomorphic, where $f_1^2 \overline{\otimes} q_2 + q_1 \overline{\otimes} f_2^2 + 2q_1 \otimes q_2 (x \otimes y) = f_1(x)^2 q_2(y) + q_1(x) f_2(y)^2 + 4q_1(x) q_2(y)$, $f_1^2 \overline{\otimes} q_2 + q_1 \overline{\otimes} f_2^2 + 2q_1 \otimes q_2 (\sum_{i=1}^n x_i \otimes y_i) = \sum_{i=1}^n (f_1(x_i)^2 q_2(y_i) + q_1(x_i) f_2(y_i)^2 + 4q_1(x_i) q_2(y_i)) + \sum_{i < j} (f_1(x_i) f_1(x_j) B_{q_2}(y_i, y_j) + B_{q_1}(x_i, x_j) f_2(y_i) f_2(y_j) + 2B_{q_1}(x_i, x_j) B_{q_2}(y_i, y_j))$, ($n > 1$), and $f_1 \otimes f_2 (\sum_{i=1}^n x_i \otimes y_i) = \sum_{i=1}^n f_1(x_i) f_2(y_i)$ for $\sum_{i=1}^n x_i \otimes y_i$ and $x \otimes y$ in $U_1 \otimes_A U_2$, (cf. (2.8) in [5]).³⁾*

Proof. By Proposition 1, there exist A -isomorphisms $\sigma_1: U_1 \rightarrow U'_1$ and $\sigma_2: U_2 \rightarrow U'_2$, and A -homomorphisms $g_1: U_1 \rightarrow A$ and $g_2: U_2 \rightarrow A$ such that $q'_1 \circ \sigma_1 = f_1 g_1 + q_1 - g_1^2$, $f'_1 \circ \sigma_1 = f_1 - 2g_1$, and $q'_2 \circ \sigma_2 = f_2 g_2 + q_2 - g_2^2$, $f'_2 \circ \sigma_2 = f_2 - 2g_2$. By the computation, we get the following:

For any element $x \otimes y$ in $U_1 \otimes_A U_2$, $(f_1'^2 \overline{\otimes} q'_2 + q'_1 \overline{\otimes} f_2'^2 + 2q'_1 \otimes q'_2) \circ (\sigma_1 \otimes \sigma_2) (x \otimes y) = (f_1(x) - 2g_1(x))^2 (f_2(y) g_2(y) + q_2(y) - g_2(y)^2) + (f_1(x) g_1(x) + q_1(x) - g_1(x)^2) (f_2(y) - 2g_2(y))^2 + 4(f_1(x) g_1(x) + q_1(x) - g_1(x)^2) (f_2(y) g_2(y) + q_2(y) - g_2(y)^2) = f_1(x) f_2(y) (f_1(x) g_2(y) + g_1(x) f_2(y) - 2g_1(x) g_2(y)) + (f_1(x)^2 q_2(y) + q_1(x) f_2(y)^2 + 4q_1(x) q_2(y)) - (f_1(x) g_2(y) + g_1(x) f_2(y) - 2g_1(x) g_2(y))^2 = [(f_1 \otimes f_2) (f_1 \otimes g_2 + g_1 \otimes f_2 - 2g_1 \otimes g_2) + (f_1^2 \overline{\otimes} q_2 + q_1 \overline{\otimes} f_2^2 + 2q_1 \otimes q_2) - (f_1 \otimes g_2 + g_1 \otimes f_2 - 2g_1 \otimes g_2)^2] (x \otimes y)$. Using the identities

$B_{q_1}(\sigma_1(x_i), \sigma_1(x_j)) = f_1(x_i) g_1(x_j) + f_1(x_j) g_1(x_i) + B_{q_1}(x_i, x_j) - 2g_1(x_i) g_1(x_j)$ and $B_{q_2}(\sigma_2(y_i), \sigma_2(y_j)) = f_2(y_i) g_2(y_j) + f_2(y_j) g_2(y_i) + B_{q_2}(y_i, y_j) - 2g_2(y_i) g_2(y_j)$ for $x_i \otimes y_i$ and $x_j \otimes y_j$ in $U_1 \otimes_A U_2$, we get as follows; $f'_1(\sigma_1(x_i)) f'_1(\sigma_1(x_j)) B_{q_2}(\sigma_2(y_i), \sigma_2(y_j)) + B_{q_1}(\sigma_1(x_i), \sigma_1(x_j)) f'_2(\sigma_2(y_i)) f'_2(\sigma_2(y_j)) + 2B_{q_1}(\sigma_1(x_i), \sigma_1(x_j)) B_{q_2}(\sigma_2(y_i), \sigma_2(y_j)) = (f_1(x_i) - 2g_1(x_i))(f_1(x_j) - 2g_1(x_j))(f_2(y_i) g_2(y_j) + f_2(y_j) g_2(y_i) + B_{q_2}(y_i, y_j) - 2g_2(y_i) g_2(y_j)) + (f_1(x_i) g_1(x_j) + f_1(x_j) g_1(x_i) + B_{q_1}(x_i, x_j) - 2g_1(x_i) g_1(x_j))(f_2(y_i) - 2g_2(y_i))(f_2(y_j) - 2g_2(y_j)) + 2(f_1(x_i) g_1(x_j) + f_1(x_j) g_1(x_i) + B_{q_1}(x_i, x_j) - 2g_1(x_i) g_1(x_j))(f_2(y_i) g_2(y_j) + f_2(y_j) g_2(y_i) + B_{q_2}(y_i, y_j) - 2g_2(y_i) g_2(y_j)) = f_1(x_i) f_2(y_i) (f_1(x_j) g_2(y_j) + g_1(x_j) f_2(y_j) - 2g_1(x_j) g_2(y_j)) + f_1(x_i) f_2(y_j) (f_1(x_j) g_2(y_i) + g_1(x_j) f_2(y_i) - 2g_1(x_j) g_2(y_i)) + f_1(x_i) f_1(x_j) B_{q_2}(y_i, y_j) + B_{q_1}(x_i, x_j)$

³⁾ $f^2 \overline{\otimes} q'$ and $f^2 \otimes q'$ are defined by $f^2 \overline{\otimes} q'(\sum x_i \otimes y_i) = \sum_i f(x_i)^2 q'(y_i) + \sum_{i < j} f(x_i) f(x_j) B_{q'}(y_i, y_j)$ and $f^2 \otimes q'(\sum x_i \otimes y_i) = \sum_i 2f(x_i)^2 q'(y_i) + \sum_{i < j} B_{f^2}(x_i, x_j) B_{q'}(y_i, y_j)$ for $\sum x_i \otimes y_i$ in $M \otimes_A M'$.

$f_2(y_i)f_2(y_j) + 2B_{q_1}(x_i, x_j)B_{q_2}(y_i, y_j) - 2(f_1(x_i)g_2(y_i) + g_1(x_i)f_2(y_i) - 2g_1(x_i)g_2(y_i))(g_1(x_j)f_2(y_j) + f_1(x_j)g_2(y_j) - 2g_1(x_j)g_2(y_j))$. Accordingly, we get $(f_1'^2 \bar{\otimes} q_2' + q_1' \bar{\otimes} f_2'^2 + 2q_1' \otimes q_2') \circ (\sigma_1 \otimes \sigma_2)(\sum_{i=1}^n x_i \otimes y_i) = [(f_1 \otimes f_2)(f_1 \otimes g_2 + g_1 \otimes f_2 - 2g_1 \otimes g_2) + (f_1^2 \bar{\otimes} q_2 + q_1 \bar{\otimes} f_2^2 + 2q_1 \otimes q_2) + (f_1 \otimes g_2 - 2g_1 \otimes g_2)^2](\sum_{i=1}^n x_i \otimes y_i)$ for all $\sum_{i=1}^n x_i \otimes y_i$ in $U_1 \otimes_A U_2$. Put $G = f_1 \otimes g_2 + g_1 \otimes f_2 - 2g_1 \otimes g_2$, then G is an A -homomorphism of $U_1 \otimes_A U_2$ to A , $\sigma_1 \otimes \sigma_2$ is an A -isomorphism of $U_1 \otimes_A U_2$ to $U_1' \otimes_A U_2'$, and these satisfy $(f_1'^2 \bar{\otimes} q_2' + q_1' \bar{\otimes} f_2'^2 + 2q_1' \otimes q_2') \circ (\sigma_1 \otimes \sigma_2) = (f_1 \otimes f_2)G + (f_1^2 \bar{\otimes} q_2 + q_1 \bar{\otimes} f_2^2 + 2q_1 \otimes q_2) + G^2$, and $(f_1' \otimes f_2') \circ (\sigma_1 \otimes \sigma_2) = f_1 \otimes f_2 - 2G$.

By Proposition 1, we have $(U_1 \otimes_A U_2, f_1 \otimes f_2, f_1^2 \bar{\otimes} q_2 + q_1 \bar{\otimes} f_2^2 + 2q_1 \otimes q_2)$ and $(U_1' \otimes_A U_2', f_1' \otimes f_2', f_1'^2 \bar{\otimes} q_2' + q_1' \bar{\otimes} f_2'^2 + 2q_1' \otimes q_2')$ are isomorphic as A -algebras.

DEFINITION. We denote by $Q(A)$ the set of all A -algebra-isomorphism classes $[U, f, q]$ of quadratic extensions (U, f, q) of A .

PROPOSITION 2. $Q(A)$ forms an abelian semi-group with unit element $[A, 1, 0]$ by the product $[U, f, q] \cdot [U', f', q'] = [U \otimes_A U', f \otimes f', f^2 \bar{\otimes} q' + q' \bar{\otimes} f'^2 + 2q \otimes q']$, where (A, a, b) denotes a quadratic extension $A \oplus Av$ such that $v^2 = av + b$, a and b in A , i.e. $f(v) = a$, $q(v) = b$.

Proof. By Lemma 3, the product in $Q(A)$ is well defined. The associative law is easily seen as follows; $([U, f, q][U', f', q'])[U'', f'', q''] = [U \otimes_A U' \otimes_A U'', f \otimes f' \otimes f'', f^2 \bar{\otimes} f'^2 \bar{\otimes} q'' + f^2 \bar{\otimes} q' \bar{\otimes} f''^2 + q \bar{\otimes} f'^2 \bar{\otimes} f''^2 + 2(q \otimes q' \bar{\otimes} f''^2 + q \bar{\otimes} f'^2 \otimes q'' + f^2 \bar{\otimes} q' \otimes q'')] + 4q \otimes q' \otimes q'' = [U, f, q]([U', f', q'] [U'', f'', q'']).⁴⁾$

DEFINITION. Let P be a finitely generated projective and faithful A -module, $f: P \rightarrow A$ an A -homomorphism and $q: P \rightarrow A$ a quadratic form. For the A -algebra $(P, f, q) = T(P)/(x \otimes x - f(x)x - q(x); x \in P)$, we consider a symmetric bilinear form $D_{f,q}: P \times P \rightarrow A$ defined by $D_{f,q}(x, y) = f(x)f(y) + 2B_q(x, y)$ for x, y in P , where $B_q(x, y) = q(x + y) - q(x) - q(y)$ for x, y in P . Then we shall call the bilinear A -module $(P, D_{f,q})$ the *discriminant* of (P, f, q) .

Remark 1. If 2 is invertible in A , then we have that (P, f, q) is a separable algebra over A if and only if $(P, D_{f,q})$ is a non-degenerate bilinear A -module, i.e. $P \rightarrow \text{Hom}_A(P, A); x \rightsquigarrow D_{f,q}(x, -)$ is an isomorphism.

⁴⁾ $(q \bar{\otimes} f^2) \otimes q' = q \otimes (f^2 \bar{\otimes} q')$, $(f^2 \bar{\otimes} q') \otimes q'' = f^2 \bar{\otimes} (q' \otimes q'')$.

Proof. $d = f^2 + 4q$ is a quadratic form of P to A , and satisfies $d(x) = f(x)^2 + 2B_q(x, x) = D_{f,q}(x, x)$. In the tensor algebra $T(P)$, we put $P' = \{x - (1/2)f(x) \in A \oplus P \subset T(P); x \in P\}$, then the map $P \rightarrow P'; x \rightsquigarrow x - (1/2)f(x)$ is an A -isomorphism. We denote by h the inverse isomorphism of it. For the ideal of $T(P)$ generated by the set $\{x \otimes x - f(x)x - q(x); x \in P\} = \{x \otimes x - d(h(x)/2); x \in P'\}$, we have $(P, f, q) = T(P)/(x \otimes x - f(x)x - q(x); x \in P) = T(P')/(x \otimes x - d(h(x)/2)); x \in P' = (P', 0, d \circ (1/2)h)$, since $T(P) = T(P')$. But, $(P', 0, d \circ (1/2)h)$ is a Clifford algebra $\text{Cl}(P', d \circ (1/2)h)$ of a quadratic module $(P', d \circ (1/2)h)$. It is known that $\text{Cl}(P', d \circ (1/2)h)$ is a separable algebra over A if and only if $(P', d \circ (1/2)h)$ is non-degenerated. Since (P, d) and $(P', d \circ (1/2)h)$ are isometric, we get this remark.

THEOREM 1. *Let U be a finitely generated projective A -module of rank 1, $f: U \rightarrow A$ an A -homomorphism, $q: U \rightarrow A$ a quadratic form, and (U, f, q) the quadratic extension of A . Then the following conditions are equivalent:*

- 1) (U, f, q) is a separable algebra over A .
- 2) $(U, D_{f,q})$ is a non-degenerate bilinear A -module.
- 3) $[U, f, q]^2 = [A, 1, 0]$.

Proof. 1) \rightleftarrows 2): To prove the equivalence of the conditions 1) and 2), we may assume that A is a local ring. Let A be the local ring. Then $U = Au$ and $(U, f, q) \approx A[X]/(X^2 - aX - b)$, where $a = f(u)$, $b = q(u)$. Hence, (U, f, q) is separable over A if and only if $a^2 + 4b = f^2 + 4q(u) = D_{f,q}(u, u)$ is invertible in A . On the other hand, $(U, D_{f,q})$ is non-degenerated if and only if $D_{f,q}(u, u)$ is invertible in A . Therefore, we obtain the equivalence.

2) \rightarrow 3): Assume that $(U, D_{f,q})$ is non-degenerate. Then the A -isomorphism $U \rightarrow \text{Hom}_A(U, A); x \rightsquigarrow D_{f,q}(x, -)$ induces an A -isomorphism $D_{f,q}: U \otimes_A U \rightarrow A; x \otimes y \rightsquigarrow D_{f,q}(x, y)$. Put $\sigma_1 = D_{f,q}$ and $g = -B_q$. Then we have $I \circ \sigma_1 = D_{f,q} = f \otimes f + 2B_q = f \otimes f - 2g$. Furthermore, we can prove the following identity:

$$(f \otimes f)g + (f^2 \bar{\otimes} q + q \bar{\otimes} f^2 + 2q \otimes q) - g^2 = 0.$$

Because, by the localizations of A and U by every maximal ideal \mathfrak{m} of A , we can check that quadratic forms $f^2 \bar{\otimes} q + q \bar{\otimes} f^2 - B_q \cdot f \otimes f: U \otimes_A U \rightarrow A$, and $2q \otimes q - B_q^2: U \otimes_A U \rightarrow A$ are equal to 0. Thus, by Proposition 1 we get $[U, f, q]^2 = [U \otimes_A U, f \otimes f, f^2 \bar{\otimes} q + q \bar{\otimes} f^2 + 2q \otimes q] =$

$[A, 1, 0]$.

3) \rightarrow 2): Let $[U, f, q]^2 = [A, 1, 0]$. To prove the condition 2) it is sufficient to show that for any maximal ideal \mathfrak{m} of A , $D_{f,q}(u, u)$ is invertible in $A_{\mathfrak{m}}$, where $U_{\mathfrak{m}} = A_{\mathfrak{m}}u$. Now, we assume A is a local ring with maximal ideal \mathfrak{m} and $U = Au$. We shall show $D_{f,q}(u, u) = f(u)^2 + 2B_q(u, u) = f(u)^2 + 4q(u) \notin \mathfrak{m}$. From $[U, f, q]^2 = [A, 1, 0]$, there exist an A -homomorphism $g: U \otimes_A U \rightarrow A$ and an A -isomorphism $\sigma_1: U \otimes_A U \rightarrow A$ such that $\sigma_1(x \otimes y) = f(x)f(y) - 2g(x \otimes y)$ and $0 = f(x)f(y)g(x \otimes y) + f(x)^2q(y) + q(x)f(y)^2 + 4q(x)q(y) + g(x \otimes y)^2$ for all $x \otimes y \in U \otimes_A U$. Especially, taking $x = y = u$, we get

$$\sigma_1(u \otimes u) = f(u)^2 - 2g(u \otimes u) \tag{3},$$

and

$$0 = f(u)^2g(u \otimes u) + 2f(u)^2q(u) + 4q(u)^2 - g(u \otimes u) \tag{4}.$$

Eliminating $f(u)^2$ from (3) and (4), we get $(\sigma_1(u \otimes u) + 2g(u \otimes u))g(u \otimes u) + 2(\sigma_1(u \otimes u) + 2g(u \otimes u))q(u) + 4q(u)^2 - g(u \otimes u)^2 = 0$, and so

$$(\sigma_1(u \otimes u) + g(u \otimes u) + 2q(u))(g(u \otimes u) + 2q(u)) = 0.$$

If $g(u \otimes u) + 2q(u)$ is contained in \mathfrak{m} , then from $\sigma_1(u \otimes u) \notin \mathfrak{m}$, $\sigma_1(u \otimes u) + g(u \otimes u) + 2q(u)$ is invertible in A . Therefore, we have $g(u \otimes u) + 2q(u) = 0$, and $D_{f,q}(u, u) = f(u)^2 + 4q(u) = f(u)^2 - 2g(u \otimes u) = \sigma_1(u \otimes u)$ is invertible in A . If $g(u \otimes u) + 2q(u) \notin \mathfrak{m}$, then $\sigma_1(u \otimes u) + g(u \otimes u) + 2q(u) = 0$. From (3) and $2\sigma_1(u \otimes u) + 2g(u \otimes u) + 4q(u) = 0$, we get $\sigma_1(u \otimes u) + f(u)^2 + 4q(u) = 0$, accordingly, $D_{f,q}(u, u) = f(u)^2 + 4q(u) = -\sigma_1(u \otimes u)$ is invertible in A .

COROLLARY 1. *The set $Q_s(A)$ of A -algebra-isomorphism classes of the separable quadratic extensions of A forms an abelian group with exponent 2.*

PROPOSITION 3. *Let (U, f, q) be a quadratic extension of A . The map $\tau_f: (U, f, q) \rightarrow (U, f, q); a + x \rightsquigarrow a + f(x) - x$ is an A -algebra-isomorphism such that $\tau_f^2 = I$. If (U, f, q) and (U', f', q') are quadratic extensions of A and $\sigma: (U, f, q) \rightarrow (U', f', q')$ is an A -algebra-isomorphism, then we have the following commutative diagram;*

$$\begin{array}{ccc}
 (U, f, q) & \xrightarrow{\sigma} & (U', f', q') \\
 \downarrow \tau_f & \circlearrowleft & \downarrow \tau'_{f'} \\
 (U, f, q) & \xrightarrow{\sigma} & (U', f', q') .
 \end{array}$$

Proof. From Proposition 1, there exist g in $\text{Hom}_A(U, A)$ and A -isomorphism $\sigma_1: U \rightarrow U'$ such that $\sigma(x) = g(x) + \sigma_1(x)$ and $f'(\sigma_1(x)) = f(x) - 2g(x)$ for all x in U . Therefore, $\tau'_{f'}(\sigma(x)) = g(x) + \tau'_{f'}(\sigma_1(x)) = g(x) + f'(\sigma_1(x)) - \sigma_1(x) = g(x) + f(x) - 2g(x) - \sigma_1(x) = f(x) - (g(x) + \sigma_1(x)) = f(x) - \sigma(x) = \sigma(f(x) - x) = \sigma(\tau_f(x))$, for all x in U .

Remark 2.

1) In Proposition 3, if we take $\sigma = I$, then $\tau_f = \tau'_{f'}$.

2) If (U, f, q) is a separable algebra over A , then τ_f is the unique A -algebra-automorphism of (U, f, q) which is not the identity.

Let $B = (U, f, q)$ and $B' = (U', f', q')$ be separable quadratic extensions of A . Then $G = \{\tau_f, I\}$ and $G' = \{\tau'_{f'}, I\}$ are the groups of automorphisms of B over A and B' over A , respectively. In [1], [3] and [4], the product $B * B'$ of quadratic extensions B and B' was defined as the fixed subalgebra $(B \otimes_A B')^{\tau_f \otimes \tau'_{f'}} = \{x \in B \otimes_A B'; \tau_f \otimes \tau'_{f'}(x) = x\}$ of $B \otimes_A B'$ by $\tau_f \otimes \tau'_{f'}$. But this product coincides with our one.

PROPOSITION 4. *Let (U, f, q) and (U', f', q') be separable quadratic extensions of A . Then we have $[(U, f, q) \otimes_A (U', f', q')]^{\tau_f \otimes \tau'_{f'}} = [U, f, q] \cdot [U', f', q']$ in $Q_s(A)$.*

Proof. For $B = (U, f, q)$ and $B' = (U', f', q')$, $B \otimes_A B'$ is expressed as a direct sum $B \otimes_A B' = A \oplus U \oplus U' \oplus U \otimes_A U'$. Put $V = \{\sum_i f(x_i)y_i + f'(y_i)x_i - 2x_i \otimes y_i \in U \oplus U' \oplus U \otimes_A U'; \text{ for all } \sum_i x_i \otimes y_i \text{ in } U \otimes_A U'\}$. Then V is an A -submodule of $B \otimes_A B'$, which is A -isomorphic to $U \otimes_A U'$ by the isomorphism $\theta: U \otimes_A U' \rightarrow V; x \otimes y \rightsquigarrow f(x)y + f'(y)x - 2x \otimes y$. It is easily seen that the A -submodule $C = A \oplus V$ of $B \otimes_A B'$ generated by V and A is contained in $B \otimes_A B'^{\tau_f \otimes \tau'_{f'}}$. To show $C = B \otimes_A B'^{\tau_f \otimes \tau'_{f'}}$, we shall prove first that the map $\theta': (U \otimes_A U', f \otimes f', f^2 \overline{\otimes} q' + q \overline{\otimes} f'^2 + 2q \otimes q') = A \oplus U \otimes_A U' \rightarrow C = A \oplus V; a + x \otimes y \rightsquigarrow a + \theta(x \otimes y)$ is an A -algebra-isomorphism. We can easily compute that for any $x \otimes y$ in $U \otimes_A U'$, $\theta'(x \otimes y)^2 = (f(x)y + f'(y)x - 2x \otimes y)^2 = f(x)^2 y^2 + f'(y)^2 x^2 + 4x^2 \otimes y^2 + 2f(x)f'(y)x \otimes y - 4f(x)x \otimes y^2 - 4f'(y)x^2 \otimes y = f(x)^2(f'(y)y + q'(y)) + f'(y)^2(f(x)x + q(x)) + 4(f(x) + q(x)) \otimes (f'(y)y + q'(y)) + 2f(x)f'(y)$

$x \otimes y - 4f(x)x \otimes (f'(y)y + q'(y)) - 4f'(y)(f(x)x + q(x)) \otimes y = f(x)f'(y)(f(x)y + f'(y)x - 2x \otimes y) + f(x)^2q'(y) + f'(y)^2q(x) + 4q(x)q'(y) = f \otimes f'(x \otimes y)\theta'(x \otimes y) + (f^2 \otimes q' + q \otimes f'^2 + 2q \otimes q')(x \otimes y) = \theta'[(f \otimes f'(x \otimes y)x \otimes y + (f^2 \otimes q' + q \otimes f'^2 + q \otimes q')(x \otimes y)] = \theta'((x \otimes y)^2)$, and for $x_i \otimes y_i, x_j \otimes y_j$ in $U \otimes_A U'$, $2\theta'(x_i \otimes y_i) \cdot \theta'(x_j \otimes y_j) = 2(f(x_i)y_i + f'(y_i)x_i - 2x_i \otimes y_i)(f(x_j)y_j + f'(y_j)x_j - 2x_j \otimes y_j) = f(x_i)f'(y_i)(f(x_j)y_j + f'(y_j)x_j - 2x_j \otimes y_j) + f(x_j)f'(y_j)(f(x_i)y_i + f'(y_i)x_i - 2x_i \otimes y_i) + f(x_i)f(x_j)B_{q'}(y_i, y_j) + f'(y_i)f'(y_j)B_q(x_i, x_j) + 2B_q(x_i, x_j)B_{q'}(y_i, y_j) = \theta'(f(x_i)f'(y_i)x_j \otimes y_j + f(x_j)f'(y_j)x_i \otimes y_i) + f(x_i)f(x_j)B_{q'}(y_i, y_j) + f'(y_i)f'(y_j)B_q(x_i, x_j) + 2B_q(x_i, x_j)B_{q'}(y_i, y_j)$.

Therefore, we have $\theta'(\sum_i x_i \otimes y_i)^2 = \theta'((\sum_i x_i \otimes y_i)^2)$ for any $\sum_i x_i \otimes y_i$ in $U \otimes_A U'$. Accordingly, θ' is an A -algebra isomorphism. Thus, C is also a separable algebra over A . Since $B \otimes_A B'$ is a finitely generated projective A -module, $B \otimes_A B'$ is also finitely generated projective over C . Therefore, C is a direct summand of $B \otimes_A B'$, and hence also a direct summand of $B \otimes_A B'^{r_f \otimes r_{f'}}$ as C -module. But, $\text{rank}(C : A) = \text{rank}(B \otimes_A B'^{r_f \otimes r_{f'}} : A) = 2$, hence we have $B \otimes_A B'^{r_f \otimes r_{f'}} = C = A \oplus V \approx (U \otimes_A U', f \otimes f', f^2 \otimes q' + q \otimes f'^2 + 2q \otimes q')$ as A -algebra.

2. Extended quadratic module.

In this section, we give a generalization of quadratic module. Let A be an arbitrary commutative ring with unit element. Let M be an A -module, $f : M \rightarrow A$ an A -homomorphism, and $q : M \rightarrow A$ a quadratic form. Then, we call the triple $\langle M, f, q \rangle$ an *extended quadratic module*

DEFINITION. Let $\langle M, f, q \rangle$ and $\langle M', f', q' \rangle$ be extended quadratic modules. If there exist an A -isomorphism $\sigma : M \rightarrow M'$ and A -homomorphism $g : M \rightarrow A$ satisfying $q' \circ \sigma = q + 2fg - 2g^2$ and $f' \circ \sigma = f - 2g$, then we call that $\langle M, f, q \rangle$ and $\langle M', f', q' \rangle$ are A -isomorphic, and denote by $(\sigma, g) : \langle M, f, q \rangle \rightarrow \langle M', f', q' \rangle$ the A -isomorphism of extended quadratic modules, or simply $\langle M, f, q \rangle \approx \langle M', f', q' \rangle$.

Then we have easily

- 1) $(I, 0)$ is identity,
- 2) $(\sigma', g')(\sigma, g) = (\sigma' \circ \sigma, g + g' \circ \sigma)$ and
- 3) $(\sigma, g)^{-1} = (\sigma^{-1}, -g \circ \sigma^{-1})$.

Thus, we can consider a category $\text{Qua}^*(A)$ in which objects are extended quadratic modules and morphisms are A -isomorphisms of extended quadratic modules. Then, $\text{Qua}^*(A)$ includes the category $\text{Qua}(A)$ of the

ordinally quadratic modules as a sub-category. Because, $(\sigma, g): \langle M, 0, q \rangle \rightarrow \langle M', 0, q' \rangle$ is an A -isomorphism in $\text{Qua}^*(A)$ if and only if $\sigma: (M, q) \rightarrow (M', q')$ is an A -isomorphism in $\text{Qua}(A)$, therefore we may regard as $\langle M, 0, q \rangle = (M, q)$ and $(\sigma, 0) = \sigma$ in $\text{Qua}(A)$.

DEFINITION. Let $\langle M, f, q \rangle$ be an extended quadratic module, and let $B_{f,q}: M \times M \rightarrow A$ be a symmetric bilinear form defined by $B_{f,q}(x, y) = f(x)f(y) + B_q(x, y)$ for x and y in M . Then, we call the bilinear module $(M, B_{f,q})$ the associated bilinear module with $\langle M, f, q \rangle$. If $(M, B_{f,q})$ is a non-degenerate bilinear module, then $\langle M, f, q \rangle$ is called a *non-degenerate extended quadratic module*.

LEMMA 4. If $(\sigma, g): \langle M, f, q \rangle \rightarrow \langle M', f', q' \rangle$ is an A -isomorphism in $\text{Qua}^*(A)$, then we have $B_{f',q'}(\sigma(x), \sigma(y)) = B_{f,q}(x, y)$ for all x and y in M , that is, $\sigma: (M, B_{f,q}) \rightarrow (M', B_{f',q'})$ is an A -isomorphism of bilinear modules.

Proof. Since the A -isomorphism $\sigma: M \rightarrow M'$ and the A -homomorphism $g: M \rightarrow A$ satisfy $f' \circ \sigma = f - 2g$ and $q' \circ \sigma = q + 2fg - 2g^2$, we have $B_{f',q'}(\sigma(x), \sigma(y)) = f'(\sigma(x))f'(\sigma(y)) + B_{q'}(\sigma(x), \sigma(y)) = (f(x) - 2g(x))(f(y) - 2g(y)) + B_q(x, y) + 2(f(x)g(y) + f(y)g(x)) - 4g(x)g(y) = f(x)f(y) + B_q(x, y) = B_{f,q}(x, y)$.

COROLLARY 2. If $\langle M, f, q \rangle \approx \langle M', f', q' \rangle$ and $\langle M, f, q \rangle$ is non-degenerate, then $\langle M', f', q' \rangle$ is also non-degenerate.

DEFINITION. Let $\langle M_1, f_1, q_1 \rangle$ and $\langle M_2, f_2, q_2 \rangle$ be extended quadratic modules. We define the orthogonal sum \perp and the tensor product \otimes of extended quadratic modules as follows:

$$\langle M_1, f_1, q_1 \rangle \perp \langle M_2, f_2, q_2 \rangle = \langle M_1 \oplus M_2, f_1 \perp f_2, q_1 \perp q_2 - f_1 \times f_2 \rangle \quad (5),$$

$$\begin{aligned} \langle M_1, f_1, q_1 \rangle \otimes \langle M_2, f_2, q_2 \rangle \\ = \langle M_1 \otimes M_2, f_1 \otimes f_2, f_1^2 \bar{\otimes} q_2 + q_1 \bar{\otimes} f_2^2 + q_1 \otimes q_2 \rangle \quad (6), \end{aligned}$$

where $f_1 \perp f_2$ is defined by the A -homomorphism $M_1 \oplus M_2 \rightarrow A$; $x_1 \oplus x_2 \rightsquigarrow f_1(x_1) + f_2(x_2)$, and $f_1 \times f_2$ the quadratic form $M_1 \oplus M_2 \rightarrow A$; $x_1 \oplus x_2 \rightsquigarrow f_1(x_1) \cdot f_2(x_2)$.

LEMMA 5. Let $\langle M_i, f_i, q_i \rangle$ and $\langle M'_i, f'_i, q'_i \rangle$ be extended quadratic modules, and $(\sigma_i, g_i): \langle M_i, f_i, q_i \rangle \rightarrow \langle M'_i, f'_i, q'_i \rangle$ an A -isomorphism in $\text{Qua}^*(A)$ for $i = 1, 2$. Then we have the following A -isomorphisms in $\text{Qua}^*(A)$;

$$(\sigma_1 \oplus \sigma_2, g_1 \perp g_2) : \langle M_1, f_1, q_1 \rangle \perp \langle M_2, f_2, q_2 \rangle \rightarrow \langle M'_1, f'_1, q'_1 \rangle \perp \langle M'_2, f'_2, q'_2 \rangle \quad (7),$$

$$\begin{aligned} (\sigma_1 \otimes \sigma_2, f_1 \otimes g_2 + g_1 \otimes f_2 - 2g_1 \otimes g_2) : \langle M_1, f_1, q_1 \rangle \otimes \langle M_2, f_2, q_2 \rangle \\ \rightarrow \langle M'_1, f'_1, q'_1 \rangle \otimes \langle M'_2, f'_2, q'_2 \rangle \end{aligned} \quad (8).$$

Proof. The proof of (7). We shall show that $(\sigma_1 \oplus \sigma_2, g_1 \perp g_2) : \langle M_1 \oplus M_2, f_1 \perp f_2, q_1 \perp q_2 - f_1 \times f_2 \rangle \rightarrow \langle M'_1 \oplus M'_2, f'_1 \perp f'_2, q'_1 \perp q'_2 - f'_1 \times f'_2 \rangle$ is an A -isomorphism in $\text{Qua}^*(A)$. For $x_1 \oplus x_2$ in $M_1 \oplus M_2$, we have

$$\begin{aligned} (q'_1 \perp q'_2 - f'_1 \times f'_2) \circ (\sigma_1 \oplus \sigma_2)(x_1 \oplus x_2) &= q'_1(\sigma_1(x_1)) + q'_2(\sigma_2(x_2)) - f'_1(\sigma_1(x_1))f'_2(\sigma_2(x_2)) \\ &= q_1(x_1) + 2f_1(x_1)g_1(x_1) - 2g_1(x_1)^2 + q_2(x_2) + 2f_2(x_2)g_2(x_2) - 2g_2(x_2)^2 - (f_1(x_1) - \\ & 2g_1(x_1))(f_2(x_2) - 2g_2(x_2)) = (q_1 \perp q_2 - f_1 \times f_2)(x_1 \oplus x_2) + 2(f_1 \perp f_2)(g_1 \perp g_2) \\ & (x_1 \oplus x_2) - 2(g_1 \perp g_2)^2(x_1 \oplus x_2), \text{ and} \end{aligned}$$

$$\begin{aligned} (f'_1 \perp f'_2) \circ (\sigma_1 \oplus \sigma_2) &= f'_1 \circ \sigma_1 \perp f'_2 \circ \sigma_2 = (f_1 - 2g_1) \perp (f_2 - 2g_2) \\ &= (f_1 \perp f_2) - 2(g_1 \perp g_2). \end{aligned}$$

The proof of (8) is obtained by similar computations the proof of Lemma 3. We omit this proof.

DEFINITION. We denote by $B_{f,q} \perp B_{f',q'}$ the associated bilinear form with $\langle M, f, q \rangle \perp \langle M', f', q' \rangle$, and by $B_{f,q} \otimes B_{f',q'}$ the associated bilinear form with $\langle M, f, q \rangle \otimes \langle M', f', q' \rangle$, that is, $B_{f,q} \perp B_{f',q'} = B_{f \perp f', (q \perp q') - (f \times f')}$ and $B_{f,q} \otimes B_{f',q'} = B_{f \otimes f', f^2 \otimes q' + q \otimes f'^2 + q \otimes q'}$.

PROPOSITION 5. *The orthogonal sum and the tensor product of extended quadratic modules $\langle M, f, q \rangle$ and $\langle M', f', q' \rangle$ induce the following identities between the associated bilinear modules with them;*

$$(M \oplus M', B_{f,q} \perp B_{f',q'}) = (M, B_{f,q}) \perp (M', B_{f',q'}) \quad (9),$$

i.e. $B_{f,q} \perp B_{f',q'}(x \oplus x', y \oplus y') = B_{f,q}(x, y) + B_{f',q'}(x', y')$ for $x \oplus x'$ and $y \oplus y'$ in $M \oplus M'$, and

$$(M \otimes M', B_{f,q} \otimes B_{f',q'}) = (M, B_{f,q}) \otimes (M', B_{f',q'}) \quad (10),$$

i.e. $B_{f,q} \otimes B_{f',q'}(\sum_i x_i \otimes x'_i, \sum_j y_j \otimes y'_j) = \sum_{i,j} B_{f,q}(x_i, y_j) \cdot B_{f',q'}(x'_i, y'_j)$ for $\sum_i x_i \otimes x'_i$ and $\sum_j y_j \otimes y'_j$ in $M \otimes M'$.

Proof. The proof of (9): $B_{f,q} \perp B_{f',q'}(x \oplus x', y \oplus y') = (f \perp f')(x \otimes x') (f \perp f')(y \otimes y') + B_{(q \perp q') - (f \times f')}(x \oplus x', y \oplus y') = (f(x) + f'(x'))(f(y) + f'(y')) + B_{q \perp q'}(x \oplus x', y \oplus y') - B_{f \times f'}(x \oplus x', y \oplus y') = f(x)f(y) + f'(x')f'(y') + f'(x')f(y) + f(x)f'(y') + B_q(x, y) + B_{q'}(x', y') - (f(x)f'(y') + f(y)f'(x')) = B_{f,q}(x, y) + B_{f',q'}(x', y')$, for any $x \oplus x'$ and $y \oplus y'$ in $M \oplus M'$.

The proof of (10): $B_{f,q} \otimes B_{f',q'}(\sum_i x_i \otimes x'_i, \sum_j y_j \otimes y'_j) = B_{f \otimes f', f^2 \otimes q' + q \otimes f'^2 + q \otimes q'}(\sum_i x_i \otimes x'_i, \sum_j y_j \otimes y'_j) = f \otimes f'(\sum_i x_i \otimes x'_i) f \otimes f'(\sum_j y_j \otimes y'_j) + B_{f^2 \otimes q' + q \otimes f'^2 + q \otimes q'}(\sum_i x_i \otimes x'_i, \sum_j y_j \otimes y'_j) + B_{q \otimes q'}(\sum_i x_i \otimes x'_i, \sum_j y_j \otimes y'_j) + B_{q \otimes q'}(\sum_i x_i \otimes x'_i, \sum_j y_j \otimes y'_j) = \sum_{i,j} (f(x_i) f(y_j) f'(x'_i) f'(y'_j) + f(x_i) f(y_j) B_{q'}(x'_i, y'_j) + B_q(x_i, y_j) f'(x'_i) f'(y'_j) + B_q(x_i, y_j) B_{q'}(x'_i, y'_j)) = \sum_{i,j} B_{f,q}(x_i, y_j) B_{f',q'}(x'_i, y'_j)$, for $\sum x_i \otimes x'_i$ and $\sum y_j \otimes y'_j$ in $M \otimes M'$.

COROLLARY 3. *If $\langle M, f, q \rangle$ and $\langle M', f', q' \rangle$ are non-degenerate extended quadratic modules, then $\langle M, f, q \rangle \perp \langle M', f', q' \rangle$ is also non-degenerate. Furthermore, if M and M' are finitely generated projective A -modules, then $\langle M, f, q \rangle \otimes \langle M', f', q' \rangle$ is non-degenerate.*

Remark 3. If 2 is invertible in the ring A , then the category $\text{Qua}^*(A)$ is equivalent to the category $\text{Qua}(A)$, i.e. for any object $\langle M, f, q \rangle$ in $\text{Qua}^*(A)$, $\langle M, f, q \rangle \approx \langle M, 0, q + (1/2)f^2 \rangle$.

Remark 4. Let $\langle M_1, f_1, q_1 \rangle$, $\langle M_2, f_2, q_2 \rangle$ and $\langle M_3, f_3, q_3 \rangle$ be extended quadratic modules. Then we get the following natural isomorphisms in $\text{Qua}^*(A)$;

- 1) $\langle M_1, f_1, q_1 \rangle \perp \langle M_2, f_2, q_2 \rangle \approx \langle M_2, f_2, q_2 \rangle \perp \langle M_1, f_1, q_1 \rangle$,
- 2) $\langle M_1, f_1, q_1 \rangle \otimes \langle M_2, f_2, q_2 \rangle \approx \langle M_2, f_2, q_2 \rangle \otimes \langle M_1, f_1, q_1 \rangle$,
- 3) $(\langle M_1, f_1, q_1 \rangle \perp \langle M_2, f_2, q_2 \rangle) \perp \langle M_3, f_3, q_3 \rangle \approx \langle M_1, f_1, q_1 \rangle \perp (\langle M_2, f_2, q_2 \rangle \perp \langle M_3, f_3, q_3 \rangle)$,
- 4) $(\langle M_1, f_1, q_1 \rangle \otimes \langle M_2, f_2, q_2 \rangle) \otimes \langle M_3, f_3, q_3 \rangle \approx \langle M_1, f_1, q_1 \rangle \otimes (\langle M_2, f_2, q_2 \rangle \otimes \langle M_3, f_3, q_3 \rangle)$,
- 5) $(\langle M_1, f_1, q_1 \rangle \perp \langle M_2, f_2, q_2 \rangle) \otimes \langle M_3, f_3, q_3 \rangle \approx (\langle M_1, f_1, q_1 \rangle \otimes \langle M_3, f_3, q_3 \rangle) \perp (\langle M_2, f_2, q_2 \rangle \otimes \langle M_3, f_3, q_3 \rangle)$,
- 6) $\langle M_1, f_1, q_1 \rangle \otimes \langle A, I, 0 \rangle \approx \langle M_1, f_1, q_1 \rangle$.

Proof. We shall show only 5). For the other isomorphisms, we can see easily. To prove it, it is enough to show the identity

$$(f_1 \perp f_2)^2 \otimes q_3 + (q_1 \perp q_2 - f_1 \times f_2) \otimes f_3^2 + (q_1 \perp q_2 - f_1 \times f_2) \otimes q_3 = (f_1^2 \otimes q_3 + q_1 \otimes f_3^2 + q_1 \otimes q_3) \perp (f_2^2 \otimes q_3 + q_2 \otimes f_3^2 + q_2 \otimes q_3) - (f_1 \otimes f_3) \times (f_2 \otimes f_3).$$

For any $\sum_i (x_i + y_i) \otimes z_i$ in $(M_1 \oplus M_2) \otimes M_3$, $(f_1 \perp f_2)^2 \otimes q_3 + (q_1 \perp q_2 - f_1 \times f_2) \otimes f_3^2 + (q_1 \perp q_2 - f_1 \times f_2) \otimes q_3 (\sum_i (x_i \oplus y_i) \otimes z_i) = \sum_i [(f_1(x_i) + f_2(y_i))^2 q_3(z_i) + (q_1(x_i) + q_2(y_i) - f_1(x_i) f_2(y_i)) f_3(z_i)^2 + 2(q_1(x_i) + q_2(y_i) - f_1(x_i) f_2(y_i)) q_3(z_i)] + \sum_{i < j} [(f_1(x_i) + f_2(y_i))(f_1(x_j) + f_2(y_j)) B_{q_3}(z_i, z_j) + (B_{q_1}(x_i, x_j) + B_{q_2}(y_i, y_j) - f_1(x_i) f_2(y_j) - f_1(x_j) f_2(y_i)) f_3(z_i) f_3(z_j) + (B_{q_1}(x_i, x_j) + B_{q_2}(y_i, y_j) - f_1(x_i) f_2(y_j))$

$$\begin{aligned}
 -f_1(x_j)f_2(y_i)B_{q_3}(z_i, z_j)] &= \sum_i [(f_1(x_i)^2q_3(z_i) + q_1(x_i)f_3(z_i)^2 + 2q_1(x_i)q_3(z_i)) + \\
 &(f_2(y_i)^2q_3(z_i)q_2(y_i)f_3(z_i)^2 + 2q_2(y_i)q_3(z_i)) - f_1(x_i)f_3(z_i)f_2(y_i)f_3(z_i)] + \sum_{i < j} [(f_1(x_i) \\
 &f_1(x_j)B_{q_3}(z_i, z_j) + B_{q_1}(x_i, x_j)f_3(z_i)f_3(z_j) + B_{q_1}(x_i, x_j)B_{q_3}(z_i, z_j)) + (f_2(y_i)f_2(y_j) \\
 &B_{q_3}(z_i, z_j) + B_{q_2}(y_i, y_j)f_3(z_i)f_3(z_j) + B_{q_2}(y_i, y_j)B_{q_3}(z_i, z_j)) - (f_1(x_i)f_3(z_i)f_2(y_j) \\
 &f_3(z_j) + f_1(x_j)f_3(z_j)f_2(y_i)f_3(z_i))] = (f_1^2 \otimes q_3 + q_1 \otimes f_3^2 + q_1 \otimes q_3) \perp (f_2^2 \otimes q_3 + \\
 &q_2 \otimes f_3^2 + q_2 \otimes q_3) - (f_1 \otimes f_3) \times (f_2 \otimes f_3)(\sum_i (x_i \oplus y_i) \otimes z_i).
 \end{aligned}$$

DEFINITION. An extended quadratic module $\langle M, f, q \rangle$ is called *hyperbolic* if the associated bilinear module $(M, B_{f,q})$ with $\langle M, f, q \rangle$ is hyperbolic, i.e. there exists an A -module N such that $M = N \oplus N'$ for some A -submodule N' , $f(N) = q(N) = 0$ and $N = N^\perp (= \{x \in M; B_{f,q}(x, N) = 0\})$.

From Proposition 5 and the well known properties on bilinear modules, we get the following proposition.

PROPOSITION 6.

- 1) If $\langle M, f, q \rangle$ and $\langle M', f', q' \rangle$ are hyperbolic, then so is also $\langle M, f, q \rangle \perp \langle M', f', q' \rangle$.
- 2) If M is a finitely generated projective A -module and $\langle M, f, q \rangle$ is hyperbolic then $\langle M, f, q \rangle$ is non-degenerate.
- 3) If $\langle M, f, q \rangle$ and $\langle M', f', q' \rangle$ are non-degenerate and $\langle M, f, q \rangle$ is hyperbolic, then $\langle M, f, q \rangle \otimes \langle M', f', q' \rangle$ is also hyperbolic.

3. Extended Witt ring $W^*(A)$.

From the argument in §2, we can construct a commutative ring $W^*(A)$. Let $\text{Qua}_p^*(A)$ be a full subcategory of $\text{Qua}^*(A)$ consisting of non-degenerate extended quadratic modules with finitely generated projective modules. In the category $\text{Qua}_p^*(A)$, as well as the construction of the Witt ring $W(A)$, we consider the full subcategory $\text{HQua}_p^*(A)$ consisting of hyperbolic extended quadratic modules. And, using the notation of K -theory in [1], we define the extended Witt ring $W^*(A)$ by $W^*(A) = \text{Coker}(K_0(\text{HQua}_p^*(A)) \rightarrow K_0(\text{Qua}_p^*(A)))$. Thus, it can be easily checked that $W^*(A)$ is a commutative ring with sum and product induced by orthogonal sum \perp and tensor product \otimes . We denote by $[\langle P, f, q \rangle]$ the class of $\langle P, f, q \rangle$ in $W^*(A)$.

THEOREM 2. *The extended Witt ring $W^*(A)$ has always the identity element $[\langle A, I, 0 \rangle]$, and there exists a ring homomorphism of the Witt ring $W(A)$ to $W^*(A)$. Then, the image of $W(A)$ becomes an ideal of*

$W^*(A)$. If 2 is invertible in A , then it is an isomorphism; $W(A) \xrightarrow{\approx} W^*(A)$.

Proof. Let $\text{Qua}_p(A)$ be the full subcategory of $\text{Qua}(A)$ consisting of non-degenerate quadratic modules (P, q) with finitely generated projective A -module P , and $\text{HQua}_p(A)$ the full subcategory of $\text{Qua}_p(A)$ whose objects are hyperbolic in $\text{Qua}_p(A)$. Consider the functor $\Phi: \text{Qua}_p(A) \rightarrow \text{Qua}_p^*(A)$; $(P, q) \rightsquigarrow \langle P, 0, q \rangle$, then we have the following commutative diagram

$$\begin{array}{ccccccc} K_0(\text{HQua}_p^*(A)) & \rightarrow & K_0(\text{Qua}_p^*(A)) & \rightarrow & W^*(A) & \rightarrow & 0 \\ & & \uparrow K_0(\Phi) & & \uparrow K_0(\Phi) & & \\ K_0(\text{HQua}_p(A)) & \rightarrow & K_0(\text{Qua}_p(A)) & \rightarrow & W(A) & \rightarrow & 0 \end{array}$$

where two rows are exact.

Thus, the ring homomorphism $K_0(\Phi)$ induces a ring homomorphism $\omega: W(A) \rightarrow W^*(A)$. Then, $\text{Im } \omega$ becomes an ideal of $W^*(A)$, for $[\langle P, f, q \rangle] [\langle P', 0, q' \rangle] = [\langle P \otimes P', 0, q \otimes q' + f^2 \overline{\otimes} q' \rangle]$ in $W^*(A)$. If 2 is invertible in A , by Remark 3, $K_0(\Phi)$ is an isomorphism, therefore, so is also $\omega: W(A) \xrightarrow{\approx} W^*(A)$.

4. The unit group of $W^*(A)$ and $Q_s(A)$.

In this section, we consider a relation between the separable quadratic extension group $Q_s(A)$ and the unit group $U(W^*(A))$ of the extended Witt ring $W^*(A)$.

THEOREM 3. *There exists a group homomorphism of $Q_s(A)$ to $U(W^*(A))$;*

$$\theta: Q_s(A) \longrightarrow U(W^*(A)); [U, f, q] \rightsquigarrow [\langle U, f, 2q \rangle].$$

Proof. Let $[U, f, q]$ be an element in $Q_s(A)$. By Theorem 1, the bilinear module $(U, D_{f,q})$, called the discriminant of $[U, f, q]$, is non-degenerate. Since $D_{f,q}(x, y) = f(x)f(y) + 2B_q(x, y) = f(x)f(y) + B_{2q}(x, y) = B_{f,2q}(x, y)$ for any x and y in U , we have $D_{f,q} = B_{f,2q}$. Therefore, $\langle U, f, 2q \rangle$ is in $\text{Qua}_p^*(A)$. Now, we shall show that θ is well defined: If $[U, f, q] = [U', f', q']$ is in $Q_s(A)$, then there exist an A -isomorphism $\sigma: U \rightarrow U'$ and an A -homomorphism $g: U \rightarrow A$ such that $q' \circ \sigma = q + fg - g^2$ and $f' \circ \sigma = f - 2g$. Then, we get $2q' \circ \sigma = 2q + 2fg - 2g^2$ and $f' \circ \sigma =$

$f - 2g$, that is, $\langle U, f, 2q \rangle \approx \langle U', f', 2q' \rangle$ in $\text{Qua}_p^*(A)$. Thus, the map $\theta: Q_s(A) \rightarrow W^*(A); [U, f, q] \rightsquigarrow [\langle U, f, 2q \rangle]$ is well defined. Furthermore, we have

$\theta([U, f, q][U', f', q']) = \theta([U \otimes U', f \otimes f', f^2 \bar{\otimes} q' + q \bar{\otimes} f'^2 + 2q \otimes q']) = [\langle U \otimes U', f \otimes f', f^2 \bar{\otimes} 2q' + 2q \bar{\otimes} f'^2 + 2q \otimes 2q' \rangle] = [\langle U, f, 2q \rangle] \cdot [\langle U', f', 2q' \rangle]$, and $\theta([A, I, 0]) = [\langle A, I, 0 \rangle]$. Accordingly, $\text{Im } \theta$ is contained in $U(W^*(A))$ and $\theta: Q_s(A) \rightarrow U(W^*(A))$ is a group homomorphism.

Remark 5.

1) if K is a field with the characteristic $\neq 2$, then $U(W^*(A)) = U(W(A)) \approx U(K)/U(K)^2$, $Q_s(K) \approx U(K)/U(K)^2$ and θ is an isomorphism.

2) If K is a field with characteristic 2, then θ is a zero homomorphism.

REFERENCES

- [1] H. Bass: Lecture on Topics in Algebraic K-theory, Tata Institute of Fundamental Research, Bombay, 1967.
- [2] P. M. Cohn: Quadratic extensions of skew fields, Proc. London Math. Soc. **11** (1961) 531-556.
- [3] T. Kanzaki: On commutator ring and Galois theory of separable algebras, Osaka J. Math. **1** (1964).
- [4] T. Kanzaki: Generalized crossed product and Brauer group, Osaka J. Math. **5** (1968) 175-188.
- [5] T. Kanzaki: On bilinear module and Witt ring over a commutative ring, Osaka J. Math. **8** (1971) 485-496.
- [6] P. A. Micali and O. E. Villamayor: Algebra de Clifford et groupe de Brauer, Ann. scient. Ec. Norm. Sup., t. **4** (1971) 285-310.
- [7] P. P. Revoy: Sur les deux premiers invariants d'une forme quadratique, Ann. scient. Ec. Norm. Sup., t. **4** (1971) 311-319.

Osaka City University

