



On the Road Coloring Problem

Author(s): Joel Friedman

Source: *Proceedings of the American Mathematical Society*, Vol. 110, No. 4 (Dec., 1990), pp. 1133-1135

Published by: [American Mathematical Society](#)

Stable URL: <http://www.jstor.org/stable/2047767>

Accessed: 08/10/2010 02:12

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ams>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Proceedings of the American Mathematical Society*.

<http://www.jstor.org>

ON THE ROAD COLORING PROBLEM

JOEL FRIEDMAN

(Communicated by Thomas H. Brylawski)

ABSTRACT. Let $G = (V, E)$ be a strongly connected, aperiodic, directed graph having outdegree 2 at each vertex. A *red-blue coloring* of G is a coloring of the edges with the colors red and blue such that each vertex has one red edge and one blue edge leaving it. Given such a coloring, we define $R: V \rightarrow V$ by $R(v) = w$ iff there is a red edge from v to w . Similarly we define $B: V \rightarrow V$. G is said to be collapsible if some composition of R 's and B 's maps V to a single vertex. The road coloring problem is to determine whether G has a collapsible coloring. It has been conjectured that all such G have a collapsible coloring. Since G has outdegree 2 everywhere and is strongly connected, the adjacency matrix, A , of G has a positive left eigenvector $w = (w(v_1), \dots, w(v_n))$ with eigenvalue 2, i.e. $wA = 2w$. Furthermore, we can assume that w 's components are integers with no common factor. We call $w(v)$ the *weight* of v . Let $W \equiv \sum_{v \in V} w(v)$, defined to be the weight of the graph. We will prove that if G has a simple cycle of length relatively prime to W , then G is collapsibly colorable.

1. INTRODUCTION

Let $G = (V, E)$ be a directed graph. G is said to be *strongly connected* if any vertex can reach any other vertex by a path in G . G is said to be *aperiodic* if V cannot be partitioned into $d > 1$ sets $V_1, \dots, V_d = V_0$ such that all edges (u, v) with $u \in V_i$ have $v \in V_{i+1}$.

Let G have outdegree 2 at each vertex. A *red-blue coloring* of G is a coloring of the edges with the colors red and blue such that each vertex has one red edge and one blue edge leaving it. Given such a coloring, we define $R: V \rightarrow V$ by $R(v) = w$ iff there is a red edge from v to w . Similarly we define $B: V \rightarrow V$. G is said to be collapsible if some composition of R 's and B 's maps V to a single vertex.

The road coloring problem is to determine whether G has a collapsible coloring. It has been conjectured that all such G have a collapsible coloring (i.e. G aperiodic, strongly connected). This problem originated in connection with [2], and appears explicitly in [1]. There it was assumed that G has no multiple edges, i.e. each vertex has edges to two distinct vertices. In [4] it was shown that a graph which has no multiple edges and a simple cycle of prime length

Received by the editors December 15, 1986 and, in revised form, July 25, 1988.
1980 *Mathematics Subject Classification* (1985 Revision). Primary 05C50.

©1990 American Mathematical Society
0002-9939/90 \$1.00 + \$.25 per page

is collapsibly colorable. In this paper we analyze a property of noncollapsible colorings; this gives further conditions for collapsible colorability.

Since G has outdegree 2 everywhere and is strongly connected, the adjacency matrix, A , of G has a positive left eigenvector $w = (w(v_1), \dots, w(v_n))$ with eigenvalue 2, i.e. $wA = 2w$. Furthermore, we can assume that w 's components are integers with no common factor. We call $w(v)$ the *weight* of v . Let $W \equiv \sum_{v \in V} w(v)$, defined to be the weight of the graph. For example, if G has indegree 2 everywhere, then $W = |V|$, the size of V . We will prove that if G has a simple cycle of length relatively prime to W , then G is collapsibly colorable.

I would like to thank Brian Marcus for posing this problem to me and for encouragement on it.

2. AN OBSERVATION

Let G be a graph as before and fix a coloring of G . For $T \subset V$, let $w(T)$, the weight of T , be the sum of the weights of the vertices in T . We say that T is collapsible if T can be mapped to a single vertex by some composition of R 's and B 's. Let T_0 be a collapsible set of maximum weight, w_0 .

Theorem 2.1. *There exist i and subsets T_1, \dots, T_{i-1} , each of which is collapsible and of weight w_0 , such that T_0, \dots, T_{i-1} is a partition of V . In particular, $w_0 i = W$.*

Proof. Let $U \subset V$, and consider its backward images, $R^{-1}U$, $B^{-1}U$. Since $wA = 2w$, we have $w(R^{-1}U) + w(B^{-1}U) = 2w(U)$. It follows that either $R^{-1}U$ or $B^{-1}U$ has greater weight than U , or both have weight equal to that of U . Also, if U is collapsible, then so are $R^{-1}U$ and $B^{-1}U$. From these observations it follows that R^{-1} and B^{-1} of any collapsible set of maximum weight is again collapsible of maximum weight.

We have $f: T_0 \rightarrow v_0$, where f is a composition of R 's and B 's and $v_0 \in V$. Suppose that T_0 is not all of V . We claim that we can extend f backwards to $g = fh$, h being a composition of R 's and B 's, so that g maps T_0 to one vertex and maps another maximum weight collapsible set to another vertex. To see this, consider f^{-1} , which maps V to subsets of V ; these subsets form a partition of V . Pick any vertex $v \notin T_0$, and let $g = fhf$, where h is any composition of R 's and B 's mapping v_0 to v . Notice that $f(v) \neq v_0$, since $v \notin f^{-1}v_0$ since T_0 is maximal, and that $f^{-1}h^{-1}T_0$ is a collapsible set of maximum weight by the preceding paragraph. It follows that g collapses two disjoint maximum weight collapsible sets. If these two sets do not comprise all of V , then we can extend g backwards to a function which collapses three disjoint maximum weight collapsible sets, one of which is T_0 , by iterating the same argument. Repeating this process enough times completes the proof of the theorem. \square

A set of vertices U is called a *minimal image* of V if U is the image of V under some composition of R 's and B 's and if U cannot be reduced in

size by any further composition of R 's and B 's. It is easy to see that any such U has size i —any set of size $> i$ must contain at least two points in one of T_0, \dots, T_{i-1} and can therefore be reduced, and any set of size $< i$ cannot be the image of a set of vertices of weight $> w_0$ times its cardinality, which is less than W .

3. COLORING WITH RED TREES

Given that G has a simple cycle of length m through vertices v_0, \dots, v_{m-1} , we can choose one outgoing edge from each other vertex so that any path through these edges leads to the cycle. Coloring these edges and the edges of the cycle red, we get a coloring of G which has a red cycle of length m and a set of red paths taking each vertex into this cycle. We call such colorings of G *colorings with red trees* (the red edges of G form a tree plus an extra edge to complete the cycle).

The following result appears in O'Brien's paper:

Theorem 3.1. *Let G have a coloring with a red tree and a red cycle of length m . Then i , the size of a minimal image, divides m .*

Proof. See [4]. \square

Combining this with Theorem 2.1 we have:

Corollary 3.2. *If G is colored with a red tree and a red cycle of length m , m relatively prime to W , then G is collapsible.*

Proof. Since i divides m and W , $i = 1$. That is to say, G is collapsible.

Corollary 3.3. *If W is a prime power, then G is collapsibly colorable.*

Proof. Since G is aperiodic, G has a cycle relatively prime to p where p is the prime with $W = p^j$. This cycle can be written as the union of simple cycles, and at least one of the simple cycles must be relatively prime to p . \square

In the case where G has indegree 2, has a prime number of vertices, and has no multiple edges, it was previously known that G is collapsibly colorable. This was due to Nelson Markley and Michael Paul, unpublished, based on ideas of G. A. Hedlund in [3].

REFERENCES

1. R. L. Adler, L. W. Goodwyn, and B. Weiss, *Equivalence of topological Markov shifts*, Israel J. Math. **27** (1977), 49–63.
2. R. L. Adler and B. Weiss, *Similarity of automorphisms of the torus*, Mem. Amer. Math. Soc., no. 98, Amer. Math. Soc., Providence, RI, 1970.
3. G. A. Hedlund, *Endomorphisms and automorphisms of the shift*, Math. Systems Theory **3** (1969), 320–375.
4. G. L. O'Brien, *The road-colouring problem*, Israel J. Math. **39** (1981), 145–154.