

On the stable rank and reducibility in algebras of real symmetric functions

R. Rupp*¹ and A. Sasane**²

¹ Fachbereich Allgemeinwissenschaften, Georg-Simon-Ohm-Fachhochschule Nürnberg, Keßlerplatz 12, D-90489 Nürnberg, Germany.

² Mathematics Department, London School of Economics, Houghton Street, London WC2A 2AE, United Kingdom.

Received April 30, 2007

Key words Real Banach algebras, Bass stable rank, topological stable rank, reducibility, stabilization.

MSC (2000) Primary 46J15; Secondary 19B10, 30H05, 93D15

Let $A_{\mathbb{R}}(\overline{\mathbb{D}})$ denote the set of functions belonging to the disc algebra having real Fourier coefficients. We show that $A_{\mathbb{R}}(\overline{\mathbb{D}})$ has Bass and topological stable ranks equal to 2, which settles the conjecture made by Brett Wick in [18]. We also give a necessary and sufficient condition for reducibility in some real algebras of functions on symmetric domains with holes, which is a generalization of the main theorem in [18]. A sufficient topological condition on the symmetric open set D is given for the corresponding real algebra $A_{\mathbb{R}}(\overline{D})$ to have Bass stable rank equal to 1.

Copyright line will be provided by the publisher

1 Introduction

The notion of stable rank of a ring (which we call Bass stable rank) was introduced by H. Bass [2] to facilitate computations in algebraic K-theory. We recall the definition of the Bass stable rank of a ring below.

Definition 1.1 Let \mathcal{A} be a commutative ring with identity 1. Let $n \in \mathbb{N}$. An element $a = (a_1, \dots, a_n) \in \mathcal{A}^n$ is called *unimodular* if there exists $b = (b_1, \dots, b_n) \in \mathcal{A}^n$ such that

$$\langle b, a \rangle := \sum_{k=1}^n b_k a_k = 1.$$

We denote by $U_n(\mathcal{A})$ the set of unimodular elements of \mathcal{A}^n .

We say that $a = (a_1, \dots, a_n) \in U_n(\mathcal{A})$ is *reducible* (in \mathcal{A}), if there exist $h_1, \dots, h_{n-1} \in \mathcal{A}$ such that $(a_1 + h_1 a_n, \dots, a_{n-1} + h_{n-1} a_n) \in U_{n-1}(\mathcal{A})$. The *Bass stable rank* of \mathcal{A} , denoted by $\text{bsr } \mathcal{A}$, is the least integer n such that every $a \in U_{n+1}(\mathcal{A})$ is reducible (and it is infinite if no such integer n exists).

The Bass stable rank is a purely algebraic notion, but when studying commutative Banach algebras of functions, analysis also plays a role. In [13], M. Rieffel introduced the notion of topological stable rank, analogous to the concept of Bass stable rank:

Definition 1.2 Let \mathcal{A} denote a commutative unital Banach algebra. The *topological stable rank* of \mathcal{A} , denoted by $\text{tsr } \mathcal{A}$, is the minimum integer n such that $U_n(\mathcal{A})$ is dense in \mathcal{A}^n (and it is infinite if no such integer exists).

Jones, Marshall and Wolff [9] showed that the Bass stable rank of the complex disc algebra $A(\overline{\mathbb{D}})$ is equal to 1, and Rieffel [13] showed that its topological stable rank is equal to 2. Recall that the complex disc algebra $A(\overline{\mathbb{D}})$ is the Banach algebra of all complex-valued functions defined on the closed unit disc $\overline{\mathbb{D}}$ that are holomorphic in the open unit disc \mathbb{D} and continuous on $\overline{\mathbb{D}}$, endowed with the supremum norm: $\|f\|_{\infty} = \sup_{z \in \overline{\mathbb{D}}} |f(z)|$.

* Part of this work was done while R. Rupp visited London School of Economics. He wishes to thank LSE for their kind hospitality and the University of Applied Sciences, Nürnberg for the sabbatical. e-mail: rudolf.rupp@fh-nuernberg.de

** Corresponding author. A. Sasane is supported by the Nuffield Grant NAL/32420. e-mail: A.J.Sasane@lse.ac.uk, Phone: +44 20 7955 6585, Fax: +44 20 7966 6877

In this article, we study the Bass/topological stable ranks, and also consider reducibility of corona pairs, in some real Banach algebras of “real symmetric” functions. We define these below. Throughout this article, we use z^* to denote the complex conjugate of z , and we use $\overline{\Omega}$ to denote the closure of the set $\Omega \subset \mathbb{C}$.

Definition 1.3 The *real disc algebra*, denoted by $A_{\mathbb{R}}(\overline{\mathbb{D}})$, is the set of all functions of $A(\overline{\mathbb{D}})$ having real Fourier coefficients. Equivalently,

$$A_{\mathbb{R}}(\overline{\mathbb{D}}) = \{f \in A(\overline{\mathbb{D}}) \mid \forall z \in \overline{\mathbb{D}}, f(z) = (f(z^*))^*\}.$$

The real disc algebra $A_{\mathbb{R}}(\overline{\mathbb{D}})$ is a real Banach algebra with the supremum norm $\|\cdot\|_{\infty}$.

More generally, if D is an open set in \mathbb{C} , then by $A(\overline{D})$ we mean the set of functions holomorphic in D that are continuous and bounded on \overline{D} . If D is *real symmetric* (that is, $z \in D$ if and only if $z^* \in D$), then we use the symbol $A_{\mathbb{R}}(\overline{D})$ to denote the set of functions f belonging to $A(\overline{D})$ that are *real symmetric*, that is, $f(z) = (f(z^*))^*$ ($z \in \overline{D}$).

If D is a real symmetric open set, then $C_{\mathbb{R}}(\overline{D})$ denotes the set of complex-valued, bounded, continuous functions f defined on \overline{D} , that satisfy $f(z) = (f(z^*))^*$ ($z \in \overline{D}$).

Brett Wick conjectured [18] that the Bass stable rank of $A_{\mathbb{R}}(\overline{\mathbb{D}})$ is equal to 2, and we prove this in Section 2, by first showing that the topological stable rank of $A_{\mathbb{R}}(\overline{\mathbb{D}})$ is 2. In Section 4 we extend the main result of B. Wick [18] to the case of subalgebras of $A_{\mathbb{R}}(\overline{\mathbb{D}})$. We also completely characterize reducible elements in algebras $A_{\mathbb{R}}(\overline{D})$ of real symmetric functions on domains with holes (under mild assumptions) in Theorem 6.4. This generalizes the main result in [18] from the case of the disc to more general domains. Finally, in Theorem 6.6, we give a sufficient topological condition on the open set D for $A_{\mathbb{R}}(\overline{D})$ to have Bass stable rank equal to 1.

2 Bass and topological stable rank of $A_{\mathbb{R}}(\overline{\mathbb{D}})$

In this section we prove that $\text{bsr } A_{\mathbb{R}}(\overline{\mathbb{D}}) = \text{tsr } A_{\mathbb{R}}(\overline{\mathbb{D}}) = 2$.

We begin by making the observation that the polynomials with real coefficients are dense in $A_{\mathbb{R}}(\overline{\mathbb{D}})$. Indeed, given $f \in A_{\mathbb{R}}(\overline{\mathbb{D}})$, f has real Fourier coefficients, which are the same as the coefficients in the Taylor expansion of the analytic function f about the point 0 in \mathbb{D} . Since f is continuous on the circle, and its negative Fourier coefficients vanish, the Cesàro means of the Fourier series for f are trigonometric polynomials with real coefficients which converge uniformly to f . The corresponding polynomials in z give the desired sequence converging uniformly to f in $A_{\mathbb{R}}(\overline{\mathbb{D}})$.

Theorem 2.1 *The topological stable rank of $A_{\mathbb{R}}(\overline{\mathbb{D}})$ is 2.*

Proof. First of all we note that $U_1(A_{\mathbb{R}}(\overline{\mathbb{D}}))$ is not dense in $A_{\mathbb{R}}(\overline{\mathbb{D}})$. Indeed, $U_1(A_{\mathbb{R}}(\overline{\mathbb{D}}))$ is the set of units in $A_{\mathbb{R}}(\overline{\mathbb{D}})$, and f is invertible as an element in $A_{\mathbb{R}}(\overline{\mathbb{D}})$ only if it has no zero in $\overline{\mathbb{D}}$. But the uniform limit of a sequence of functions from the disc algebra which are never zero in \mathbb{D} is either identically zero or has no zeros in \mathbb{D} ; see [1, Theorem 2, p. 178]. So taking any function with finitely many zeros in \mathbb{D} , say z , we have a contradiction. So $\text{tsr } A_{\mathbb{R}}(\overline{\mathbb{D}}) > 1$.

Next we show that $U_2(A_{\mathbb{R}}(\overline{\mathbb{D}}))$ is dense in $A_{\mathbb{R}}(\overline{\mathbb{D}})^2$. Take $(f, g) \in A_{\mathbb{R}}(\overline{\mathbb{D}})^2$ and approximate f, g by polynomials p, q , respectively, having real coefficients. Since $p \in \mathbb{R}[z]$, we have the following product representation for p :

$$p(z) = C \prod (z - r_j) \prod (z^2 + s_j z + t_j),$$

where C, r_j, s_j, t_j are real numbers. If p and q have a common root in $\overline{\mathbb{D}}$, then we replace r_j, s_j, t_j by $r_j + \epsilon, s_j + \epsilon, t_j + \epsilon$ with a sufficiently small real ϵ so that the new polynomial \tilde{p} has no common root with q in $\overline{\mathbb{D}}$, and so $(\tilde{p}, q) \in U_2(A_{\mathbb{R}}(\overline{\mathbb{D}}))$ is near (f, g) . Consequently $\text{tsr } A_{\mathbb{R}}(\overline{\mathbb{D}}) \leq 2$. \square

We recall the following result [4, Theorem 3, p. 293]:

Proposition 2.2 *Let \mathcal{A} be a commutative unital real (or complex) Banach algebra. If $U_n(\mathcal{A})$ is a dense subset of \mathcal{A}^n , then $\text{bsr } \mathcal{A} \leq n$.*

Theorem 2.3 *The Bass stable rank of $A_{\mathbb{R}}(\overline{\mathbb{D}})$ is 2.*

Proof. First we show that $\text{bsr } A_{\mathbb{R}}(\overline{\mathbb{D}}) > 1$. Consider $a := (z, 1 - z^2) \in A_{\mathbb{R}}(\overline{\mathbb{D}})^2$. The element a is unimodular, since with $b := (z, 1) \in A_{\mathbb{R}}(\overline{\mathbb{D}})^2$, we have $\langle b, a \rangle = 1$. However a is not reducible. Indeed, otherwise there exists an element $h \in A_{\mathbb{R}}(\overline{\mathbb{D}})$ such that with $f(z) := z + h(z)(1 - z^2)$, $z \in \overline{\mathbb{D}}$, f is an invertible element of $A_{\mathbb{R}}(\overline{\mathbb{D}})$. However, $f(-1) = -1$ and $f(1) = 1$, and so by the intermediate value theorem $f(c) = 0$ for some $c \in (-1, 1)$, contradicting the invertibility of f .

From Theorem 2.1 it follows that $U_2(A_{\mathbb{R}}(\overline{\mathbb{D}}))$ is dense in $A_{\mathbb{R}}(\overline{\mathbb{D}})^2$, and so by Proposition 2.2, we obtain that $\text{bsr } A_{\mathbb{R}}(\overline{\mathbb{D}}) \leq 2$. This completes the proof. \square

Remark 2.4

1. Brett Wick conjectured in [18] that the Bass stable rank of $A_{\mathbb{R}}(\overline{\mathbb{D}})$ is equal to 2; the above result settles this conjecture.
2. Bass and topological stable ranks of $A_{\mathbb{R}}(\overline{\mathbb{D}})$ play an important role in *control theory* in the problem of stabilization of linear systems. We refer the reader to [12] and [17] for background on the connection between stable rank and control theory.

3 Preliminaries

3.1 Reducibility in general real Banach algebras

We adapt the definition from [14, Definition 1.1] to the case of *real* Banach algebras as follows:

Definition 3.1 Let \mathcal{A} be a real commutative Banach algebra with unit element denoted by 1. Suppose that $(f, g) \in \mathcal{A}^2$ and $\lambda, \mu \in \mathbb{R}$. The real numbers λ, μ are called *equivalent* if there exist elements $h, k \in \mathcal{A}$ such that

$$f - \lambda + hg = (f - \mu) \exp(k).$$

It is not hard to see that for fixed (f, g) this indeed gives an equivalence relation. We denote the equivalence class of λ by $[\lambda]$.

Theorem 3.2 *If $(f - \mu, g) \in U_2(\mathcal{A})$, then $[\mu]$ is open in \mathbb{R} .*

The proof is the same as that of [14, Proposition 1.2], but all numbers have to be real.

Theorem 3.3 *Let $(f, g) \in U_2(\mathcal{A})$ and suppose that for some positive ϵ the interval $(-\infty, \epsilon)$ belongs to the real inversion set*

$$I_{\mathbb{R}}(f, g) := \{\lambda \in \mathbb{R} \mid (f - \lambda, g) \in U_2(\mathcal{A})\}.$$

Then (f, g) is reducible in \mathcal{A} .

The proof is similar to that of [14, Proposition 1.3]: Take a real number M such that $M > \|f\|$. Then there exists $l \in \mathcal{A}$ such that $f - (-M) = \exp(l)$. Then $(f - (-M), g) \in U_2(\mathcal{A})$ and, by assumption, $-M$ and 0 belong to the same connected component of $I_{\mathbb{R}}(f, g)$. But then Theorem 3.2 implies that $\mu = -M$ and $\lambda = 0$ are equivalent. (Otherwise the open connected set $(-\infty, \epsilon)$ would split into disjoint open sets, namely certain equivalence classes.) Thus there exist $h, k \in \mathcal{A}$ such that $f + hg = (f - (-M)) \exp(k) = \exp(k + l)$.

3.2 Some notation and terminology.

When we consider domains with holes in sections 5 and 6, the following notation will be convenient.

Notation 3.4 Let D denote a bounded symmetric domain in \mathbb{C} with n holes, having a boundary that is a union of pairwise disjoint Jordan curves. The outer boundary curve is denoted by Γ_{n+1} . From these n holes, bounded by pairwise disjoint Jordan curves Γ_j , we have r holes intersecting \mathbb{R} and $2m$ which do not intersect \mathbb{R} . Here $n = r + 2m$. For notational reasons the Jordan curves Γ_j belonging to the upper half plane are indexed by $j = r + 1, \dots, r + m$, while the Jordan curves belonging to the lower half plane are indexed by $j = r + m + 1, \dots, r + 2m$. These curves Γ_{r+m+j} are the reflection of Γ_{r+j} . Let C_j be a hole of D ($j = 1, \dots, r, r + 1, \dots, r + 2m$). Choose for each $j \in \{1, \dots, r\}$ a point $x_j \in \mathbb{R} \cap C_j$, and for $j \in \{r + 1, \dots, r + m\}$ let $z_j \in C_j$. Finally let $S = \{x_1, \dots, x_r, z_1, \dots, z_m, z_1^*, \dots, z_m^*\}$. See Figure 1. To ensure that the domain D is always to the left of the orientation, we assume that the Jordan curves $\Gamma_1, \dots, \Gamma_n$ surrounding the holes are negatively oriented, whereas the outer boundary curve Γ_{n+1} is positively oriented.




Fig. 1 The domain D with $n = r + 2m$ holes.

Definition 3.5 If D denotes an open subset of \mathbb{C} , then we say the *corona theorem* holds for $\mathcal{A} (\subset A_{\mathbb{R}}(\overline{D}))$ if the following is true for all $n \in \mathbb{N}$:

$(f_1, \dots, f_n) \in U_n(\mathcal{A})$ if and only if there exists $\delta > 0$ such that $\sum_{j=1}^n |f_j(z)| \geq \delta$ ($z \in \overline{D}$), that is, if and only if the functions f_1, \dots, f_n have no common zero in \overline{D} .

Definition 3.6 For functions $g \in A_{\mathbb{R}}(\overline{D})$ the *zero set* Z_g of g is

$$Z_g := \{z \in \overline{D} \mid g(z) = 0\},$$

and for $\delta > 0$ the *level set* $Z(\delta)$ is

$$Z(\delta) := \{z \in \overline{D} \mid |g(z)| \leq \delta\}.$$

Of course the inclusion $Z_g \subset Z(\delta)$ holds.

Following B. Wick [18], we will use the following terminology:

Definition 3.7 Let $f, g \in A_{\mathbb{R}}(\overline{D})$. The function f is said to be *positive on real zeros of g* (abbreviated as f is *POZ of g*), if f has the same sign at all real zeros of g .

For example, consider $(f, g) := (z, 1 - z^2) \in A_{\mathbb{R}}(\overline{\mathbb{D}})^2$. Then f is not POZ of g .

3.3 Technical lemmata.

In this subsection, we will prove two technical lemmata which will be used in the sequel. The first one is well-known among the workers in the field. For the sake of completeness we include a proof.

Lemma 3.8 *For every function $g \in A_{\mathbb{R}}(\overline{\mathbb{D}})$, the complement $\mathbb{C} \setminus Z(\delta)$ of the level set $Z(\delta)$ is connected. Moreover, the complement $\mathbb{C} \setminus Z_g$ of the zero set Z_g is also connected.*

Proof. For constant functions g the assertions are trivially true. So we may assume that g is non constant. By the very definition $\mathbb{C} \setminus \overline{\mathbb{D}} \subset \mathbb{C} \setminus Z(\delta)$, and so if $\mathbb{C} \setminus Z(\delta)$ is not connected, there exists a bounded component $G \subset \overline{\mathbb{D}}$. Being in the complement of the level set, we must have $|g(z)| \geq \delta$ for all $z \in \partial G \subset \overline{\mathbb{D}}$. On the other hand, $|g(z)| \leq \delta$ for all $z \in \partial G \subset \overline{\mathbb{D}}$, because $\partial G \subset \partial(\mathbb{C} \setminus Z(\delta)) = \partial Z(\delta) \subset Z(\delta)$. This gives $|g(z)| = \delta$ for all $z \in \partial G$ by the maximum modulus theorem, implying $|g(z)| \leq \delta$ for all $z \in G$, a contradiction. Hence no such bounded component of the complement of $Z(\delta)$ can exist. That $\mathbb{C} \setminus Z_g$ is connected follows from

$$\mathbb{C} \setminus Z_g = \bigcup_{\delta > 0} (\mathbb{C} \setminus Z(\delta))$$

and the fact that $\bigcap_{\delta > 0} (\mathbb{C} \setminus Z(\delta)) \neq \emptyset$ (indeed, $\mathbb{C} \setminus \overline{\mathbb{D}} \subset \mathbb{C} \setminus Z(\delta)$ for every $\delta > 0$). □

In order to facilitate handling zero sets, we prove the following result, in which we enclose the zero set by finitely many closed sets.

Lemma 3.9 *Let $g \in A_{\mathbb{R}}(\overline{\mathbb{D}})$ be such that it has at least one zero in $\overline{\mathbb{D}}$, but it does not vanish identically. Then for all $\delta > 0$, there exist finitely many closed sets $H_1, \dots, H_N \subset \overline{\mathbb{D}}$, lying symmetrically with respect to the real axis, that is, $H_j = H_k^*$ for certain j, k , with the following properties:*

1. $Z_g \subset \cup_{j=1}^N H_j \subset Z(\delta)$.
2. $H_j \cap H_k = \emptyset$ ($j \neq k$).
3. 1° *If no real zero of g belongs to H_j , then $H_j \cap \mathbb{R} = \emptyset$, H_j belongs entirely to the upper (respectively lower) half plane, and $H_j = H_k^*$ for some $j \neq k$.*
 2° *If at least one real zero x_0 of g belongs to H_j (that is, $x_0 \in Z_g \cap H_j \cap \mathbb{R}$), then $H_j = H_j^*$ holds and H_j is connected.*

Proof. The zero set Z_g is compact, hence finitely many components K_j , $j = 1, \dots, M$, of the relatively open set $H := \{z \in \overline{\mathbb{D}} \mid |g(z)| < \delta\}$ will suffice to cover Z_g . Note that these components are open because H is locally connected. Since \overline{H} is symmetric with respect to the real axis, its components are symmetric as well. Unfortunately, the closures $\overline{K_j}$ need not be disjoint. However, we may take the closed connected components of $\cup_{j=1}^M \overline{K_j}$; at most there are M such components. These components are symmetric as well.

To ensure all the three assertions hold, we must eventually truncate the closed sets $\overline{K_j}$:

- 1° *If no real zero of g is in $\overline{K_j}$, then $|g(z)| \geq \rho_j > 0$ for all $z \in (\overline{K_j} \cap \mathbb{R}) \times (|\operatorname{Im}(z)| \leq \delta_j)$. Hence no zero of g belongs to $z \in (\overline{K_j} \cap \mathbb{R}) \times (|\operatorname{Im}(z)| \leq \delta_j)$. We truncate as follows: $H_j := \overline{K_j} \cap (|\operatorname{Im}(z)| \geq \delta_j)$. The closed set $\overline{K_j}$ splits in two closed sets belonging entirely to the upper (respectively lower) half plane.*
- 2° *If at least one real zero of g belongs to $\overline{K_j}$, then we don't truncate, that is, $H_j := \overline{K_j}$. By symmetry we have $H_j = H_j^*$, and $H_j = \overline{K_j}$ is connected, because K_j is.*

By construction all the zeros of g belong to exactly one closed set H_j , $j = 1, \dots, N$. □

4 Reducibility in real symmetric algebras on the disc

In this section we generalize the main result of B. Wick [18] to subalgebras \mathcal{A} of $A_{\mathbb{R}}(\overline{\mathbb{D}})$.

In real algebras $\mathcal{A} \subset A_{\mathbb{R}}(\overline{\mathbb{D}})$ where the corona theorem holds, the real inversion set from Theorem 3.3 is given by

$$I_{\mathbb{R}}(f, g) = \mathbb{R} \setminus f(Z_g).$$

That the corona theorem holds for $A_{\mathbb{R}}(\overline{\mathbb{D}})$ follows easily from the corona theorem for the complex algebra $A(\overline{\mathbb{D}})$ by symmetrization of the solution. We refer the reader to [10] for a constructive proof (not using any Gelfand theory nor Banach algebra theory) of the corona theorem for certain subalgebras of $A(\overline{\mathbb{D}})$ and certain domains D (including, of course, \mathbb{D}).

Theorem 4.1 (Units) *Let \mathcal{A} denote one of the algebras $A_{\mathbb{R}}(\overline{\mathbb{D}}), C_{\mathbb{R}}(\overline{\mathbb{D}})$ respectively. For any unit $u \in \mathcal{A}^{-1}$ either u or $-u$ can be expressed as*

$$u = \exp(h),$$

where $h \in \mathcal{A}$.

Proof. First of all we prove the theorem in case $\mathcal{A} = C_{\mathbb{R}}(\overline{\mathbb{D}})$. Choose a closed disc $U \supset \overline{\mathbb{D}}$, small enough so that a continuous extension of u to U has no zeros in \overline{U} . Using a theorem of Borsuk [3, Corollary 4.33], there exists a continuous logarithm h on $U \supset \overline{\mathbb{D}}$. However, this function h need not be symmetric. Because the unit u is symmetric we derive

$$u(z) = \exp(h(z)) = \exp((h(z^*))^*), \quad z \in \overline{\mathbb{D}}.$$

Because $\overline{\mathbb{D}}$ is connected and h is continuous in $\overline{\mathbb{D}}$, there exists an integer k such that

$$h(z) = (h(z^*))^* + 2k\pi i.$$

Restricting to the interval $[-1, 1]$ gives $\operatorname{Im} h(x) = k\pi$. Since u is a unit, $u(z)$ is either a positive or negative real number when $z \in [-1, 1]$.

- 1° If $u(x) > 0$, then we have $u(x) = \exp h(x) > 0$, that is, the integer k is even. But then $h - k\pi i$ is a symmetric continuous logarithm of u .
- 2° If $u(x) < 0$, then we just look at the unit $-u$.

Hence there exists $h \in C_{\mathbb{R}}(\overline{\mathbb{D}})$ such that $u = \exp(h)$.

The remaining case $\mathcal{A} = A_{\mathbb{R}}(\overline{\mathbb{D}})$ now follows from the first case and the implicit function theorem as follows: By the holomorphic inverse function theorem applied to $z \mapsto \exp(z)$, we see that it has a local holomorphic inverse around each point z_0 , say g_{z_0} . Thus $z \mapsto h(z) = g_{z_0}(u(z))$ is holomorphic near z_0 as well. \square

Theorem 4.2 *Let \mathcal{A} denote a subalgebra of $A_{\mathbb{R}}(\overline{\mathbb{D}})$ containing all real polynomials such that the corona theorem holds for \mathcal{A} . The following are equivalent for any unimodular pair $(f, g) \in U_2(\mathcal{A})$:*

1. *There exists a continuous and zero free extension F of f from the zero set Z_g to $F \in C_{\mathbb{R}}(\overline{\mathbb{D}})^{-1}$.*
2. *(f, g) is reducible in \mathcal{A} , that is, there exists a unit $u \in \mathcal{A}^{-1}$ and there exists a $h \in \mathcal{A}$ such that $f + hg = u$.*

Proof. The implication (2) \Rightarrow (1) is obvious: indeed if there exist $h \in \mathcal{A}$ and a unit $u \in \mathcal{A}^{-1}$ such that $f + hg = u$, then u serves as the desired zero free extension of f from the zero set Z_g to $u \in C_{\mathbb{R}}(\overline{\mathbb{D}})^{-1}$.

(1) \Rightarrow (2): Using Theorem 4.1 for units in $C_{\mathbb{R}}(\overline{\mathbb{D}})$, we write either F or $-F$ as

$$F = \exp(K),$$

where $K \in C_{\mathbb{R}}(\overline{\mathbb{D}})$. For either f or $-f$ this gives

$$f(z) = \exp(K(z)) \quad (z \in Z_g),$$

and so f (respectively $-f$) is in fact an exponential on the zero set Z_g .

If g is the zero function then f must be invertible in \mathcal{A} , because the pair $(f, 0)$ was assumed to be unimodular. So $(f, 0)$ is reducible in \mathcal{A} , and the unit u is just the function f . Hence we may assume that g is not the zero function, and so the interior of the zero set Z_g is empty by the identity theorem. By Lemma 3.8 the complement $\mathbb{C} \setminus Z_g$ of the zero set is connected. Using Mergelyan's theorem there exist polynomials q_n converging uniformly on Z_g to K . Because K is symmetric, we can also approximate by the symmetrization of q_n , that is, the real polynomials \tilde{q}_n given by

$$\tilde{q}_n(z) := \frac{q_n(z) + (q_n(z^*))^*}{2}$$

converge uniformly to K too. Pick a real polynomial q such that

$$\operatorname{Re}(f(z) \exp(-q(z))) > 1/2 \quad (z \in Z_g).$$

Using the corona theorem, all pairs $(f \exp(-q) - \lambda, g)$ are unimodular for $\lambda \in (-\infty, 1/2)$. For the real inversion set from Theorem 3.3 this gives the inclusion

$$I_{\mathbb{R}}(f \exp(-q), g) = \mathbb{R} \setminus (f \exp(-q))(Z_g) \supset (-\infty, 1/2).$$

Again Theorem 3.3 shows that $(f \exp(-q), g)$ is reducible in $A_{\mathbb{R}}(\overline{\mathbb{D}})$, hence (f, g) , that is, there exists a unit $U \in A_{\mathbb{R}}(\overline{\mathbb{D}})^{-1}$ and $H \in A_{\mathbb{R}}(\overline{\mathbb{D}})$ such that

$$f + Hg = U.$$

U being a unit, we must have

$$|U(z)| > \delta > 0 \quad (z \in \overline{\mathbb{D}}).$$

Now the real polynomials are dense in $A_{\mathbb{R}}(\overline{\mathbb{D}})$. Take real polynomials $h \in \mathcal{A}$ near H such that

$$|U(z) - (H(z) - h(z))g(z)| > \delta/2 > 0 \quad (z \in \overline{\mathbb{D}}).$$

We conclude that

$$f + hg = U - (H - h)g$$

belongs to the algebra \mathcal{A} and has no zeros in $\overline{\mathbb{D}}$ and so it is invertible, proving the reducibility of (f, g) in \mathcal{A} . \square

Theorem 4.3 *Let \mathcal{A} denote a subalgebra of $A_{\mathbb{R}}(\overline{\mathbb{D}})$ containing all real polynomials such that the corona theorem holds for \mathcal{A} . The following assertions are equivalent for any unimodular pair $(f, g) \in \mathcal{A}^2$:*

1. (f, g) is reducible in \mathcal{A} .
2. f is POZ of g .

Proof. (1) \Rightarrow (2): Suppose that (f, g) is reducible in \mathcal{A} . Then there exist $h, u \in \mathcal{A}$, $u \in \mathcal{A}^{-1}$ such that

$$f + h \cdot g = u.$$

By Theorem 4.1 either u or $-u$ can be written as

$$u = \exp(k),$$

for a function $k \in A_{\mathbb{R}}(\overline{\mathbb{D}})$. We arrive at $f + h \cdot g = \exp(k)$, respectively the same equation with $-\exp(k)$ instead of $\exp(k)$. Hence the function f is POZ of g .

(2) \Rightarrow (1): Now assume that f is POZ of g . If g has no zero at all in $\overline{\mathbb{D}}$, then we have $|g(z)| \geq \rho > 0$ for all $z \in \overline{\mathbb{D}}$. But then $M := (1 + \|f\|_{\infty})/\rho$ gives

$$|f(z) + Mg(z)| > 1 + \|f\|_{\infty} - \|f\|_{\infty} = 1,$$

and so $f + Mg$ is invertible in A ; hence the pair (f, g) is reducible. We may also assume that g is not the zero function. Otherwise f itself is invertible and again the pair (f, g) would be reducible. So our assumption is: g has at least one zero in $\overline{\mathbb{D}}$ but is not identically zero. In order to use Theorem 4.2, we must show that there exists a continuous, zero free extension F of f from the zero set Z_g to $\overline{\mathbb{D}}$.

Because (f, g) is unimodular there exists $\delta > 0$ such that $|f(z)| + |g(z)| \geq \delta$ for all $z \in \overline{\mathbb{D}}$. By Lemma 3.9 (with $\delta/2$ instead of δ), there exist finitely many pairwise disjoint closed sets $H_1, \dots, H_N \subset \overline{\mathbb{D}}$ lying symmetrically with respect to the real axis, such that: $Z_g \subset \cup_{j=1}^N H_j$ and $|g(z)| \leq \delta/2$ holds there. Hence $|f(z)| \geq \delta/2$ in the union of this sets. Moreover, we have a continuous logarithm of f on $Z(\delta/2)$: To prove this we quote a theorem of Borsuk, see [3, Corollary 4.33]: Every continuous, zero-free function on $Z(\delta)$ has a continuous logarithm on $Z(\delta)$ if and only if $\mathbb{C} \setminus Z(\delta)$ is connected. This is the case by Lemma 3.8. In particular, there exist functions l_j , continuous in the closed sets $H_j \subset Z(\delta)$, such that

$$f(z) = \exp(l_j(z)), \quad z \in H_j, \quad j = 1, \dots, N.$$

By assertion (3) of Lemma 3.9, if no real zero of g belongs to H_j then $H_j \cap \mathbb{R} = \emptyset$. Moreover, H_j belongs entirely to the upper (respectively lower) half plane. The desired logarithm is very easy to obtain for these sets, because they don't intersect the real line. By symmetry we have $H_j = H_k^*$ for some $j \neq k$. So we may redefine $l_j(z) = (l_k(z^*))^*$.

Thus only the case of a real zero x_0 of g belonging to H_j remains to be discussed. In this case H_j is connected. Since f is POZ of g , we may assume that $f(x_0) > 0$ holds for all real zeros of g . Because f is real symmetric we derive

$$f(z) = \exp(l_j(z)) = \exp((l_j(z^*))^*) \quad (z \in H_j = H_j^*).$$

Since H_j is connected and l_j is continuous in H_j , there exists an integer k such that

$$l_j(z) = (l_j(z^*))^* + 2k\pi i.$$

Restricting to the real zero $x_0 \in H_j \cap \mathbb{R}$ of g gives $\text{Im } l_j(x_0) = k\pi$. As $f(x_0) = \exp(l_j(x_0)) > 0$, the integer k must be even. Now $l_j - k\pi i$ is the desired symmetric logarithm of f on $H_j = H_j^*$.

By Tietze's Theorem we can find a continuous function l on $\overline{\mathbb{D}}$ such that

$$l(z) = l_j(z) \quad z \in H_j, \quad j = 1, \dots, N.$$

The desired logarithm is now given by symmetrization in $\overline{\mathbb{D}}$:

$$L(z) := \frac{l(z) + (l(z^*))^*}{2}.$$

Recall that we either have $l_j(z) = (l_k(z^*))^*$ for $H_j = H_k^*$, $j \neq k$, or else $l_j(z) = (l_j(z^*))^*$ and $H_j = H_j^*$. We end up with $F = \exp(L)$, where we have $\exp(L(z)) = f(z)$ ($z \in Z_g$). \square

5 Units in $A_{\mathbb{R}}(\overline{D})$ and $C_{\mathbb{R}}(\overline{D})$

We recall the notation of the winding number $n(\Gamma; z)$ from [3, Definition 4.2]:

Let Γ denote a closed loop given by a continuous parametrization $t \mapsto \zeta(t)$ ($a \leq t \leq b$), and z denote a point outside Γ . Then there exists a continuous logarithm h of $\zeta(t) - z$, that is

$$\zeta(t) - z = \exp(h(t)), \quad a \leq t \leq b.$$

The winding number $n(\Gamma; z)$ is defined to be

$$n(\Gamma; z) := \frac{h(b) - h(a)}{2\pi i} = \frac{\phi(b) - \phi(a)}{2\pi},$$

where ϕ denotes the imaginary part of the logarithm h . From this definition the following facts are easily seen:

(F1) Let f, g denote zero free continuous functions near the closed loop Γ . Then we can form the closed loops $f(\Gamma)$, $g(\Gamma)$, $(f \cdot g)(\Gamma)$ by their parametrizations $f(\zeta(t))$, $g(\zeta(t))$, $(f \cdot g)(\zeta(t))$, respectively. Since f and g never vanish on Γ , we conclude

$$n((f \cdot g)(\Gamma); 0) = n(f(\Gamma); 0) + n(g(\Gamma); 0).$$

Also $n(\exp(f)(\Gamma); 0) = 0$.

(F2) The curves Γ_{r+m+j} are the reflection of Γ_{r+j} with *reversed* orientation, and so

$$n(f(\Gamma_{r+m+j}); 0) = n(f(\Gamma_{r+j}); 0)$$

holds for all continuous symmetric functions, zero free near Γ_{r+j} .

Theorem 5.1 (Product theorem for units) *Let the notations be as in Notation 3.4. Let \mathcal{A} denote one of the algebras $A_{\mathbb{R}}(\overline{D})$, $C_{\mathbb{R}}(\overline{D})$. For any unit $u \in \mathcal{A}^{-1}$ there exist integers $n_1, \dots, n_r, n_{r+1}, \dots, n_{r+m}$ and a function $h \in \mathcal{A}$ such that the following structure theorem holds: Either u or $-u$ can be factored as*

$$u = p \cdot \exp(h),$$

where $p \in \mathcal{A}^{-1}$, $h \in \mathcal{A}$ and the unit p is given by

$$p(z) = \prod_{j=1}^r (z - x_j)^{n_j} \cdot \prod_{j=1}^m (z^2 - 2(\text{Re}(z_j))z + |z_j|^2)^{n_{r+j}}.$$

Proof. Choose a small compact neighborhood $U \supset \overline{D}$, so small that a continuous extension of u to U has no zeros in U . Each hole of U belongs to exactly one hole of D . The factorization for u in U follows then from [3, Theorem 4.59]. To be precise:

$$u = p \cdot \exp(h),$$

where $p \in \mathcal{A}^{-1}$, h is analytic in $U \supset \overline{D}$, and the unit p is given by

$$p(z) = \prod_{j=1}^r (z - x_j)^{n_j} \prod_{j=1}^m (z - z_j)^{n_{r+j}} \prod_{j=1}^m (z - z_j^*)^{n_{r+m+j}}.$$

With these products and the facts above we can compute the integers n_k ($k = 1, \dots, r + 2m$):

$$n(u(\Gamma_k); 0) = \sum_{j=1}^r n_j \cdot n(\Gamma_k; x_j) + \sum_{j=1}^m n_{r+j} \cdot n(\Gamma_k; z_j) + \sum_{j=1}^m n_{r+m+j} \cdot n(\Gamma_k; z_j^*) + n(\exp(h)(\Gamma_k); 0),$$

that is, $n(u(\Gamma_k); 0) = n_k$ ($k = 1, \dots, r + 2m$). Recall that the curves Γ_{r+m+j} are the reflection of Γ_{r+j} . Observe that reflection has *reversed* orientation. Using the fact F2 and the symmetric choice of our points, we derive

$$n_{r+m+j} = n(u(\Gamma_{r+m+j}); 0) = n(u(\Gamma_{r+j}); 0) = n_{r+j} \quad (j = 1, \dots, m).$$

We conclude that

$$(z - z_j^*)^{n_{r+m+j}} \cdot (z - z_j)^{n_{r+j}} = (z^2 - 2 \cdot \operatorname{Re}(z_j) \cdot z + |z_j|^2)^{n_{r+j}} \quad (j = 1, \dots, m).$$

This proves the product representation. We now show that the logarithm h can be chosen to be symmetric. Using the symmetry of the functions u and p , we conclude that

$$\exp(h(z^*)) = \exp(h(z))$$

holds in the connected set \overline{D} . Thus there is an integer k such that $h(z) = (h(z^*))^* + 2k\pi i$. Hence $\operatorname{Im}(h(x)) = k\pi$ ($x \in \mathbb{R} \cap \overline{D}$). Take a point $x_0 \in \mathbb{R} \cap \overline{D}$, such that $x_0 > \max\{x_1, \dots, x_r\}$. Then $p(x_0) > 0$, and we consider the two cases:

- 1° If $u(x_0) > 0$, then we have $\exp h(x_0) = u(x_0)/p(x_0) > 0$. Since $\operatorname{Im}(h(x_0)) = k\pi$ it follows that k is even. But then $h - k\pi i$ is a symmetric logarithm of u/p .
- 2° If $u(x_0) < 0$, then we just look at the unit $-u$.

□

6 Reducibility in algebras of real symmetric functions

In this section we generalize our technical lemmata to the case of certain finitely connected domains.

Lemma 6.1 *Let D denote a domain as described in Notation 3.4. For every non constant function $g \in A_{\mathbb{R}}(\overline{D})$, the complement $\mathbb{C} \setminus Z(\delta)$ of the level set $Z(\delta)$ is connected for all sufficiently small $\delta > 0$. Moreover, the complement $\mathbb{C} \setminus Z_g$ of the zero set Z_g is also connected.*

Proof. We will first prove that the complement $\mathbb{C} \setminus Z_g$ of the zero set is connected. If g is not identically zero, then by [15, Theorem 3.1], the zero set $Z_g \subset \overline{D}$ is totally disconnected, and so its covering dimension is zero. Hence its open complement $\mathbb{C} \setminus Z_g$ is connected, see [8, Theorem IV.4].

Now we prove that $\mathbb{C} \setminus Z(\delta)$ is connected. We connect each hole C_j by pairwise disjoint cross-cuts $Q_j \subset \overline{D}$, $j = 1, \dots, n$, connecting Γ_j to Γ_{j+1} such that $g(z) \neq 0$ for all $z \in \bigcup_{j=1}^n Q_j$. (This can best be done mapping \overline{D} homeomorphically onto the closure of a circular domain, see for example [11]. Note that even in this situation, the zero set remains totally disconnected.)

Assume that δ is sufficiently small, that is $\delta < \delta_0/2$, where

$$\delta_0 := \min \left\{ |g(z)|, z \in \bigcup_{j=1}^n Q_j \right\}.$$

From Topology we know that $D \setminus \bigcup_{j=1}^n (Q_j \cup C_j)$ is a simply connected domain. The restriction to the choice δ gives

$$Z(\delta) \subset \overline{D} \setminus \bigcup_{j=1}^n (Q_j \cup C_j).$$

If $\mathbb{C} \setminus Z(\delta)$ is not connected, then there exists a bounded component G . In contrast to the simply connected case \mathbb{D} , we may have two cases:

1° $G \subset \overline{D}$ or

2° $G \cap (\mathbb{C} \setminus \overline{D}) \neq \emptyset$, that is, there exists $z_0 \in G$ belonging to a hole, say C : so $z_0 \in C$. But then the hole C is contained in G . Using our cross-cuts we can find a path in $\mathbb{C} \setminus Z(\delta)$ connecting C to the outer boundary Γ_{n+1} of D and beyond. So the starting point of this path belongs to \overline{G} , whereas the endpoint does not. Hence there exists a boundary point $w \in \partial G$ belonging to $\bigcup_{j=1}^n Q_j \subset \overline{D}$. Since $\partial G \subset \partial(\mathbb{C} \setminus Z(\delta)) = \partial Z(\delta) \subset Z(\delta)$ we must have $|g(w)| \leq \delta$, contradicting the choice of $\delta < \frac{\delta_0}{2} = \frac{1}{2} \min\{|g(z)|, z \in \bigcup_{j=1}^n Q_j\}$.

Thus only the first case $G \subset \overline{D}$ remains to be dealt with. But this is done exactly as in the proof of Lemma 3.8. Hence no such bounded component of the complement of $Z(\delta)$ can exist. \square

Lemma 6.2 *Let D denote a domain as described in Notation 3.4 and let $g \in A_{\mathbb{R}}(\overline{D})$ be such that it has at least one zero in \overline{D} , but it does not vanish identically. Then for all $\delta > 0$, there exist finitely many closed sets $H_1, \dots, H_N \subset \overline{D}$, lying symmetrically with respect to the real axis, that is, $H_j = H_k^*$ for certain j, k , with the following properties:*

1. $Z_g \subset \bigcup_{j=1}^N H_j \subset Z(\delta)$.
2. $H_j \cap H_k = \emptyset$ ($j \neq k$).
3. 1° If no real zero of g belongs to H_j , then $H_j \cap \mathbb{R} = \emptyset$, H_j belongs entirely to the upper (respectively lower) half plane, and $H_j = H_k^*$ for some $j \neq k$.
2° If at least one real zero x_0 of g belongs to H_j (that is, $x_0 \in Z_g \cap H_j \cap \mathbb{R}$), then $H_j = H_j^*$ holds and H_j is connected.

Proof. The proof proceeds exactly as in the proof of Lemma 3.9. Note that the components of the level set H are open because $H := \{z \in \overline{D} \mid |g(z)| < \delta\}$ are locally connected by the Jordan curve theorem. We just replace the use of Lemma 3.8 by Lemma 6.1. \square

Theorem 6.3 *Let the notations be as in 3.4 above. Let \mathcal{A} denote a subalgebra of $A_{\mathbb{R}}(\overline{D})$ containing all real polynomials such that the corona theorem holds for \mathcal{A} . Let $(f, g) \in \mathcal{A}^2$ be a unimodular pair. Then the following assertions are equivalent:*

1. There exists a continuous and zero free extension F of f from the zero set Z_g to $F \in C_{\mathbb{R}}(\overline{D})^{-1}$.
2. (f, g) is reducible in \mathcal{A} , that is, there exists a unit $u \in \mathcal{A}^{-1}$ and $h \in \mathcal{A}$ such that $f + hg = u$.

Proof. The implication (2) \Rightarrow (1) is obvious: indeed if there exist $h \in \mathcal{A}$ and a unit $u \in \mathcal{A}^{-1}$ such that $f + hg = u$, then u serves as the desired zero free extension of f from the zero set Z_g to $u \in C_{\mathbb{R}}(\overline{D})^{-1}$.

(1) \Rightarrow (2): Using the factorization theorem 5.1 for units in $C_{\mathbb{R}}(\overline{D})$, we can factor either F or $-F$ as

$$F = p \cdot \exp(K),$$

where $p \in \mathcal{A}^{-1}$ (because \mathcal{A} contains all polynomials and the corona theorem holds) and $K \in C_{\mathbb{R}}(\overline{D})$. For either f or $-f$ this gives

$$\frac{f(z)}{p(z)} = \exp(K(z)) \quad (z \in Z_g),$$

and so the fraction is in fact an exponential on the zero set Z_g . The rest of the proof is now entirely analogous to the corresponding case in the proof of Theorem 4.2. \square

Recall the Notation 3.4: Let D denote a symmetric domain in \mathbb{C} with n holes. From these n holes, bounded by pairwise disjoint Jordan curves, we have r holes intersecting \mathbb{R} and $2m$ which do not intersect \mathbb{R} . Here $n = r + 2m$.

Let C_j be a hole of D ($j = 1, \dots, r, r+1, \dots, r+2m$). Choose for each $j \in \{1, \dots, r\}$ a point $x_j \in \mathbb{R} \cap C_j$, and for $j \in \{r+1, \dots, r+m\}$ let $z_j \in C_j$.

Finally let $S = \{x_1, \dots, x_r, z_1, \dots, z_m, z_1^*, \dots, z_m^*\}$. See Figure 1. For such domains we associate the following family of 2^r polynomials:

$$\mathcal{P} := \left\{ p(z) := \prod_{j=1}^r (z - x_j)^{m_j} \mid m_j \in \{0, 1\} \right\}. \quad (1)$$

As an example, consider an annulus with center at the origin, and let us choose $x_1 = 0$. Then \mathcal{P} has only two polynomials, namely $p_1(z) := 1$ and $p_2(z) := z$.

In [18], Brett Wick showed that a unimodular pair (f, g) in $A_{\mathbb{R}}(\overline{\mathbb{D}})^2$ is reducible in $A_{\mathbb{R}}(\overline{\mathbb{D}})$ if and only if f is POZ of g . We generalize this result from the case of \mathbb{D} to our domains D .

Theorem 6.4 *As in Notation 3.4, let D denote a symmetric domain with n holes, bounded by pairwise disjoint Jordan curves, and let \mathcal{P} denote the associated family of 2^r polynomials (1). Let \mathcal{A} denote a subalgebra of $A_{\mathbb{R}}(\overline{D})$ containing all real polynomials such that the corona theorem holds for \mathcal{A} . The following assertions are equivalent for a unimodular pair $(f, g) \in \mathcal{A}^2$:*

1. (f, g) is reducible in \mathcal{A} .
2. For at least one polynomial $p \in \mathcal{P}$ the product $p \cdot f$ is POZ of g .

Proof. (1) \Rightarrow (2): Suppose that (f, g) is reducible in \mathcal{A} . Then there exist $h, u \in \mathcal{A}$, $u \in \mathcal{A}^{-1}$ such that

$$f + h \cdot g = u.$$

By the factorization theorem for units (Theorem 5.1) in $A_{\mathbb{R}}(\overline{D})$, there exists an invertible polynomial $P \in \mathcal{A}$ such that either u or $-u$ can be factored as

$$u = P \cdot \exp(k),$$

for a function $k \in A_{\mathbb{R}}(\overline{D})$. We arrive at $f + h \cdot g = P \cdot \exp(k)$, respectively the same equation with $-P$ instead of P . Hence the function $P \cdot f$ is POZ of g . But the sign of $P(x)$ depends only on the exponents modulo 2 of its linear factors $x - x_j$. So we can find a polynomial $p \in \mathcal{P}$ such that $p \cdot f$ is POZ of g .

(2) \Rightarrow (1): Now assume that for a polynomial $p \in \mathcal{P}$ the product $p \cdot f$ is POZ of g . Since $p \in \mathcal{A}^{-1}$, it is enough to show that the corona pair (\tilde{f}, g) is reducible, where $\tilde{f} := f/p$. The rest of the proof is entirely analogous to the corresponding case in the proof of Theorem 4.3. \square

Remark 6.5 The family \mathcal{P} obviously depends on the choice of points in the associated set S , hence so does the second assertion in Theorem 6.4, whereas the question of reducibility is independent of the particular choice of S . This mystery can be cleared by Eilenberg's theorem [3, Exercise 4.36, p. 100]:

For any two points a, b belonging to the same hole of D there exists a continuous logarithm $\log \frac{z-a}{z-b}$ which we think of as extended continuously to \mathbb{C} . Thus the product representation of the unit in the proof of Theorem 6.4 won't change much as long as both points belong to the same hole.

The proof of Theorem 2.1 can be generalized to the real Banach algebras $A_{\mathbb{R}}(\overline{D})$, where D is as in Notation 3.4. By Mergelyan's theorem, (real) rational functions are dense in this Banach algebra, and so the proof alters only slightly. So we conclude that the Bass stable rank of these algebras is less than or equal 2. However, our domains D contain an open interval $(a, b) \subset \mathbb{R}$. Consider the unimodular pair $(z - \frac{a+b}{2}, (z-a)(z-b))$. The necessary condition that f is POZ of g is violated, and so this pair is not reducible, hence the stable rank of $A_{\mathbb{R}}(\overline{D})$ is 2. Are there situations where the stable rank of the real algebras is one? Obviously we must allow open sets instead of domains, since there are no symmetric domains such that $D \cap \mathbb{R} = \emptyset$.

Question: For which bounded symmetric open sets $D \subset \mathbb{C}$ do we have $\text{bsr } A_{\mathbb{R}}(\overline{D}) = 1$?

If the stable rank of $A_{\mathbb{R}}(\overline{D})$ is 1, then the open set D necessarily must fulfill the following requirement: $\overline{D} \cap \mathbb{R}$ is either empty or totally disconnected.

If $\overline{D} \cap \mathbb{R}$ is not empty and not totally disconnected, then it contains an open interval $(a, b) \subset \mathbb{R}$. The unimodular pair $(z - \frac{a+b}{2}, (z-a)(z-b))$ is again not reducible by the intermediate value theorem.

Theorem 6.6 *Let $D \subset \mathbb{C}$ be a bounded symmetric open set such that $D^+ := D \cap \{\text{Im}(z) > 0\}$ is a finitely connected domain and $\overline{D} \cap \mathbb{R}$ is either empty or a totally disconnected set of linear measure zero. Then the Bass stable rank of $A_{\mathbb{R}}(\overline{D})$ is 1.*

Proof. Using [5, Theorem 3.11], we see that the complex Banach algebra $A(\overline{D^+})$ has Bass stable rank one, because by Mergelyan's theorem the rational functions with poles off $\overline{D^+}$ are dense in $A(\overline{D^+})$. Again the corona theorem holds in $A(\overline{D})$ by Arens's theorem, hence in $A_{\mathbb{R}}(\overline{D})$, see for example [6, Theorem 1.9, p. 31]. Let (f, g) be a unimodular pair in $A_{\mathbb{R}}(\overline{D})$. If $\overline{D} \cap \mathbb{R} = \emptyset$, we proceed as follows: the unimodular pair (f, g) is reducible in $A(\overline{D^+})$, that is, there exist $u, h \in A(\overline{D^+})$, u invertible in $A(\overline{D^+})$ such that $f + hg = u$. Now we reflect u, h to the lower half plane. Since $\overline{D} \cap \mathbb{R} = \emptyset$ this reflection is well-defined, hence (f, g) is reducible in $A_{\mathbb{R}}(\overline{D})$.

So we may assume that $\overline{D} \cap \mathbb{R}$ is not empty and is a totally disconnected set of linear measure zero.

1° $f(x) \neq 0$ for all $x \in \overline{D} \cap \mathbb{R}$.

Take a peak-function p in the upper half plane for the totally disconnected set $\overline{D} \cap \mathbb{R}$ of linear measure zero. The existence of such a peak function follows from Rudin's theorem (see for instance [7, p. 81]) by conformally mapping the unit disc onto the upper half plane. Then the function $q := 1 - p \in A(\overline{D^+})$ vanishes at z_0 if and only if $z_0 \in \overline{D} \cap \mathbb{R}$. The corona theorem for $A(\overline{D^+})$ and 1° now implies that the pair $(f, q \cdot g)$ is unimodular in $A(\overline{D^+})$, and so it is reducible in $A(\overline{D^+})$. Thus there exists $h \in A(\overline{D^+})$ such that $f + h \cdot (qg)$ has no zeros in $\overline{D^+}$. We define the function $H \in A_{\mathbb{R}}(\overline{D})$ by reflection: $H(z) := h(z)q(z)$ in case $z \in \overline{D^+}$ and $H(z) := (h(z^*)q(z^*))^*$ in case $z \in \overline{D} \setminus \overline{D^+}$. Since q vanishes identically on $\overline{D} \cap \mathbb{R}$ this reflection is well-defined. We conclude that $f + H \cdot g$ has no zeros in \overline{D} and so it is invertible by the corona theorem. Therefore the pair (f, g) is reducible in $A_{\mathbb{R}}(\overline{D})$.

2° $f(x) = 0$ for some $x \in \overline{D} \cap \mathbb{R}$.

We approximate f uniformly by real rational functions f_n on \overline{D} , and again use Mergelyan's theorem and symmetry. Since $\overline{D} \cap \mathbb{R}$ is totally disconnected, we can perturb the finitely many zeros of f_n slightly (respecting symmetry) such that f_n has no zeros in $\overline{D} \cap \mathbb{R}$. Using 1° we see that the pairs (f_n, g) are reducible in $A_{\mathbb{R}}(\overline{D})$, and so (f, g) is reducible in $A_{\mathbb{R}}(\overline{D})$, see for example [5, Lemma 3.7].

□

Acknowledgement: The authors thank the referees for their numerous comments which improved the exposition of the paper.

References

- [1] L.V. Ahlfors, Complex Analysis, 3rd Edition (McGraw-Hill, New York, 1979).
- [2] H. Bass, K-theory and stable algebra, Publications Mathématiques de L'I.H.É.S., **22**, 5–60 (1964).
- [3] R.B. Burckel, An Introduction to Classical Complex Analysis, Vol. 1 (Birkhäuser, New York-London, 1979).
- [4] G. Corach and A.R. Larotonda, Stable range in Banach algebras, Journal of Pure and Applied Algebra, **32**, 289–300 (1984).
- [5] G. Corach and F.D. Suárez, Stable range in holomorphic function algebras, Illinois Journal of Mathematics, **29**, 627–639 (1985).
- [6] T.W. Gamelin, Uniform Algebras (Prentice-Hall, N.J., 1969).
- [7] K. Hoffman, Banach Spaces of Analytic Functions (Dover, New York, 1962).
- [8] W. Hurewicz and H.W. Wallman, Dimension Theory (Princeton University Press, Princeton, N.J., 1941).
- [9] P.W. Jones, D. Marshall and T. Wolff, Stable rank of the disc algebra, Proceeding of the American Mathematical Society, no. 4, **96**, 603–604 (1986).
- [10] R. Mortini and R. Rupp, A constructive proof of the Nullstellensatz for subalgebras of $A(K)$, Séminaire de mathématique de Luxembourg, Trav. Math., **3**, 45–49 (1991).
- [11] Chr. Pommerenke, Univalent Functions (Vandenhoeck and Ruprecht, Göttingen, 1975).
- [12] A. Quadrat, On a general structure of the stabilizing controllers based on stable range, SIAM Journal on Control and Optimization, no. 6, **42**, 2264–2285 (2004).
- [13] M.A. Rieffel, Dimension and stable rank in the K-theory of C^* -algebras, Proceedings of the London Mathematical Society, no.2, **46**, 301–333 (1983).
- [14] R. Rupp, Stable rank in Banach algebras, Function spaces, Lecture Notes in Pure and Applied Mathematics, **136**, 357–365 (Dekker, New York, 1992).
- [15] R. Rupp, Stable rank of subalgebras of the disc algebra, Proceedings of the American Mathematical Society, **108**, 137–142 (1990).
- [16] R. Rupp, Stable rank and the $\bar{\partial}$ -equation, Canadian Mathematical Bulletin, **34**, 113–118 (1991).
- [17] M. Vidyasagar, Control System Synthesis: A Factorization Approach, MIT Press Series in Signal Processing, Optimization, and Control, **7** (MIT Press, Cambridge, MA, 1985).
- [18] B. Wick, A note about stabilization in the real disc algebra, To appear in Mathematische Nachrichten (2007).