

Mario Petrich

On the structure of a class of commutative semigroups

Czechoslovak Mathematical Journal, Vol. 14 (1964), No. 1, 147–153

Persistent URL: <http://dml.cz/dmlcz/100607>

Terms of use:

© Institute of Mathematics AS CR, 1964

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE STRUCTURE OF A CLASS OF COMMUTATIVE SEMIGROUPS

MARIO PETRICH, Maryland (USA)

(Received March 18, 1963)

The purpose of the paper is to clarify the structure of a special type of commutative semigroups to which several authors have been led by studying decompositions of general semigroups.

1. Introduction and summary. In this paper we investigate the structure of a class of semigroups which we call N -semigroups. An N -semigroup is a commutative non-potent archimedean cancellative semigroup. E. HEWITT and H. S. ZUCKERMAN [2] have shown that if G is any commutative semigroup, then there is a maximal separative homomorphic image G' of G , and that a member H_x of the maximal semilattice decomposition of G' is either a group or an N -semigroup. T. TAMURA [5] has given a characterization of N -semigroups. T. TAMURA and N. KIMURA [4] have shown that a member of the maximal semilattice decomposition of an arbitrary commutative semigroup is archimedean and has at most one idempotent. Š. SCHWARZ [6] has established certain properties of decompositions of a semigroup similar to those already mentioned.

In section 2 we define N -semigroups and discuss some properties of commutative semigroups in connection with it. Then in section 3 we establish a property of N -semigroups with a finite number of generators. In section 4 we find the structure of N -semigroups with two generators. Finally in section 5 we give a classification and several examples of N -semigroups.

A semigroup is a non-empty set on which an associative multiplication is defined. We will discuss only commutative semigroups. Throughout the whole paper S will denote an arbitrary commutative semigroup unless stated otherwise. We follow the notation and terminology of A. H. CLIFFORD and G. B. PRESTON [1] for all concepts not defined in the paper. By $x^m y^n$ with $m = 0$ and $n > 0$, we mean y^n .

The writer wishes to thank Professor *Edwin Hewitt* for mentioning this problem to him, and Professor *Herbert S. Zuckerman* for his help in preparation of this paper.

2. Definitions and properties. We first define an N -semigroup and then discuss some properties of semigroups in connection with it.

Definition. S will be called an N -semigroup if it has the following properties:

(A) for every $x, y, z \in S$, $xz = yz$ implies $x = y$ (S is cancellative);
 (B) for every $x, y \in S$, $x^n = ay$ for some $a \in S$ and some natural number n (S is archimedean);

(C) S has no idempotents (S is nonpotent).

Proposition 1. *If S satisfies condition (B) and*

(D) *for every $x, y \in S$, $x^2 = y^2 = xy$ implies $x = y$ (S is separative), then S also satisfies (A).*

Proof. Let S satisfy conditions (B) and (D) and suppose that $xz = yz$ for some $x, y, z \in S$. Then $x^m = az$ and $y^n = bz$ for some $a, b \in S$ and some m, n . Hence

$$(1) \quad x^{m+1} = (az)x = a(xz) = a(yz) = (az)y = x^m y,$$

$$(2) \quad y^{n+1} = (bz)y = b(yz) = b(xz) = (bz)x = y^n x.$$

If in (1) $m > 1$, then

$$x^{2m-2}xy = x^{m-2}x^{m+1}y = x^{m-2}x^m y^2 = x^{2m-2}y^2,$$

$$x^{2m-2}xy = x^{m-1}x^m y = x^{m-1}x^{m+1} = x^{2m}.$$

Consequently $(x^{m-1}y)^2 = (x^m)^2 = x^m(x^{m-1}y)$ and thus $x^m = x^{m-1}y$. After $m - 1$ steps, we obtain $x^2 = xy$. Similarly from (2), we obtain $y^2 = xy$ and therefore $x = y$.

Corollary. *In the definition of an N -semigroup, we can substitute condition (A) by the weaker condition (D).*

The proofs of the following statements are either contained in the works mentioned at the beginning of the paper or in [3].

Proposition 2. *If S satisfies (B), then it contains at most one idempotent.*

Proposition 3. *If S satisfies (A) and (B) and does not satisfy (C), then it is a group.*

Theorem 1. *S contains no prime (proper semiprime) ideals if and only if S satisfies (B).*

Theorem 2. *Each member of the maximal semilattice decomposition of S satisfies (B).*

Theorem 3. *S satisfies (D) if and only if S is a semilattice of semigroups each of which satisfies (A).*

3. Finitely generated N -semigroups. The set of positive integers under addition is an N -semigroup generated by the element 1. It is evident that this is the only cyclic N -semigroup. The following theorem establishes a property of N -semigroups with a finite number of generators.

Theorem 4. A finitely generated N -semigroup S satisfies the following condition:

(E) for every $x, y \in S$, there are natural numbers p and q such that $x^p = y^q$.

Proof. Let a_1, a_2, \dots, a_n be the set of generators of S . We first show that condition (E) holds for a_1, a_2, \dots, a_n . We do this by mathematical induction on the number k defined as follows. A power of a fixed generator can be written as a product of powers of any $n - k$ of the remaining generators with the power of any specified generator positive, $1 \leq k < n$.

The proof for $k = 1$. Consider the generator a_1 ; the other cases are similar. For $m > 1$, we have

$$a_1^t = (a_1^{t_1} a_2^{t_2} \dots a_n^{t_n})^m$$

for some t, t_1, t_2, \dots, t_n with $t > 1$ and $\sum_{i=1}^n t_i > 0$. Here $t > t_1$, for otherwise we would arrive at a contradiction after cancellation. Hence

$$a_1^{t-t_1} = a_2^{t_2} \dots a_m^{t_m+1} \dots a_n^{t_n}.$$

Suppose now that the condition stated at the beginning of the proof is satisfied for some $k, 1 \leq k < n$. We again consider only the case of the generator a_1 , the other cases being similar. We show that the condition in question is also valid for $k + 1$. By hypothesis we have

$$(1) \quad \begin{aligned} a_1^p &= a_{k+1}^{p_{k+1}} a_{k+2}^{p_{k+2}} \dots a_n^{p_n}, \\ &\dots \\ a_{k+1}^q &= a_1^{q_1} a_{k+2}^{q_{k+2}} \dots a_n^{q_n} \end{aligned}$$

where $p_n > 0$. We obtain

$$a_1^{pq} = a_{k+1}^{p_{k+1}q} a_{k+2}^{p_{k+2}q} \dots a_n^{p_nq} = a_1^{q_1 p_{k+1}} a_{k+2}^{q_{k+2} p_{k+1}} \dots a_n^{q_n p_{k+1}} a_{k+2}^{p_{k+2}q} \dots a_n^{p_nq}$$

whence

$$a_1^{pq - q_1 p_{k+1}} = a_{k+2}^{q_{k+2} p_{k+1} + p_{k+2}q} \dots a_n^{q_n p_{k+1} + p_nq}$$

since necessarily $pq > q_1 p_{k+1}$ and also $p_n q > 0$. The general case is proved by considering in (1) any $n - k$ generators different from a_1 which merely amounts to a change of notation. This concludes the proof of induction.

We have in particular $a_1^{m_i} = a_i^{s_i}$ for $i = 2, 3, \dots, n$ where $m_i, s_i > 1$. Let $a_1^{k_1} a_2^{k_2} \dots a_n^{k_n}, a_1^{l_1} a_2^{l_2} \dots a_n^{l_n} \in S$ be arbitrary. Then

$$\begin{aligned} &(a_1^{k_1} a_2^{k_2} \dots a_n^{k_n})^{s_2 s_3 \dots s_n (l_1 s_2 s_3 \dots s_n + m_2 l_2 s_3 \dots s_n + \dots + m_n s_2 s_3 \dots s_{n-1} l_n)} = \\ &= a_1^{(k_1 s_2 s_3 \dots s_n + m_2 k_2 s_3 \dots s_n + \dots + m_n s_2 s_3 \dots s_{n-1} k_n) (l_1 s_2 s_3 \dots s_n + m_2 l_2 s_3 \dots s_n + \dots + m_n s_2 s_3 \dots s_{n-1} l_n)} = \\ &= (a_1^{l_1} a_2^{l_2} \dots a_n^{l_n})^{s_2 s_3 \dots s_n (k_1 s_2 s_3 \dots s_n + m_2 k_2 s_3 \dots s_n + \dots + m_n s_2 s_3 \dots s_{n-1} k_n)}, \end{aligned}$$

which completes the proof.

4. N -semigroups with two generators. In this section we find the structure and give a concrete realization of N -semigroups with two generators. We do not consider N -semigroups with more than two generators because the characterization given in this paper becomes too involved in such a case. We first introduce some notations.

Notation. Let S be an N -semigroup with two generators, say a_1 and a_2 . Let m_1 and m_2 be the smallest positive integers such that $a_1^{m_1} = ua_2$ and $a_2^{m_2} = va_1$ for some $u, v \in S$. We suppose that $m_1 \leq m_2$ and denote S by $N(m_1, m_2)$.

Since S is cancellative, a_2 is a generator of S , and m_1 is minimal, we have $u = a_2^{k_2}$ for some $k_2 > 0$, and thus $a_1^{m_1} = a_2^{k_2+1}$. Similarly $a_2^{m_2} = a_1^{k_1+1}$ for some $k_1 > 0$. By minimality of m_1 and m_2 , $m_1 \leq k_1 + 1$ and $m_2 \leq k_2 + 1$. If $k_2 + 1 > m_2$, then

$$a_1^{m_1} = a_2^{k_2+1} = a_2^{m_2} a_2^{k_2+1-m_2} = a_1^{k_1+1} a_2^{k_2+1-m_2}$$

which is impossible since $m_1 \leq k_1 + 1$. Thus $m_2 = k_2 + 1$, that is, $a_1^{m_1} = a_2^{m_2}$.

Notation. Let m_1 and m_2 be integers such that $2 \leq m_1 \leq m_2$. A set S will be denoted by (m_1, m_2) -s.g. if

$$S = \{(k_1, k_2) \mid k_1 = 0, 1, 2, \dots, \quad k_2 = 0, 1, 2, \dots, m_2 - 1, \quad k_1 + k_2 > 0\}$$

with multiplication

$$(k_1, k_2)(l_1, l_2) = (k_1 + l_1 + jm_1, k_2 + l_2 - jm_2)$$

where j is the integer such that $0 \leq k_2 + l_2 - jm_2 < m_2$, and $(k_1, k_2) = (l_1, l_2)$ implies $k_1 = l_1$ and $k_2 = l_2$.

The following theorem gives a simple characterization of N -semigroups with two generators:

Theorem 5. *Let S be a set. Then $S = N(m_1, m_2)$ if and only if $S = (m_1, m_2)$ -s.g.*

Proof. We first prove *necessity*. Thus let $S = N(m_1, m_2)$ with the generators a_1 and a_2 such that $a_1^{m_1} = a_2^{m_2}$. If x is an element of S , then $x = a_1^{k_1} a_2^{k_2}$ for some non-negative integers k_1 and k_2 such that $k_1 + k_2 > 0$. We have $0 \leq k_2 - jm_2 < m_2$ for some non-negative integer j , and hence

$$a_1^{k_1} a_2^{k_2} = a_1^{k_1} a_2^{jm_2 + (k_2 - jm_2)} = a_1^{k_1 + jm_1} a_2^{k_2 - jm_2}.$$

Thus every element of S can be written in the form of an element of (m_1, m_2) -s.g. with a suitable change of notation. One checks similarly that the multiplication of S coincides with that of (m_1, m_2) -s.g. under the restriction that $0 \leq k_2 < m_2$ where $a_1^{k_1} a_2^{k_2} \in S$. Suppose that $a_1^{k_1} a_2^{k_2} = a_1^{l_1} a_2^{l_2}$ with $0 \leq k_2, l_2 < m_2$. If $k_1 > l_1$, then $a_1^{k_1 - l_1} a_2^{k_2} = a_2^{l_2}$ and thus necessarily $k_2 < l_2$. Consequently $a_1^{k_1 - l_1} = a_2^{l_2 - k_2}$ and thus $l_2 - k_2 \geq m_2$ by minimality of m_2 . But this contradicts the hypothesis that $l_2 < m_2$. The case $k_1 < l_1$ is symmetric. Hence $k_1 = l_1$ and consequently $k_2 = l_2$. Therefore $S = (m_1, m_2)$ -s.g.

We next prove *sufficiency*. Let $S = (m_1, m_2)$ -s.g. It is clear that S is closed under its multiplication and is commutative. We verify the postulates for $N(m_1, m_2)$.

Associativity. It is easily seen that both $[(k_1, k_2)(l_1, l_2)](r_1, r_2)$ and $(k_1, k_2) \cdot [(l_1, l_2)(r_1, r_2)]$ are equal to $(k_1 + l_1 + r_1 + im_1, k_2 + l_2 + r_2 - im_2)$ where i is the integer such that $0 \leq k_2 + l_2 + r_2 - im_2 < m_2$. Hence the associative law holds.

Condition (A). If $(k_1, k_2)(r_1, r_2) = (l_1, l_2)(r_1, r_2)$, then

$$(k_1 + r_1 + im_1, k_2 + r_2 - im_2) = (l_1 + r_1 + jm_1, l_2 + r_2 - jm_2)$$

where i and j are the integers such that $0 \leq k_2 + r_2 - im_2 < m_2$ and $0 \leq l_2 + r_2 - jm_2 < m_2$. Hence

$$(1) \quad k_1 + r_1 + im_1 = l_1 + r_1 + jm_1,$$

$$(2) \quad k_2 + r_2 - im_2 = l_2 + r_2 - jm_2.$$

From (2) we obtain $k_2 - l_2 = (i - j)m_2$ which implies that $i = j$ and consequently $k_2 = l_2$. But $i = j$ in (1) yields $k_1 = l_1$. Therefore $(k_1, k_2) = (l_1, l_2)$ and the cancellation law holds in S .

Condition (B). Now let (k_1, k_2) and (l_1, l_2) be any elements of S . If $k_1 > 0$, then let $n = m_1 + l_1 + 1$ and q be the non-negative integer such that $0 \leq nk_2 - m_2q < m_2$. If $k_1 = 0$, then let $q = m_1 + l_1 + 1$ and n be the integer satisfying the inequality $m_2q/k_2 \leq n < (m_2q/k_2) + 1$ (in this case $k_2 > 0$). Moreover let

$$j = \begin{cases} 1 & \text{if } nk_2 - m_2q - l_2 < 0, \\ 0 & \text{if } nk_2 - m_2q - l_2 \geq 0. \end{cases}$$

It follows easily from the definitions of n , q , and j that

$$(a) \quad nk_1 + m_1q - l_1 - jm_1 > (m_1 + l_1) - l_1 - jm_1 \geq 0;$$

$$(b) \quad nk_1 + m_1q > 0;$$

$$(c) \quad 0 \leq nk_2 - m_2q < m_2;$$

$$(d) \quad 0 \leq nk_2 - m_2q - l_2 + jm_2 < m_2.$$

We therefore have

$$\begin{aligned} (nk_1 + m_1q - l_1 - jm_1, nk_2 - m_2q - l_2 + jm_2)(l_1, l_2) &= \\ &= (nk_1 + m_1q, nk_2 - m_2q) = (k_1, k_2)^n. \end{aligned}$$

Condition (C). Suppose now that $(k_1, k_2)^2 = (k_1, k_2)$ for some $(k_1, k_2) \in S$. Then

$$(2k_1 + jm_1, 2k_2 - jm_2) = (k_1, k_2)$$

where j is the integer such that $0 \leq 2k_2 - jm_2 < m_2$. Hence $2k_1 + jm_1 = k_1$ and $2k_2 - jm_2 = k_2$. Consequently $k_2 - jm_2 = 0$ which implies that $j = 0$ and hence also $k_2 = 0$. But $j = 0$ in the first equation yields $k_1 = 0$, which is impossible. Therefore S has no idempotents.

$S = N(m_1, m_2)$. We have already proved that S is an N -semigroup. It is clear that $(1, 0)$ and $(0, 1)$ are the generators of S . Let n be the smallest positive integer such that $(1, 0)^n = (k_1, k_2)(0, 1)$ for some $(k_1, k_2) \in S$. Then $k_1 = 0$ by minimality of n . Hence $(1, 0)^n = (0, k_2)(0, 1)$ and thus $(n, 0) = (jm_1, k_2 + 1 - jm_2)$ where j is the integer such that $0 \leq k_2 + 1 - jm_2 < m_2$. Therefore $n = jm_1$ and $0 = k_2 + 1 - jm_2$. Consequently $j = 1$ and thus $n = m_1$ and $k_2 = m_2 - 1$.

Similarly we see that if n is the smallest positive integer such that $(0, 1)^n = (k_1, k_2)(1, 0)$ for some $(k_1, k_2) \in S$, then $n = m_2$. Since $m_1 \leq m_2$, we have proved that $S = N(m_1, m_2)$.

Remark. From the definition of (m_1, m_2) -s.g. it follows easily that (m_1, m_2) -s.g. is isomorphic to the free commutative semigroup on two generators, say a_1 and a_2 , with the defining relation $a_1^{m_1} = a_2^{m_2}$. This furnishes a second characterization of $N(m_1, m_2)$ by virtue of Theorem 5.

We next give a concrete realization of (m_1, m_2) -s.g. First we introduce some notation.

Notation. For integers m_1 and m_2 such that $2 \leq m_1 \leq m_2$, let $C(m_1, m_2)$ be the subsemigroup of the group of non-zero complex numbers, generated by the two elements

$$a_1 = 2^{1/m_1} e^{(2\pi i)/m_1} \quad \text{and} \quad a_2 = 2^{1/m_2} e^{(4\pi i)/m_2}.$$

Then we have the following result:

Theorem 6. *The semigroups $C(m_1, m_2)$ and (m_1, m_2) -s.g. are isomorphic.*

Proof. First note that $a_1 \neq a_2$ because $2 \leq m_1 \leq m_2$. Since $C(m_1, m_2)$ is a sub-semigroup of a commutative group (non-zero complex numbers), by the remark above it suffices to show that $a_1^{k_1} = a_2^{k_2}$ implies $k_1/k_2 = m_1/m_2$ and $k_1 \geq m_1$. If $a_1^{k_1} = a_2^{k_2}$, that is,

$$2^{k_1/m_1} e^{[(2\pi i)/(m_1)]k_1} = 2^{k_2/m_2} e^{[(4\pi i)/(m_2)]k_2},$$

then by equating moduli, we obtain $k_1/m_1 = k_2/m_2$. But then

$$e^{2\pi i[2(k_1/m_1) - (k_1/m_1)]} = 1$$


and consequently $k_1 \geq m_1$. The theorem follows.

The following is essentially a resume of some results of this section:

Theorem 7. *The semigroup $C(m_1, m_2)$ is an N -semigroup with two-generators and $C(m_1, m_2) = C(m'_1, m'_2)$ only if $m_1 = m'_1$ and $m_2 = m'_2$. Conversely, every N -semigroup with two generators is isomorphic to the semigroup $C(m_1, m_2)$ for some (unique) integers m_1 and m_2 , $2 \leq m_1 \leq m_2$.*

5. Classification and examples of N -semigroups. We classify N -semigroups according to whether they satisfy condition (E) of Theorem 4, whether they are finitely generated, and whether they contain indecomposable elements. We say that an element x of S is indecomposable if $x \neq yz$ for all $y, z \in S$. We also give an example

for each group of N -semigroups. All the examples given are subsemigroups of the additive semigroup of positive real numbers. The numbers in brackets denote the generators of S ; the letter a denotes any positive real number.



References

- [1] Clifford A. H. and G. B. Preston: The Algebraic Theory of Semigroups. Vol. I. American Math. Society, 1961.
- [2] Hewitt Edwin and H. S. Zuckerman: The l_1 -algebra of a commutative semigroup. Trans. A.M.S., 83 (1956), 70—97.
- [3] Petrich Mario: The greatest semilattice decomposition of a semigroup (to appear).
- [4] Tamura T. and N. Kimura: On decompositions of a commutative semigroup. Kōdai Math. Sem. Rep., 4 (1954), 109—112.
- [5] Tamura T.: Commutative nonpotent archimedean semigroup with cancellation law I. J. Gakugei Tokushima Univ., 8 (1957), 5—11.
- [6] Schwarz Štefan: O pologrupách splňujúcich zoslabené pravidla krátania. Matematicko-Fyzikálny Časopis, S.A.V. 6 (1956), 149—158.

Резюме

О СТРОЕНИИ ОПРЕДЕЛЕННОГО КЛАССА КОММУТАТИВНЫХ ПОЛУГРУПП

МАРИО ПЕТРИХ (Mario Petrich), Мариланд (США)

Коммутативная полугруппа S называется N -полугруппой, если 1. В S имеет место правило сокращения; 2. для каждой пары $x, y \in S$ существует $a \in S$ и целое число $n > 0$ так, что $x^n = ay$; 3. S не имеет идемпотентов. Работа посвящена изучению строения N -полугрупп. Именно, описана структура всех N -полугрупп, обладающих двумя генераторами.