


On the Sulfation Pattern of Polysaccharides in the Extracellular Matrix of Sheep with Chondrodysplasia

Cartilage
2(1) 36–39
© The Author(s) 2011
Reprints and permission:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/1947603510377465
http://cart.sagepub.com


S.A. Piripi¹, M.A.K. Williams^{2,3}, and K.G. Thompson¹

Abstract

Objective: Chondroitin sulfate is the major sulfated polysaccharide attached to the core protein, aggrecan, in the hyaline cartilage matrix. Sulfation of the cartilage matrix polysaccharide is vital for normal matrix integrity and compressive stiffness of the tissue and is therefore crucial to normal cartilage formation and consequently to endochondral ossification. Several forms of chondrodysplasia, a condition resulting in clear macroscopic deficiencies in the mechanical properties of the cartilage and characterized by reduced levels of sulfate, have been identified in both human beings and animals. **Design:** In this study, the authors used capillary electrophoresis to investigate the sulfation state of extracted chondroitin sulfate polymers. **Results:** Significantly, cartilage from affected sheep had a lower ratio of the chondroitin-derived enzymatically liberated disaccharides Δ di-mono4S to Δ di-mono6S, demonstrating reduced levels of chondroitin 4-sulfate, but not chondroitin 6-sulfate, in chondrodysplastic sheep compared to age-matched controls at all ages measured. **Conclusion:** This supports the hypothesis that a difference in chondroitin sulfate disaccharides is detectable in affected newborn lambs prior to the development of lesions.

Keywords

cartilage sulfation, capillary electrophoresis, chondrodysplasia

Animals Used and Samples Collected

Several forms of chondrodysplasia, a condition resulting in clear macroscopic deficiencies in the mechanical properties of the cartilage and characterized by reduced levels of sulfate, have been identified in both human beings and animals.^{1–7} In this work an entire humerus was collected from sheep that had undergone postmortem examination and was stored at -20°C for 6 to 24 months. Near-full-thickness articular cartilage shavings (10–15 mg) were harvested from the central humeral heads of 8 animals of various ages. Due to the age-related differences in articular cartilage thickness and composition, sulfation profiles are expected to vary with age of the animal: an issue that we attempted to ameliorate by using age-matched controls. Samples 1 and 2 were from obligate affected newborn lambs (the form of chondrodysplasia investigated in this study is a heritable disorder with a simple autosomal recessive mode of inheritance, so although there are no macroscopic or microscopic lesions identifiable in newborn lambs, lambs with both parents affected by the disorder were considered to be obligately affected), samples 3 and 4 were from control newborn lambs, sample 5 was from a 1-month-old lamb with severe chondrodysplasia, sample 6 was from a 1-month-old control lamb, sample 7 was from a mature chondrodysplastic ram, and sample 8 was from a mature control sheep. The animals used as controls were

not taken from the same back-cross breeding trial that produced the affected animals for the study. This was because although lambs with both parents affected could be considered obligately affected, there was no way to confidently determine nonaffected newborn lambs. The control animals were genetically unrelated but age matched and from the same farm. Although unfortunately the breed was not recorded in this instance, the results reported herein are highly suggestive and should stimulate further work.

Chondroitin Disaccharide Extraction

The method for extraction of chondroitin disaccharides from articular cartilage was adapted from similar previous studies.^{8,9} Briefly, the glycosidic bond linking sulfated glycosaminoglycans to aggrecan was cleaved by incubation

¹Institute of Veterinary, Animal & Biomedical Sciences, Massey University, Palmerston North, New Zealand

²Institute of Fundamental Sciences, Massey University, Palmerston North, New Zealand

³MacDiarmid Institute for Nanotechnology and Advanced Materials, Wellington, New Zealand

Corresponding Author:

Martin Williams, Private Bag 11 222 Palmerston North, 4442
New Zealand
Email: m.williams@massey.ac.nz

with 100 μL of 0.5M NaOH per mg of sample at 4°C, with agitation overnight. The NaOH was then removed by minidialysis against distilled water in 3.5-kDa molecular weight cutoff dialysis cassettes (Slide-A-Lyzer, Pierce Biotechnology, Rockford, IL) for 48 hours at 4°C. Samples were then centrifuged at 13,000 g for 3 minutes and the supernatant freeze-dried and then rehydrated in 25 μL deionized water. In order to digest the extracted polysaccharides into disaccharides, 10 μL of resuspended supernatant was incubated with 10 μL chondroitinase ABC (from *Proteus vulgaris*, Sigma-Aldrich Corp., St. Louis, MO; 0.05 mU/ μL in 200 mM Tris base at pH 7.3) for 1 hour at 37°C. Digested samples were filtered through 3-kDa nominal molecular weight limit centrifugal filter units (Microcon ultracel YM-3, Amicon Bioseparations, Millipore, Billerica, MA) at 13,000 g for 90 minutes, and the filtrate containing the disaccharides was further diluted with 20 μL distilled water.

Capillary Electrophoresis

Experiments were performed using an automated capillary electrophoresis (CE) system (HP 3D), equipped with a diode array detector, based on a previously reported method.^{10,11} Briefly, electrophoresis was carried out at reverse polarity and 25°C in a fused silica capillary of internal diameter of 50 μm and a total length of 46.5 cm. Phosphate buffer (15 mM) at pH 3.0 was used as a CE background electrolyte, with UV detection of the unsaturated bond resulting from enzymatic digestion absorbing at 232 nm. Samples were loaded hydrodynamically at 5,000 Pa for 10 seconds and typically electrophoresed across a potential difference of 20 kV. Standard samples of 2-acetamido-2-deoxy-3-O-(4-deoxy- α -L-threo-hex-4-enopyranosyluronic acid)-4-O-sulfo-D-galactose (known as chondroitin 4-sulfate disaccharide, chondroitin sulfate A disaccharide, or $\Delta\text{di-mono4S}$) and 2-acetamido-2-deoxy-3-O-(4-deoxy- α -L-threo-hex-4-enopyranosyluronic acid)-6-O-sulfo-D-galactose (known as chondroitin 6-sulfate disaccharide, chondroitin sulfate C or $\Delta\text{di-mono6S}$) (Sigma-Aldrich Corp.) were used to identify ionic species and to construct standard curves for quantification. All samples were run at least 4 times and the peaks integrated using HP Chemstation software (Agilent Technologies, Santa Clara, CA). Where a particular disaccharide exhibited a double peak, both anomeric forms were pooled. Normalized¹² peak areas were subsequently used, via a calibration of absorptivity using the standard samples, to produce concentrations in mg/L.

Analysis

In order to compensate for slight differences in injection volume, the comparison of the results from the different




Figure 1. Typical electropherogram of chondroitin disaccharides from ovine articular cartilage. **(A)** Double peaks result from anomeric forms of hexosamine. **(B)** Different ages of background buffer alter the resolution of these anomeric forms. 6S indicates $\Delta\text{di-mono6S}$, 4S indicates $\Delta\text{di-mono4S}$, and OS represents unsulfated chondroitin disaccharide.

samples was carried out by taking the concentration of $\Delta\text{di-mono4S}$ and dividing by the concentration of $\Delta\text{di-mono6S}$, giving a ratio for each sample. Statistical analyses were performed using S-Plus 8.0 student edition (Insightful Corp., Seattle, WA, 2007). The analysis of variance (ANOVA) model was as follows: ratio of $\Delta\text{di-mono4S}$ to $\Delta\text{di-mono6S}$ \sim Age + Status, where Age was the age of the animal from which cartilage was taken, and Status was the presence or absence of chondrodysplasia. The Student t test was used to compare individual animals.

A typical electropherogram is shown in **Figure 1**, demonstrating peaks caused by $\Delta\text{di-mono4S}$, $\Delta\text{di-mono6S}$, and a peak at the expected location for unsulfated chondroitin disaccharide, as demonstrated by Karamanos *et al.*¹¹ Double peaks were routinely observed, owing to the different anomeric forms of the hexosamine present at the reducing terminal of the disaccharide. Ten milligrams of each sample was run and mean concentrations of disaccharide for each of the tested animals are listed in **Table 1**, expressed as ng/10-mg sample. Concentrations of $\Delta\text{di-mono4S}$ were markedly lower in cartilage from chondrodysplastic sheep of all ages compared with controls, but there was no such difference in $\Delta\text{di-mono6S}$ concentrations. The ratios of $\Delta\text{di-mono4S}$ to $\Delta\text{di-mono6S}$ were also lower in chondrodysplastic sheep than in unrelated control animals in all age groups ($P \leq 0.0001$; **Fig. 2**). In addition, the ratio of $\Delta\text{di-mono4S}$ to $\Delta\text{di-mono6S}$ was markedly lower in the mature control than in the younger controls and in the mature affected ram than in the younger affected lambs

Table 1. Concentrations of Chondroitin Sulfate Disaccharides from Chondrodysplastic and Control Texel Sheep of Varying Ages

Animal	Status	Age	Δ di-mono6S	Δ di-mono4S
1	Affected	Newborn	2.42 <i>n</i> = 5, SD = 0.20	4.86 <i>n</i> = 5, SD = 0.38
2	Affected	Newborn	2.88 <i>n</i> = 5, SD = 0.29	5.52 <i>n</i> = 5, SD = 0.71
3	Control	Newborn	2.61 <i>n</i> = 4, SD = 0.45	7.43 <i>n</i> = 4, SD = 1.47
4	Control	Newborn	3.87 <i>n</i> = 4, SD = 0.62	9.11 <i>n</i> = 4, SD = 1.84
5	Affected	1 month	4.12 <i>n</i> = 5, SD = 0.79	6.19 <i>n</i> = 5, SD = 1.44
6	Control	1 month	2.86 <i>n</i> = 5, SD = 0.53	7.14 <i>n</i> = 5, SD = 1.66
7	Affected	Mature	3.83 <i>n</i> = 4, SD = 0.55	3.61 <i>n</i> = 4, SD = 0.64
8	Control	Mature	4.45 <i>n</i> = 5, SD = 0.78	5.63 <i>n</i> = 5, SD = 1.22

Concentrations are expressed as ng per 10 mg of sample. *n* = the number of repeat measurements on the sample.

($P \leq 0.0001$). The concentration of unsulfated chondroitin disaccharide did not vary significantly with disease status.

Although the ratio of C4S to C6S decreases with age, alterations due to degenerative joint disease are variable, with some previous reports of no change and other reports of an increased ratio of C4S to C6S. The older affected animals in this study demonstrated angular limb deformities with resultant misloading of joints, and therefore it is possible that the change in chondroitin sulfation was a reflection of secondary degenerative joint disease. The change in sulfation, however, was a decrease rather than an increase in the ratio of C4S to C6S, and the presence of an altered sulfation profile in the newborn lambs that showed no gross or microscopic joint lesions supports a primary alteration in chondroitin sulfation.

A decreased ratio of Δ di-mono4S to Δ di-mono6S has been recorded in human chondrodysplasias resulting from sulfate transport disorders due to reduced activity of the diastrophic dysplasia sulfate transporter (DTDST)¹³ and in mutations in the gene encoding chondroitin 4-sulfotransferase-1, *CHST11*.¹⁴ This finding indicates that *CHST11*, which maps to ovine chromosome 3 at approximately 221.4cM, is potentially a candidate gene for involvement in the condition. A naturally occurring decrease in the ratio of Δ di-mono4S to Δ di-mono6S has previously been described in aging articular cartilage in humans,^{9,15} and indeed a similar change was evident in the articular cartilage of mature sheep in this study.

The decreased ratio of Δ di-mono4S to Δ di-mono6S in newborn chondrodysplastic lambs indicates that although affected lambs cannot be distinguished phenotypically




Figure 2. Ratios of Δ di-mono4S to Δ di-mono6S in chondrodysplastic and control Texel sheep of different ages. Gray plots are chondrodysplastic animals; white plots are control animals. The ratios are lower in affected animals of all ages ($P \leq 0.0001$).

from normal or heterozygous lambs at birth, the underlying cartilage abnormality is present. Capillary electrophoresis techniques may therefore be useful in identifying the phenotype of newborn animals that are yet to demonstrate gross or microscopic lesions of chondrodysplasia. Although more samples would be required to confirm the true diagnostic power of the ratio of cartilage Δ di-mono4S to Δ di-mono6S, this study indicates that the information would at least be useful in future genetic studies as quantitative trait data associated with the chondrodysplastic phenotype.

The newborn affected lambs, which had no gross or microscopic lesions, and the mature affected ram, which had mild lesions of chondrodysplasia, all had ratios of Δ di-mono4S to Δ di-mono6S closer to the values for control animals than did the 1-month-old affected lamb, which demonstrated severe lesions of the disease. This suggested that the magnitude of the difference in the ratio of Δ di-mono4S to Δ di-mono6S may have been related to the severity of lesions. A study including a greater number of animals of different ages and representing the full spectrum of severity in this disease would be required to investigate this further. Such a study could be based on capillary electrophoresis of serial cartilage biopsies because of the small sample size (10 mg of cartilage) required for this technique.

Although the number of animals studied was relatively small, there are statistically significant, even striking, differences between affected sheep and control animals. Whatever the precise origins of the disease, it is clear that the manifest changes in sulfation at one particular position on the sugar

ring produce large changes in the macroscopic mechanical properties of the resultant cartilage. Understanding in detail how such molecular change is mediated, presumably through modifications of biopolymeric assembly, would also be an interesting topic of further work.

Funding

The authors received no financial support for the research or authorship of this article.

Declaration of Conflicting Interests

The authors declare no potential conflicts of interests with respect to the authorship and publication of this article.

References

1. Krueger RCJ, Fields TA, Hildreth J, Schwartz NB. Chick cartilage chondroitin sulfate proteoglycan core protein: I. Generation and characterization of peptides and specificity for glycosaminoglycan attachment. *J Biol Chem.* 1990;265:12075-87.
2. Sandy JD, Plaas AHK, Rosenberg L. Structure, function and metabolism of cartilage proteoglycans. In: Koopman WJ, editor. *Arthritis and allied conditions: a textbook of rheumatology.* 13th ed. Philadelphia: Williams & Wilkins; 1997. p. 229-44.
3. Rossi A, Bonaventure J, Delezoide AL, Cetta G, Superti-Furga A. Undersulfation of proteoglycans synthesized by chondrocytes from a patient with achondrogenesis type 1b homozygous for an 1483p substitution in the diastrophic dysplasia sulfate transporter. *J Biol Chem.* 1996;271:18456-64.
4. Rossi A, Superti-Furga A. Mutations in the diastrophic dysplasia sulfate transporter (DTDST) gene (SLC26A2): 22 novel mutations, mutation review, associated skeletal phenotypes, and diagnostic relevance. *Hum Mutat.* 2001;17:159-71.
5. Forlino A, Piazza R, Torre SD, Tatangelo L, Bonafe L, Gualeni B, et al. A diastrophic dysplasia sulfate transporter (SLC26A2) mutant mouse: morphological and biochemical characterization of the resulting chondrodysplasia phenotype. *Hum Mol Genet.* 2005;14:859-71.
6. Haque MF, King LM, Krakow D, Cantor RM, Rusiniak ME, Swank RT, et al. Mutations in orthologous genes in human spondyloepimetaphyseal dysplasia and the brachymorphic mouse. *Nat Genet.* 1998;20:157-62.
7. Leach RMJ, Muenster AM. Studies on the role of manganese in bone formation: I. Effect on the mucopolysaccharidal content of chick bone. *J Nutr.* 1962;78:51.
8. Price FM, Levick JR, Mason RM. Glycosaminoglycan concentration in synovium and other tissues of rabbit knee in relation to synovial hydraulic resistance. *J Physiol.* 1996;495:803-20.
9. Bayliss MT, Osborne D, Woodhouse S, Davidson C. Sulfation of chondroitin sulfate in human articular cartilage: the effect of age, topographical position, and zone of cartilage on tissue composition. *J Biol Chem.* 1999;274:15892-900.
10. Theocharis AD, Theocharis DA. High-performance capillary electrophoretic analysis of hyaluronan and galactosaminoglycan-disaccharides in gastrointestinal carcinomas: differential disaccharide composition as a possible tool-indicator for malignancies. *Biomed Chromatogr.* 2002;16:157-61.
11. Karamanos NK, Axelsson S, Vanky P, Tzanakakis GN, Hjerpe A. Determination of hyaluronan and galactosaminoglycan disaccharides by high-performance capillary electrophoresis at the attomole level: applications to analyses of tissue and cell culture proteoglycans. *J Chromatogr.* 1995;696A:295-305.
12. Goodall DM, Williams SJ, Lloyd DK. Quantitative aspects of capillary electrophoresis. *Trends Anal Chem.* 1991;10:272-9.
13. Rossi A, Kaitila I, Wilcox WR, Rimo DL, Steinmann B, Cetta G, et al. Proteoglycan sulfation in cartilage and cell cultures from patients with sulfate transporter chondrodysplasias: relationship to clinical severity and indications on the role of intracellular sulfate production. *Matrix Biol.* 1998;17:361-9.
14. Klüppel M, Wight TN, Chan C, Hinek A, Wrana JL. Maintenance of chondroitin sulfation balance by chondroitin-4-sulfotransferase 1 is required for chondrocyte development and growth factor signaling during cartilage morphogenesis. *Development.* 2005;132:3989-4003.
15. Lauder RM, Huckerby TN, Brown GM, Bayliss MT, Nieduszynski IA. Age-related changes in the sulfation of the chondroitin sulfate linkage region from human articular cartilage aggrecan. *Biochem J.* 2001;358:523-8.