

On the sums of continued fractions

by

BOHUSLAV DIVIŠ (Columbus, Ohio)

*Dedicated to the memory of my teacher
Professor Vojtěch Jarník*

Let α be an irrational number and $\alpha = (g_0; g_1, g_2, \dots)$ its regular continued fraction expansion. For each natural number N we shall denote by F_N the set of all irrational numbers α , for which $g_j \leq N$ ($j = 1, 2, 3, \dots$). M. Hall Jr. [1] has proved that each real number β can be written in the form $\beta = a_1 + a_2$, where $a_j \in F_4$ ($j = 1, 2$). The purpose of this paper is to show that each real number can be expressed as the sum of three elements of the set F_3 or four elements of the set F_2 . Moreover, we show that a lesser number of summands, in general, does not suffice.

THEOREM 1. *Let β be a real number. Then there exist three numbers $a_j \in F_3$ ($j = 1, 2, 3$) such that $\beta = a_1 + a_2 + a_3$. Also there exist real numbers β' such that $\beta' \neq a'_1 + a'_2$ for any pair (a'_1, a'_2) , $a'_i \in F_3$ ($i = 1, 2$).*

THEOREM 2. *Let β be a real number. Then there exist four numbers $a_j \in F_2$ ($j = 1, 2, 3, 4$) such that $\beta = a_1 + a_2 + a_3 + a_4$. Also there exist real numbers β' such that $\beta' \neq a'_1 + a'_2 + a'_3$ for any triple (a'_1, a'_2, a'_3) , $a'_i \in F_2$ ($i = 1, 2, 3$).*

Now, let $k \geq 2$ be a natural number. Let A_1, A_2, \dots, A_k be non-empty sets of real numbers. We shall call their Schnirelman sum (notation $A_1 \dot{+} A_2 \dot{+} \dots \dot{+} A_k$) the set of all numbers of the form $\alpha_1 + \alpha_2 + \dots + \alpha_k$, where $\alpha_j \in A_j$ ($j = 1, 2, \dots, k$). Theorems 1 and 2 then follow from Theorems 1' and 2', respectively.

THEOREM 1'.

$$(1') \quad F_2 \dot{+} F_3 \dot{+} F_3 = (-\infty, +\infty),$$

$$(2') \quad F_3 \dot{+} F_3 \neq (-\infty, +\infty).$$

THEOREM 2'.

$$(3') \quad F_2 \dot{+} F_2 \dot{+} F_2 \dot{+} F_2 = (-\infty, +\infty),$$

$$(4') \quad F_2 \dot{+} F_2 \dot{+} F_2 \neq (-\infty, +\infty).$$

For integral numbers $k \geq 0$ and for natural numbers N we denote by the symbol $F_N(n_0, n_1, \dots, n_k)$ the set of all $a \in F_N$, $a = (g_0; g_1, g_2, \dots)$, for which $g_i = n_i$ ($i = 0, 1, \dots, k$). Theorems 1' and 2' then follow from Theorems 1'' and 2''.

THEOREM 1''.

$$(1'') \quad F_3(0) + F_3(0) + F_3(0) = \left[\frac{\sqrt{21}-3}{2}, 3 \frac{\sqrt{21}-3}{2} \right],$$

$$(2'') \quad F_3(0) + F_3(0) \subset \left[\frac{\sqrt{21}-3}{2}, \frac{15-\sqrt{21}}{17} \right] \cup \left[\frac{4\sqrt{21}-9}{15}, \sqrt{21}-3 \right].$$

THEOREM 2''.

$$(3'') \quad F_2(0) + F_2(0) + F_2(0) + F_2(0) = [2(\sqrt{3}-1), 4(\sqrt{3}-1)],$$

$$(4'') \quad F_2(0) + F_2(0) + F_2(0) \subset \left[3 \frac{\sqrt{3}-1}{2}, 3-\sqrt{3} \right] \cup \left[\frac{4\sqrt{3}-3}{3}, 3(\sqrt{3}-1) \right].$$

We have obviously

$$\begin{aligned} F_3 + F_3 + F_3 &= \bigcup_{\substack{n_0, n'_0, n''_0 \\ \text{int.}}} (F_3(n_0) + F_3(n'_0) + F_3(n''_0)) \\ &= \bigcup_{\substack{n_0, n'_0, n''_0 \\ \text{int.}}} (F_3(0) + F_3(0) + F_3(0) + \{n_0 + n'_0 + n''_0\}) \\ &= \bigcup_{n \text{ int.}} (F_3(0) + F_3(0) + F_3(0) + \{n\}) \end{aligned}$$

and the length of the interval $\left[\frac{\sqrt{21}-3}{2}, 3 \frac{\sqrt{21}-3}{2} \right]$ is $\sqrt{21}-3 > 1$.

Hence, (1') follows from (1''). Similarly, we have

$$F_2 + F_2 = \bigcup_{n \text{ int.}} (F_2(0) + F_2(0) + \{n\}).$$

Since $\frac{4\sqrt{21}-9}{15} > \frac{15-\sqrt{21}}{17}$,

$$\max(F_3(0) + F_3(0) + \{-1\}) = \sqrt{21}-4 < \frac{15-\sqrt{21}}{17},$$

$$\min(F_3(0) + F_3(0) + \{1\}) = \frac{\sqrt{21}-3}{3} + 1 > \frac{4\sqrt{21}-9}{15},$$

the inclusion (2'') implies (2'). In an analogous manner we get very easily that (3'') implies (3').

Finally, since $\frac{4\sqrt{3}-3}{3} > 3-\sqrt{3}$,

$$\max(F_2(0) + F_2(0) + F_2(0) + \{-1\}) = 3(\sqrt{3}-1) - 1 < 3-\sqrt{3},$$

$$\min(F_2(0) + F_2(0) + F_2(0) + \{1\}) = 3 \frac{\sqrt{3}-1}{2} + 1 > \frac{4\sqrt{3}-3}{3},$$

the inclusion (4'') implies (4').

We start with the proof of inclusions (2'') and (4'').

Proof of (2''). We have obviously $F_3(0) = F_3(0, 1) \cup F_3(0, 2) \cup F_3(0, 3)$ and also

$$F_3(0, 3) \subset [(0; 3, \overline{1, 3}), (0; 3, \overline{3, 1})] = \left[\frac{\sqrt{21}-3}{6}, \frac{15-\sqrt{21}}{34} \right] \stackrel{\text{df}}{=} K_1,$$

$$\begin{aligned} F_3(0, 2) \cup F_3(0, 1) &\subset [(0; 2, \overline{1, 3}), (0; 1, \overline{3, 1})] \\ &= \left[\frac{\sqrt{21}-1}{10}, \frac{\sqrt{21}-3}{2} \right] \stackrel{\text{df}}{=} K_2. \end{aligned}$$

From this follows that $F_3(0) \subset K_1 \cup K_2$ and hence

$$\begin{aligned} F_3(0) + F_3(0) &\subset (K_1 + K_1) \cup (K_1 + K_2) \cup (K_2 + K_2) \\ &= \left[\frac{\sqrt{21}-3}{3}, \frac{15-\sqrt{21}}{17} \right] \cup \left[\frac{4\sqrt{21}-9}{15}, \sqrt{21}-3 \right]. \end{aligned}$$

Proof of (4''). Obviously, $F_2(0) = F_2(0, 1) \cup F_2(0, 2)$ and also

$$F_2(0, 2) \subset [(0; 2, \overline{1, 2}), (0; 2, \overline{2, 1})] = \left[\frac{\sqrt{3}-1}{2}, \frac{3-\sqrt{3}}{3} \right] \stackrel{\text{df}}{=} K_3,$$

$$F_2(0, 1) \subset [(0; 1, \overline{1, 2}), (0; 1, \overline{2, 1})] = \left[\frac{\sqrt{3}}{3}, \sqrt{3}-1 \right] \stackrel{\text{df}}{=} K_4.$$

From this follows that $F_2(0) \subset K_3 \cup K_4$ and hence

$$\begin{aligned} F_2(0) + F_2(0) + F_2(0) &\subset (K_3 + K_3 + K_3) \cup (K_3 + K_3 + K_4) \cup (K_3 + K_4 + K_4) \cup (K_4 + K_4 + K_4) \\ &= \left[3 \frac{\sqrt{3}-1}{2}, 3-\sqrt{3} \right] \cup \left[\frac{4\sqrt{3}-3}{3}, 3(\sqrt{3}-1) \right]. \end{aligned}$$

We shall prove relations (1'), (3'') as an application of the following more general considerations.

Let I_0 be a non-degenerate compact interval with endpoints a, b , i.e. $I_0 = [a, b]$. We shall call I_0 an interval of order 0. We delete from I_0 an open interval (c, d) , such that $a < c < d < b$. We get thus two closed intervals $I_1^{(1)} = [a, c]$, $I_1^{(2)} = [d, b]$, which we shall call intervals of order 1. Generally, when we have already constructed all intervals $I_n^{(i_1 \dots i_n)}$ of order n for some $n \geq 0$ we get intervals $I_{n+1}^{(i_1 \dots i_{n+1})}$, $I_{n+1}^{(i_1 \dots i_n 2)}$ of order $n+1$ by deleting from each interval of order n an open non-empty interval, in such a way that the resulting intervals of order $n+1$ are non-degenerate. If we denote by K_n ($n \geq 0$) the union of all intervals of order n , then $\bigcap_{n \geq 0} K_n \stackrel{\text{df}}{=} L(I_0)$ is a closed, non-countable set contained in I_0 and with measure less than the length of I_0 . $L(I_0)$ contains either a non-empty interval or is nowhere dense in I_0 .

In the following, we shall denote by $\mu(A)$ the measure (length) of an interval A .

DEFINITION. We shall say that $L(I_0)$ satisfies the k -condition, if for arbitrary $n \geq 0$, the lengths of the intervals $I_n^{(i_1 \dots i_n)}$, $I_{n+1}^{(i_1 \dots i_{n+1})}$ and $I_{n+1}^{(i_1 \dots i_n 2)}$ satisfy the following two inequalities:

$$\mu(I_n^{(i_1 \dots i_n)}) \leq k \cdot \mu(I_{n+1}^{(i_1 \dots i_{n+1})}) + \mu(I_{n+1}^{(i_1 \dots i_n 2)}),$$

$$\mu(I_n^{(i_1 \dots i_n)}) \leq \mu(I_{n+1}^{(i_1 \dots i_{n+1})}) + k \cdot \mu(I_{n+1}^{(i_1 \dots i_n 2)}).$$

The following assertion, in a different formulation was proved by M. Hall Jr. [1].

ASSERTION 1. Let A, B be non-degenerate compact intervals and $L(A), L(B)$ satisfy the 2-condition. If $\frac{1}{2} \leq \mu(A)/\mu(B) \leq 3$, then

$$L(A) + L(B) = A + B.$$

For the proof of (1'') we shall need an analogue of Assertion 1 for three "summands":

ASSERTION 2. Let A, B, C be non-degenerate compact intervals and $L(A), L(B), L(C)$ satisfy the 3-condition. If $\mu(A) + \mu(B) + \mu(C) \leq 6 \min(\mu(A), \mu(B), \mu(C))$, then

$$L(A) + L(B) + L(C) = A + B + C.$$

For the proof of (3'') we shall need an analogue of Assertion 1 for four "summands":

ASSERTION 3. Let A, B, C, D be non-degenerate compact intervals and $L(A), L(B), L(C), L(D)$ satisfy the 4-condition. If $\mu(A) + \mu(B) + \mu(C) + \mu(D) \leq 8 \min(\mu(A), \mu(B), \mu(C), \mu(D))$, then

$$L(A) + L(B) + L(C) + L(D) = A + B + C + D.$$

Remark. The similarities of Assertions 2 and 3 with Assertion 1 will become clearer, when we rewrite the condition $\frac{1}{2} \leq \mu(A)/\mu(B) \leq 3$ from Assertion 1 to the equivalent form $\mu(A) + \mu(B) \leq 4 \min(\mu(A), \mu(B))$.

Instead of Assertions 2 and 3 we shall formulate and prove more general Assertions 2' and 3'.

ASSERTION 2'. Let $A = [A_L, A_R]$, $B = [B_L, B_R]$, $C = [C_L, C_R]$ be non-degenerate compact intervals. Let $L(A), L(B), L(C)$ satisfy the 3-condition. Then

$$[A_L + B_L + C_L, A_L + B_L + C_L + 3 \min(\mu(A), \mu(B), \mu(C))] \subset L(A) + L(B) + L(C),$$

$$[A_R + B_R + C_R - 3 \min(\mu(A), \mu(B), \mu(C)), A_R + B_R + C_R] \subset L(A) + L(B) + L(C).$$

ASSERTION 3'. Let $A = [A_L, A_R]$, $B = [B_L, B_R]$, $C = [C_L, C_R]$, $D = [D_L, D_R]$ be non-degenerate compact intervals. Let $L(A), L(B), L(C), L(D)$ satisfy the 4-condition. Then

$$[A_L + B_L + C_L + D_L, A_L + B_L + C_L + D_L + 4 \min(\mu(A), \mu(B), \mu(C), \mu(D))] \subset L(A) + L(B) + L(C) + L(D),$$

$$[A_R + B_R + C_R + D_R - 4 \min(\mu(A), \mu(B), \mu(C), \mu(D)), A_R + B_R + C_R + D_R] \subset L(A) + L(B) + L(C) + L(D).$$

The proofs of Assertions 2' and 3' will make use of Lemmas 1 and 2 below. Before proceeding, some new definitions are needed.

DEFINITION. Let $A_j = [x_j, x_j + a_j]$ ($j = 1, 2, \dots, n$) be n ($n \geq 2$) non-degenerate compact intervals. The interval

$$\overline{(A_1, A_2, \dots, A_n)} = \left[\sum_{j=1}^n x_j, \sum_{j=1}^n x_j + n \min_{1 \leq j \leq n} a_j \right]$$

will be called their *lower associate*, the interval

$$\overline{(A_1, A_2, \dots, A_n)} = \left[\sum_{j=1}^n (x_j + a_j) - n \min_{1 \leq j \leq n} a_j, \sum_{j=1}^n (x_j + a_j) \right]$$

will be called their *upper associate*.

LEMMA 1. Let a_j ($j = 1, 2, 3$) be real numbers. Let a_j ($j = 1, 2, 3$), b_1, c_1 be positive numbers, such that $a_1 \geq a_2 \geq a_3$, $a_1 > b_1 + c_1$,

$$(5) \quad a_1 \leq b_1 + 3c_1,$$

$$(6) \quad a_1 \leq 3b_1 + c_1.$$

Let us put

$$A_j = [x_j, x_j + a_j] \quad (j = 1, 2, 3),$$

$$A_{11} = [x_1, x_1 + b_1],$$

$$A_{12} = [x_1 + a_1 - c_1, x_1 + a_1].$$

Then

$$(7) \quad \overline{(A_1, A_2, A_3)} \cup \overline{(A_1, A_2, A_3)} \subset \bigcup_{j=1}^2 \overline{((A_{1j}, A_2, A_3) \cup (A_{1j}, A_2, A_3))}.$$

Proof. Because the situation is completely invariant with respect to the numbers x_j ($j = 1, 2, 3$), we may restrict ourselves to the case $x_1 = x_2 = x_3 = 0$. Moreover, it suffices to prove the inclusion (7) only for $\overline{(A_1, A_2, A_3)}$ due to the symmetry between the letters b, c and the lower and upper associates.

By the very definition and by the assumptions of the lemma we have

$$\overline{(A_1, A_2, A_3)} = [0, 3a_3], \quad \overline{(A_{11}, A_2, A_3)} = [0, 3 \min(a_3, b_1)].$$

Hence, if $b_1 \geq a_3$, we are done. Let in the following $b_1 < a_3$, i.e. $\overline{(A_{11}, A_2, A_3)} = [0, 3b_1]$. We have

$$\overline{(A_{12}, A_2, A_3)} = [a_1 - c_1, a_1 - c_1 + 3 \min(c_1, a_3)]$$

and using (6):

$$[0, a_1 - c_1 + 3 \min(c_1, a_3)] \subset \overline{(A_{11}, A_2, A_3)} \cup \overline{(A_{12}, A_2, A_3)}.$$

Hence, if $c_1 \geq a_3$, we are done. Let in the following also $c_1 < a_3$, i.e.

$$(8) \quad [0, a_1 + 2c_1] \subset \overline{(A_{11}, A_2, A_3)} \cup \overline{(A_{12}, A_2, A_3)}.$$

Now, we have

$$\overline{(A_{11}, A_2, A_3)} = [a_2 + a_3 - 2b_1, a_2 + a_3 + b_1],$$

$$\overline{(A_{12}, A_2, A_3)} = [a_1 + a_2 + a_3 - 3c_1, a_1 + a_2 + a_3].$$

Using (5), we get

$$(9) \quad [a_2 + a_3 - 2b_1, a_1 + a_2 + a_3] \subset \overline{(A_{11}, A_2, A_3)} \cup \overline{(A_{12}, A_2, A_3)}.$$

From the relations (8), (9) we obtain that

$$[0, a_1 + a_2 + a_3] \subset \bigcup_{j=1}^2 \overline{((A_{1j}, A_2, A_3) \cup (A_{1j}, A_2, A_3))},$$

since $a_2 + a_3 - 2b_1 \leq 2a_1 - 2b_1 \leq a_1 + 2c_1$ by (5), (6).

LEMMA 2. Let x_j ($j = 1, 2, 3, 4$) be real numbers. Let a_j ($j = 1, 2, 3, 4$), b_j, c_j ($j = 1, 2, 3$) be positive numbers such that

$$(10) \quad a_4 = \min_{1 \leq j \leq 4} a_j,$$

$$(11) \quad a_j > b_j + c_j \quad (j = 1, 2, 3),$$

$$(12) \quad a_j \leq 4b_j + c_j \quad (j = 1, 2, 3),$$

$$(13) \quad a_j \leq b_j + 4c_j \quad (j = 1, 2, 3).$$

Let us denote

$$A_{j0} = [x_j, x_j + a_j] \quad (j = 1, 2, 3, 4),$$

$$A_{j1} = [x_j, x_j + b_j] \quad (j = 1, 2, 3),$$

$$A_{j2} = [x_j + a_j - c_j, x_j + a_j] \quad (j = 1, 2, 3).$$

Then

$$(14) \quad \overline{(A_{10}, A_{20}, A_{30}, A_{40})} \cup \overline{(A_{10}, A_{20}, A_{30}, A_{40})} \subset \bigcup_{\substack{0 \leq j, k, l \leq 2 \\ j+k+l \geq 1}} \overline{((A_{1j}, A_{2k}, A_{3l}, A_{40}) \cup (A_{1j}, A_{2k}, A_{3l}, A_{40}))}.$$

Proof. For the same reasons as in the proof of Lemma 1 we shall restrict ourselves only to the proof of the inclusion (14) for the lower associate $\overline{(A_{10}, A_{20}, A_{30}, A_{40})}$ and, without loss of generality, we shall suppose that $x_1 = x_2 = x_3 = x_4 = 0$. Hence we have

$$\overline{(A_{10}, A_{20}, A_{30}, A_{40})} = [0, 4a_4].$$

If $b_1 \geq a_4$, then

$$\overline{(A_{11}, A_{20}, A_{30}, A_{40})} = [0, 4a_4]$$

and we are done. The cases $b_2 \geq a_4, b_3 \geq a_4$ can be handled similarly. Let in the following

$$(15) \quad \max_{1 \leq j \leq 3} b_j < a_4,$$

which means that

$$(16) \quad \overline{(A_{11}, A_{20}, A_{30}, A_{40})} \cup \overline{(A_{10}, A_{21}, A_{30}, A_{40})} \cup \overline{(A_{10}, A_{20}, A_{31}, A_{40})} = [0, 4 \max_{1 \leq j \leq 3} b_j].$$

Now, $\overline{(A_{12}, A_{20}, A_{30}, A_{40})} = [a_1 - c_1, a_1 - c_1 + 4 \min(c_1, a_4)]$ and by (12), this interval overlaps the interval (16). If $c_1 \geq a_4$, then we are done and also in the cases $c_2 \geq a_4$ or $c_3 \geq a_4$, by an analogous argument. Hence, we shall suppose in the following that

$$(17) \quad \max_{1 \leq j \leq 3} c_j < a_4.$$

It follows from the considerations above, the lower associates which occur on the right-hand side of the inclusion (14), cover the interval

$$(18) \quad \left[0, \max\left\{\max_{1 \leq j \leq 3} 4b_j; \max_{1 \leq j \leq 3} (a_j + 3c_j)\right\}\right].$$

Completely analogously we could show, that the upper associates which occur on the right-hand side of the inclusion (14), cover the interval

$$(19) \quad \left[\sum_{j=1}^4 a_j - \max\left\{\max_{1 \leq j \leq 3} 4c_j; \max_{1 \leq j \leq 3} (a_j + 3b_j)\right\}, \sum_{j=1}^4 a_j\right].$$

Unfortunately, in the general case, the union of the intervals (18) and (19) does not cover the interval $[0, 4a_4]$; however, fortunately, we still have not used all the associates on the right-hand side of the inclusion (14).

We shall show that the interval $(A_{12}, A_{22}, A_{32}, A_{40})$ overlaps the interval (19). This statement is equivalent to the inequality

$$\sum_{j=1}^3 a_j - \sum_{j=1}^3 c_j + 4 \min_{1 \leq j \leq 3} c_j \geq \sum_{j=1}^4 a_j - \max\left\{\max_{1 \leq j \leq 3} 4c_j; \max_{1 \leq j \leq 3} (a_j + 3b_j)\right\},$$

which is equivalent to

$$(20) \quad \max\left\{\max_{1 \leq j \leq 3} 4c_j; \max_{1 \leq j \leq 3} (a_j + 3b_j)\right\} + 4 \min_{1 \leq j \leq 3} c_j \geq a_4 + \sum_{j=1}^3 c_j.$$

The last inequality can be easily proved using the relations (10), (12) and (13). Namely, we have

$$\begin{aligned} & \max\left\{\max_{1 \leq j \leq 3} 4c_j; \max_{1 \leq j \leq 3} (a_j + 3b_j)\right\} + 4 \min_{1 \leq j \leq 3} c_j \\ & \geq \frac{1}{2} \max_{1 \leq j \leq 3} 4c_j + \frac{1}{2} \max_{1 \leq j \leq 3} (a_j + 3b_j) + 4 \min_{1 \leq j \leq 3} c_j \\ & \geq \sum_{j=1}^3 c_j + \frac{1}{2} \min_{1 \leq j \leq 3} a_j + \frac{3}{2} \max_{1 \leq j \leq 3} b_j + 3 \min_{1 \leq j \leq 3} c_j \\ & \geq \sum_{j=1}^3 c_j + \frac{1}{2} \min_{1 \leq j \leq 3} a_j + \frac{9}{10} \min_{1 \leq j \leq 3} a_j > a_4 + \sum_{j=1}^3 c_j. \end{aligned}$$

Quite analogously we could show that the interval $(A_{12}, A_{21}, A_{31}, A_{40})$ overlaps the interval (18). Hence, under the assumptions (15) and (17), the following intervals will certainly be covered:

$$(21) \quad \left[0, \max\left\{\max_{1 \leq j \leq 3} 4b_j; \max_{1 \leq j \leq 3} (a_j + 3c_j); a_4 + \sum_{j=1}^3 b_j\right\}\right],$$

$$(22) \quad \left[\sum_{j=1}^4 a_j - \max\left\{\max_{1 \leq j \leq 3} 4c_j; \max_{1 \leq j \leq 3} (a_j + 3b_j); a_4 + \sum_{j=1}^3 c_j\right\}, \sum_{j=1}^4 a_j\right].$$

Unfortunately, even the union of the intervals (21), (22) need not cover the interval $[0, 4a_4]$. But we still have not made use of all the intervals on the right-hand side of the inclusion (14). Moreover, we shall show that the whole interval $\left[0, \sum_{j=1}^4 a_j\right]$ is covered.

Case A. Let

$$b_j \leq c_j \quad (j = 1, 2, 3) \quad \text{and} \quad c_1 \geq c_2 \geq c_3.$$

We show that the interval $(A_{12}, A_{22}, A_{30}, A_{40})$ overlaps the interval (21).

If not, we have

$$(24) \quad a_1 - c_1 + a_2 - c_2 > a_1 + 3c_1, \quad a_1 - c_1 + a_2 - c_2 > a_2 + 3c_2.$$

Adding the inequalities (24) we would get

$$a_1 + a_2 > 5c_1 + 5c_2 \geq c_1 + 4b_1 + c_2 + 4b_2,$$

contradicting the assumptions (12), (13). Similarly we get that also the intervals $(A_{12}, A_{20}, A_{32}, A_{40})$, $(A_{10}, A_{22}, A_{32}, A_{40})$ overlap the interval (21). Hence, certainly the interval

$$(25) \quad \left[0, \max\{a_1 - c_1 + a_2 + 3c_2; a_1 - c_1 + a_3 + 3c_3\}\right]$$

is covered. Now, we show that the interval $(A_{12}, A_{22}, A_{32}, A_{40})$ overlaps (25). If not, we have

$$(26) \quad \begin{aligned} a_1 - c_1 + a_2 - c_2 + a_3 - c_3 &> a_1 - c_1 + a_2 + 3c_2, \\ a_1 - c_1 + a_2 - c_2 + a_3 - c_3 &> a_1 - c_1 + a_3 + 3c_3. \end{aligned}$$

Adding the inequalities (26) we would get

$$a_2 + a_3 > 5c_2 + 5c_3 \geq c_2 + 4b_2 + c_3 + 4b_3,$$

thus contradicting (12) and (13). Now, the interval $(A_{12}, A_{22}, A_{32}, A_{40})$ is contained in (22) and hence we have shown that (25) overlaps (22), i.e. the whole interval $\left[0, \sum_{j=1}^4 a_j\right]$ is covered.

Case B. Let

$$(27) \quad b_j \leq c_j \quad (j = 1, 2), \quad b_3 > c_3, \quad c_1 \geq c_2.$$

Again as in case A, the interval $(A_{12}, A_{22}, A_{30}, A_{40})$ overlaps the interval (21). We shall distinguish three subcases.

Subcase B1. Let

$$(28) \quad b_3 \geq c_1.$$

Then the interval $(A_{12}, A_{22}, A_{30}, A_{40})$ overlaps the interval (22), since (by (27), (28), (12), (13) and (10))

$$\begin{aligned} a_1 - c_1 + a_2 + 3c_2 &\geq a_1 + a_2 + 2c_1 + 3c_2 - 3b_2 \geq a_1 + a_2 + 2b_2 + 3c_2 - 3b_2 \\ &\geq a_1 + a_2 + a_4 - 3b_3 = \sum_{j=1}^4 a_j - (a_3 + 3b_3). \end{aligned}$$

Thus the whole interval $[0, \sum_{j=1}^4 a_j]$ is covered.

Subcase B2. Let

$$(29) \quad b_3 < c_1, \quad 4c_2 + c_3 \geq a_3.$$

Then the interval $(A_{12}, A_{22}, A_{30}, A_{40})$ overlaps the interval (22), since (by (29))

$$a_1 - c_1 + a_2 + 3c_2 \geq a_1 - c_1 + a_2 - c_2 + a_3 - c_3 = \sum_{j=1}^4 a_j - \left(a_4 + \sum_{j=1}^3 c_j \right).$$

Thus the whole interval $[0, \sum_{j=1}^4 a_j]$ is covered.

Subcase B3. Let

$$(30) \quad b_3 < c_1, \quad 4c_2 + c_3 < a_3.$$

The assumption $4c_2 + c_3 < a_3$ implies (using (12))

$$4c_2 + c_3 < a_3 \leq 4b_2 + c_3,$$

thus

$$(31) \quad c_2 < b_2,$$

hence (using (27))

$$(32) \quad b_2 < b_3,$$

thus (by (13), (31))

$$(33) \quad a_2 \leq b_2 + 4c_2 < b_2 + 4b_2.$$

We shall show that the interval $(A_{10}, A_{21}, A_{31}, A_{40})$ overlaps the interval (21). We have (using (32), (10), (30), (12), (13), (27))

$$\begin{aligned} \sum_{j=1}^4 a_j - (a_2 - b_2 + a_3 - b_3 + 4 \min b_j) &= a_1 + a_4 + b_3 - 3b_2 \\ &\leq a_1 + a_2 + c_1 - 3b_2 \leq a_1 + c_1 + 2c_2 \leq a_1 + 3c_1. \end{aligned}$$

Finally, we show that the interval $(A_{10}, A_{21}, A_{31}, A_{40})$ overlaps the interval (22). We have (by (33))

$$\sum_{j=1}^4 a_j - (a_2 - b_2 + a_3 - b_3) \geq \sum_{j=1}^4 a_j - (a_3 + 3b_3).$$

Thus again the whole interval $[0, \sum_{j=1}^4 a_j]$ is covered.

Case C. We have neither case A nor B. Then by a suitable renumbering or change of the role of the letters b and c , we can reduce the situation to one of the cases A, B.

Proof of Assertion 2'. The intervals on the left-hand sides of the inclusions in Assertion 2' are associates (A, B, C) and $(\bar{A}, \bar{B}, \bar{C})$. If a number δ belongs to one of them, then (by Lemma 1) δ belongs to an infinite sequence of nested closed associates $(A^{(i)}, B^{(i)}, C^{(i)})$ ($i = 1, 2, \dots$) (upper or lower) such that $A^{(i+1)} \subset A^{(i)}, B^{(i+1)} \subset B^{(i)}, C^{(i+1)} \subset C^{(i)}$ ($i = 1, 2, \dots$) and we cannot have equality in all three cases. Also, $A^{(1)} = A, B^{(1)} = B, C^{(1)} = C$ and each of the intervals $A^{(i)}$ or $B^{(i)}$ or $C^{(i)}$ contains points of $L(A)$ or $L(B)$ or $L(C)$, respectively. If the sets $L(A), L(B)$ and $L(C)$ do not contain any interval, then, since 1) the intervals $A^{(i)}, B^{(i)}, C^{(i)}$ are closed, 2) always the shortest one of them remains undivided at each step (unless they have the same length), 3) the associates $(A^{(i)}, B^{(i)}, C^{(i)})$ are closed, 4) $(A^{(i)}, B^{(i)}, C^{(i)}) \subset A^{(i)} + B^{(i)} + C^{(i)}$ ($i = 1, 2, 3, \dots$), the intersections $\bigcap_i A^{(i)}, \bigcap_i B^{(i)}, \bigcap_i C^{(i)}$ contain just one point and we have

$$\begin{aligned} \bigcap_i A^{(i)} &= \{\alpha\} \subset L(A), \quad \bigcap_i B^{(i)} = \{\beta\} \subset L(B), \quad \bigcap_i C^{(i)} = \{\gamma\} \subset L(C), \\ a + \beta + \gamma &= \delta, \quad \{\delta\} = \bigcap_i (A^{(i)}, B^{(i)}, C^{(i)}). \end{aligned}$$

If some of the sets $L(A), L(B), L(C)$ contains an interval, e.g. $L(A)$ say, then we can delete from $L(A)$ countably many open intervals such that the resulting set $L'(A)$ does not contain any interval and such that, by a suitable ordering of the steps in the construction of the set $L'(A)$ from the interval A by the scheme above, the 3-condition will be satisfied. This can be achieved easily e.g. so, that from each interval contained in $L(A)$, we delete countably many open intervals in a manner analogous to the construction of the Cantor discontinuum (as far as the ratios of the lengths of the corresponding intervals are concerned). Then we construct the set $L'(A)$ in the following way. We shall delete from the interval A all the intervals which have been deleted by the construction of the set $L(A)$ and all the intervals which were deleted from the intervals contained in the set $L(A)$ in such an order, that at each step we delete from each interval to be divided a longest possible to be deleted.

In the same manner we construct the sets $L'(B), L'(C)$. If $L(B)$ does not contain any interval, then, of course, $L'(B) = L(B)$; similarly in the case of $L'(C)$. From the first part of this proof then follows that $\delta = \alpha' + \beta' + \gamma'$, where $\alpha' \in L'(A) \subset L(A), \beta' \in L'(B) \subset L(B), \gamma' \in L'(C) \subset L(C)$.

Proof of Assertion 3'. The proof is completely analogous to the proof of Assertion 2'.

Proof of (1'') from Theorem 1''. We shall show that

$$F_3(0) = L\left(\left[\frac{\sqrt{21}-3}{6}, \frac{\sqrt{21}-3}{2}\right]\right)$$

and $F_3(0)$ satisfies the 3-condition.

We put $I_0 = \left[\frac{1}{6}(\sqrt{21}-3), \frac{1}{2}(\sqrt{21}-3)\right] = [(0; \overline{3, 1}), (0; \overline{1, 3})]$. From I_0 we delete the open interval $((0; \overline{2, 3, 1}), (0; \overline{1, 1, 3}))$, whose endpoints belong to the set $F_3(0)$, but which itself contains no point of $F_3(0)$. We get two closed intervals $I_1^{(1)} = [(0; \overline{3, 1, 3}), (0; \overline{2, 3, 1})]$, $I_1^{(2)} = [(0; \overline{1, 1, 3}), (0; \overline{1, 3, 1})]$, such that their union covers $F_3(0)$. From the interval $I_1^{(1)}$ we delete the open interval $((0; \overline{3, 3, 1}), (0; \overline{2, 1, 3}))$, from the interval $I_1^{(2)}$ we delete the open interval $((0; \overline{1, 1, 1, 3}), (0; \overline{1, 2, 3, 1}))$. Thus we get four closed intervals

$$\begin{aligned} I_2^{(1,1)} &= [(0; \overline{3, 1, 3}), (0; \overline{3, 3, 1})], & I_2^{(1,2)} &= [(0; \overline{2, 1, 3}), (0; \overline{2, 3, 1})], \\ I_2^{(2,1)} &= [(0; \overline{1, 1, 3, 1}), (0; \overline{1, 1, 1, 3})], \\ I_2^{(2,2)} &= [(0; \overline{1, 2, 3, 1}), (0; \overline{1, 3, 1, 3})], \end{aligned}$$

whose endpoints belong to $F_3(0)$ and such that the union covers $F_3(0)$. We show generally that anyone of the intervals $I_n^{(i_1 \dots i_n)}$ will have endpoints of one of the following two types:

$$(34) \quad (0; \overline{a_1, \dots, a_r, 1, 3}), \quad (0; \overline{a_1, \dots, a_r, 3, 1}),$$

$$(35) \quad (0; \overline{a_1, \dots, a_r, 3, 1, 3}), \quad (0; \overline{a_1, \dots, a_r, 2, 3, 1}).$$

Here, of course, the points on the left can be both left and right endpoints; this depends on the parity of r . We remark first of all that I_0 has endpoints of the type (34) (with $r = 0$). When we have an interval with endpoints of the type (34), then after deleting the open interval with endpoints $(0; \overline{a_1, \dots, a_r, 1, 1, 3}), (0; \overline{a_1, \dots, a_r, 2, 3, 1})$ we get two closed intervals, the one with endpoints $(0; \overline{a_1, \dots, a_r, 1, 3, 1}), (0; \overline{a_1, \dots, a_r, 1, 1, 3})$ (of type (34)), the other with endpoints $(0; \overline{a_1, \dots, a_r, 2, 3, 1}), (0; \overline{a_1, \dots, a_r, 3, 1, 3})$ (of type (35)). When we have an interval with endpoints of the type (35), then after deleting the open interval with endpoints $(0; \overline{a_1, \dots, a_r, 3, 3, 1}), (0; \overline{a_1, \dots, a_r, 2, 1, 3})$ we get two closed intervals, the one with endpoints $(0; \overline{a_1, \dots, a_r, 3, 1, 3}), (0; \overline{a_1, \dots, a_r, 3, 3, 1})$ (of type (34)), the other with endpoints $(0; \overline{a_1, \dots, a_r, 2, 1, 3}), (0; \overline{a_1, \dots, a_r, 2, 3, 1})$ (of type (34)). It is easily seen that by this process we get from the interval $[\frac{1}{6}(\sqrt{21}-3), \frac{1}{2}(\sqrt{21}-3)]$ just the set $F_3(0)$.

Case A. The division of an interval with endpoints of type (34). We must prove the following:

$$\begin{aligned} (36) \quad & |(0; \overline{a_1, \dots, a_r, 1, 3}) - (0; \overline{a_1, \dots, a_r, 3, 1})| \\ & \leq |(0; \overline{a_1, \dots, a_r, 1, 3, 1}) - (0; \overline{a_1, \dots, a_r, 1, 1, 3})| + \\ & \quad + 3|(0; \overline{a_1, \dots, a_r, 2, 3, 1}) - (0; \overline{a_1, \dots, a_r, 3, 1, 3})|, \\ (37) \quad & |(0; \overline{a_1, \dots, a_r, 1, 3}) - (0; \overline{a_1, \dots, a_r, 3, 1})| \\ & \leq 3|(0; \overline{a_1, \dots, a_r, 1, 3, 1}) - (0; \overline{a_1, \dots, a_r, 1, 1, 3})| + \\ & \quad + |(0; \overline{a_1, \dots, a_r, 2, 3, 1}) - (0; \overline{a_1, \dots, a_r, 3, 1, 3})|. \end{aligned}$$

The inequalities (36) and (37), respectively, are easily seen to be equivalent to the following inequalities, (36') and (37'), respectively.

$$\begin{aligned} (36') \quad & \left| \frac{(0; \overline{a_1, \dots, a_r, 1, 1, 3}) - (0; \overline{a_1, \dots, a_r, 2, 3, 1})}{(0; \overline{a_1, \dots, a_r, 2, 3, 1}) - (0; \overline{a_1, \dots, a_r, 3, 1, 3})} \right| \leq 2, \\ (37') \quad & \left| \frac{(0; \overline{a_1, \dots, a_r, 1, 1, 3}) - (0; \overline{a_1, \dots, a_r, 2, 3, 1})}{(0; \overline{a_1, \dots, a_r, 1, 3, 1}) - (0; \overline{a_1, \dots, a_r, 1, 1, 3})} \right| \leq 2. \end{aligned}$$

In the following, we shall denote by $x_{r-1}/y_{r-1}, x_r/y_r$ the last two convergents of the number $(0; \overline{a_1, \dots, a_r})$ (for $r = 0$ we set $x_{-1} = 1, y_{-1} = 0$). As well known, we have

$$(38) \quad |x_r y_{r-1} - x_{r-1} y_r| = 1.$$

We shall put

$$(39) \quad \xi = \overline{(1; 3)} = \frac{\sqrt{21}+3}{6},$$

so that $\overline{(3; 1)} = 3\xi, 3\xi^2 = 3\xi+1$; and

$$(40) \quad t = \frac{y_{r-1}}{y_r},$$

so that $0 \leq t \leq 1$.

Each of the continued fractions occurring on the left-hand sides of (36'), (37') can be expressed in terms of $x_{r-1}, x_r, y_{r-1}, y_r, \xi$. We have

$$\begin{aligned} (41) \quad & (0; \overline{a_1, \dots, a_r, 1, 1, 3}) = \frac{(3\xi-2)x_r + x_{r-1}}{(3\xi-2)y_r + y_{r-1}}, \\ & (0; \overline{a_1, \dots, a_r, 2, 3, 1}) = \frac{(\xi+1)x_r + x_{r-1}}{(\xi+1)y_r + y_{r-1}}, \\ & (0; \overline{a_1, \dots, a_r, 3, 1, 3}) = \frac{3\xi x_r + x_{r-1}}{3\xi y_r + y_{r-1}}, \\ & (0; \overline{a_1, \dots, a_r, 1, 3, 1}) = \frac{\xi x_r + x_{r-1}}{\xi y_r + y_{r-1}}. \end{aligned}$$

When we substitute these expressions into the inequalities (36'), (37'), we can easily see (using (38)) that the relations (36'), (37') are equivalent to the following inequalities:

$$(36'') \quad \frac{3-2\xi}{2\xi-1} \frac{3\xi+t}{3\xi-2+t} \leq 2,$$

$$(37'') \quad \frac{3-2\xi}{2\xi-2} \frac{\xi+t}{\xi+1+t} \leq 2.$$

Since $0 \leq t \leq 1$, the expression on the left-hand side of (36'') is at most

$$\frac{3-2\xi}{2\xi-1} \frac{3\xi}{3\xi-2} = \frac{\sqrt{21}}{7} < 2.$$

Similarly, the expression on the left-hand side of (37'') is at most

$$\frac{3-2\xi}{2\xi-2} \frac{\xi+1}{\xi+2} = \frac{9\sqrt{21}+1}{68} < 2.$$

Case B. The division of an interval with endpoints of type (35). As in case A, it is sufficient to show that the following inequalities are satisfied:

$$(42) \quad \left| \frac{(0; a_1, \dots, a_r, \overline{3, 3, 1}) - (0; a_1, \dots, a_r, \overline{2, 1, 3})}{(0; a_1, \dots, a_r, \overline{3, 1, 3}) - (0; a_1, \dots, a_r, \overline{3, 3, 1})} \right| \leq 2,$$

$$(43) \quad \left| \frac{(0; a_1, \dots, a_r, \overline{3, 3, 1}) - (0; a_1, \dots, a_r, \overline{2, 1, 3})}{(0; a_1, \dots, a_r, \overline{2, 1, 3}) - (0; a_1, \dots, a_r, \overline{2, 3, 1})} \right| \leq 2.$$

Using the same notation as in case A, we have:

$$(44) \quad (0; a_1, \dots, a_r, \overline{2, 1, 3}) = \frac{(3\xi-1)x_r + x_{r-1}}{(3\xi-1)y_r + y_{r-1}},$$

$$(0; a_1, \dots, a_r, \overline{3, 3, 1}) = \frac{(\xi+2)x_r + x_{r-1}}{(\xi+2)y_r + y_{r-1}}.$$

After using (41), (44) and (38) in (42) and (43), we see easily that the inequalities (42), (43) are equivalent to the inequalities

$$(42') \quad \frac{3-2\xi}{2\xi-2} \frac{\xi+2+t}{3\xi-1+t} \leq 2,$$

$$(43') \quad \frac{3-2\xi}{2\xi-2} \frac{\xi+1+t}{\xi+2+t} \leq 2.$$

Since $0 \leq t \leq 1$, the expression on the left-hand side of (42') is at most

$$\frac{3-2\xi}{2\xi-2} \frac{\xi+2}{3\xi-1} = \frac{36-\sqrt{21}}{30} < 2.$$

Notice that the expression on the left-hand side of (42') is greater than 1 for each t with $0 \leq t \leq 1$; hence using this procedure it is not possible to prove that $F_3(0) + F_3(0)$ contains any interval. Finally, the expression on the left-hand side of (43') is at most

$$\frac{3-2\xi}{2\xi-2} \frac{\xi+2}{\xi+3} < 2.$$

Proof of (3'') from Theorem 2''. We shall show that

$$F_2(0) = L\left(\left[\frac{\sqrt{3}-1}{2}, \sqrt{3}-1\right]\right)$$

and that $F_2(0)$ satisfies the 4-condition.

We put

$$I_0 = [\frac{1}{2}(\sqrt{3}-1), \sqrt{3}-1] = [(0; \overline{2, 1}), (0; \overline{1, 2})].$$

We show that each of the intervals $I_n^{(i_1 \dots i_n)}$ will have endpoints of the type

$$(45) \quad (0; a_1, \dots, a_r, \overline{1, 2}), \quad (0; a_1, \dots, a_r, \overline{2, 1}).$$

We remark, first of all, that I_0 has endpoints of type (45) (with $r=0$). If we have an interval with endpoints of type (45), then after deleting of the open interval with endpoints

$$(0; a_1, \dots, a_r, \overline{1, 1, 2}), \quad (0; a_1, \dots, a_r, \overline{2, 2, 1})$$

we get two closed intervals, one with endpoints $(0; a_1, \dots, a_r, \overline{1, 2, 1})$, $(0; a_1, \dots, a_r, \overline{1, 1, 2})$, the other with endpoints $(0; a_1, \dots, a_r, \overline{2, 2, 1})$, $(0; a_1, \dots, a_r, \overline{2, 1, 2})$, thus both of type (45).

Analogously, it suffices to show that the following inequalities hold:

$$(46) \quad \left| \frac{(0; a_1, \dots, a_r, \overline{1, 1, 2}) - (0; a_1, \dots, a_r, \overline{2, 2, 1})}{(0; a_1, \dots, a_r, \overline{1, 2, 1}) - (0; a_1, \dots, a_r, \overline{1, 1, 2})} \right| \leq 3,$$

$$(47) \quad \left| \frac{(0; a_1, \dots, a_r, \overline{1, 1, 2}) - (0; a_1, \dots, a_r, \overline{2, 2, 1})}{(0; a_1, \dots, a_r, \overline{2, 2, 1}) - (0; a_1, \dots, a_r, \overline{2, 1, 2})} \right| \leq 3.$$

We shall use the symbols $x_r, y_r, x_{r-1}, y_{r-1}, t$ in the same meaning as in the preceding proof. Instead of ξ we introduce

$$\eta = \overline{(1; 2)} = \frac{1 + \sqrt{3}}{2}.$$

Then we can write

$$(48) \quad \begin{aligned} (0; a_1, \dots, a_r, 1, \overline{1, 2}) &= \frac{(2\eta - 1)x_r + x_{r-1}}{(2\eta - 1)y_r + y_{r-1}}, \\ (0; a_1, \dots, a_r, 2, \overline{2, 1}) &= \frac{(\eta + 1)x_r + x_{r-1}}{(\eta + 1)y_r + y_{r-1}}, \\ (0; a_1, \dots, a_r, 1, \overline{2, 1}) &= \frac{\eta x_r + x_{r-1}}{\eta y_r + y_{r-1}}, \\ (0; a_1, \dots, a_r, 2, \overline{1, 2}) &= \frac{2\eta x_r + x_{r-1}}{2\eta y_r + y_{r-1}}. \end{aligned}$$

After using (48) in (46) and (47), we can convince ourselves that the inequalities (46), (47) are equivalent to

$$(46') \quad \frac{2 - \eta}{\eta - 1} \frac{\eta + t}{\eta + 1 + t} \leq 3,$$

$$(47') \quad \frac{2 - \eta}{\eta - 1} \frac{2\eta + t}{2\eta - 1 + t} \leq 3.$$

The expression on the left-hand side of (46') attains its maximum in the interval $0 \leq t \leq 1$ at $t = 1$ and the maximum is obviously less than 3. The expression on the left-hand side of (47') attains its maximum at $t = 0$:

$$\frac{2 - \eta}{\eta - 1} \frac{2\eta}{2\eta - 1} = \sqrt{3} + 1 < 3.$$

Notice that the expression on the left-hand side of (47') is greater than 2 if $0 \leq t \leq 1$: hence it is not possible to show by this procedure that $F_2(0) + F_2(0) + F_2(0)$ contains any interval.

Analogous results concerning the products of continued fractions will be published later.

Note added in proof. The results of this paper were obtained in the seminar of Prof. Jarník in 1968. In connection with similar questions, T. W. Cusick and R. A. Lee have arrived recently at general statements about sums of discontinui which make it possible to give another proof of our results (see [2] and [3]).

Bibliography

- [1] M. Hall, Jr., *On the sum and product of continued fractions*, Annals of Math. 48 (1947), pp. 966-993.
- [2] T. W. Cusick and R. A. Lee, *Sums of sets of continued fractions*, Proc. AMS 30 (1971), pp. 241-246.
- [3] T. W. Cusick, *On M. Hall's continued fraction theorem*, Proc. AMS (to appear, March 1973).

DEPARTMENT OF MATHEMATICS
OHIO STATE UNIVERSITY
Columbus, Ohio

Received on 5. 11. 1971

(235)