

On the Uniqueness of the Equilibrium State for Plane Rotators

J. Bricmont and J. R. Fontaine

Institut de Physique Théorique, U.C.L., Université de Louvain, B-1348 Louvain-la-Neuve, Belgium

L. J. Landau

Department of Mathematics, Bedford College, London NW1 4NS, England

Abstract. We study the classical statistical mechanics of the plane rotator, and show that there is a unique translation invariant equilibrium state in zero external field, if there is no spontaneous magnetization. Moreover, this state is then extremal in the equilibrium states. In particular there is a unique phase for the two dimensional rotator, and a unique phase for the three dimensional rotator above the critical temperature. It is also shown that in a sufficiently large external field the Lee-Yang theorem implies uniqueness of the equilibrium state.

1. Introduction

Some new results have been obtained recently concerning the classical statistical mechanics of the plane rotator model, defined by the Hamiltonian

$$H = -\beta \sum J_{ij} \sigma_i \cdot \sigma_j \quad J_{ij} \geq 0 \quad (1.1)$$

where σ_i is a two-dimensional vector of unit length. For $d \geq 3$, it has been proven [6] that spontaneous magnetization occurs for large β . For $d = 2$, it is well-known that there is no spontaneous magnetization for sufficiently short range interactions [17]; moreover Dobrushin and Shlosman have proven that for finite range interactions all the equilibrium states are invariant under the action of $SO(2)$ on the configuration space [2]. One may ask: does the absence of spontaneous magnetization imply uniqueness of the equilibrium state, as for example in the Ising model [16, 19]? We show in the present paper that it implies at least uniqueness of the translation invariant equilibrium state and that the latter cannot be decomposed into non-invariant equilibrium states. Compared with the similar problem in the Ising model [16, 19], our method looks more complicated; the reason is that for this model, we don't have correlation inequalities comparable to the F.K.G. inequalities [5] which are valid for all the different boundary conditions. Instead of considering boundary conditions, we characterize invariant equilibrium states as tangents to the pressure [9]. We introduce various perturbations to the pressure and control the derivatives of these perturbed pressures with correlation inequalities.

In a sufficiently large external field, the correlation inequalities are sufficient to control all boundary conditions, and together with the Lee-Yang theorem gives uniqueness of the equilibrium state.

The extension to models where σ_i has more than two components seems difficult because of the lack of the necessary correlation inequalities.

Acknowledgements. It is a pleasure to thank F. Dunlop and J. Slawny for helpful discussions, and A. Messager, S. Miracle, and Ch. E. Pfister for providing us with Lemma 3.2 [24] which leads to a simplification of our original proof of the extremality of the unique phase.

2. The Model

At each point i of the lattice \mathbb{Z}^d is associated a spin variable $\sigma_i \in \mathbb{R}^2$. We use two parametrizations, given by:

$$\sigma_i = (s_i, t_i) = (r_i \cos \phi_i, r_i \sin \phi_i)$$

where

$$s_i, t_i \in \mathbb{R}, \quad r_i \in \mathbb{R}_+, \quad \phi_i \in [0, 2\pi[.$$

The a priori probability distribution for σ_i is assumed rotation invariant:

$$dv_0(\sigma_i) = d\lambda(r_i) d\phi_i$$

and the spin is bounded: $|\sigma_i| \leq b$ with probability 1. In addition the measure v_0 satisfies Condition A, which is important for the development of correlation inequalities:

Condition A.

$$\forall \alpha, \beta, \gamma, \delta \in \mathbb{N}$$

$$\int (s+s')^\alpha (s-s')^\beta (t+t')^\gamma (t-t')^\delta dv_0(s, t) dv_0(s', t') \geq 0.$$

Remark. 1) The following measures (suitably normalized) are known to satisfy Condition A [1, 3, 12, 18, 22]:

i) $\delta(r-b) dr d\phi \quad b > 0$ (fixed length)

ii) $\chi(r \in [0, b]) r dr d\phi \quad b > 0$ (uniform distribution)

iii) $\exp(-ar^4 + cr^2) \chi(r \in [0, b]) r dr d\phi \quad a, b > 0.$

The interaction between spins is given by a translation invariant, ferromagnetic pair interaction

$$H = - \sum J_{ij} \sigma_i \cdot \sigma_j$$

where

$$(2.1)$$

$$J_{ij} = J(i-j) \geq 0.$$

We consider the case of finite range interactions:

$$J_{ij} = 0 \quad \text{if } |i-j| > L.$$

However our results are not sensitive to this restriction and carry over to long range interactions with sufficiently rapid fall-off.

If A is a finite region, an exterior configuration ω is a specification of the spins in A_c , the complement of A , and a boundary condition is a probability measure $\mu_{bc}(\omega)$ on the exterior configurations. A particularly useful boundary condition is denoted s -b.c. for which all spins in A_c are in the s -direction with maximum value:

$$\text{for all } i \in A_c \quad \sigma_i = \left(s_i = \max_{\text{supp } \lambda} r, t_i = 0 \right).$$

The boundary condition t -b.c. is obtained by the interchange of s and t variables.

An exterior configuration ω induces an effective Hamiltonian for the region A given by

$$H_{A,\omega} = - \sum_{i,j \in A} J_{ij} \sigma_i \cdot \sigma_j - \sum_{\substack{i \in A \\ j \in A_c}} J_{ij} \sigma_i \cdot \sigma_j(\omega)$$

where $\sigma_j(\omega)$ takes the value specified by ω .

The joint probability distribution of the spins in the region A is given by the Boltzmann factor

$$d\mu_{A,\omega} = Z_{A,\omega}^{-1} \exp(-H_{A,\omega}) \prod_{i \in A} dv_0(\sigma_i)$$

where

$$Z_{A,\omega} = \int \exp(-H_{A,\omega}) \prod_{i \in A} dv_0(\sigma_i).$$

Expectations with respect to $\mu_{A,\omega}$ will be denoted $\langle \rangle_{A,\omega}$. For a general boundary condition

$$\langle \rangle_{A,bc} = \int \langle \rangle_{A,\omega} d\mu_{bc}(\omega).$$

The pressure

$$p_{A,\omega} = |A|^{-1} \ln Z_{A,\omega}$$

and for a general boundary condition

$$p_{A,bc} = \int p_{A,\omega} d\mu_{bc}(\omega).$$

The pressure

$$p = \lim_{A \nearrow \mathbb{Z}^d} p_{A,bc}$$

and is independent of the boundary condition [20].

An equilibrium state is an infinite volume limit of finite region states $\langle \rangle_{A,bc}$ with some boundary condition, or equivalently a probability measure on the infinite volume configuration space satisfying the DLR equations [9, 13]. We will generally use the symbol $\langle \rangle$ for a finite region state and the symbol q for an infinite volume equilibrium state. The set of equilibrium states for the Hamiltonian H is denoted $\mathcal{A}(H)$. A phase is an equilibrium state invariant under the lattice translations, and the set of phases is denoted $\mathcal{A}_T(H)$.

In an external field hn the Hamiltonian (2.1) is modified to

$$\begin{aligned} H_h &= - \sum J_{ij} \sigma_i \cdot \sigma_j - \sum hn \cdot \sigma_i \\ &= - \sum J_{ij} (s_i s_j + t_i t_j) - \sum (h_s s_i + h_t t_i). \end{aligned} \tag{2.2}$$

Definition. The spontaneous magnetization M is defined by

$$M = \lim_{h \searrow 0} \varrho_h(\mathbf{n} \cdot \boldsymbol{\sigma}_i)$$

where

$$\varrho_h \in \Delta_I(H_h).$$

Remark. By translation invariance of the phase ϱ_h , M is independent of i ; by rotation covariance, M is independent of the direction \mathbf{n} . Using the fact that the pressure is a convex function of h and the relation between phases and tangents to the graph of the pressure [9] it is seen that M is the right-derivative of the pressure with respect to h at $h=0$ and is independent of the choice of phase $\varrho_h \in \Delta_I(H_h)$.

Our basic results can now be formulated:

2.1. Theorem. *If the spontaneous magnetization $M=0$ then there is a unique phase in zero external field. Moreover, this phase is then extremal in the equilibrium states. This phase is also the unique quasi-periodic state.*

2.2. Theorem. *Let $d\lambda(r)=\delta(r-b)dr$. Then for a large enough external field the equilibrium state is unique.*

2.3. Corollary. *If the lattice dimension $d=2$, there is a unique phase in zero external field. If $d=3$, there is a unique phase for $T>T_c$ where T_c is the critical temperature for spontaneous magnetization.*

We point out that for simplicity of presentation we have not given each theorem or lemma in its most general form. We discuss extensions in Section 6.

3. Inequalities

Let A be a finite subset of \mathbb{Z}^d with multiplicities, and define

$$s_A = \prod_{i \in A} s_i.$$

For example $s_i^2 s_j = s_A$ where $A = \{i, i, j\}$.

$|A|$ denotes the (finite) cardinality of A . Similarly we define t_A and r_A . If N is a function from \mathbb{Z}^d to \mathbb{Z} which is equal to zero except at a finite number of points, we define

$$N\phi = \sum n_i \phi_i.$$

Note that if $|A|$ is even, $s_A \pm t_A$ may be expressed in terms of cosines.

$$s_A + t_A = 2^{(1-|A|)} r_A \sum_M \cos M\phi \tag{3.1}$$

and similarly for $s_A - t_A$: We decompose $s_A \pm t_A$ into a sum of products of $r_i r_j$ ($\cos \phi_i \cos \phi_j \pm \sin \phi_i \sin \phi_j = r_i r_j \cos(\phi_i \mp \phi_j)$) and then use the identity $\cos N\phi \cos M\phi = \frac{1}{2} [\cos(N+M)\phi + \cos(N-M)\phi]$.

We denote by $\text{supp } A$ the set A without multiplicities:

$$\text{supp } A = \{i \in \mathbb{Z}^d : i \in A\}.$$

Also

$$\text{supp } N = \{i \in \mathbb{Z}^d : n_i \neq 0\}.$$

We define the algebra \mathfrak{A} as the set of finite linear combinations of the functions $s_A t_B$, and the even and odd subspaces $\mathfrak{A}_e, \mathfrak{A}_o$ as linear combinations of $\{r_A \cos N\phi\}$, respectively $\{r_A \sin N\phi\}$. Equivalently these subspaces are defined as linear combinations of $\{s_A t_B\}$ with $|B|$ even, respectively odd. The subspaces may equally well be defined according to transformation properties under the transformation $t_i \rightarrow -t_i \forall i$ ($\phi_i \rightarrow -\phi_i \forall i$). Consider the interaction (2.2) with the external field in the positive s -direction. The Hamiltonian has the form

$$H = - \sum J_{ij} r_i r_j \cos(\phi_i - \phi_j) - h \sum r_i \cos \phi_i.$$

With s -b.c. the effective interaction in the region A has the form

$$H_{A,s} = - \sum_{i,j \in A} J_{ij} r_i r_j \cos(\phi_i - \phi_j) - \sum_{i \in A} \alpha_i r_i \cos \phi_i$$

where $\alpha_i \geq 0$. Then Ginibre's inequalities hold [Appendix, Theorems A1, A4]

$$\langle r_A \cos N\phi r_B \cos M\phi \rangle_{s,A} \geq \langle r_A \cos N\phi \rangle_{s,A} \langle r_B \cos M\phi \rangle_{s,A}. \tag{3.2}$$

Now consider the interaction (2.2) with the external field such that $h_s, h_t \geq 0$. Then the generalized Griffiths' inequalities hold [Theorems A2, A3]

$$\begin{aligned} \langle s_A s_B \rangle_{s,A} &\geq \langle s_A \rangle_{s,A} \langle s_B \rangle_{s,A} \\ \langle t_A t_B \rangle_{s,A} &\geq \langle t_A \rangle_{s,A} \langle t_B \rangle_{s,A} \\ 0 &\leq \langle s_A t_B \rangle_{s,A} \leq \langle s_A \rangle_{s,A} \langle t_B \rangle_{s,A}. \end{aligned} \tag{3.3}$$

In addition there are comparison inequalities relating different states [Theorems A3, A4].

We derive here some basic results which will be useful in the proofs of Theorems 2.1 and 2.2. The discussion will be given for the interaction (2.1).

3.1. Theorem. $q_s \equiv \lim_{A \nearrow \mathbb{Z}^d} \langle \cdot \rangle_{s,A}$ exists by monotonicity and is an extremal equilibrium state.

Remark. We note a useful property of the state q_s . By monotonicity in both h and A it follows that $M = q_s(s_i)$.

The proof of Theorem 3.1 is based on ¹.

3.2. Lemma. Let ω be any exterior configuration for the region A . Then

$$\langle r_A \cos M\phi \rangle_{s,A} \geq \langle r_A \cos M\phi \rangle_{\omega,A}.$$

Proof. We must show

$$\int (r_A \cos N\phi - r'_A \cos N\phi') d\mu_s(\{r_i, \phi_i\}) d\mu_\omega(\{r'_i, \phi'_i\}) \geq 0.$$

¹ Lemma 3.2 is due to A. Messager, S. Miracle, and Ch. E. Pfister. It simplifies our original proof of Theorem 3.1

This is proved in the same way as Ginibre's inequalities [7] once the effective Hamiltonians are combined as

$$-H_{\Lambda,s} - H_{\Lambda,\omega} = \sum J_{ij} [r_i r_j \cos(\phi_i - \phi_j) + r'_i r'_j \cos(\phi'_i - \phi'_j)] + \sum \alpha_i r_i \cos \phi_i + \alpha'_i r'_i \cos(\phi'_i + \psi_i)$$

where

$$|\alpha'_i| \leq \alpha_i.$$

This can be written as a sum with positive coefficients of products of $(r_i \pm r'_i)$, $\cos\left(\frac{\phi'_i + \phi_i + \psi}{2}\right) \cos\left(\frac{\phi'_i - \phi_i + \psi}{2}\right)$ and similar terms with sine instead of cosine. $(r_A \cos M\phi - r'_A \cos M\phi')$ can be written similarly.

Then one expands the exponential in series and, due to the above decompositions, each term is a product of integrals over r_i, r'_i and over ϕ_i, ϕ'_i . The integral over ϕ_i, ϕ'_i factorizes into a product of two integrals of two identical functions, one in the variables $(\phi'_i + \phi_i)$ and the other in $(\phi'_i - \phi_i)$. Being a square, this expression is positive. The integral over the variables r_i, r'_i reduces to a product of integrals of the type:

$$\int (r_i + r'_i)^{a_i} (r_i - r'_i)^{b_i} d\lambda_i(r_i) d\lambda_i(r'_i).$$

This integral obviously vanishes if b_i is odd and is positive if b_i is even. □

Definition. We say the state ϱ is *clustering* if for all $f, g \in \mathfrak{A}$

$$\varrho(f\tau_a g) - \varrho(f)\varrho(\tau_a g) \rightarrow 0 \quad \text{as } |a| \rightarrow \infty$$

where τ_a denotes a translation by the amount a .

Proof of Theorem 3.1. From Lemma 3.2 it follows that $\langle r_A \cos M\phi \rangle_{s,A} (\geq 0)$ decreases as $\Lambda \nearrow \mathbb{Z}^d$. Thus the limit exists. Since $\langle r_A \sin M\phi \rangle_{s,A} = 0$ the state ϱ_s is well-defined. Again from Lemma 3.2 it follows that

$$\varrho(r_A \cos N\phi) \leq \varrho_s(r_A \cos N\phi) \quad \text{for all } \varrho \in \Delta(H),$$

which shows that ϱ_s is extremal as a state on the even subspace \mathfrak{A}_e .

The extremality of ϱ_s on the even subspace \mathfrak{A}_e implies that ϱ_s is clustering on \mathfrak{A}_e . Indeed, since $\Delta(H)$ is a metrizable Choquet simplex [9], there is a unique probability measure ω_s carried by the extremal elements of $\Delta(H)$ such that

$$\varrho_s(f) = \int \sigma(f) d\omega_s(\sigma), \quad \forall f \in \mathfrak{A}.$$

By Lemma 3.2 it follows that $\varrho_s \upharpoonright \mathfrak{A}_e = \sigma \upharpoonright \mathfrak{A}_e \omega_s$ -a.e. Since σ is clustering, ϱ_s is clustering on \mathfrak{A}_e . By correlation inequalities it follows that ϱ_s is clustering on the full algebra \mathfrak{A} : (Theorem A5 or [3])

$$|\varrho_s(s_A t_B s_C t_D) - \varrho_s(s_A t_B) \varrho_s(s_C t_D)| \leq (\varrho_s(s_A s_B s_C s_D) - \varrho_s(s_A s_B) \varrho_s(s_C s_D)) \text{ if } |B| \text{ is even}$$

and

$$\varrho_s(s_A t_B s_C t_D)^2 \leq (\varrho_s(s_A s_B s_C s_D)^2 - \varrho_s(s_A s_B)^2 \varrho_s(s_C s_D)^2) \text{ if } |B| \text{ is odd.}$$

² Lemma 3.2 holds for all exterior configurations and hence for any boundary condition b . Take b as the boundary condition induced by s -b.c. in Λ'_i where $\Lambda' \supset \Lambda$

By construction, the restriction of ϱ_s to the odd subspace \mathfrak{A}_0 is identically zero. Thus the extremality of ϱ_s follows from the following lemma.

3.3. Lemma. (a) Let ϱ, σ be two states on \mathfrak{A} satisfying

- (i) ϱ, σ are clustering ;
- (ii) $\varrho/\mathfrak{A}_e = \sigma/\mathfrak{A}_e$;
- (iii) $\varrho/\mathfrak{A}_0 = 0$,

then $\lim_{i \rightarrow \infty} \sigma(\tau_i(f)) = 0, \forall f \in \mathfrak{A}_0$.

(b) Let ω be a translation invariant probability measure on a set E of states σ , satisfying $\lim_{i \rightarrow \infty} \sigma(\tau_i(f)) = 0$ ω -a.e. for some $f \in \mathfrak{A}$.

Then $\sigma(f) = 0$ ω -a.e.

(c) Let $\varrho = \int_E \sigma d\omega(\sigma)$ be a decomposition of the invariant state ϱ into a set of states E , with the measure ω translation invariant. Suppose ϱ and ω -a.e. σ satisfy the hypothesis in (a). Then $\varrho = \sigma$ ω -a.e.

Proof. (a) By (i) and (iii)

$$\lim_{i \rightarrow \infty} \varrho(f\tau_i(g)) = 0 \quad \forall f, g \in \mathfrak{A}_0$$

and by (ii)

$$\lim_{i \rightarrow \infty} \sigma(f\tau_i(g)) = 0.$$

Therefore by (i) $\sigma(f) \lim_{i \rightarrow \infty} \sigma(\tau_i(g)) = 0 \quad \forall f, g \in \mathfrak{A}_0$ and the conclusion follows.

(b) Since ω is translation invariant, $\int_E |\sigma(f)| d\omega(\sigma) = \int_E |\sigma(\tau_i(f))| d\omega(\sigma), \forall i \in \mathbb{Z}^d$ and by dominated convergence, $(|\sigma(\tau_i(f))|) \leq \|f\|$

$$\lim_{i \rightarrow \infty} \int_E |\sigma(\tau_i(f))| d\omega(\sigma) = 0.$$

So $\sigma(f) = 0$ ω -a.e.

(c) By points (a) and (b) $\sigma(f) = 0$ ω -a.e. for each $f \in \mathfrak{A}_0$. Since \mathfrak{A}_0 is separable, we conclude $\sigma/\mathfrak{A}_0 = 0$ ω -a.e. and $\sigma = \varrho$ ω -a.e. \square

We now begin the argument leading to the proof of Theorem 2.1.

There is a useful “bootstrap” principle which allows one to conclude that certain higher order correlation functions are zero if certain lower order ones are zero. In particular we shall conclude from $M = 0$ and Lemma 3.2 that the equilibrium states have certain symmetry properties.

3.4. Theorem. Let ϱ be any weak* limit point of the finite volume states $\langle \cdot \rangle_A$ as $A \nearrow \mathbb{Z}^d$, where the effective Hamiltonian for the region A has the form

$$-H_A = \sum_{i,j \in A} J_{ij}(s_i s_j + t_i t_j) + \sum_{i \in A} \alpha_i s_i + \beta_i t_i \tag{3.4}$$

with $\alpha_i \geq |\beta_i|$.

Then if $\varrho(s_i - t_i) = 0$ for all i , it follows that $\varrho(s_A - t_A) = 0$ for all A .

Proof. Define the random variables $x_i = 2^{-\frac{1}{2}}(s_i + t_i)$, $y_i = 2^{-\frac{1}{2}}(s_i - t_i)$. The effective Hamiltonian in terms of the variables x, y is

$$\sum J_{ij}(x_i x_j + y_i y_j) + 2^{-\frac{1}{2}} \sum (\alpha_i + \beta_i)x_i + (\alpha_i - \beta_i)y_i.$$

Also the a priori measure for (x, y) is the same as for (s, t) by rotation invariance.

Therefore the generalized Griffiths' inequalities are valid (Theorem A2) and these extend to the limit point ϱ :

$$0 \leq \varrho(x_B y_C) \leq \varrho(x_B)\varrho(y_C). \tag{3.5}$$

We first show: Given n (odd), if $\varrho(y_A) = 0$ for all $|A|$ (odd) $< n$ then $\varrho(s_A) = \varrho(t_A)$ for all $|A| < n$.

Indeed³

$$\varrho(s_A - t_A) = 2^{1 - \frac{1}{2}|A|} \sum_{\substack{B \cup C = A \\ |C| \text{ odd}}} \varrho(x_B y_C) = 0 \quad \text{if } |A| < n$$

by (3.5) and the hypothesis $\varrho(y_C) = 0$.

We next show: $\varrho(y_A) = 0$ for all $|A|$ odd. Indeed, by hypothesis $\varrho(s_i) = \varrho(t_i)$ for all i and so $\varrho(y_i) = 0$ for all i . The proof proceeds by induction. Let n (odd) be given and suppose $\varrho(y_A) = 0$ for all $|A|$ (odd) $< n$. To show $\varrho(y_A) = 0$ if $|A| = n$ we write

$$\begin{aligned} \varrho(y_A) &= 2^{-\frac{1}{2}|A|} \sum_{B \cup C = A} \varrho(s_B t_C) (-1)^{|C|} \\ &= 2^{-\frac{1}{2}|A|} \left\{ \varrho(s_A - t_A) + \sum_{\substack{B \cup C = A \\ |B| \text{ odd} \\ |B| < |A|}} \varrho(s_B t_C) - \varrho(s_C t_B) \right\}. \end{aligned}$$

But

$$\begin{aligned} |\varrho(s_B t_C) - \varrho(s_C t_B)| &= |\varrho(s_B t_C) - \varrho'(s_B t_C)| \\ &\leq \varrho(s_B)\varrho'(t_C) - \varrho'(s_B)\varrho(t_C) \\ &= \varrho(s_B)\varrho(s_C) - \varrho(t_B)\varrho(t_C) \end{aligned}$$

by a comparison inequality (A6) where ϱ' denotes the state obtained from ϱ by the interchange of s and t variables⁴.

Since $\varrho(s_B) = \varrho(t_B)$, $\varrho(s_C) = \varrho(t_C)$ by the induction hypothesis, we have

$$\varrho(y_A) = 2^{-\frac{1}{2}|A|} \varrho(s_A - t_A) = 2^{1 - |A|} \sum_{\substack{B \cup C = A \\ |C| \text{ odd}}} \varrho(x_B y_C) = 2^{1 - |A|} \varrho(y_A)$$

which implies $\varrho(y_A) = 0$. \square

³ In the summation over subsets of A we distinguish different occurrences of the same lattice point (multiplicities). Otherwise combinatorial factors should be included in the expression

⁴ Note that for ϱ' the effective Hamiltonian for the region A has the form

$$-H'_A = \sum_{i, j \in A} J_{ij}(s_i s_j + t_i t_j) + \sum_{i \in A} \alpha'_i s_i + \beta'_i t_i$$

where $\alpha'_i = \beta_i$, $\beta'_i = \alpha_i$. Since $\alpha_i \geq |\beta_i|$ the hypotheses of comparison Theorem A3 are satisfied

4. Uniqueness of the Phase

We consider the plane rotator in zero external field and suppose there is no spontaneous magnetization. We will show that there is a unique phase. We use the equivalence between phases and tangents to the graph of the pressure [9]. For $f \in \mathfrak{A}$ we consider the perturbation λf to the Hamiltonian H . That is, we consider

$$H_\lambda = H - \lambda \sum \tau_i f$$

where we sum over the lattice translates of f . (Equilibrium states for H_λ are defined as in Section 2, via finite volume states with some effective Hamiltonian $H_{\lambda, A}$, or directly by the DLR equations appropriate for H_λ .) If we can show that the pressure p_λ is differentiable at $\lambda=0$ it follows that all invariant equilibrium states take the same value on f . This may be formulated as

4.1. Lemma. *Let there exist a sequence of positive numbers $(\lambda_n)_{n \in \mathbb{N}}$ and another one of negative numbers $(\lambda'_n)_{n \in \mathbb{N}}$, both converging to zero and invariant equilibrium states $q_{\lambda'_n}, q_{\lambda_n}$ of $H_{\lambda'_n}, H_{\lambda_n}$ such that $\lim_{n \rightarrow \infty} q_{\lambda'_n}(f) = \lim_{n \rightarrow \infty} q_{\lambda_n}(f)$.*

Then all the $q \in \Delta_I(H)$ take the same value on f ($= \lim_{n \rightarrow \infty} q_{\lambda_n}(f)$).

Proof. The equivalence between invariant equilibrium states and tangents to the graph of the pressure [9], together with the hypothesis and the convexity of P_λ shows that P_λ is differentiable at $\lambda=0$ and this is equivalent to the conclusion [9, 13]. \square

We shall apply Lemma 4.1 to various perturbations and first to $\{s_A, t_A\}$. Once we have shown uniqueness on these functions (Lemma 4.2) we use this result to get uniqueness on the even subspace \mathfrak{A}_e .

4.2. Lemma. *All phases take the same value on s_A and on t_A .*

Proof. The proof proceeds in three steps.

a) $q_s(s_A) = q_s(t_A)$ for all A , and $q_s(s_A) = 0$ if $|A|$ odd:

Since $M=0$ Lemma 4.1 implies $q_s(s_i) = 0$. Since $q_s(t_i) = 0$ by construction, we have $q_s(s_i - t_i) = 0$. By the “bootstrap” Theorem 3.4, $q_s(s_A - t_A) = 0$ for all A . Since $q_s(t_A) = 0$ if $|A|$ odd by construction, we may also conclude $q_s(s_A) = 0$ if $|A|$ odd.

b) For all $q \in \Delta(H)$ $q(s_A) = q(t_A)$ for all A and $q(s_A) = 0$ if $|A|$ odd:

By Lemma 3.2 $q(s_A - t_A) \leq q_s(s_A - t_A) = 0$ if $|A|$ even [expanding in terms of cosines—Eq. (3.1)]. Thus by the symmetry of interchanging s and t variables, we conclude

$$q(s_A - t_A) = 0 \quad \text{for all } q \in \Delta(H), |A| \text{ even.}$$

If $|A|$ odd, $q(s_A) \leq q_s(s_A) = 0$. By the symmetry $s \rightarrow -s$, we conclude

$$q(s_A) = 0 \quad \text{for all } q \in \Delta(H), |A| \text{ odd.}$$

By the symmetry of interchanging s and t variables we now conclude

$$q(t_A) = 0 \quad \text{for all } q \in \Delta(H), |A| \text{ odd.}$$

c) All phases take the same value on s_A and on t_A :

Consider the perturbation $\pm \lambda s_A$, $\lambda > 0$.

Let $\varrho_{+\lambda}$ be the (translation invariant) limit (Theorem 3.1) as $\Lambda \nearrow \mathbb{Z}^d$ of $\langle \cdot \rangle_{s,+\lambda,\Lambda}$ and $\varrho_+ = \lim_{\lambda \searrow 0} \varrho_{+\lambda}$ (Griffiths' inequalities). Let $\tilde{\varrho}_{-\lambda}$ be any weak* limit point as $\Lambda \nearrow \mathbb{Z}^d$ of $\langle \cdot \rangle_{s,-\lambda,\Lambda}$ and let $\varrho_{-\lambda}$ be an average over translations of $\tilde{\varrho}_{-\lambda}$. Let ϱ_- be a weak* limit point of $\varrho_{-\lambda}$ as $\lambda \rightarrow 0$. Since by a comparison inequality [Eq. (A4)]

$$\langle s_B \rangle_{s,\lambda,\Lambda} \geq \langle s_B \rangle_{s,-\lambda,\Lambda}$$

it follows that

$$\varrho_+(s_B) \geq \varrho_-(s_B)$$

Similarly

$$\varrho_+(t_B) \leq \varrho_-(t_B).$$

Since $\varrho_{\pm} \in \Delta_I(H)$, it follows by point (b) that $\varrho_{\pm}(s_B - t_B) = 0$ and so $\varrho_+(s_A) = \varrho_-(s_A)$. By Lemma 4.1 all phases agree on s_A . By the symmetry of interchanging s and t variables all phases agree on t_A . \square

4.3. Lemma. *All phases take the same value on the even subspace \mathfrak{A}_e .*

Proof. a) Consider the perturbation $\pm \lambda r_A \cos M\phi$. From a comparison inequality (Theorem A4) we have

$$\langle r_B \cos N\phi \rangle_{s,\lambda,\Lambda} \geq \langle r_B \cos N\phi \rangle_{s,-\lambda,\Lambda} \quad \text{for } \lambda > 0.$$

Taking weak* limit points as in Lemma 4.2 we obtain states ϱ_{\pm} which satisfy

$$\varrho_+(r_B \cos N\phi) \geq \varrho_-(r_B \cos N\phi). \tag{4.1}$$

b) If $|B|$ is even, inequality (4.1) implies

$$\begin{aligned} \varrho_-(s_A t_B) &= \varrho_-(s_A(s_B + t_B)) - \varrho_-(s_A s_B) \\ &\leq \varrho_+(s_A(s_B + t_B)) - \varrho_-(s_A s_B) \end{aligned}$$

and

$$\begin{aligned} \varrho_-(s_A t_B) &= -\varrho_-(s_A(s_B - t_B)) + \varrho_-(s_A s_B) \\ &\geq -\varrho_+(s_A(s_B - t_B)) + \varrho_-(s_A s_B). \end{aligned}$$

Since $\varrho_+(s_A s_B) = \varrho_-(s_A s_B)$ by Lemma 4.2 the above inequalities imply

$$\varrho_-(s_A t_B) \leq \varrho_+(s_A t_B)$$

and

$$\varrho_-(s_A t_B) \geq \varrho_+(s_A t_B).$$

Thus $\varrho_-(s_A t_B) = \varrho_+(s_A t_B)$. In particular $\varrho_+(r_A \cos M\phi) = \varrho_-(r_A \cos M\phi)$. Therefore all phases agree on the even subspace \mathfrak{A}_e by Lemma 4.1. \square

Proof of Theorem 2.1. From Lemma 4.3 $\varrho(f) = \varrho_s(f)$ for all $f \in \mathfrak{A}_e$, and by construction $\varrho_s(f) = 0$ for all f in the odd sub-space \mathfrak{A}_o . If there were a phase ϱ such that, $\varrho(f) \neq 0$ for some $f \in \mathfrak{A}_o$, then defining ϱ' from ϱ by the transformation

$\phi_i \rightarrow -\phi_i \forall i$ we would have a nontrivial decomposition of $\varrho_s = \frac{1}{2}(\varrho + \varrho')$ which contradicts the extremality of ϱ_s (Theorem 3.1). Thus $\varrho(f) = 0$ for all $\varrho \in \Delta_I(H)$ if $f \in \mathfrak{A}_0$. Thus ϱ_s is the unique phase, and by Theorem 3.1 this phase is an extremal equilibrium state. That ϱ_s is also the unique quasi-periodic state follows from the extremality of the unique phase, as in [21, Appendix C]. \square

Proof of Corollary 2.3. If the lattice dimension $d = 2$, we know by the theorem of Mermin [17] that $\lim_{h \searrow 0} \langle \sigma \cdot n \rangle_{h, \text{periodic}} = 0$ with periodic boundary conditions. This implies that $M = 0$ and the corollary follows from Theorem 2.1. If $d = 3$ we define T_c as the lowest temperature such that $M = 0$ for all $T > T_c$. (T_c is finite by the high temperature cluster expansion and $T_c \neq 0$ by [6].) Theorem 2.1 implies the uniqueness of the phase for $T > T_c$. \square

5. Uniqueness of the Equilibrium State

It will be shown that for sufficiently large external field, the plane rotator with fixed length ($d\lambda = \delta(r - b)dr$) has a unique equilibrium state. We define two phases ϱ_M and ϱ_m which suitably bound all equilibrium states. Proving that $\varrho_M = \varrho_m$ then gives the uniqueness of the equilibrium state. We take the external field interaction

$$-\sum (h_s s_i + h_t t_i)$$

for the case $h_s = h_t = h$. (By rotation covariance the result is true for the external field in an arbitrary direction.)

The effective Hamiltonian for the region A has the form (3.4) where α_i, β_i depend on the exterior configuration. If $h \geq b \sum_j J_{ij}$ then $\alpha_i, \beta_i \geq 0$ for all exterior configurations. This allows the use of correlation inequalities for all equilibrium states.

Let

$$A_i = \max_{\omega \in A_c} \alpha_i = \max_{\omega \in A_c} \beta_i$$

$$B_i = \min_{\omega \in A_c} \alpha_i = \min_{\omega \in A_c} \beta_i$$

and define the states $\langle \cdot \rangle_{M,A}$ (resp. $\langle \cdot \rangle_{m,A}$) by taking $\alpha_i = A_i, \beta_i = B_i$ (resp. $\alpha_i = B_i, \beta_i = A_i$).

Note that $\langle \cdot \rangle_{m,A}$ is obtained from $\langle \cdot \rangle_{M,A}$ by the interchange of s and t variables. Then from comparison inequalities (A4, A5) for any exterior configuration ω

$$0 \leq \langle s_B \rangle_{m,A} \leq \langle s_B \rangle_{\omega,A} \leq \langle s_B \rangle_{M,A}$$

$$0 \leq \langle t_B \rangle_{M,A} \leq \langle t_B \rangle_{\omega,A} \leq \langle t_B \rangle_{m,A}$$
(5.1)

and

$$|\langle s_A t_B \rangle_{M,A} - \langle s_A t_B \rangle_{\omega,A}| \leq \langle s_A \rangle_{M,A} \langle t_B \rangle_{\omega,A} - \langle s_A \rangle_{\omega,A} \langle t_B \rangle_{M,A}$$
(5.2)

from (A6).

5.1. Lemma. *The following infinite volume limits exist.*

$$\varrho_M \equiv \lim_{A \nearrow \mathbb{Z}^d} \langle \cdot \rangle_{M,A}, \quad \varrho_m \equiv \lim_{A \nearrow \mathbb{Z}^d} \langle \cdot \rangle_{m,A}.$$

Proof. It follows from Equations (5.1) that⁵

- $\langle s_A \rangle_{M,A}$ decreases as A increases,
- $\langle t_A \rangle_{M,A}$ increases as A increases.

Thus the limits as $A \nearrow \mathbb{Z}^d$ exist. From (5.2) it now follows that $\langle s_A t_B \rangle_{M,A}$ converges as $A \nearrow \mathbb{Z}^d$. By the interchange of s and t variables the same holds for $\langle \cdot \rangle_{m,A}$. \square

Proof of Theorem 2.2. The Lee-Yang theorem for this model has been deduced by Dunlop and Newman [4] from a theorem of Suzuki and Fisher [23]. Although it is not stated in this form in [4], one can deduce from [23] analyticity of the pressure in h_s with h_t fixed, real, and non-zero and analyticity in h_t with h_s fixed, real and non-zero. Here h_s (resp. h_t) is the field in the s -direction (resp. the t -direction). (In our case we are interested in a neighbourhood of $h_s = h_t = h$.) Therefore for all $h \neq 0$, $\varrho(s_i)$ and $\varrho(t_i)$ are independent of $\varrho \in \Delta_1(H)$.

In particular $\varrho_M(s_i) = \varrho_M(t_i)$. The “bootstrap” Theorem 3.5 then implies $\varrho_M(s_A) = \varrho_M(t_A)$ for all A . By the symmetry of interchanging s and t variables we conclude that $\varrho_M(s_A) = \varrho_m(s_A)$. Now by Equations (5.1) we conclude that all equilibrium states take the same value on s_A ; similarly for t_A . By Equation (5.2) all equilibrium states take the same value on $s_A t_B$. \square

6. Generalizations

In this section we discuss extensions of results derived in preceding sections.

Using a modification of Condition A, Messenger et al. [24] have obtained uniqueness of the phase in any nonzero external field with the Lee-Yang theorem (e.g. fixed length rotator).

Theorems 2.1, 2.2, and Corollary 2.3 extend to long-range interactions with sufficiently rapid fall-off.

For $d=2$, Corollary 2.3 is valid with J_{ij} such that $\sum_{j \in \mathbb{Z}^2} |J_{ij}|^2 < \infty$ i.e. for J_{ij} decreasing like $|i-j|^{-\alpha}$ with $\alpha > 4$. Kunz and Pfister [11] have shown that if J_{ij} decreases like $|i-j|^{-\alpha}$ with $2 < \alpha < 4$, spontaneous magnetization occurs at low temperatures. They remark that in the borderline case, $\alpha=4$, there is no spontaneous magnetization. The conclusion of Corollary 2.3 should hold in this case too.

Theorem 2.2 is valid for any a priori measure satisfying Condition A and such that the pressure is differentiable in h_s and h_t (the s - and t -components of the external field). In particular a weak version of the Lee-Yang property would be sufficient.

We will denote by “generalized interaction” a Hamiltonian of the form

$$-H = \sum_A J(A)(s_A + t_A)$$

where $J(A) \geq 0$ and $|A|$ is even.

⁵ Inequalities (5.1) and (5.2) hold for any exterior configuration and hence for any boundary condition b . Take b induced by $\langle \cdot \rangle_{M,A'}$ where $A' \supset A$

Lemma 3.2 and Theorem 3.1 depend only on Ginibre’s inequalities [Theorem A1].

Lemma 3.3 is a general result for states on a separable algebra.

Lemma 4.1 is a general statistical mechanical result.

Given that $\varrho_s(s_A) = \varrho_s(t_A) \forall A$ and $\varrho_s(s_A) = 0$ for $|A|$ odd, then Lemma 4.2b) follows for any generalized interaction, c) follows for a translation invariant generalized interaction, as does Lemma 4.3.

Given the correlation inequalities of Dunlop [3], one can extend all the above results to get results on the Heisenberg model and on 4-component models. The correlation inequalities are not sufficient to get uniqueness of the translation invariant equilibrium state on all the correlation functions but only on a subset of those.

It has already been noticed that Griffiths’ inequality for rotators (Theorem A2—A3) has some analogy with Lebowitz’ inequality [15] for Ising models [3, 1, 22]. In support of this, one may note that using only Lebowitz’ inequality (in a similar way to the use of Theorem A3 in this paper), and assuming that the equilibrium states are invariant under the spin-flip symmetry, at zero external field, one can show that the equilibrium state is unique. Of course, in this case, the use of FKG inequalities [5] gives much stronger results [16, 19].

Appendix: Correlation Inequalities

We first recall Ginibre’s and (generalized) Griffiths’ inequalities:

Theorem A1. (Ginibre [7], Dunlop-Newman [4].) *Let*

$$d\mu = Z_A^{-1} \exp \sum_{\substack{\text{supp } A \subset A \\ \text{supp } M \subset A}} (J(A, M)r_A \cos M\phi) \prod_{i \in A} d\lambda_i(r_i) d\phi_i$$

with $J(A, M) \geq 0$.

$\lambda_i(r_i)$ any measure on R_+ of compact support. Then

$$\langle r_A \cos M\phi r_B \cos N\phi \rangle \geq \langle r_A \cos M\phi \rangle \langle r_B \cos N\phi \rangle. \tag{A1}$$

Theorem A2. (Generalized Griffiths’ inequality [1, 3, 12, 18, 22].) *Let*

$$d\mu = Z_A^{-1} \exp \left(\sum_{A \subset A} J_1(A)s_A + J_2(A)t_A \right) \prod_{i \in A} dv_i(s_i, t_i)$$

$$J_1(A) \geq 0$$

$$J_2(A) \geq 0.$$

$dv_i(s_i, t_i)$ satisfy Condition A and are of compact support. Then

$$\langle s_A s_B \rangle \geq \langle s_A \rangle \langle s_B \rangle \tag{A2}$$

$$0 \leq \langle s_A t_B \rangle \leq \langle s_A \rangle \langle t_B \rangle. \tag{A3}$$

Proof. Stated in this form, the proof is in [1, 3]. \square

We want to extend these inequalities to some cases where J_1 or J_2 are not necessarily positive. This is an extension of a remark of Griffiths [8] and of a recent inequality of Lebowitz [14].

Theorem A3. *Let*

$$d\mu = Z_A^{-1} \exp\left(\sum_{\text{supp } A \subset A} J_1(A)s_A + J_2(A)t_A\right) \prod_{i \in A} dv_i(s_i t_i)$$

$$d\mu' = Z'_A{}^{-1} \exp\left(\sum_{\text{supp } A \subset A} J'_1(A)s_A + J'_2(A)t_A\right) \prod_{i \in A} dv_i(s_i t_i).$$

dv_i satisfy Condition A and are of compact support.

$$|J'_1(A)| \leq J_1(A)$$

$$|J'_2(A)| \leq J_2(A).$$

Then

$$(i) \quad |\langle s_A \rangle'| \leq \langle s_A \rangle \tag{A4}$$

$$(ii) \quad |\langle t_A \rangle'| \leq \langle t_A \rangle' \tag{A5}$$

$$(iii) \quad |\langle s_A t_B \rangle' - \langle s_A t_B \rangle'| \leq \langle s_A \rangle \langle t_B \rangle' - \langle s_A \rangle' \langle t_B \rangle. \tag{A6}$$

Proof. We use the method of duplicate variables [7]. We consider the integral

$$\int (s_A - \varepsilon_1 s'_A)(t'_B - \varepsilon_2 t_B) d\mu(\{s_i t_i\}) d\mu'(\{s'_i t'_i\}) \tag{*}$$

where $\varepsilon_i = \pm 1$. Now

$$\begin{aligned} & J_1(A)s_A + J_2(A)t_A + J'_1(A)s'_A + J'_2(A)t'_A \\ &= \frac{1}{2} [(J_1(A) + J'_1(A))(s_A + s'_A) + (J_1(A) - J'_1(A))(s_A - s'_A) \\ & \quad + (J'_2(A) + J_2(A))(t_A + t'_A) + (J'_2(A) - J_2(A))(t'_A - t_A)]. \end{aligned}$$

By hypothesis all the coefficients

$$\begin{aligned} & J_1(A) + J'_1(A), \quad J_1(A) - J'_1(A) \\ & J'_2(A) + J_2(A), \quad J'_2(A) - J_2(A) \end{aligned} \text{ are positive.}$$

We expand the exponential in series and develop in each term the factors $s_A + s'_A$, $s_A - s'_A$, etc. with the iterated formula:

$$x_1 x_2 \pm y_1 y_2 = \frac{1}{2} ((x_1 + y_1)(x_2 \pm y_2) + (x_1 - y_1)(x_2 \mp y_2)).$$

We get a series with positive coefficients of products of integrals which are all positive by Condition A.

Thus the integral (*) is ≥ 0 , which gives

$$\varepsilon_1 \langle s_A t_B \rangle' + \varepsilon_2 \langle s_A t_B \rangle \leq \langle s_A \rangle \langle t_B \rangle' + \varepsilon_1 \varepsilon_2 \langle s_A \rangle' \langle t_B \rangle.$$

This gives

$$|\langle s_A t_B \rangle + \varepsilon \langle s_A t_B \rangle'| \leq \langle s_A \rangle \langle t_B \rangle' + \varepsilon \langle s_A \rangle' \langle t_B \rangle$$

where $\varepsilon = \pm 1$.

- (i) follows taking $B = \phi, \quad \varepsilon = \pm 1$
- (ii) follows taking $A = \phi, \quad \varepsilon = \pm 1$
- (iii) follows taking $\varepsilon = -1. \quad \square$

Combining this method with Ginibre’s method of proving his inequalities, we have:

Theorem A4. *Let*

$$d\mu = Z_A^{-1} \exp\left(\sum_{\substack{\text{supp } M \subset A \\ \text{supp } A \subset A}} J(A, M)r_A \cos M\phi\right) \prod d\lambda_i(r_i)d\phi_i$$

$$d\mu' = Z_A^{-1} \exp\left(\sum_{\substack{\text{supp } M \subset A \\ \text{supp } A \subset A}} J(A, M)r'_A \cos M\phi\right) \prod d\lambda_i(r_i)d\phi_i.$$

If for all $M, A \quad J(M, A) \geq |J(M, A)'$ then

$$\langle r_B \cos N\phi \rangle \geq \langle r'_B \cos N\phi \rangle'.$$

Proof.

$$\int (r_A \cos M\phi - r'_A \cos M\phi') d\mu(\{r_i, \phi_i\}) d\mu(\{r'_i, \phi'_i\}) \geq 0$$

by the same method as above. The inequalities used in the proof of Theorem 3.1 are derived as follows.

Theorem A5. (Generalized Dunlop-Newman inequalities [3].) *Let ϱ be a state satisfying Ginibre’s inequalities (Theorem A1) and such that $\varrho(s_{A^c}t_B) = 0$ if $|B|$ is odd. Then*

- i) $|\varrho(s_A t_B s_C t_D) - \varrho(s_A t_B)\varrho(s_C t_D)|$
 $\leq \varrho(s_A s_B s_C s_D) - \varrho(s_A s_B)\varrho(s_C t_D) \quad \text{if } |B| \text{ is even};$
- ii) $\varrho(s_A t_B s_C t_D)^2 \leq \varrho(s_A s_B s_C s_D)^2 - \varrho(s_A s_B)^2 \varrho(s_C s_D)^2$
 if $|B|$ is odd.

Proof. We assume that $|D|$ is even in i) and odd in ii) otherwise the l.h.s. vanishes.

i) Let $l_i = \pm 1$. Using Ginibre’s inequalities and Equation (3.1), we have

$$\varrho(s_A(s_B + l_1 t_B)s_C(s_D + l_2 t_D)) \geq \varrho(s_A(s_B + l_1 t_B))\varrho(s_C(s_D + l_2 t_D))$$

which gives

$$\begin{aligned} & -l_1 l_2 (\varrho(s_A t_B s_C t_D) - \varrho(s_A t_B)\varrho(s_C t_D)) \\ & \leq \varrho(s_A s_B s_C s_D) - \varrho(s_A s_B)\varrho(s_C s_D) + l_1 (\varrho(s_A t_B s_C s_D) \\ & - \varrho(s_A t_B)\varrho(s_C s_D)) + l_2 (\varrho(s_A s_B s_C t_D) - \varrho(s_A s_B)\varrho(s_C t_D)). \end{aligned}$$

Take the case $l_1 = 1, l_2 = 1$, add it to the case $l_1 = -1, l_2 = -1$ and divide by 2:

$$-(\varrho(s_A t_B s_C t_D) - \varrho(s_A t_B)\varrho(s_C t_D)) \leq \varrho(s_A s_B s_C s_D) - \varrho(s_A s_B)\varrho(s_C s_D).$$

With $l_1 = +1, l_2 = -1$ added to $l_1 = -1, l_2 = +1$, we get the same result with the sign instead of $-$ in front of the l.h.s. So i) is proved.

ii) We use duplicate variables and apply i). Then

$$\begin{aligned} & |q(s_A t_B s_C t_D)^2 - q(s_A t_B)^2 q(s_C t_D)^2| \\ & \leq q(s_A s_B s_C s_D)^2 - q(s_A s_B)^2 q(s_C s_D)^2 \end{aligned}$$

but $q(s_A t_B) = q(s_C t_D) = 0$ since $|B|, |D|$ are odd. \square

References

1. Bricmont, J.: Correlation inequalities for two-component fields. *Ann. Soc. Sc. Brux.* **90**, 245—252 (1976)
2. Dobrushin, R.L., Shlosman, S.B.: Absence of breakdown of continuous symmetry in two-dimensional models of statistical physics. *Commun. math. Phys.* **42**, 31—40 (1975)
3. Dunlop, F.: Correlation inequalities for multicomponent rotators. *Commun. math. Phys.* **49**, 247—256 (1976)
4. Dunlop, F., Newman, C.M.: Multicomponent field theories and classical rotators. *Commun. math. Phys.* **44**, 223—235 (1974)
5. Fortuin, C.M., Ginibre, J., Kastelyn, P.W.: Correlation inequalities on some partially ordered sets. *Commun. math. Phys.* **22**, 89—103 (1971)
6. Fröhlich, J., Simon, B., Spencer, T.: Infrared bounds, phase transitions and continuous symmetry breaking. *Commun. math. Phys.* **50**, 79—85 (1976)
7. Ginibre, J.: General formulation of Griffiths' inequalities. *Commun. math. Phys.* **16**, 310—328 (1970)
8. Griffiths, R.B.: Phase transitions. In: *Statistical mechanics and quantum field theory* (Les Houches), pp. 241—279. New York, London, Paris: Gordon and Breach 1970
9. Israël, R.: Thesis, Princeton University
10. Israël, R.B.: High-temperature analyticity in classical lattice systems. *Commun. math. Phys.* **50**, 245—257 (1976)
11. Kunz, H., Pfister, Ch.Ed.: First order phase transitions in the plane rotator ferromagnetic model in two dimensions. *Commun. math. Phys.* **46**, 245—251 (1976)
12. Kunz, H., Pfister, Ch.Ed., Vuilleumot, P.A.: Inequalities for some classical spin vector models. Bielefeld preprint
13. Lanford, O.E., Ruelle, D.: Observables at infinity and states with short range correlations in statistical mechanics. *Commun. math. Phys.* **13**, 194—215 (1969)
14. Lebowitz, J.L.: Coexistence of phases in Ising ferromagnets. Preprint
15. Lebowitz, J.L.: GHS and other inequalities. *Commun. math. Phys.* **35**, 87—92 (1974)
16. Lebowitz, J.L., Martin-Löf, A.: On the uniqueness of the equilibrium state for Ising spin systems. *Commun. math. Phys.* **25**, 276—282 (1976)
17. Mermin, N.D.: Absence of ordering in certain classical systems. *J. Math. Phys.* **6**, 1061—1064 (1967)
18. Monroe, J.L.: Correlation inequalities for two-dimensional vector spin system. *J. Math. Phys.* **16**, 1809—1812 (1975)
19. Ruelle, D.: On the use of "small external fields" in the problem of symmetry breakdown in statistical mechanics. *Ann. Phys.* **69**, 364—374 (1972)
20. Ruelle, D.: *Statistical mechanics*. New York: Benjamin 1969
21. Holsztynski, W., Slawny, J.: Phase transitions in ferromagnetic spin systems at low temperatures. Preprint
22. Sylvester, G.S.: Thesis, Harvard University
23. Suzuki, M., Fisher, M.: Zeros of the partition function for the Heisenberg, Ferroelectric and general Ising models. *J. Math. Phys.* **12**, 235—246 (1971)
24. Messager, A., Miracle, S., Pfister, Ch.E.: to be published

Communicated by E. Lieb

Received March 25, 1977; in revised form July 6, 1977