

ON THE ZEROS OF SOLUTIONS OF A CLASS OF SECOND ORDER LINEAR DIFFERENTIAL EQUATIONS*†§

JIN TU AND XIANG-DONG YANG

Abstract

In this paper, we investigate the exponent of convergence of the zero-sequence of solutions of the second order linear differential equation

$$f'' + \left(\sum_{j=1}^l Q_j(z) e^{P_j(z)} \right) f = 0,$$

where $P_j(z)$ ($j = 1, 2, \dots, l \geq 3$) are polynomials of degree $n \geq 1$, $Q_j(z)$ are entire functions of order less than n , and obtain some results which improve and generalize the previous results in [8, 9, 13].

1. Introduction and results

We shall assume that reader is familiar with the fundamental results and the standard notations of the Nevanlinna's value distribution theory of meromorphic functions (see [7, 10]). We will use the notation $\rho(f)$ to denote the order of growth of meromorphic function $f(z)$, $\lambda(f)$ to denote the exponent of convergence of the zero-sequence of $f(z)$.

For second order linear differential equation

$$(1.1) \quad f'' + A(z)f = 0,$$

where $A(z)$ is an entire function, many authors have investigated the growth and the convergence of the zero-sequence of solutions of (1.1), and have achieved many results (see [1, 2, 3, 11]). When $A(z) = e^{P_1(z)} + e^{P_2(z)} + Q_0(z)$, for the following second order linear differential equation

$$(1.2) \quad f'' + (e^{P_1(z)} + e^{P_2(z)} + Q_0(z))f = 0,$$

*2000 *Mathematics Subject Classification*: 30D35, 34A20.

†Key words: linear differential equation, entire function, zero.

§The project is by the Youth Foundation of Education Bureau of Jiangxi Province in China (Grant No. GJJ09463), the Natural Science Foundation of Jiangxi Province in China (Grant No. 2009GQS0013), Basic Research Foundation of Yunnan Province in China (Grant No. 2009ZC013X) and Basic Research Foundation of Yunnan Education Bureau in China (Grant No. 09Y0079).

Received July 22, 2008; revised November 4, 2009.

where $P_1(z), P_2(z)$ are non-constant polynomials

$$P_1(z) = \zeta_1 z^n + \dots, \quad P_2(z) = \zeta_2 z^m + \dots, \quad \zeta_1 \zeta_2 \neq 0, \quad (n, m \in \mathbb{N}).$$

and $Q_0(z)$ is an entire function of order less than $\max\{n, m\}$. If $e^{P_1(z)}$ and $e^{P_2(z)}$ are linearly independent, K. Ishizaki and K. Tohge have studied the exponent of convergence of the zero-sequence of solutions of (1.2) and obtained the following results.

THEOREM A ([9]). *Suppose that $n = m$, and that $\zeta_1 \neq \zeta_2$ in (1.2). If $\frac{\zeta_1}{\zeta_2}$ is non-real, then for any solution $f \not\equiv 0$ of (1.2), we have $\lambda(f) = \infty$.*

THEOREM B ([8]). *Suppose that $n = m$, and that $\frac{\zeta_1}{\zeta_2} = \rho > 0$ in (1.2). If $0 < \rho < \frac{1}{2}$ or $Q_0(z) \equiv 0, \frac{3}{4} < \rho < 1$, then for any solution $f \not\equiv 0$ of (1.2), we have $\lambda(f) \geq n$.*

When $A(z) = Q_1(z)e^{P_1(z)} + Q_2(z)e^{P_2(z)} + Q_3(z)e^{P_3(z)}$, for the following second order linear differential equation

$$(1.3) \quad f'' + (Q_1(z)e^{P_1(z)} + Q_2(z)e^{P_2(z)} + Q_3(z)e^{P_3(z)})f = 0,$$

in 2007, J. Tu and Z. X. Chen studied the exponent of convergence of the zero-sequence of solutions of (1.3) and obtain the following results.

THEOREM C ([13]). *Let $Q_1(z), Q_2(z), Q_3(z)$ be entire functions of order less than n , and $P_1(z), P_2(z), P_3(z)$ be polynomials of degree $n \geq 1$,*

$$P_1(z) = \zeta_1 z^n + \dots, \quad P_2(z) = \zeta_2 z^n + \dots, \quad P_3(z) = \zeta_3 z^n + \dots,$$

where $\zeta_1, \zeta_2, \zeta_3$ are complex numbers.

(i) *If $\frac{\zeta_1}{\zeta_2}$ is non-real, $0 < \lambda = \frac{\zeta_3}{\zeta_2} < \frac{1}{2}$, then for any solution $f \not\equiv 0$ of (1.3), we have $\lambda(f) = \infty$.*

(ii) *If $0 < \frac{\zeta_2}{\zeta_1} < \frac{1}{4}, 0 < \lambda = \frac{\zeta_3}{\zeta_2} < 1$, then for any solution $f \not\equiv 0$ of (1.3), we have $\lambda(f) \geq n$.*

Then a natural question is: what is the case if $A(z) = \sum_{j=1}^l Q_j e^{P_j(z)}$ ($l \geq 3$)? Can we get the same results as Theorem C?

In this paper, we investigate the exponent of convergence of the zero-sequence of solutions of the following equation

$$(1.4) \quad f'' + \left(\sum_{j=1}^l Q_j(z)e^{P_j(z)} \right) f = 0,$$

and obtain the following results which improve and generalize the results in [8, 9, 13].

THEOREM 1. Let $Q_1(z), Q_2(z), \dots, Q_l(z)$ ($l \geq 3$) be entire functions of order less than n , and $P_1(z), P_2(z), \dots, P_l(z)$ ($l \geq 3$) be polynomials of degree $n \geq 1$,

$$P_1(z) = \zeta_1 z^n + \dots, \quad P_2(z) = \zeta_2 z^n + \dots, \dots, \quad P_l(z) = \zeta_l z^n + \dots,$$

where $\zeta_1, \zeta_2, \dots, \zeta_l$ are complex numbers.

(i) If $\frac{\zeta_1}{\zeta_2}$ is non-real, $0 < \lambda_j = \frac{\zeta_j}{\zeta_2} < \frac{1}{2}$ ($j = 3, \dots, l$), then any solution $f \neq 0$ of (1.4) satisfies $\lambda(f) = \infty$.

(ii) If $0 < \rho = \frac{\zeta_2}{\zeta_1} < \frac{1}{4}$, $\lambda_j = \frac{\zeta_j}{\zeta_2} > 0$ and $\sum_{j=3}^l \lambda_j < 1$, then any solution $f \neq 0$ of (1.4) satisfies $\lambda(f) \geq n$.

2. Notations and lemmas

To prove the theorem, we need some notations and a series of lemmas. Let $P_j(z)$ ($j = 1, \dots, l$) be polynomials of degree $n \geq 1$, where $P_j(z) = (\alpha_j + i\beta_j)z^n + \dots$, $\alpha_j, \beta_j \in \mathbf{R}$.

Define

$$\delta(P_j, \theta) = \delta_j(\theta) = \alpha_j \cos n\theta - \beta_j \sin n\theta, \quad \theta \in [0, 2\pi) \quad (j = 1, \dots, l),$$

$$S_j^+ = \{\theta \mid \delta_j(\theta) > 0\}, \quad S_j^- = \{\theta \mid \delta_j(\theta) < 0\} \quad (j = 1, \dots, l).$$

Let $f(z)$ be a meromorphic function in the complex plane, throughout the paper, $S(r, f)$ will be used to denoted any quantity that satisfies $S(r, f) = o\{T(r, f)\}$ as $r \rightarrow \infty$, outside possibly an exceptional set of r values of finite linear measure. We will use M to denote a positive constant throughout this paper, not always the same at each occurrence. We call a meromorphic function $a(z)$ a small function of $f(z)$ if $T(r, a(z)) = S(r, f)$. A differential polynomial $P(f)$ in f is a polynomial in f and its derivatives with small functions of f as the coefficients (see [7]).

LEMMA 1 [5]. Suppose that $f(z)$ is meromorphic and transcendental in the plane and that

$$(2.1) \quad f^n(z)P(f) = Q(f),$$

where $P(f), Q(f)$ are differential polynomials in f with small functions of f as the coefficients and the degree of $Q(f)$ is at most n . Then

$$(2.2) \quad m(r, P(f)) = S(r, f).$$

LEMMA 2 [6]. Let $f(z)$ be a transcendental meromorphic function with $\rho(f) = \rho < \infty$, $\Gamma = \{(k_1, j_1), \dots, (k_m, j_m)\}$ be a finite set of distinct pairs of integers which satisfy $k_i > j_i \geq 0$ for $i = 1, \dots, m$. And let $\varepsilon > 0$ be a given constant, then there exists a set $E \subset [0, 2\pi)$ which has linear measure zero, such that if $\varphi \in [0, 2\pi) \setminus E$,

there is a constant $R_1 = R_1(\varphi) > 1$, such that for all z satisfying $\arg z = \varphi$ and $|z| = r > R_1$ and for all $(k, j) \in \Gamma$, we have

$$(2.3) \quad \left| \frac{f^{(k)}(z)}{f^{(j)}(z)} \right| \leq |z|^{(k-j)(\sigma-1+\varepsilon)}.$$

LEMMA 3 [12]. Suppose that $P(z) = (\alpha + \beta i)z^n + \dots$ (α, β are real numbers, $|\alpha| + |\beta| \neq 0$) is a polynomial with degree $n \geq 1$, that $A(z) (\neq 0)$ is an entire function with $\rho(A) < n$. Set $g(z) = A(z)e^{P(z)}$, $z = re^{i\theta}$, $\delta(P, \theta) = \alpha \cos n\theta - \beta \sin n\theta$. Then for any given $\varepsilon > 0$, there exists a set $H_1 \subset [0, 2\pi)$ that has the linear measure zero, such that for any $\theta \in [0, 2\pi) \setminus (H_1 \cup H_2)$, there is a constant $R_2 > 0$ such that for $|z| = r > R_2$, we have

(i) If $\delta(P, \theta) > 0$, then

$$(2.4) \quad \exp\{(1 - \varepsilon)\delta(P, \theta)r^n\} < |g(re^{i\theta})| < \exp\{(1 + \varepsilon)\delta(P, \theta)r^n\};$$

(ii) If $\delta(P, \theta) < 0$, then

$$(2.5) \quad \exp\{(1 + \varepsilon)\delta(P, \theta)r^n\} < |g(re^{i\theta})| < \exp\{(1 - \varepsilon)\delta(P, \theta)r^n\},$$

where $H_2 = \{\theta \in [0, 2\pi); \delta(P, \theta) = 0\}$ is a finite set.

Remark. Lemma 3 also holds when $A(z)$ is a meromorphic function with $\rho(A) < n$.

LEMMA 4 [4]. Let $f(z)$ be an entire function of order $\rho(f) = \alpha < +\infty$. Then for any given $\varepsilon > 0$, there is a set $E \subset [1, \infty)$ that has finite linear measure such that for all z satisfying $|z| = r \notin [0, 1] \cup E$, we have

$$(2.6) \quad \exp\{-r^{\alpha+\varepsilon}\} \leq |f(z)| \leq \exp\{r^{\alpha+\varepsilon}\}.$$

LEMMA 5. Let $P_j(z)$ ($j = 1, \dots, l$) be polynomials of degree $n \geq 1$,

$$P_1(z) = \zeta z^n + B_1(z), \quad P_2(z) = \rho_2 \zeta z^n + B_2(z), \quad \dots, \quad P_l(z) = \rho_l \zeta z^n + B_l(z),$$

where $\zeta = \alpha + \beta i$, $\alpha, \beta \in \mathbf{R}$, $|\alpha| + |\beta| \neq 0$, $0 < \rho_j < 1$, $j = 2, \dots, l$, $B_1(z), \dots, B_l(z)$ are polynomials of degree at most $n - 1$. Let $Q_1(z) \neq 0$, $Q_2(z), \dots, Q_l(z)$ be entire functions of order less than n , then for any given $\varepsilon > 0$, there exist a set E with finite linear measure and a constant $\xi (n - 1 < \xi < n)$ such that

$$(2.7) \quad m(r, Q_1 e^{P_1} + Q_2 e^{P_2} + \dots + Q_l e^{P_l}) \\ \geq (1 - \varepsilon)m(r, e^{P_1}) + O(r^\xi), \quad r \rightarrow \infty, \quad (r \notin E).$$

Proof. By definition, for sufficiently large r , we have

$$(2.8) \quad m(r, e^{P_1}) = \frac{1}{2\pi} \int_0^{2\pi} \log^+ |e^{P_1(re^{i\theta})}| d\theta = \frac{1}{2\pi} \int_{S_1^+} \log^+ |e^{P_1(re^{i\theta})}| d\theta \\ = \frac{|\zeta| r^n}{\pi} + O(r^{n-1}).$$

If $\theta \in S_1^-$, then $\delta(P_j, \theta) < 0$ ($j = 2, \dots, l$), by Lemma 3 and Lemma 4, for any given $\varepsilon > 0$ and for sufficiently large r , we have

$$(2.9) \quad |Q_1 e^{P_1(re^{i\theta})} + Q_2 e^{P_2(re^{i\theta})} + \dots + Q_l e^{P_l(re^{i\theta})}| \leq \sum_{j=1}^l \exp\{(1 - 2\varepsilon)\delta(P_j, \theta)r^n\} \leq 1.$$

If $\theta \in S_1^+$, since $0 < \rho_j < 1$ ($j = 2, \dots, l$), by Lemma 3 and Lemma 4, there exist a set E with finite linear measure, for any given $\varepsilon > 0$ and for sufficiently large r , we have

$$(2.10) \quad \begin{aligned} & |Q_1 + Q_2 e^{P_2(re^{i\theta}) - P_1(re^{i\theta})} + \dots + Q_l e^{P_l(re^{i\theta}) - P_1(re^{i\theta})}| \\ & \geq |Q_1| - |Q_2 e^{P_2(re^{i\theta}) - P_1(re^{i\theta})}| - \dots - |Q_l e^{P_l(re^{i\theta}) - P_1(re^{i\theta})}| \\ & \geq \frac{1}{2} \exp\{-r^{\sigma(Q_1) + \varepsilon}\} \geq \exp\{-r^\xi\}, \quad (r \notin E), \end{aligned}$$

where $\rho(Q_1) < \xi < n$. By (2.8)–(2.10), we have

$$(2.11) \quad \begin{aligned} & m(r, Q_1 e^{P_1} + Q_2 e^{P_2} + \dots + Q_l e^{P_l}) \\ & = \frac{1}{2\pi} \int_0^{2\pi} \log^+ |Q_1 e^{P_1(re^{i\theta})} + Q_2 e^{P_2(re^{i\theta})} + \dots + Q_l e^{P_l(re^{i\theta})}| d\theta \\ & = \frac{1}{2\pi} \int_{S_1^+} \log^+ (|e^{P_1(re^{i\theta})}| |Q_1 + Q_2 e^{P_2(re^{i\theta}) - P_1(re^{i\theta})} \\ & \quad + \dots + Q_l e^{P_l(re^{i\theta}) - P_1(re^{i\theta})}|) d\theta \\ & = \frac{(1 - \varepsilon)|\zeta|r^n}{\pi} - O(r^\xi), \quad (r \notin E). \end{aligned}$$

By (2.8) and (2.11), we obtain (2.7).

3. Proof of Theorem 1 (i)

Since $\zeta_j = \lambda_j \zeta_2$, $\lambda_j > 0$, $j = 3, \dots, l$, we have $S_2^+ = S_3^+ = \dots = S_l^+$, $S_2^- = S_3^- \dots = S_l^-$. We see that S_j^+ and S_j^- have n components S_{jq}^+ and S_{jq}^- respectively ($j = 1, \dots, l$; $q = 1, 2, \dots, n$). Hence we write

$$S_j^+ = \bigcup_{q=1}^n S_{jq}^+, \quad S_j^- = \bigcup_{q=1}^n S_{jq}^- \quad (j = 1, 2, \dots, l).$$

Let $f \not\equiv 0$ be a solution of (1.4). Suppose that $\lambda(f) < \infty$. Write $f = \pi e^h$, where π is the canonical product from zeros of f , and h is an entire function. From our hypothesis, we have $\sigma(\pi) = \lambda(\pi) < \infty$. From (1.4), we get

$$(3.1) \quad (h')^2 = -h'' - 2\frac{\pi'}{\pi}h' - \frac{\pi''}{\pi} - Q_1 e^{P_1} - Q_2 e^{P_2} - \dots - Q_l e^{P_l}.$$

Eliminating e^{P_1} from (3.1) and set $\frac{Q'_1}{Q_1} + P'_1 = R$, we have

$$(3.2) \quad 2U_1h' = -h''' + \left(R - 2\frac{\pi'}{\pi}\right)h'' + 2\left(R\frac{\pi'}{\pi} - \left(\frac{\pi'}{\pi}\right)'\right)h' + R\frac{\pi''}{\pi} - \left(\frac{\pi''}{\pi}\right)' + \sum_{j=2}^l (RQ_j - Q'_j - Q_jP'_j)e^{P_j},$$

$$(3.3) \quad U_1 = h'' - \frac{1}{2}Rh'.$$

Eliminating e^{P_2} from (3.1) and set $\frac{Q'_2}{Q_2} + P'_2 = T$, we have

$$(3.4) \quad 2U_2h' = -h''' + \left(T - 2\frac{\pi'}{\pi}\right)h'' + 2\left(T\frac{\pi'}{\pi} - \left(\frac{\pi'}{\pi}\right)'\right)h' + T\frac{\pi''}{\pi} - \left(\frac{\pi''}{\pi}\right)' + (TQ_1 - Q'_1 - Q_1P'_1)e^{P_1} + \sum_{j=3}^l (RQ_j - Q'_j - Q_jP'_j)e^{P_j},$$

where

$$(3.5) \quad U_2 = h'' - \frac{1}{2}Th'.$$

We next proceed to prove that $\rho(U_1) \leq n$ and $\rho(U_2) \leq n$. Since $\max\{\rho(Q_j), j = 1, \dots, l\} < n$, we choose constants ξ_1, ξ_2, ξ_3 satisfying $\max\{\rho(Q_j), j = 1, \dots, l\} < \xi_1 < \xi_2 < \xi_3 < n$, then we have

$$|Q_j(re^{i\theta})| \leq \exp\{r^{\xi_1}\}, \quad T(r, Q_j) = m(r, Q_j) \leq r^{\xi_1}, \quad (j = 1, \dots, l)$$

for sufficiently large r and for any $\theta \in [0, 2\pi)$. We apply Lemma 1 to (3.1), for any given $\varepsilon > 0$, we have

$$T(r, h') = m(r, h') \leq m\left(r, \frac{\pi''}{\pi}\right) + m\left(r, \frac{\pi'}{\pi}\right) + m(r, Q_1e^{P_1(z)} + Q_2e^{P_2(z)} + \dots + Q_l e^{P_l(z)}) + S(r, h') \leq O(r^{n+\varepsilon}) + S(r, h'),$$

which implies $\rho(h') \leq n$. It follows from (3.3) and (3.5) that $\rho(U_1) \leq n$ and $\rho(U_2) \leq n$ respectively.

We next show that there exists a set $E_0 \subset [0, 2\pi)$ with $m(E_0) = 0$ such that if $\theta \in S_2^- \setminus E_0$, then

$$(3.6) \quad |U_1(re^{i\theta})| \leq O(e^{r^{\xi_2}}), \quad \text{as } r \rightarrow \infty, \theta \notin E_0,$$

where E_0 denote a set of linear measure zero, not always the same at each occurrence. If $|h'(re^{i\theta})| < 1$, by Lemma 2 and (3.3), we have

$$(3.7) \quad |U_1(re^{i\theta})| \leq \left| \frac{h''(re^{i\theta})}{h'(re^{i\theta})} \right| + \frac{1}{2} |R(re^{i\theta})| \leq O(r^M) \quad \text{as } r \rightarrow \infty, \theta \notin E_0.$$

If $|h'(re^{i\theta})| \geq 1$, then from (3.2), we get

$$(3.8) \quad \begin{aligned} |2U_1(re^{i\theta})| &\leq \left| \frac{h'''(re^{i\theta})}{h'(re^{i\theta})} \right| + \left(|R(re^{i\theta})| + 2 \left| \frac{\pi'(re^{i\theta})}{\pi(re^{i\theta})} \right| \right) \left| \frac{h''(re^{i\theta})}{h'(re^{i\theta})} \right| \\ &\quad + 2 \left(|R(re^{i\theta})| \left| \frac{\pi'(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi''(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi'(re^{i\theta})}{\pi(re^{i\theta})} \right|^2 \right) \\ &\quad + |R(re^{i\theta})| \left| \frac{\pi''(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi'''(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi''(re^{i\theta})\pi'(re^{i\theta})}{\pi(re^{i\theta})^2} \right| \\ &\quad + \sum_{j=2}^l (|R(re^{i\theta})Q_j(re^{i\theta})| + |Q'_j(re^{i\theta})| \\ &\quad + |Q_j(re^{i\theta})P'_j(re^{i\theta})|) |e^{P_j(re^{i\theta})}| \\ &\leq O(e^{r^{\xi_2}}), \quad \text{as } r \rightarrow \infty, \theta \in S_2^- \setminus E_0. \end{aligned}$$

Since Q and h' are of finite order, combining (3.7) and (3.8), we obtain (3.6).

In the following, we prove that for any $\theta \in [0, 2\pi)$,

$$(3.9) \quad |U_1(re^{i\theta})| \leq O(e^{r^{\xi_3}}), \quad \text{as } r \rightarrow \infty.$$

We note that there exist $\bar{\theta}_j$ ($j = 1, 2, \dots, l$) satisfying $\delta_j(\theta) = 0$ on the rays $\arg z = \bar{\theta}_j + \frac{q\pi}{n}$, where $q = 0, \dots, 2n - 1$, which form $2n$ sectors of opening $\frac{\pi}{n}$ respectively. Without loss of generality, we may assume that $\bar{\theta}_j \in \left[0, \frac{\pi}{n}\right)$. Since $\lambda_j = \frac{\zeta_j}{\zeta_2} > 0$ ($j = 3, \dots, l$), we have $\bar{\theta}_j = \bar{\theta}_2$ ($j = 3, \dots, l$). Set $\bar{\theta}_{jq} = \bar{\theta}_j + \frac{q\pi}{n}$, $j = 1, 2$, if there are some integers q_1 and q_2 such that $\bar{\theta}_{1q_1} = \bar{\theta}_{2q_2}$, then $\bar{\theta}_1 - \bar{\theta}_2 + (q_1 - q_2)\frac{\pi}{n} = 0$, we have that $\tan n\bar{\theta}_j = \frac{\alpha_j}{\beta_j}$, $j = 1, 2$. Which gives

$$0 = \tan(n\bar{\theta}_1 - n\bar{\theta}_2 + (q_1 - q_2)\pi) = \frac{\alpha_1\beta_2 - \alpha_2\beta_1}{\alpha_1\alpha_2 + \beta_1\beta_2} = -\Im m \frac{\zeta_1}{\zeta_2}.$$

This contradicts the assumption that $\frac{\zeta_1}{\zeta_2}$ is non-real. Hence we see that each component of S_1^+ and S_2^+ contains a component of $S_1^+ \cap S_2^+$. The boundaries of

the components of $S_1^+ \cap S_2^+$ are some of the rays $\arg z = \bar{\theta}_{jq}$, we fix a component of $S_1^+ \cap S_2^+$, say S^* . We may write

$$S^* = \{\theta \in S_1^+ \cap S_2^+ : \theta_1^* < \theta < \theta_2^*, \delta_1(\theta_1^*) = \delta_2(\theta_2^*) = 0\}$$

or

$$S^* = \{\theta \in S_1^+ \cap S_2^+ : \theta_2^* < \theta < \theta_1^*, \delta_1(\theta_1^*) = \delta_2(\theta_2^*) = 0\}.$$

Furthermore, we define

$$D_{12} = \{\theta \in S_1^+ \cap S_2^+ : \delta_1(\theta) > (2\lambda + 2)\delta_2(\theta)\},$$

$$D_{21} = \left\{ \theta \in S_1^+ \cap S_2^+ : \delta_2(\theta) > \frac{\lambda + 1}{\lambda} \delta_1(\theta) \right\},$$

where $\lambda = \max\{\lambda_j : j = 3, \dots, l\} < \frac{1}{2}$. Since each component of S_1^+ and S_2^+ is a sector of opening $\frac{\pi}{n}$, the rays $\arg z = \theta_1^*$ and $\arg z = \theta_2^*$ are contained in S_2^+ and S_1^+ respectively. We prove the first case, the proof of the second case can be obtained similarly. Hence there exist $\eta_1 > 0, \eta_2 > 0$ such that

$$\{\theta : \theta_1^* < \theta < \theta_1^* + \eta_1\} \subset D_{21}, \quad \{\theta : \theta_2^* - \eta_2 < \theta < \theta_2^*\} \subset D_{12}.$$

Hence there exists a $\theta \in (S_{2k}^+ \cap D_{12}) \setminus E_0$ for any $k = 1, 2, \dots, n$. Set $0 < (2\lambda + 2)\delta_2 < \rho_2 < \rho_1 < \delta_1, 0 < \varepsilon_{11} < 1 - \frac{\rho_1}{\delta_1}, 0 < \varepsilon_{12} < \frac{\rho_2}{2\delta_2} - 1, 0 < \varepsilon_{1j} < \frac{\rho_2}{2\lambda_j\delta_2} - 1, (j = 3, \dots, l)$, by Lemma 3, we have

$$\begin{aligned} (3.10) \quad & |Q_1 e^{P_1(re^{i\theta})} + Q_2 e^{P_2(re^{i\theta})} + \dots + Q_l e^{P_l(re^{i\theta})}| \\ & \geq |Q_1 e^{P_1(re^{i\theta})}| \left| 1 - \frac{Q_2}{Q_1} e^{P_2(re^{i\theta}) - P_1(re^{i\theta})} - \dots - \frac{Q_l}{Q_1} e^{P_l(re^{i\theta}) - P_1(re^{i\theta})} \right| \\ & \geq (1 - o(1)) e^{(1 - \varepsilon_{11})\delta_1 r^n} \geq (1 - o(1)) e^{\rho_1 r^n}, \quad \text{as } r \rightarrow \infty. \end{aligned}$$

We assume that there exists an unbounded sequence $\{r_m\}_{m=1}^\infty$ such that $0 < |h'(r_m e^{i\theta})| \leq 1$. From (3.1), (3.10) and Lemma 2, we get for an $N_1 \in \mathbf{N}$

$$\begin{aligned} e^{\rho_1 r_m^n} (1 - o(1)) & \leq 1 + \left| \frac{h''(r_m e^{i\theta})}{h'(r_m e^{i\theta})} \right| + 2 \left| \frac{\pi'(r_m e^{i\theta})}{\pi(r_m e^{i\theta})} \right| + \left| \frac{\pi''(r_m e^{i\theta})}{\pi(r_m e^{i\theta})} \right| \\ & \leq r_m^{N_1}, \quad \text{as } m \rightarrow \infty. \end{aligned}$$

Which is absurd. Hence we may assume that $|h'(re^{i\theta})| \geq 1$ for sufficiently large r . It follows from (3.1) and Lemma 2, for an $N_2 \in \mathbf{N}$

$$\begin{aligned} (3.11) \quad & |Q_1 e^{P_1(re^{i\theta})} + Q_2 e^{P_2(re^{i\theta})} + \dots + Q_l e^{P_l(re^{i\theta})}| \\ & \leq |h'(re^{i\theta})|^2 \left(1 + \left| \frac{h''(re^{i\theta})}{h'(re^{i\theta})} \right| + 2 \left| \frac{\pi'(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi''(re^{i\theta})}{\pi(re^{i\theta})} \right| \right) \\ & \leq |h'(re^{i\theta})|^2 (1 + O(r^{N_2})), \quad \text{as } r \rightarrow \infty. \end{aligned}$$

Thus, by (3.10) and (3.11) and for sufficiently large r , we have

$$(3.12) \quad |h'(re^{i\theta})| \geq e^{(1/2)\rho_2 r^n}.$$

From Lemma 2, (3.2) and (3.12), we get

$$(3.13) \quad \begin{aligned} |2U_1(re^{i\theta})| &\leq \left| \frac{h'''(re^{i\theta})}{h'(re^{i\theta})} \right| + \left(|R(re^{i\theta})| + 2 \left| \frac{\pi'(re^{i\theta})}{\pi(re^{i\theta})} \right| \right) \left| \frac{h''(re^{i\theta})}{h'(re^{i\theta})} \right| \\ &\quad + 2 \left(|R(re^{i\theta})| \left| \frac{\pi'(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi''(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi'(re^{i\theta})}{\pi(re^{i\theta})} \right|^2 \right) \\ &\quad + |R(re^{i\theta})| \left| \frac{\pi''(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi'''(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi''(re^{i\theta})\pi'(re^{i\theta})}{\pi(re^{i\theta})^2} \right| \\ &\quad + \sum_{j=2}^l (|R(re^{i\theta})Q_j(re^{i\theta})| + |Q_j'(re^{i\theta})| \\ &\quad + |Q_j(re^{i\theta})P_j'(re^{i\theta})|) \left| \frac{e^{P_j(re^{i\theta})}}{h'(re^{i\theta})} \right| \\ &\leq O(r^{N_2}) + (1 + o(1)) \exp \left\{ \left(\delta_2(1 + \varepsilon_{12}) - \frac{\rho_2}{2} \right) r^n \right\} \\ &\quad + \sum_{j=3}^l (1 + o(1)) \exp \left\{ \left(\lambda_j \delta_2(1 + \varepsilon_{1j}) - \frac{\rho_2}{2} \right) r^n \right\}, \quad \text{as } r \rightarrow \infty. \end{aligned}$$

Since $\delta_2(1 + \varepsilon_{12}) - \frac{\rho_2}{2} < 0$, $\lambda_j \delta_2(1 + \varepsilon_{1j}) - \frac{\rho_2}{2} < 0$ ($j = 3, \dots, l$), it gives that for an $N_3 \in \mathbb{N}$ and for sufficiently large r , we have

$$(3.14) \quad |U_1(re^{i\theta})| \leq r^{N_3}.$$

Now we fix a $\gamma (= \gamma_{2k}) \in (S_{2k}^+ \cap D_{21}) \setminus E_0$, $k = 1, 2, \dots, n$. Then we find $\gamma_1, \gamma_2 \in S_2^- \setminus E_0$, $\gamma_1 < \gamma < \gamma_2$ such that $\gamma - \gamma_1 < \frac{\pi}{n}$, $\gamma_2 - \gamma < \frac{\pi}{n}$. We first show that (3.9) holds for any $\theta \in [\gamma_1, \gamma]$. Write $\gamma - \gamma_1 = \frac{\pi}{n + \tau_1}$, $\tau_1 > 0$, since $\rho(U_1) \leq n$, we have that $|U_1(re^{i\theta})| \leq e^{r^{n+\tau_2}}$, $0 < \tau_2 < \tau_1$ for sufficiently large r . Set $g(z) = U_1(z) / \exp((ze^{-((\gamma+\gamma_1)/2)i})^{\xi_3})$, then $g(z)$ is analytic in the region $\{z : \gamma_1 \leq \arg z \leq \gamma\}$. Since $\gamma_1 \leq \arg z = \theta \leq \gamma$, $\gamma - \gamma_1 < \frac{\pi}{n}$, we infer that $\cos(\arg((ze^{-((\gamma+\gamma_1)/2)i})^{\xi_3})) \geq K$ for some $K > 0$. In fact,

$$-\frac{\pi}{2} < -\frac{\pi \xi_3}{2n} \leq -\xi_3 \frac{\gamma - \gamma_1}{2} \leq \arg((ze^{-((\gamma+\gamma_1)/2)i})^{\xi_3}) \leq \xi_3 \frac{\gamma - \gamma_1}{2} \leq \frac{\pi \xi_3}{2n} < \frac{\pi}{2}.$$

Hence for $\gamma_1 < \theta < \gamma$,

$$|g(re^{i\theta})| \leq \left| \frac{U_1(re^{i\theta})}{e^{Kr^{\xi_3}}} \right| \leq O(e^{r^{n+\tau_2}}), \quad \text{as } r \rightarrow \infty.$$

It follows from (3.6) and (3.14) that for some $M > 0$, as $r \rightarrow \infty$

$$|g(re^{i\gamma_1})| \leq \frac{O(e^{r^{\xi_2}})}{e^{Kr^{\xi_3}}} \leq M$$

and

$$|g(re^{i\gamma})| \leq \frac{O(r^{N_3})}{e^{Kr^{\xi_3}}} \leq M.$$

By the Phragmen-Lindelöf theorem, we obtain (3.9). Similarly we see that (3.9) holds for any $\theta \in [\gamma, \gamma_2]$. Hence we conclude that (3.9) holds for any $\theta \in [0, 2\pi)$.

We next need to prove that for any $\theta \in [0, 2\pi)$,

$$(3.15) \quad |U_2(re^{i\theta})| \leq O(e^{r^{\xi_3}}), \quad \text{as } r \rightarrow \infty.$$

By recalling the previous reasoning in (3.6) and (3.8), we can also obtain that there exists a set $E_0 \subset [0, 2\pi)$ with $m(E_0) = 0$ such that if $\theta \in S_1^+ \cap S_2^- \setminus E_0$, then

$$(3.16) \quad |U_2(re^{i\theta})| \leq O(e^{r^{\xi_2}}), \quad \text{as } r \rightarrow \infty.$$

By the similar proof in (3.9), there exists a $\theta \in (S_{1k}^+ \cap D_{21}) \setminus E_0$ for any $k = 1, 2, \dots, n$. Set $0 < (2\lambda + 2)\delta_1 < 2\lambda\delta_2 < \rho_4 < \rho_3 < \delta_2$, $0 < \varepsilon_{21} < 1 - \frac{\rho_3}{\delta_2}$, $0 < \varepsilon_{22} < \frac{\rho_4}{2\delta_1} - 1$, $0 < \varepsilon_{2j} < \frac{\rho_4}{2\lambda_j\delta_2} - 1$, ($j = 3, \dots, l$). By Lemma 3, we have

$$(3.17) \quad \begin{aligned} & |Q_1 e^{P_1(re^{i\theta})} + Q_2 e^{P_2(re^{i\theta})} + \dots + Q_l e^{P_l(re^{i\theta})}| \\ & \geq |Q_2 e^{P_2(re^{i\theta})}| \left| 1 - \frac{Q_1}{Q_2} e^{P_1(re^{i\theta}) - P_2(re^{i\theta})} - \dots - \frac{Q_l}{Q_2} e^{P_l(re^{i\theta}) - P_2(re^{i\theta})} \right| \\ & \geq (1 - o(1)) e^{(1 - \varepsilon_{21})\delta_2 r^n} \geq (1 - o(1)) e^{\rho_3 r^n}, \quad \text{as } r \rightarrow \infty. \end{aligned}$$

We assume that there exists an unbounded sequence $\{r_m\}_{m=1}^\infty$ such that $0 < |h'(r_m e^{i\theta})| \leq 1$. From (3.1), (3.17) and Lemma 2, we get for an $N_4 \in \mathbf{N}$

$$\begin{aligned} e^{\rho_3 r_m^n} (1 - o(1)) & \leq 1 + \left| \frac{h''(r_m e^{i\theta})}{h'(r_m e^{i\theta})} \right| + 2 \left| \frac{\pi'(r_m e^{i\theta})}{\pi(r_m e^{i\theta})} \right| + \left| \frac{\pi''(r_m e^{i\theta})}{\pi(r_m e^{i\theta})} \right| \\ & \leq r_m^{N_4}, \quad \text{as } m \rightarrow \infty. \end{aligned}$$

This is absurd. Hence we may assume that $|h'(re^{i\theta})| \geq 1$ for sufficiently large r . It follows from (3.1) and Lemma 2, for an $N_5 \in \mathbf{N}$

$$\begin{aligned}
 (3.18) \quad & |Q_1 e^{P_1(re^{i\theta})} + Q_2 e^{P_2(re^{i\theta})} + \dots + Q_l e^{P_l(re^{i\theta})}| \\
 & \leq |h'(re^{i\theta})|^2 \left(1 + \left| \frac{h''(re^{i\theta})}{h'(re^{i\theta})} \right| + 2 \left| \frac{\pi'(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi''(re^{i\theta})}{\pi(re^{i\theta})} \right| \right) \\
 & \leq |h'(re^{i\theta})|^2 (1 + O(r^{N_5})), \quad \text{as } r \rightarrow \infty.
 \end{aligned}$$

Combining (3.17) and (3.18), we obtain for sufficiently large r

$$(3.19) \quad |h'(re^{i\theta})| \geq e^{(1/2)\rho_4 r^n}.$$

It follows from (3.4) and (3.19) that

$$\begin{aligned}
 (3.20) \quad & |2U_2(re^{i\theta})| \\
 & \leq \left| \frac{h'''(re^{i\theta})}{h'(re^{i\theta})} \right| + \left(|T(re^{i\theta})| + 2 \left| \frac{\pi'(re^{i\theta})}{\pi(re^{i\theta})} \right| \right) \left| \frac{h''(re^{i\theta})}{h'(re^{i\theta})} \right| \\
 & \quad + 2 \left(|T(re^{i\theta})| \left| \frac{\pi'(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi''(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi'(re^{i\theta})}{\pi(re^{i\theta})} \right|^2 \right) \\
 & \quad + |T(re^{i\theta})| \left| \frac{\pi''(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi'''(re^{i\theta})}{\pi(re^{i\theta})} \right| + \left| \frac{\pi''(re^{i\theta})\pi'(re^{i\theta})}{\pi(re^{i\theta})^2} \right| \\
 & \quad + (|T(re^{i\theta})Q_1(re^{i\theta})| + |Q_1'(re^{i\theta})| + |Q_1(re^{i\theta})P_1'(re^{i\theta})|) \left| \frac{e^{P_1(re^{i\theta})}}{h'(re^{i\theta})} \right| \\
 & \quad + \sum_{j=3}^l (|T(re^{i\theta})Q_j(re^{i\theta})| + |Q_j'(re^{i\theta})| + |Q_j(re^{i\theta})P_j'(re^{i\theta})|) \left| \frac{e^{P_j(re^{i\theta})}}{h'(re^{i\theta})} \right| \\
 & \leq O(r^{N_5}) + (1 + o(1)) \exp \left\{ \left(\delta_1(1 + \varepsilon_{22}) - \frac{\rho_4}{2} \right) r^n \right\} \\
 & \quad + \sum_{j=3}^l (1 + o(1)) \exp \left\{ \left(\lambda_j \delta_2(1 + \varepsilon_{2j}) - \frac{\rho_4}{2} \right) r^n \right\}, \quad \text{as } r \rightarrow \infty.
 \end{aligned}$$

Since $\delta_1(1 + \varepsilon_{22}) - \frac{\rho_4}{2} < 0$, $\lambda_j \delta_2(1 + \varepsilon_{2j}) - \frac{\rho_4}{2} < 0$ ($j = 3, \dots, l$), it gives that for an $N_6 \in \mathbf{N}$ and for sufficiently large r ,

$$(3.21) \quad |U_2(re^{i\theta})| \leq r^{N_6}.$$

Now we fix a $\gamma' (= \gamma'_{2k}) \in (S_{2k}^+ \cap D_{12}) \setminus E_0$, $k = 1, 2, \dots, n$. Then we find $\gamma_3, \gamma_4 \in S_1^- \cap S_2^- \setminus E_0$, $\gamma_3 < \gamma' < \gamma_4$ such that $\gamma' - \gamma_3 < \frac{\pi}{n}$, $\gamma_4 - \gamma' < \frac{\pi}{n}$. By the same reasoning in (3.14), for any $\gamma_3 \leq \theta \leq \gamma_4$, we have

$$(3.22) \quad |U_2(re^{i\theta})| \leq O(e^{r^{\varepsilon_3}}), \quad \text{as } r \rightarrow \infty.$$

Hence we conclude that (3.15) holds for any $\theta \in [0, 2\pi)$.

To complete the proof of Theorem 1 (i), by (3.2) and (3.5), we have

$$(3.23) \quad U_1 - U_2 = \frac{1}{2}h'(T - R),$$

since $\max\{\rho(Q_j), j = 1, 2, \dots, l\} < \xi_2 < \xi_3$, by the theorem on the logarithmic derivative and by (3.1), (3.9), (3.15), (3.23), we have

$$(3.24) \quad \begin{aligned} m(r, Q_1 e^{P_1(z)} + Q_2 e^{P_2(z)} + \dots + Q_l e^{P_l(z)}) \\ \leq 2m(r, h') + O(\log r) \leq 2m(r, U_1 - U_2) + O(\log r) \\ \leq O(r^{\xi_3}), \quad \text{as } r \rightarrow \infty. \end{aligned}$$

Since $\frac{\xi_1}{\xi_2}$ is non-real, $S_1^+ \cap S_2^-$ contains an interval $I = [\varphi_1, \varphi_2]$ satisfying $\min_{\theta \in I} \delta_1(\theta) = s > 0$. By Lemma 3, there exists a constant $R_2(\theta) (> 0)$ such that for any $\theta \in I$ and for any given $\varepsilon > 0$, we have for sufficiently large $r \geq R_2(\theta)$

$$\begin{aligned} |Q_1 e^{P_1(re^{i\theta})}| &\geq \exp((1 - \varepsilon)\delta_1 r^n), \\ |Q_2 e^{P_2(re^{i\theta})}| &\leq \exp((1 - \varepsilon)\delta_2 r^n), \\ |Q_j e^{P_j(re^{i\theta})}| &\leq \exp((1 - \varepsilon)\lambda_j \delta_2 r^n), \quad (j = 3, \dots, l). \end{aligned}$$

Hence,

$$(3.25) \quad \begin{aligned} m(r, Q_1 e^{P_1(z)} + Q_2 e^{P_2(z)} + \dots + Q_l e^{P_l(z)}) \\ \geq \int_{\varphi_1}^{\varphi_2} \log^+ |Q_1 e^{P_1(re^{i\theta})} + Q_2 e^{P_2(re^{i\theta})} + \dots + Q_l e^{P_l(re^{i\theta})}| d\theta \\ \geq \int_{\varphi_1}^{\varphi_2} (1 - o(1)) \log^+ |Q_1 e^{P_1(re^{i\theta})}| d\theta \\ \geq \int_{\varphi_1}^{\varphi_2} (1 - o(1))(1 - \varepsilon)sr^n d\theta \\ \geq (1 - o(1))(1 - \varepsilon)sr^n(\varphi_2 - \varphi_1), \quad \text{as } r \rightarrow \infty. \end{aligned}$$

Combining (3.24) and (3.25) and recalling that $\xi_3 < n$, we get a contradiction. Hence, $\lambda(f) = \infty$.

4. Proof of Theorem 1 (ii)

Let $f \not\equiv 0$ be a solution of (1.4). Write $f = \pi e^h$, suppose that $\lambda(f) < n$. From our hypothesis, we have $\rho(\pi) = \lambda(\pi) < n$. Eliminating e^{P_1} from (3.1) and recalling that $R = \frac{Q_1'}{Q_1} + P_1'$, we have

$$(3.26) \quad 2Uh' = -h''' + \left(R - 2\frac{\pi'}{\pi}\right)h'' + 2\left(R\frac{\pi'}{\pi} - \left(\frac{\pi'}{\pi}\right)'\right)h' + R\frac{\pi''}{\pi} - \left(\frac{\pi''}{\pi}\right)' + \sum_{j=2}^l (RQ_j - Q_j' - Q_jP_j')e^{P_j},$$

where

$$(3.27) \quad U = h'' - \frac{1}{2}Rh'.$$

From (3.26) and (3.27), we get

$$C_1(z)h' = C_0(z),$$

where

$$(3.28) \quad C_0(z) = -U' + \frac{1}{2}RU - 2\frac{\pi'}{\pi}U + R\frac{\pi''}{\pi} - \frac{\pi'''}{\pi} + \frac{\pi''\pi'}{\pi^2} + \sum_{j=2}^l (RQ_j - Q_j' - Q_jP_j')e^{P_j},$$

$$(3.29) \quad C_1(z) = 2U + \frac{1}{2}R' - \frac{1}{4}R^2 - R\frac{\pi'}{\pi} + 2\frac{\pi''}{\pi} - 2\left(\frac{\pi'}{\pi}\right)'^2.$$

We next show that $C_0(z) \equiv 0$ and $C_1(z) \equiv 0$. If $C_0(z) \not\equiv 0$, $C_1(z) \not\equiv 0$, by Nevanlinna's first fundamental theorem, we obtain

$$T(r, h') \leq T(r, C_0) + T(r, C_1) + o(1).$$

Set $\max\{\rho(Q_j) \ (j = 1, \dots, l), \lambda(f)\} < \xi_2 < \xi_3 < n$, from (3.1), we obtain

$$(3.30) \quad T(r, Q_1e^{P_1(z)} + Q_2e^{P_2(z)} + \dots + Q_le^{P_l(z)}) \leq 2T(r, h') + O(\log r).$$

By Lemma 5, we have

$$(3.31) \quad m(r, Q_1e^{P_1(z)} + Q_2e^{P_2(z)} + \dots + Q_le^{P_l(z)}) \geq (1 - \varepsilon)m(r, e^{P_1}) + O(r^{\xi_3}), \quad r \rightarrow \infty, (r \notin E),$$

where E has finite linear measure. From (3.30) and (3.31), we obtain

$$(3.32) \quad T(r, h') \geq \frac{1 - \varepsilon}{2}T(r, e^{P_1}) + O(r^{\xi_3}), \quad r \rightarrow \infty, (r \notin E).$$

Since $0 < \rho = \frac{\zeta_2}{\zeta_1} < \frac{1}{4}$, $\lambda_j = \frac{\zeta_j}{\zeta_2} > 0 \ (j = 3, \dots, l)$, $\sum_{j=3}^l \lambda_j < 1$, we get

$$\delta(P_2, \theta) = \rho\delta(P_1, \theta), \quad S_{1k}^+ = S_{2k}^+ = \dots = S_{lk}^+, \\ S_{1k}^- = S_{2k}^- = \dots = S_{lk}^-, \quad (k = 1, \dots, n).$$

By the same reasoning in (3.7) and (3.8), we have

$$(3.33) \quad |U(re^{i\theta})| \leq O(e^{r^{\xi_2}}), \quad \text{as } r \rightarrow \infty$$

for any $\theta \in S_1^- \setminus E_0$, $m(E_0) = 0$. Also by the same reasoning in (3.9)–(3.13), we have

$$(3.34) \quad |U(re^{i\theta})| \leq r^{N_3}, \quad \text{as } r \rightarrow \infty$$

for any $\theta \in S_1^+ \setminus E_0$, $m(E_0) = 0$. Since $\rho(U) \leq n$, by the Phragmen-Lindelöf theorem, we have

$$(3.35) \quad |U(re^{i\theta})| \leq O(e^{r^{\xi_3}}), \quad \text{as } r \rightarrow \infty$$

for any $\theta \in [0, 2\pi)$. In the following, we estimate $T(r, C_0)$ and $T(r, C_1)$.

$$\begin{aligned} T(r, C_0) &\leq T\left(r, U' - \frac{1}{2}RU + 2\frac{\pi'}{\pi}U\right) + T\left(r, R\frac{\pi''}{\pi} - \frac{\pi'''}{\pi} + \frac{\pi''\pi'}{\pi^2}\right) \\ &\quad + \sum_{j=2}^l T(r, RQ_j - Q_j' - Q_jP_j') + \sum_{j=2}^l T(r, e^{P_j}). \end{aligned}$$

Since $\max\{\rho(Q_j) \ (j = 1, \dots, l), \rho(R), \rho(\pi)\} < n$, we have

$$(3.36) \quad \begin{aligned} T(r, C_0) &\leq \sum_{j=2}^l T(r, e^{P_j}) + O(r^{\xi_3}) = \left(1 + \sum_{j=3}^l \lambda_j\right) T(r, e^{P_2}) + O(r^{\xi_3}) \\ &\leq \left(1 + \sum_{j=3}^l \lambda_j\right) \rho T(r, e^{P_1}) + O(r^{\xi_3}), \quad \text{as } r \rightarrow \infty. \end{aligned}$$

From (3.29) and (3.35), we have

$$(3.37) \quad T(r, C_1) \leq O(r^{\xi_3}), \quad \text{as } r \rightarrow \infty.$$

From (3.30), (3.32), (3.36) and (3.37), we get

$$(3.38) \quad \begin{aligned} &\frac{1-\varepsilon}{2} T(r, e^{P_1}) + O(r^{\xi_3}) \\ &\leq T(r, h') \leq \left(1 + \sum_{j=3}^l \lambda_j\right) \rho T(r, e^{P_1}) + O(r^{\xi_3}), \quad r \rightarrow \infty, \ (r \notin E). \end{aligned}$$

Thus (3.38) implies

$$\left(\frac{1-\varepsilon}{2} - \left(1 + \sum_{j=3}^l \lambda_j\right) \rho - o(1)\right) T(r, e^{P_1}) \leq 0, \quad r \rightarrow \infty, \ (r \notin E).$$

Since $0 < \rho = \frac{\zeta_2}{\zeta_1} < \frac{1}{4}$, $0 < \sum_{j=3}^l \lambda_j < 1$, we get a contradiction. Hence $C_0(z) \equiv C_1(z) \equiv 0$. From (3.28), we obtain

$$(3.39) \quad \sum_{j=2}^l (RQ_j - Q'_j - Q_j P'_j) e^{P_j} = U' - \frac{1}{2}RU + 2\frac{\pi'}{\pi}U - R\frac{\pi''}{\pi} + \frac{\pi'''}{\pi} - \frac{\pi''\pi'}{\pi^2}.$$

We assume that $\sum_{j=2}^l (RQ_j - Q'_j - Q_j P'_j) e^{P_j} \neq 0$, if $\sum_{j=2}^l (RQ_j - Q'_j - Q_j P'_j) e^{P_j} \equiv 0$, since $\lambda_j = \frac{\zeta_j}{\zeta_2} > 0$ ($j = 3, \dots, l$) and $0 < \sum_{j=3}^l \lambda_j < 1$, by Lemma 3 and by a simple calculation, this is a contradiction. From (3.39), by Lemma 5, we obtain

$$(3.40) \quad (1 - \varepsilon)T(r, e^{P_2}) + O(r^{\xi_3}) \leq \sum_{j=2}^l T(r, (RQ_j - Q'_j - Q_j P'_j) e^{P_j}) \\ \leq T\left(r, U' - \frac{1}{2}RU\right) + T(r, U) + T(r, R) \\ + T\left(r, \frac{\pi'}{\pi}\right) + T\left(r, \frac{\pi''}{\pi}\right) + T\left(r, \frac{\pi'''}{\pi}\right) + o(1) \\ \leq O(r^{\xi_3}), \quad r \rightarrow \infty, (r \notin E).$$

From (3.40), we have $\rho(e^{P_2}) < \xi_3 < n$, we get a contradiction. Hence $\lambda(f) \geq n$. Thus, we complete the proof of Theorem 1.

Acknowledgement. Authors are thankful to the referee for valuable suggestions to improve our paper.

REFERENCES

- [1] S. BANK AND I. LAINE, On the oscillation theory of $f'' + Af = 0$ where A is entire, Trans. Amer. Math. Soc. **273** (1982), 352–363.
- [2] S. BANK, I. LAINE AND J. K. LANGLEY, On the frequency of zeros of solutions of second order linear differential equations, Results Math. **10** (1986), 8–24.
- [3] S. BANK AND J. K. LANGLEY, On the oscillation of certain linear differential equation in the complex domain, Proc. Edinburgh Math. Soc. (2) **30** (1987), 455–469.
- [4] Z. X. CHEN, On the hyper order of solutions of some second order linear differential equations, Acta Mathematica Sinica (1) **18B** (2002), 79–88.
- [5] J. CLUNIE, On integral and meromorphic functions, J. London Math. Soc. **37** (1962), 17–27.
- [6] G. GUNDERSEN, Estimates for the logarithmic derivative of a meromorphic function, plus similar estimates, J. London Math. Soc. (2) **37** (1988), 88–104.
- [7] W. K. HAYMAN, Meromorphic functions, Clarendon Press, Oxford, 1964.
- [8] K. ISHIZAKI, An oscillation result for a certain linear differential equation of second order, Hokkaido Math. J. **26** (1997), 421–434.
- [9] K. ISHIZAKI AND K. TOHGE, On the complex oscillation of some linear differential equations, J. Math. Anal. Appl. **206** (1997), 503–517.

- [10] I. LAINE, Nevanlinna theory and complex differential equations, W. de Gruyter, Berlin, 1993.
- [11] J. K. LANGLEY, On complex oscillation and a problem of Ozawa, *Kodai Math. J.* **9** (1986), 430–439.
- [12] A. I. MARKUSHEVICH, Theory of functions of a complex variable **2**, Englewood Cliffs, N. J., Prentice-Hall, 1965.
- [13] J. TU AND Z. X. CHEN, Zeros of solutions of certain second order linear differential equation, *J. Math. Anal. Appl.* **332** (2007), 279–291.

Jin Tu
COLLEGE OF MATHEMATICS AND INFORMATION SCIENCES
JIANGXI NORMAL UNIVERSITY
NANCHANG, 330022
P.R. CHINA
E-mail: tujin2008@sina.com

Xiang-Dong Yang
DEPARTMENT OF MATHEMATICS
KUNMING UNIVERSITY OF SCIENCE AND TECHNOLOGY
KUNMING, 650093
P.R. CHINA
E-mail: yangsddp@126.com