

## Topological semigroups of matrix units

Oleg V. Gutik, Kateryna P. Pavlyk

Communicated by B. V. Novikov

**ABSTRACT.** We prove that the semigroup of matrix units is stable. Compact, countably compact and pseudocompact topologies  $\tau$  on the infinite semigroup of matrix units  $B_\lambda$  such that  $(B_\lambda, \tau)$  is a semitopological (inverse) semigroup are described. We prove the following properties of an infinite topological semigroup of matrix units. On the infinite semigroup of matrix units there exists no semigroup pseudocompact topology. Any continuous homomorphism from the infinite topological semigroup of matrix units into a compact topological semigroup is annihilating. The semigroup of matrix units is algebraically  $h$ -closed in the class of topological inverse semigroups. Some  $H$ -closed minimal semigroup topologies on the infinite semigroup of matrix units are considered.

In this paper all topological spaces are Hausdorff.

A semigroup is a set with a binary associative operation. The semigroup operation is called a *multiplication*. A semigroup  $S$  is called *inverse* if for any  $x \in S$  there exists a unique  $y \in S$  such that  $xyx = x$  and  $yx y = y$ . An element  $y$  of  $S$  is called *inverse* to  $x$  and is denoted by  $x^{-1}$ . If  $S$  is an inverse semigroup, then the map which takes  $x \in S$  to the inverse element of  $x$  is called the *inversion*.

A topological space  $S$  that is algebraically semigroup with a separately continuous semigroup operation is called a *semitopological semigroup*. If the multiplication on  $S$  is jointly continuous, then  $S$  is called a *topological semigroup*.

---

**2000 Mathematics Subject Classification:** 20M15, 20M18, 22A15, 54A10, 54C25, 54D25, 54D35, 54H10.

**Key words and phrases:** *semigroup of matrix units, semitopological semigroup, topological semigroup, topological inverse semigroup,  $H$ -closed semigroup, absolutely  $H$ -closed semigroup, algebraically  $h$ -closed semigroup, Bohr compactification, minimal topological semigroup, minimal semigroup topology.*

A *topological (semitopological) inverse semigroup* is a topological (semitopological) semigroup  $S$  that is algebraically an inverse semigroup with continuous inversion. Obviously, any topological (inverse) semigroup is a semitopological (inverse) semigroup.

If  $\tau$  is a topology on a (inverse) semigroup  $S$  such that  $(S, \tau)$  is a topological (inverse) semigroup, then  $\tau$  is called a (*inverse*) *semigroup topology* on  $S$ .

We follow the terminology of [6, 7, 11, 16], and [22].

If  $S$  is a semigroup, then by  $E(S)$  we denote the subset of idempotents of  $S$ . By  $\omega$  we denote the first infinite ordinal. Further, we identify all cardinals with their corresponding initial ordinals.

Let  $S$  be a semigroup and  $I_\lambda$  be a set of cardinality  $\lambda \geq 2$ . On the set  $B_\lambda(S) = I_\lambda \times S^1 \times I_\lambda \cup \{0\}$  we define the semigroup operation  $' \cdot '$  as follows

$$(\alpha, a, \beta) \cdot (\gamma, b, \delta) = \begin{cases} (\alpha, ab, \delta), & \text{if } \beta = \gamma, \\ 0, & \text{if } \beta \neq \gamma, \end{cases}$$

and  $(\alpha, a, \beta) \cdot 0 = 0 \cdot (\alpha, a, \beta) = 0 \cdot 0 = 0$  for  $\alpha, \beta, \gamma, \delta \in I_\lambda$ ,  $a, b \in S^1$ . The semigroup  $B_\lambda(S)$  is called a *Brandt-Howie semigroup of the weight  $\lambda$  over  $S$*  [13] or a *Brandt  $\lambda$ -extension of the semigroup  $S$*  [14]. Obviously  $B_\lambda(S)$  is the Rees matrix semigroup  $M^0(S^1; I_\lambda, I_\lambda, \mathcal{M})$ , where  $\mathcal{M}$  is the  $I_\lambda \times I_\lambda$  identity matrix. If a semigroup  $S$  is trivial (i.e. if  $S$  contains only one element), then  $B_\lambda(S)$  is the *semigroup of  $I_\lambda \times I_\lambda$ -matrix units* [7], which we shall denote by  $B_\lambda$ .

A semigroup  $\mathcal{B}(p, q)$  generated by elements  $p$  and  $q$  which satisfy the condition  $pq = 1$  is called *bicyclic*. The bicyclic semigroup plays the important role in the Algebraic Theory of Semigroups and in the Theory of Topological Semigroups. For example the well-known O. Andersen's result [1] states that a (0-) simple semigroup is completely (0-) simple if and only if it does not contain the bicyclic semigroup (see Theorem 2.54 of [7]). L. W. Anderson, R. P. Hunter and R. J. Koch in [2] proved that the bicyclic semigroup cannot be embedded into a stable semigroup. Also any  $\Gamma$ -compact topological semigroup (and hence compact topological semigroup) does not contain the bicyclic semigroup [15] and therefore every (0-) simple  $\Gamma$ -compact topological semigroup is completely (0-) simple.

In this paper we discuss semigroup topologies on the semigroup of matrix units. At the beginning we shall prove that the semigroup of matrix units is stable. Further we shall show that on any semigroup of matrix units  $B_\lambda$  there exists a unique compact topology  $\tau$  such that  $(B_\lambda, \tau)$  is a semitopological semigroup. Also we shall prove that on the infinite semigroup of matrix units there exists no semigroup pseudocompact topology

and this implies the structure theorem for 0-simple compact topological inverse semigroups. Moreover, any continuous homomorphism from the infinite topological semigroup of matrix units into a compact topological semigroup is annihilating. Also we shall prove that if a topological inverse semigroup  $S$  contains a semigroup of matrix units  $B_\lambda$ , then  $B_\lambda$  is a closed subsemigroup of  $S$ , i.e.  $B_\lambda$  is algebraically  $h$ -closed in the class of topological inverse semigroups. Some  $H$ -closed minimal semigroup topologies on the infinite semigroup of matrix units will be considered.

**Lemma 1.** *Let  $a, b \in S^1$ ,  $\alpha, \beta \in I_\lambda$ . Then the following conditions are equivalent:*

$$(i) \quad aS^1 \subseteq bS^1 \quad [S^1a \subseteq S^1b];$$

$$(ii) \quad (\alpha, a, \beta)B_\lambda(S) \subseteq (\alpha, b, \beta)B_\lambda(S) \quad [B_\lambda(S)(\alpha, a, \beta) \subseteq B_\lambda(S)(\alpha, b, \beta)].$$

*Proof.* (i) $\Rightarrow$ (ii).

$$\begin{aligned} (\alpha, a, \beta)B_\lambda(S) &= \\ &= \bigcup_{\gamma \in I_\lambda} (\alpha, aS^1, \gamma) \cup \{0\} \subseteq \bigcup_{\gamma \in I_\lambda} (\alpha, bS^1, \gamma) \cup \{0\} = (\alpha, b, \beta)B_\lambda(S). \end{aligned}$$

(ii) $\Rightarrow$ (i). Let  $(\alpha, a, \beta)B_\lambda(S) \subseteq (\alpha, b, \beta)B_\lambda(S)$ , then

$$\bigcup_{\gamma \in I_\lambda} (\alpha, aS^1, \gamma) \cup \{0\} \subseteq \bigcup_{\gamma \in I_\lambda} (\alpha, bS^1, \gamma) \cup \{0\}$$

and hence  $\bigcup_{\gamma \in I_\lambda} (\alpha, aS^1, \gamma) \subseteq \bigcup_{\gamma \in I_\lambda} (\alpha, bS^1, \gamma)$ . Therefore,  $aS^1 \subseteq bS^1$ .

The proof of equivalency of the dual conditions is similar.  $\square$

A semigroup  $S$  is called *stable* if and only if

$$(i) \quad \text{for } a, b \in S, Sa \subseteq Sab \text{ implies } Sa = Sab;$$

$$(ii) \quad \text{for } c, d \in S, cS \subseteq dcS \text{ implies } cS = dcS.$$

Stable semigroups were first investigated by R. J. Koch and A. D. Wallace in [17]. A semigroup  $S$  is called *weakly stable* if  $S^1$  is stable [20]. Every stable semigroup  $S$  is weakly stable [17]. If  $S$  is a regular semigroup, then the converse holds. L. O'Carroll [20] proved that the converse does not hold in general.

**Theorem 1.** *A semigroup  $S$  is weakly stable if and only if  $B_\lambda(S)$  is stable for each cardinal  $\lambda \geq 2$ .*

*Proof.* ( $\Leftarrow$ ) Suppose  $aS^1 \subseteq baS^1$  [ $S^1a \subseteq S^1ab$ ] for  $a, b \in S^1$ . By Lemma 1 we get

$$(\alpha, a, \beta)B_\lambda(S) \subseteq (\alpha, b, \alpha)(\alpha, a, \beta)B_\lambda(S) = (\alpha, ba, \beta)B_\lambda(S)$$

$$[B_\lambda(S)(\alpha, a, \beta) \subseteq B_\lambda(S)(\alpha, a, \beta)(\beta, b, \beta) = B_\lambda(S)(\alpha, ab, \beta)]$$

for all  $\alpha, \beta \in I_\lambda$ . Since the semigroup  $B_\lambda(S)$  is stable for each cardinal  $\lambda \geq 2$ , then

$$(\alpha, a, \beta)B_\lambda(S) = (\alpha, b, \alpha)(\alpha, a, \beta)B_\lambda(S) = (\alpha, ba, \beta)B_\lambda(S)$$

$$[B_\lambda(S)(\alpha, a, \beta) = B_\lambda(S)(\alpha, a, \beta)(\beta, b, \beta) = B_\lambda(S)(\alpha, ab, \beta)]$$

and by Lemma 1,  $aS^1 = baS^1$  [ $S^1a = S^1ab$ ]. Therefore, the semigroup  $S$  is weakly stable.

( $\Rightarrow$ ) Let  $\lambda \geq 2$ . Suppose  $(\alpha, a, \beta)B_\lambda(S) \subseteq (\gamma, b, \delta)(\alpha, a, \beta)B_\lambda(S)$  for  $\alpha, \beta, \gamma, \delta \in I_\lambda$ ,  $a, b \in S^1$ . Obviously,  $\alpha = \gamma = \delta$ . Thus  $(\alpha, a, \beta)B_\lambda(S) \subseteq (\alpha, ba, \beta)B_\lambda(S)$ . By Lemma 1 we get  $aS^1 \subseteq baS^1$ . Since  $S$  is a weakly stable semigroup, then  $aS^1 = baS^1$ . Then by Lemma 1 we get

$$\begin{aligned} (\alpha, a, \beta)B_\lambda(S) &= \\ &= (\alpha, ba, \beta)B_\lambda(S) = (\alpha, b, \alpha)(\alpha, a, \beta)B_\lambda(S) = (\gamma, b, \delta)(\alpha, a, \beta)B_\lambda(S). \end{aligned}$$

The proof of the dual statement is similar.

Therefore, the semigroup  $B_\lambda(S)$  is stable for all cardinals  $\lambda \geq 2$ .  $\square$

**Corollary 1.** *For every cardinal  $\lambda \geq 2$  the semigroup  $B_\lambda$  is stable.*

In [10] C. Eberhart and J. Selden proved that the bicyclic semigroup  $\mathcal{B}(p, q)$  admits only the discrete Hausdorff semigroup topology. M. O. Bertman and T. T. West generalized this result and showed that any Hausdorff topology  $\tau$  on  $\mathcal{B}(p, q)$  such that  $(\mathcal{B}(p, q), \tau)$  is a semitopological semigroup is discrete [5]. Lemma 2 implies that the semigroup of matrix units has similar properties.

Further by 0 we denote the zero of the semigroup  $B_\lambda$ .

**Lemma 2.** *Let  $\tau$  be a topology on  $B_\lambda$  such  $(B_\lambda, \tau)$  is a semitopological semigroup. Then any nonzero element of  $B_\lambda$  is an isolated point of  $(B_\lambda, \tau)$ .*

*Proof.* Since  $(B_\lambda, \tau)$  is a semitopological semigroup, every left internal translation  $l_s: B_\lambda \rightarrow B_\lambda$  and every right internal translation  $r_s: B_\lambda \rightarrow B_\lambda$  are continuous maps for any  $s \in B_\lambda$ . Thus for  $s = (\alpha, \alpha) \in B_\lambda$  the sets  $l_s^{-1}(0) = \{0\} \cup \{(\gamma, \beta) \mid \gamma \in I_\lambda \setminus \{\alpha\}, \beta \in I_\lambda\}$  and  $r_s^{-1}(0) =$

$\{0\} \cup \{(\gamma, \beta) \mid \gamma \in I_\lambda, \beta \in I_\lambda \setminus \{\alpha\}\}$  are closed in  $(B_\lambda, \tau)$ , and hence the sets  $B_\lambda \setminus l_s^{-1}(0) = \{(\alpha, \gamma) \mid \gamma \in I_\lambda\}$  and  $B_\lambda \setminus r_s^{-1}(0) = \{(\gamma, \alpha) \mid \gamma \in I_\lambda\}$  are open in  $(B_\lambda, \tau)$  for any  $\alpha \in I_\lambda$ . Therefore any nonzero element of  $B_\lambda$  is an isolated point of  $(B_\lambda, \tau)$ .  $\square$

In [5] M. O. Bertman and T. T. West showed that the bicyclic semigroup is embedded into a compact semitopological semigroup. The next example shows that on the infinite semigroup of matrix units  $B_\lambda$  there exists a topology  $\tau_c$  such that  $(B_\lambda, \tau_c)$  is a compact semitopological inverse semigroup.

**Example 1.** Let  $\lambda \geq \omega$ . A topology  $\tau_c$  on  $B_\lambda$  is defined as follows:

- a) all nonzero elements of  $B_\lambda$  are isolated points in  $B_\lambda$ ;
- b)  $\mathcal{B}(0) = \{A \subseteq B_\lambda \mid 0 \in A \text{ and } |B_\lambda \setminus A| < \omega\}$  is the base of the topology  $\tau_c$  at the point  $0 \in B_\lambda$ .

**Lemma 3.**  $(B_\lambda, \tau_c)$  is a compact semitopological inverse semigroup.

*Proof.* Obviously,  $\tau_c$  is a compact topology on  $B_\lambda$ .

For any  $U = B_\lambda \setminus \{(\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\} \in \mathcal{B}(0)$ , where  $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_n \in I_\lambda$  we have

- 1)  $U_1 \cdot \{(\alpha, \beta)\} = \{0\} \cup \{(\gamma, \beta) \mid \gamma \in I_\lambda \setminus \{\alpha_1, \dots, \alpha_n\}\} \subseteq U$ , where  $U_1 = U \setminus \{(\alpha_1, \alpha_1), \dots, (\alpha_n, \alpha_n)\} \in \mathcal{B}(0)$ ;
- 2)  $\{(\alpha, \beta)\} \cdot U_2 = \{0\} \cup \{(\gamma, \beta) \mid \gamma \in I_\lambda \setminus \{\beta_1, \dots, \beta_n\}\} \subseteq U$ , where  $U_2 = U \setminus \{(\beta_1, \beta_1), \dots, (\beta_n, \beta_n)\} \in \mathcal{B}(0)$ ;
- 3)  $\{(\alpha, \beta)\} \cdot \{(\gamma, \delta)\} = \{0\} \subseteq U$  if  $\beta \neq \gamma$ ;
- 4)  $\{0\} \cdot U = \{0\} \subseteq U$  and  $U \cdot \{0\} = \{0\} \subseteq U$ ;
- 5)  $\{(\alpha, \beta)\} \cdot \{(\beta, \gamma)\} = \{(\alpha, \gamma)\}$ ;
- 6)  $(U_3)^{-1} \subseteq U$ , where  $U_3 = B_\lambda \setminus \{(\beta_1, \alpha_1), \dots, (\beta_n, \alpha_n)\} \in \mathcal{B}(0)$ .

Therefore,  $(B_\lambda, \tau_c)$  is a compact semitopological inverse semigroup.  $\square$

**Remark 1.** In [21] A. B. Paalman-de-Miranda proved that the zero of a compact completely 0-simple topological semigroup  $S$  is an isolated point in  $S$ . Example 1 implies that the zero of a completely 0-simple compact semitopological inverse semigroup  $(B_\lambda, \tau)$  is not necessarily an isolated point in  $(B_\lambda, \tau)$ .

Lemmas 2 and 3 imply

**Corollary 2.** *If  $\lambda \geq \omega$  then there exists no other topology  $\tau$  on  $B_\lambda$  different from  $\tau_c$  such that  $(B_\lambda, \tau)$  is a compact semitopological semigroup.*

A topological space  $X$  is called *countably compact* if any countable open cover of  $X$  contains a finite subcover [11]. A topological space  $X$  is called *pseudocompact* [*discretely pseudocompact*] if any locally finite [discrete] collection of open subsets of  $X$  is finite. Obviously any countably compact space and any discretely pseudocompact space are pseudocompact.

**Theorem 2.** *Let  $\lambda \geq \omega$  and let  $\tau$  be a topology on the semigroup of matrix units  $B_\lambda$  such that  $(B_\lambda, \tau)$  is a semitopological semigroup. Then the following statements are equivalent:*

- (i)  $(B_\lambda, \tau)$  is a compact semitopological semigroup;
- (ii)  $(B_\lambda, \tau)$  is a countably compact semitopological semigroup;
- (iii)  $(B_\lambda, \tau)$  is a discretely pseudocompact semitopological semigroup;
- (iv)  $(B_\lambda, \tau)$  is a pseudocompact semitopological semigroup;
- (v)  $(B_\lambda, \tau)$  is topologically isomorphic to  $(B_\lambda, \tau_c)$ .

*Proof.* The implications (i)  $\Rightarrow$  (ii), (i)  $\Rightarrow$  (iii), (ii)  $\Rightarrow$  (iv) and (iii)  $\Rightarrow$  (iv) are trivial.

(iv)  $\Rightarrow$  (i) Suppose there exists a topology  $\tau$  on the infinite semigroup of matrix units  $B_\lambda$  such that  $(B_\lambda, \tau)$  is a pseudocompact non-compact semitopological semigroup. Then there exists an open cover  $\mathcal{U} = \{U_\alpha\}_{\alpha \in \mathcal{A}}$  which contains no finite subcover. Let  $U_{\alpha_1} \in \mathcal{U}$  such that  $U_{\alpha_1} \ni 0$ . Then the set  $B_\lambda \setminus U_{\alpha_1}$  is infinite. We put

$$\mathcal{U}^* = \{U_{\alpha_1}\} \cup \{x \mid x \in B_\lambda \setminus U_{\alpha_1}\}.$$

Then  $\mathcal{U}^*$  is an infinite locally finite family, which contradicts the pseudocompactness of the topological space  $(B_\lambda, \tau)$ .

Corollary 2 implies the equivalency (i)  $\Leftrightarrow$  (v). □

Since the bicyclic semigroup  $\mathcal{B}(p, q)$  admits only discrete semigroup topology [10],  $\mathcal{B}(p, q)$  admits no compact (countably compact, pseudocompact) semigroup topology. The next proposition is a similar result for the infinite semigroup of matrix units and it follows from Theorem 2.

**Proposition 1.** *If  $\lambda \geq \omega$ , then there exists no compact (countably compact, pseudocompact) semigroup topology on  $B_\lambda$ .*

A topological semigroup  $S$  is called  $\Gamma$ -compact if

$$\Gamma(x) = \overline{\{x, x^2, x^3, \dots, x^n, \dots\}}$$

is a compact subsemigroup of  $S$  for every  $x \in S$ . Obviously every compact semigroup is  $\Gamma$ -compact. J. A. Hildebrandt and R. J. Koch proved that every  $\Gamma$ -compact topological semigroup and hence compact topological semigroup does not contain the bicyclic semigroup [15]. Since for any element  $a$  of the semigroup of matrix units we have either  $aa = a$  or  $aa = 0$ , the semigroup of matrix units is  $\Gamma$ -compact.

**Question 1.** *Does there exist a compact topological inverse semigroup which contains the semigroup  $B_\omega$ ?*

In this paper we give a negative answer to Question 1. Moreover, we show that if  $\lambda \geq \omega$ , then every continuous homomorphism of the topological semigroup  $B_\lambda$  into a compact topological semigroup is annihilating and  $B_\lambda$  as a topological inverse semigroup is absolutely  $H$ -closed in the class of topological inverse semigroups.

**Lemma 4.** *Let  $T$  be a dense subsemigroup of a topological semigroup  $S$  and  $0$  be the zero of  $T$ . Then  $0$  is the zero of  $S$ .*

*Proof.* Suppose that there exists  $a \in S \setminus T$  such that  $0 \cdot a = b \neq 0$ . Then for every open neighbourhood  $U(b) \not\cong 0$  in  $S$  there exists an open neighbourhood  $V(a) \not\cong 0$  in  $S$  such that  $0 \cdot V(a) \subseteq U(b)$ . But  $|V(a) \cap T| \geq \omega$ , and hence  $0 \in 0 \cdot V(a) \subseteq U(b)$ , a contradiction with the choice of  $U(b)$ . Therefore  $0 \cdot a = 0$  for all  $a \in S \setminus T$ .

The proof of the equality  $a \cdot 0 = 0$  is similar.  $\square$

**Lemma 5.** *If  $A$  is non-singleton in  $B_\lambda$ , then  $0 \in A \cdot A$ .*

*Proof.* Suppose  $|A| = 2$ . Obviously if  $0 \in A$ , then  $0 \in A \cdot A$ .

Let  $0 \notin A$  and  $A = \{(\alpha, \beta), (\gamma, \delta)\}$ . If  $(\alpha, \beta), (\gamma, \delta)$  are idempotents of  $B_\lambda$ , then  $\alpha = \beta, \gamma = \delta, \beta \neq \gamma$ , and hence  $0 = (\alpha, \beta) \cdot (\gamma, \delta)$ . If the set  $A$  contains a non-idempotent element  $(\alpha, \beta)$ , then  $\alpha \neq \beta$  and  $0 = (\alpha, \beta) \cdot (\alpha, \beta) \in A \cdot A$ .  $\square$

**Lemma 6.** *Let  $\lambda \geq \omega$  and let  $B_\lambda$  be a dense subsemigroup of a topological semigroup  $S$ . Then  $a \cdot a = 0$  for all  $a \in S \setminus B_\lambda$ .*

*Proof.* Suppose  $a \cdot a = b \neq 0$  for some  $a \in S \setminus B_\lambda$ . Then for any open neighbourhood  $U(b) \not\cong 0$  in  $S$  there exists an open neighbourhood  $V(a) \not\cong 0$  in  $S$  such that  $V(a) \cdot V(a) \subseteq U(b)$ . But  $|V(a) \cap B_\lambda| \geq \omega$  and by Lemma 5,  $0 \in V(a) \cdot V(a) \subseteq U(b)$ , a contradiction.  $\square$

**Theorem 3.** *If  $\lambda \geq \omega$ , then there exists no semigroup topology  $\tau$  on  $B_\lambda$  such that  $(B_\lambda, \tau)$  is embedded into a compact topological semigroup.*

*Proof.* Suppose, on the contrary, that there exists a semigroup topology  $\tau$  on  $B_\lambda$  such that  $(B_\lambda, \tau)$  is a subsemigroup of some compact topological semigroup  $S$ . By Proposition 1,  $B_\lambda$  is not a closed subsemigroup of  $S$ . Without loss of generality we assume that  $B_\lambda$  is a dense subsemigroup of  $S$ . We denote  $X = B_\lambda \setminus \{0\}$ . By Lemma 2,  $X$  is a discrete subspace of  $B_\lambda$  and hence  $X$  is a locally compact subspace in  $B_\lambda$ . Thus, by Theorem 3.5.8 [11],  $\mathcal{J} = S \setminus X$  is a closed subspace of  $S$ . Therefore,  $X$  is a discrete subspace of  $S$ .

Further, we shall show that  $\mathcal{J}$  is an ideal of the semigroup  $S$ . By Lemmas 5 and 6 it is sufficient to prove that  $ax, xa, ab \in \mathcal{J}$  for every  $x \in B_\lambda \setminus \{0\}$ ,  $a, b \in \mathcal{J} \setminus \{0\}$ .

Assume that there exist  $x \in B_\lambda \setminus \{0\}$  and  $a \in \mathcal{J} \setminus \{0\}$  such that  $ax = c \notin \mathcal{J}$ . Then  $c$  is an isolated point in  $S$ . Thus for every open neighbourhood  $U(a) \not\ni 0$  at least one of the following conditions holds

- (i)  $|(U(a) \setminus \{a\}) \cdot x| \geq \omega$ ,
- (ii)  $0 \in (U(a) \setminus \{a\}) \cdot x$ .

But  $c$  is an isolated point in  $S$ , a contradiction. The proof for  $xa$  is similar.

Suppose  $ab = c \notin \mathcal{J}$  for some  $a, b \in \mathcal{J}$ . Then  $c$  is an isolated point in  $S$ . For every open neighbourhoods  $U(a) \not\ni 0$  and  $U(b) \not\ni 0$  at least one of the following conditions holds

- (iii)  $|(U(a) \setminus \{a\}) \cdot (U(b) \setminus \{b\})| \geq \omega$ ,
- (iv)  $0 \in (U(a) \setminus \{a\}) \cdot (U(b) \setminus \{b\})$ .

But  $c$  is an isolated point in  $S$ , a contradiction. Thus,  $ab = c \in \mathcal{J}$ .

Therefore,  $\mathcal{J}$  is a compact ideal of  $S$ .

By Theorem A.2.23 [16] the Rees quotient-semigroup  $S/\mathcal{J}$  is a compact topological semigroup. But the semigroup  $S/\mathcal{J}$  is algebraically isomorphic to  $B_\lambda$ , a contradiction with Proposition 1.  $\square$

A semigroup  $S$  is called *congruence-free* (*congruence-simple*, *h-simple*) if it has only two congruences: identical and universal [23]. Such semigroups E. S. Lyapin [18] and L. M. Gluskin [12] called *simple*. Obviously, a semigroup  $S$  is congruence-free if and only if every homomorphism  $h$  of  $S$  into an arbitrary semigroup  $T$  is an isomorphism "into" or is an annihilating homomorphism (i. e. there exists  $c \in T$  such that  $h(a) = c$  for all  $a \in S$ ).

Theorem 1 [12] implies



**Corollary 3.** *The semigroup  $B_\lambda$  is congruence-free for every cardinal  $\lambda \geq 2$ .*

Theorem 3 and Corollary 3 imply

**Proposition 2.** *Let  $\lambda \geq \omega$ . Then every continuous homomorphism of the topological semigroup  $B_\lambda$  into a compact topological semigroup is annihilating.*

Recall [8] that a *Bohr compactification* of a topological semigroup  $S$  is a pair  $(\beta, B(S))$  such that  $B(S)$  is a compact topological semigroup,  $\beta: S \rightarrow B(S)$  is a continuous homomorphism, and if  $g: S \rightarrow T$  is a continuous homomorphism of  $S$  into a compact semigroup  $T$ , then there exists a unique continuous homomorphism  $f: B(S) \rightarrow T$  such that the diagram

$$\begin{array}{ccc} S & \xrightarrow{\beta} & B(S) \\ g \downarrow & \searrow f & \\ T & & \end{array}$$

commutes.

Let  $S$  be a topological semigroup. Let  $\{(T_\alpha, \varphi_\alpha)\}_{\alpha \in \mathcal{A}}$  be a family of pairs of compact topological semigroups and continuous homomorphisms  $\varphi_\alpha: S \rightarrow T_\alpha$ , respectively, such that  $\varphi_\alpha(S)$  is a dense subsemigroup of  $T_\alpha$  for any  $\alpha \in \mathcal{A}$ . Then  $B(S)$  is a subsemigroup of  $\prod_{\alpha \in \mathcal{A}} T_\alpha$  (see the proofs of Lemma 2.43 and Theorem 2.44 [6, Vol. 1]), where  $|\mathcal{A}| \leq 2^{2^{|S|}}$ . Therefore Proposition 2 implies

**Corollary 4.** *If  $\lambda \geq \omega$ , then the Bohr compactification of the topological semigroup  $B_\lambda$  is a trivial semigroup.*

**Theorem 4.** *Let  $\lambda$  be a cardinal  $\geq 2$  and  $B_\lambda$  be a subsemigroup of a topological inverse semigroup  $S$ . Then  $B_\lambda$  is a closed subsemigroup of  $S$ .*

*Proof.* If  $\lambda < \omega$  then  $B_\omega$  is finite and hence  $B_\lambda$  is a closed subsemigroup of  $S$ .

Suppose  $\lambda \geq \omega$ .

Let  $\overline{B_\lambda} = S_1$  in  $S$ . Then by Proposition II.2 [10]  $S_1$  is a topological inverse semigroup and by Lemma 4 the zero  $0$  of the semigroup  $B_\lambda$  is the zero of the semigroup  $S_1$ .

Let  $b$  be any element of  $S_1 \setminus B_\lambda$ . We consider two cases:  $b \in E(S_1)$  and  $b \in S_1 \setminus E(S_1)$ .

- 1) Let  $b \in E(S_1)$ . Then for every open neighbourhood  $W(b) \not\ni 0$  there exists an open neighbourhood  $U(b) \not\ni 0$  such that  $U(b) \cdot U(b) \subseteq$

$W(b)$ . Since  $|U(b) \cap B_\lambda| \geq \omega$ , there exist  $\alpha, \beta, \gamma, \delta \in I_\lambda$  such that  $(\alpha, \beta), (\gamma, \delta) \in U(b)$ , and  $\beta \neq \gamma$  or  $\alpha \neq \delta$ . Then  $0 \in U(b) \cdot U(b) \subseteq W(b)$ , a contradiction. Therefore,  $E(S_1) = E(B_\lambda)$ .

- 2) Let  $b \in S_1 \setminus E(S_1)$ . Then  $b^{-1} \in S_1 \setminus E(S_1)$ . Since 0 is the zero of the topological semigroup  $S_1$ , then  $b \cdot b^{-1} \neq 0$  and  $b^{-1} \cdot b \neq 0$ . Otherwise, if  $b \cdot b^{-1} = 0$  or  $b^{-1} \cdot b = 0$ , then  $b = b \cdot b^{-1} \cdot b = 0 \cdot b = 0$  or  $b = b \cdot b^{-1} \cdot b = b \cdot 0 = 0$ , a contradiction with  $b \in S_1 \setminus E(S_1)$ .

Therefore, there exist  $e, f \in E(S_1) = E(B_\lambda)$  such that  $b \cdot b^{-1} = e$ ,  $b^{-1} \cdot b = f$  and  $e \neq f$ . Let  $W(e)$  and  $W(f)$  be open neighbourhoods of  $e$  and  $f$  in  $S_1$ , respectively, such that  $0 \notin W(e)$  and  $0 \notin W(f)$ . Then there exist disjoint open neighbourhoods  $U(b) \not\ni 0$  and  $U(b^{-1}) \not\ni 0$  such that  $U(b) \cdot U(b^{-1}) \subseteq W(e)$  and  $U(b^{-1}) \cdot U(b) \subseteq W(f)$ . Since  $|U(b) \cap B_\lambda| \geq \omega$  and  $|U(b^{-1}) \cap B_\lambda| \geq \omega$ , there exist  $(\alpha, \beta) \in U(b)$  and  $(\gamma, \delta) \in U(b^{-1})$  such that  $\beta \neq \gamma$  or  $\alpha \neq \delta$ . Therefore,  $0 \in U(b) \cdot U(b^{-1}) \subseteq W(e)$  or  $0 \in U(b^{-1}) \cdot U(b) \subseteq W(f)$ , a contradiction with  $0 \notin W(e)$  and  $0 \notin W(f)$ .

If  $e = f$  the proof of the statement is similar.

The obtained contradictions imply the statement of the theorem.  $\square$

Since a compact topological semigroup is stable (see Theorem 3.31 [6]) a compact 0-simple topological inverse semigroup  $S$  is completely 0-simple and by Theorem 3.9 [7]  $S$  is algebraically isomorphic to the Brandt  $\lambda$ -extension of a group. Theorems 3 and 4 imply

**Corollary 5.** *Let  $S$  be a compact 0-simple topological inverse semigroup. Then  $E(S)$  is finite.*

**Definition 1** ([25]). Let  $\mathcal{S}$  be a class of topological semigroups. A semigroup  $S \in \mathcal{S}$  is called *H-closed in  $\mathcal{S}$* , if  $S$  is a closed subsemigroup of any topological semigroup  $T \in \mathcal{S}$  which contains  $S$  as subsemigroup. If  $\mathcal{S}$  coincides with the class of all topological semigroups, then the semigroup  $S$  is called *H-closed*.

We remark that in [25] *H-closed* semigroups are called *maximal*.

Theorem 4 implies

**Corollary 6.** *Let  $\lambda$  be a cardinal  $\geq 2$  and  $\tau$  be a semigroup inverse topology on  $B_\lambda$ . Then  $(B_\lambda, \tau)$  is H-closed in the class of topological inverse semigroups.*

**Definition 2** ([26]). Let  $\mathcal{S}$  be a class of topological semigroups. A topological semigroup  $S \in \mathcal{S}$  is called *absolutely H-closed in the class  $\mathcal{S}$*  if any continuous homomorphic image of  $S$  into  $T \in \mathcal{S}$  is *H-closed in  $\mathcal{S}$* . If  $\mathcal{S}$  coincides with the class of all topological semigroups, then the semigroup  $S$  is called *absolutely H-closed*.

Corollary 3 and Theorem 4 imply

**Corollary 7.** *Let  $\lambda$  be a cardinal  $\geq 2$  and  $\tau$  be a semigroup inverse topology on  $B_\lambda$ . Then  $(B_\lambda, \tau)$  is absolutely  $H$ -closed in the class of topological inverse semigroups.*

Let  $\mathcal{S}$  be a class of topological semigroups. A semigroup  $S$  is called *algebraically  $h$ -closed in  $\mathcal{S}$*  if  $S$  with discrete topology  $d$  is absolutely  $H$ -closed in  $\mathcal{S}$  and  $(S, d) \in \mathcal{S}$ . If  $\mathcal{S}$  coincides with the class of all topological semigroups, then the semigroup  $S$  is called *algebraically  $h$ -closed*.

Absolutely  $H$ -closed semigroups and algebraically  $h$ -closed semigroups were introduced by J. W. Stepp in [26]. There they were called *absolutely maximal* and *algebraic maximal*, respectively.

Corollary 7 implies

**Proposition 3.** *For any cardinal  $\lambda \geq 2$  the semigroup  $B_\lambda$  is algebraically  $h$ -closed in the class of topological inverse semigroups.*

The following example shows that  $B_\omega$  with the discrete topology is not  $H$ -closed.

**Example 2.** Let  $B_\omega = I_\omega \times I_\omega \cup \{0\}$  be the semigroup of matrix units and  $a \notin B_\omega$ . Let  $S = B_\omega \cup \{a\}$ . We put

$$a \cdot a = a \cdot 0 = 0 \cdot a = a \cdot (\alpha, \beta) = (\alpha, \beta) \cdot a = 0$$

for all  $(\alpha, \beta) \in B_\omega \setminus \{0\}$ .

Further we enumerate the elements of the set  $I_\omega$  by natural numbers. Let  $A_n = \{(2k-1, 2k) \mid k \geq n\}$  for each  $n \in \mathbb{N}$ . A topology  $\tau$  on  $S$  is defined as follows:

- 1) all points of  $B_\omega$  are isolated in  $S$ ;
- 2)  $\mathcal{B}(a) = \{U_n(a) = \{a\} \cup A_n \mid n \in \mathbb{N}\}$  is the base of the topology  $\tau$  at the point  $a \in S$ .

Then

- a)  $\{(l, m)\} \cdot U_n(a) = U_n(a) \cdot \{(l, m)\} = \{0\}$  for all  $(l, m) \in B_\omega \setminus \{0\}$ ,  $n \geq \max\{l, m\}$ ;
- b)  $U_n(a) \cdot U_n(a) = U_n(a) \cdot \{0\} = \{0\} \cdot U_n(a) = \{0\}$  for any  $n \in \mathbb{N}$ ;
- c)  $U_n(a)$  is a compact subset of  $S$  for each  $n \in \mathbb{N}$ .

Therefore  $(S, \tau)$  is a locally compact topological semigroup. Obviously  $B_\omega$  is not a closed subset of  $(S, \tau)$ .

**Remark 2.** Let  $\lambda_1$  and  $\lambda_2$  be cardinals and  $\lambda_1 \leq \lambda_2$ . Then  $B_{\lambda_1}$  is a subsemigroup of  $B_{\lambda_2}$ . Example 2 implies that for any infinite cardinal  $\lambda$  the semigroup  $B_\lambda$  is not  $H$ -closed.

**Definition 3.** A Hausdorff topological (inverse) semigroup  $(S, \tau)$  is said to be *minimal* if no Hausdorff semigroup (inverse) topology on  $S$  is strictly contained in  $\tau$ . If  $(S, \tau)$  is minimal topological (inverse) semigroup, then  $\tau$  is called *minimal semigroup (inverse) topology*.

The concept of minimal topological groups was introduced independently in the early 1970's by Doitchinov [9] and Stephenson [24]. Both authors were motivated by the theory of minimal topological spaces, which was well understood at that time (cf. [4]). More than 20 years earlier L. Nachbin [19] had studied minimality in the context of division rings, and B. Banaschewski [3] investigated minimality in the more general setting of topological algebras.

**Question 2** (T. O. Banach). *Is it true that for any cardinal  $\lambda \geq \omega$  the semigroup  $B_\lambda$  admits minimal (inverse) semigroup topology?*

For each  $\alpha, \beta \in I_\lambda$  we define

$$V_\alpha = B_\lambda \setminus \{(\alpha, \gamma) \mid \gamma \in I_\lambda\} \quad \text{and} \quad H_\beta = B_\lambda \setminus \{(\gamma, \beta) \mid \gamma \in I_\lambda\}.$$

Put

$$U^{\alpha_1, \dots, \alpha_n} = \bigcap_{i=1}^n V_{\alpha_i}, \quad U_{\beta_1, \dots, \beta_m} = \bigcap_{i=1}^m H_{\beta_i} \quad \text{and}$$

$$U_{\beta_1, \dots, \beta_m}^{\alpha_1, \dots, \alpha_n} = U^{\alpha_1, \dots, \alpha_n} \cap U_{\beta_1, \dots, \beta_m},$$

where  $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_m \in I_\lambda$ ,  $n, m \in \mathbb{N}$ . Further we define the following families

$$\mathcal{B}_{mv} = \{U^{\alpha_1, \dots, \alpha_n} \mid \alpha_1, \dots, \alpha_n \in I_\lambda, n \in \mathbb{N}\} \cup \{(\alpha, \beta) \mid \alpha, \beta \in I_\lambda\},$$

$$\mathcal{B}_{mh} = \{U_{\beta_1, \dots, \beta_m} \mid \beta_1, \dots, \beta_m \in I_\lambda, m \in \mathbb{N}\} \cup \{(\alpha, \beta) \mid \alpha, \beta \in I_\lambda\},$$

$$\mathcal{B}_{mi} = \{U_{\beta_1, \dots, \beta_m}^{\alpha_1, \dots, \alpha_n} \mid \alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_m \in I_\lambda, n, m \in \mathbb{N}\} \cup \{(\alpha, \beta) \mid \alpha, \beta \in I_\lambda\}.$$

Obviously, the conditions (BP1)–(BP3) [11] hold for families  $\mathcal{B}_{mv}$ ,  $\mathcal{B}_{mh}$  and  $\mathcal{B}_{mi}$ , and hence  $\mathcal{B}_{mv}$ ,  $\mathcal{B}_{mh}$  and  $\mathcal{B}_{mi}$  are bases on  $B_\lambda$  of topologies  $\tau_{mv}$ ,  $\tau_{mh}$  and  $\tau_{mi}$ , respectively.

**Lemma 7.** *Let  $\lambda$  be an infinite cardinal. Then*

- (i)  $(B_\lambda, \tau_{mv})$  is a topological semigroup;

(ii)  $(B_\lambda, \tau_{mh})$  is a topological semigroup;

(iii)  $(B_\lambda, \tau_{mi})$  is a topological inverse semigroup.

*Proof.* (i) Since the following conditions hold

$$U^{\alpha_1, \dots, \alpha_n} \cdot U^{\alpha_1, \dots, \alpha_n} \subseteq U^{\alpha_1, \dots, \alpha_n}, \quad (\alpha, \beta) \cdot U^{\alpha_1, \dots, \alpha_n, \beta} = \{0\} \subseteq U^{\alpha_1, \dots, \alpha_n},$$

$$U^{\alpha_1, \dots, \alpha_n} \cdot (\alpha, \beta) = \{0\} \cup \{(\gamma, \delta) \mid \gamma \in I_\lambda \setminus \{\alpha_1, \dots, \alpha_n\}\} \subseteq U^{\alpha_1, \dots, \alpha_n}$$

for every open neighbourhood  $U^{\alpha_1, \dots, \alpha_n}$  of the zero of  $B_\lambda$  and for any  $(\alpha, \beta) \in B_\lambda \setminus \{0\}$ ,  $(B_\lambda, \tau_{mv})$  is a topological semigroup.

The proof of statement (ii) is similar to the proof of item (i).

(iii) For every open neighbourhood  $U^{\alpha_1, \dots, \alpha_n}_{\beta_1, \dots, \beta_m}$  of the zero of  $B_\lambda$  and for any  $(\beta, \alpha) \in B_\lambda \setminus \{0\}$  we have:

$$U^{\alpha_1, \dots, \alpha_n}_{\beta_1, \dots, \beta_m} \cdot U^{\alpha_1, \dots, \alpha_n}_{\beta_1, \dots, \beta_m} \subseteq U^{\alpha_1, \dots, \alpha_n}_{\beta_1, \dots, \beta_m}, \quad (\beta, \alpha) \cdot U^{\alpha_1, \dots, \alpha_n}_{\alpha, \beta_1, \dots, \beta_m} = \{0\} \subseteq U^{\alpha_1, \dots, \alpha_n}_{\beta_1, \dots, \beta_m},$$

$$U^{\beta, \alpha_1, \dots, \alpha_n}_{\beta_1, \dots, \beta_m} \cdot (\beta, \alpha) = \{0\} \subseteq U^{\alpha_1, \dots, \alpha_n}_{\beta_1, \dots, \beta_m}, \quad \left( U^{\beta_1, \dots, \beta_m}_{\alpha_1, \dots, \alpha_n} \right)^{-1} \subseteq U^{\alpha_1, \dots, \alpha_n}_{\beta_1, \dots, \beta_m}.$$

Therefore,  $(B_\lambda, \tau_{mi})$  is a topological inverse semigroup.  $\square$

We remark that  $\tau_{mv}$ ,  $\tau_{mh}$  and  $\tau_{mi}$  are not locally compact topologies on  $B_\lambda$  for  $\lambda \geq \omega$ .

For  $A \subseteq I_\lambda$  and  $a \in I_\lambda$  we denote  $A^\alpha = \{(\alpha, \beta) \in B_\lambda \mid \beta \in A\}$  and  $A_\alpha = \{(\beta, \alpha) \in B_\lambda \mid \beta \in A\}$ .

**Lemma 8.** *Let  $\lambda$  be an infinite cardinal,  $B_\lambda$  be a topological semigroup, and  $A^\alpha [A_\alpha]$  be a closed subset in  $B_\lambda$  for some  $\alpha \in I_\lambda$ . Then  $A^\beta [A_\beta]$  is a closed subset of  $B_\lambda$  for any  $\beta \in I_\lambda$ .*

*Proof.* Since  $B_\lambda$  is a topological semigroup, then the map  $\lambda_{(\alpha, \beta)}: B_\lambda \rightarrow B_\lambda$  [ $\rho_{(\beta, \alpha)}: B_\lambda \rightarrow B_\lambda$ ] defined by the formula  $\lambda_{(\alpha, \beta)}(x) = (\alpha, \beta) \cdot x$  [ $\rho_{(\beta, \alpha)}(x) = x \cdot (\beta, \alpha)$ ] is continuous. Therefore  $A^\beta = (\lambda_{(\alpha, \beta)})^{-1}(A^\alpha)$  [ $A_\beta = (\rho_{(\beta, \alpha)})^{-1}(A_\alpha)$ ] is a closed subset of  $B_\lambda$ .  $\square$

For any  $A \subseteq I_\lambda$ ,  $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_m \in I_\lambda$ ,  $n, m \in \mathbb{N}$  we denote

$$U^{\alpha_1, \dots, \alpha_n}(A) = U^{\alpha_1, \dots, \alpha_n} \cup \{(\alpha_i, x) \mid x \in A, i = 1, \dots, n\},$$

$$U_{\beta_1, \dots, \beta_m}(A) = U_{\beta_1, \dots, \beta_m} \cup \{(x, \beta_i) \mid x \in A, i = 1, \dots, m\},$$

$$U(\alpha_1, \dots, \alpha_n; A) = U^{\alpha_1, \dots, \alpha_n}(A) \cap U_{\alpha_1, \dots, \alpha_n}(A).$$

The following theorem gives a positive answer to Question 2.

**Theorem 5.** *Let  $\lambda$  be an infinite cardinal. Then the following conclusions hold:*

- (i)  $\tau_{mv}$  is a minimal semigroup topology on  $B_\lambda$ ;
- (ii)  $\tau_{mh}$  is a minimal semigroup topology on  $B_\lambda$ ;
- (iii)  $\tau_{mi}$  is the coarsest semigroup inverse topology on  $B_\lambda$ , and hence is minimal semigroup inverse.

*Proof.* (i) Suppose that there exists a Hausdorff semigroup topology  $\tau_0$  on  $B_\lambda$  which is coarser than  $\tau_{mv}$ . Let  $V_0$  be an element of a base of the topology  $\tau_0$  at the zero of  $B_\lambda$ . Then by Lemma 8 there exist an infinite subset  $A$  in  $I_\lambda$  and  $\alpha_1, \dots, \alpha_m \in I_\lambda$  ( $m \in \mathbb{N}$ ) such that  $U^{\alpha_1, \dots, \alpha_m}(A) \subseteq V_0$ . For any  $\beta \in I_\lambda$ ,  $\gamma \in I_\lambda \setminus \{\alpha_1, \dots, \alpha_m\}$  the following conditions hold:

- a)  $(\alpha_i, \beta) = (\alpha_i, \delta) \cdot (\delta, \beta)$ , where  $\delta \in A \setminus \{\alpha_1, \dots, \alpha_m\}$  and, obviously,  $(\alpha_i, \delta), (\delta, \beta) \in U^{\alpha_1, \dots, \alpha_m}(A)$  ( $i = 1, \dots, m$ );
- b)  $(\gamma, \beta) = (\gamma, \gamma) \cdot (\gamma, \beta)$ , and  $(\gamma, \gamma), (\gamma, \beta) \in U^{\alpha_1, \dots, \alpha_m}(A)$ .

Thus,

$$B_\lambda = U^{\alpha_1, \dots, \alpha_m}(A) \cdot U^{\alpha_1, \dots, \alpha_m}(A) \subseteq V_0 \cdot V_0$$

for any element  $V_0$  of a base of the topology  $\tau_0$  at the zero of  $B_\lambda$ . This gives a contradiction with the continuity of the semigroup operation in  $(B_\lambda, \tau_0)$ . Therefore  $(B_\lambda, \tau_{mv})$  is a minimal topological semigroup.

The proof of item (ii) is similar to the proof of item (i).

(iii) Let  $\tau$  be any Hausdorff semigroup inverse topology on  $B_\lambda$ . We define the maps:  $\varphi: B_\lambda \rightarrow E(B_\lambda)$  and  $\psi: B_\lambda \rightarrow E(B_\lambda)$  by formulae  $\varphi(x) = xx^{-1}$  and  $\psi(x) = x^{-1}x$ . Since the topology  $\tau$  is Hausdorff then the sets  $\varphi^{-1}((\alpha, \alpha)) = \{(\alpha, \gamma) \mid \gamma \in I_\lambda\}$  and  $\psi^{-1}((\alpha, \alpha)) = \{(\gamma, \alpha) \mid \gamma \in I_\lambda\}$  are closed for each  $\alpha \in I_\lambda$ , and hence  $U_{\beta_1, \dots, \beta_m}^{\alpha_1, \dots, \alpha_n} \in \tau$  for all  $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_m \in I_\lambda$ . Therefore  $\tau_{mi} \subseteq \tau$ .  $\square$

**Theorem 6.** *Let  $\lambda$  be an infinite cardinal. Then  $(B_\lambda, \tau_{mv})$ ,  $(B_\lambda, \tau_{mh})$ ,  $(B_\lambda, \tau_{mi})$  are  $H$ -closed topological semigroups.*

*Proof.* We shall show that the semigroup  $(B_\lambda, \tau_{mi})$  is  $H$ -closed. The proofs of  $H$ -closedness of the semigroups  $(B_\lambda, \tau_{mh})$  and  $(B_\lambda, \tau_{mv})$  are similar.

Suppose that there exists a topological semigroup  $S$  which contains  $(B_\lambda, \tau_{mi})$  as a non-closed subsemigroup. Then there exists  $x \in \overline{B_\lambda} \setminus B_\lambda \subseteq S$ . By Lemma 4,  $x \cdot 0 = 0 \cdot x = 0$ . Then for every open neighbourhood  $W(0)$  in  $S$  there exist open neighbourhoods  $U(0)$ ,  $V(0)$ , and  $V(x)$  in  $S$  such that  $V(0) \cap V(x) = \emptyset$ ,  $U(0) \cap V(x) = \emptyset$ ,  $V(0) \subseteq W(0)$ ,  $U(0) \subseteq$

$W(0), V(x) \cdot V(0) \subseteq U(0)$ , and  $V(0) \cdot V(x) \subseteq U(0)$ . We can suppose that  $U_{\beta_1, \dots, \beta_m}^{\alpha_1, \dots, \alpha_n} = U(0) \cap B_\lambda$  for some  $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_m \in I_\lambda$ .

Since  $|V(x) \cap B_\lambda| \geq \omega$ , one of the following conditions holds:

- 1) the set  $B_{i_0} = V(x) \cap \{(\alpha_{i_0}, \gamma) \mid \gamma \in I_\lambda\}$  is infinite for some  $i_0 \in \{1, \dots, n\}$ ;
- 2) the set  $B^{j_0} = V(x) \cap \{(\gamma, \alpha_{j_0}) \mid \gamma \in I_\lambda\}$  is infinite for some  $j_0 \in \{1, \dots, m\}$ .

In the first case we put

$$\Gamma_{i_0} = \{\gamma \in I_\lambda \mid (\alpha_{i_0}, \gamma) \in V(x)\}.$$

Then the set  $\{(\gamma, \gamma) \mid \gamma \in \Gamma_{i_0}\} \cap U_{\delta_1, \dots, \delta_l}^{\gamma_1, \dots, \gamma_k}$  is infinite for any basic neighbourhood  $U_{\delta_1, \dots, \delta_l}^{\gamma_1, \dots, \gamma_k}$ ,  $\gamma_1, \dots, \gamma_k, \delta_1, \dots, \delta_l \in I_\lambda$ . Thus

$$B_{i_0} \cdot U_{\delta_1, \dots, \delta_l}^{\gamma_1, \dots, \gamma_k} \not\subseteq U_{\beta_1, \dots, \beta_m}^{\alpha_1, \dots, \alpha_n},$$

a contradiction with  $V(x) \cdot V(0) \subseteq U(0)$ .

In the other case we put

$$\Gamma^{j_0} = \{\gamma \in I_\lambda \mid (\gamma, \alpha_{j_0}) \in V(x)\}.$$

Then the set  $\{(\gamma, \gamma) \mid \gamma \in \Gamma^{j_0}\} \cap U_{\delta_1, \dots, \delta_l}^{\gamma_1, \dots, \gamma_k}$  is infinite for every basic neighbourhood  $U_{\delta_1, \dots, \delta_l}^{\gamma_1, \dots, \gamma_k}$ ,  $\gamma_1, \dots, \gamma_k, \delta_1, \dots, \delta_l \in I_\lambda$ . Hence

$$U_{\delta_1, \dots, \delta_l}^{\gamma_1, \dots, \gamma_k} \cdot B^{j_0} \not\subseteq U_{\beta_1, \dots, \beta_m}^{\alpha_1, \dots, \alpha_n},$$

a contradiction with  $V(0) \cdot V(x) \subseteq U(0)$ .

Therefore the topological semigroup  $(B_\lambda, \tau_{mi})$  is  $H$ -closed.  $\square$

Theorem 6 and Corollary 3 imply

**Corollary 8.** *Let  $\lambda$  be an infinite cardinal. Then  $(B_\lambda, \tau_{mv})$  and  $(B_\lambda, \tau_{mh})$  are absolutely  $H$ -closed topological semigroups.*

**Theorem 7.** *For every cardinal  $\lambda \geq \omega$  any continuous homomorphism from  $(B_\lambda, \tau_{mv})$   $[(B_\lambda, \tau_{mh})]$  into a locally compact topological semigroup  $S$  is annihilating.*

*Proof.* Let  $h: (B_\lambda, \tau_{mv}) \rightarrow S$  be a continuous homomorphism. If  $h$  is not annihilating, then by Corollary 3,  $h$  is algebraic isomorphism, and hence, since  $(B_\lambda, \tau_{mv})$  is a minimal topological semigroup,  $h: B_\lambda \rightarrow S$  is a topological embedding.

By Theorem 6,  $h(B_\lambda)$  is a closed subsemigroup of  $S$  and by Theorem 3.3.8 [11],  $h(B_\lambda)$  is a locally compact topological semigroup. This is a contradiction with the fact that  $\tau_{mv}$  is not a locally compact semigroup topology on  $B_\lambda$ .

The proof of the theorem for the semigroup  $(B_\lambda, \tau_{mh})$  is similar.  $\square$

**Acknowledgement.** The authors are deeply grateful to T. Banach and I. Guran for useful discussions of the results of this paper.

**Add to Proof.** The authors express their sincere thanks to the referee for very careful reading the manuscript and valuable remarks improving the presentation.

### References

- [1] O. Andersen, *Ein Bericht über die Struktur abstrakter Halbgruppen*, PhD Thesis, Hamburg, 1952.
- [2] L. W. Anderson, R. P. Hunter, and R. J. Koch, *Some results on stability in semigroups*, Trans. Amer. Math. Soc. **117**:5 (1964) 521–529.
- [3] B. Banaschewski, *Minimal topological algebras*, Math. Ann. **211** (1974), 107–114.
- [4] M. P. Berri, J. R. Porter, and R. M. Stephenson, Jr., *A survey of minimal topological spaces*, "General Topology and its Relations to Modern Analysis and Algebra". Proc. (1968) Kanpur Topol. Conf. Ed.: S. P. Franklin, Z. Frolik, and V. Koutnik. Academic. Praha (1971), 93–114.
- [5] M. O. Bertman, and T. T. West, *Conditionally compact bicyclic semitopological semigroups*, Proc. Roy. Irish Acad. **A76**:21–23 (1976), 219–226.
- [6] J. H. Carruth, J. A. Hildebrandt, and R. J. Koch, "The Theory of Topological Semigroups," I, II, Marcell Dekker, Inc., New York and Basel, 1983 and 1986.
- [7] A. H. Clifford, and G. B. Preston, "The Algebraic Theory of Semigroups," I, II, Amer. Math. Soc., Surveys 7, 1961 and 1967.
- [8] K. DeLeeuw, and I. Glicksberg, *Almost-periodic functions on semigroups*, Acta Math. **105** (1961), 99–140.
- [9] D. Doitchinov, *Produits de groupes topologiques minimaux*, Bull. Sci. Math. (2) **97** (1972), 59–64.
- [10] C. Eberhart, and J. Selden, *On the closure of the bicyclic semigroup*, Trans. Amer. Math. Soc. **144** (1969), 115–126.
- [11] R. Engelking, "General Topology," PWN, Warszawa, 2nd ed., 1986.
- [12] L. M. Gluskin, *Simple semigroups with zero*, Doklady Akademii Nauk SSSR **103**:1 (1955), 5–8, in Russian.
- [13] O. V. Gutik, *On Howie semigroup*, Mat. Metody Phys.-Mech. Polya **42**:4 (1999), 127–132, in Ukrainian.
- [14] O. V. Gutik, and K. P. Pavlyk, *The H-closedness of topological inverse semigroups and topological Brandt  $\lambda$ -extensions*, XVI Open Scientific and Technical Conference of Young Scientists and Specialists of the Karpenko Physico-Mechanical Institute of NASU, May 16–18, 2001, Lviv, Materials. Lviv (2001), 240–243.
- [15] J. A. Hildebrandt, and R. J. Koch, *Swelling actions of  $\Gamma$ -compact semigroups*, Semigroup Forum **33**:1 (1986) 65–85.



- [16] K. H. Hofmann, and P. S. Mostert, “Elements of Compact Semigroups,” Charles E. Merrill Books, Inc., Columbus, Ohio, 1966.
- [17] R. J. Koch, and A. D. Wallace, *Stability in semigroups*, Duke Math. J. **24** (1957), 193–195.
- [18] E. S. Lyapin, *Simple commutative associative systems*, Izvestiya Akademii Nauk SSSR, Ser. Mat. **14**:3 (1950), 275–285, in Russian.
- [19] L. Nachbin, *On strictly minimal topological division rings*, Bull. Amer. Math. Soc. **55** (1949), 1128–1136.
- [20] L. O’Carroll, *Counterexamples in stable semigroups*, Trans. Amer. Math. Soc. **146** (1969), 377–386.
- [21] A. B. Paalman-de-Miranda, “Topological Semigroup,” Mathematical Centre Tracts. Vol. 11. Mathematisch Centrum, Amsterdam, 1964.
- [22] W. Ruppert, “Compact Semitopological Semigroups; An Intrinsic Theory,” Lecture Notes in Mathematics, Vol. 1079. Springer, Berlin, 1984.
- [23] B. M. Schein, *Homomorphisms and subdirect decompositions of semigroups*, Pacific J. Math. **17**:3 (1966), 529–547.
- [24] R. M. Stephenson, Jr., *Minimal topological groups*, Math. Ann. **192** (1971), 193–195.
- [25] J. W. Stepp, *A note on maximal locally compact semigroups*, Proc. Amer. Math. Soc. **20**:1 (1969), 251–253.
- [26] J. W. Stepp, *Algebraic maximal semilattices*, Pacific J. Math. **58**:1 (1975), 243–248.

## CONTACT INFORMATION

**O. V. Gutik**

Department of Algebra, Pidstryhach Institute for Applied Problems of Mechanics and Mathematics of National Academy of Sciences, Naukova 3b, Lviv, 79060, Ukraine, and Department of Mathematics, Ivan Franko Lviv National University, Universytetska 1, Lviv, 79000, Ukraine  
*E-Mail:* ovgutik@yahoo.com,  
topology@franko.lviv.ua

**K. P. Pavlyk**

Department of Algebra, Pidstrygach Institute for Applied Problems of Mechanics and Mathematics of National Academy of Sciences, Naukova 3b, Lviv, 79060, Ukraine  
*E-Mail:* katyapavlyk@rambler.ru

Received by the editors: 16.06.2005  
and final form in 15.09.2005.