

ON TOTALLY REAL BISECTIONAL CURVATURE

CHORNG-SHI HOUH

ABSTRACT. A Kaehler manifold of dimension ≥ 3 is a complex space form if and only if it has constant totally real bisectional curvature.

Goldberg and Kobayashi [2] introduced the notion of holomorphic bisectional curvature on a Kaehler manifold. It is determined by two holomorphic planes. In this paper, we consider a Kaehler manifold with constant totally real (or called antiholomorphic) bisectional curvature, which is determined by an antiholomorphic plane and its image by the complex structure. Namely it is defined by $R(X, JX; Y, JY)$ for a totally real section $\{X, Y\}$. A complex space form is a Kaehler manifold of constant holomorphic sectional curvature. It turns out that a complex space form can be characterized by having constant totally real bisectional curvature.

1. Let M be a real $2n$ -dimensional Kaehler manifold with complex structure J and Riemann metric g . Let R be the curvature tensor field of M . Then we have $R(JX, JY) = R(X, Y)$ and $R(X, Y)JZ = JR(X, Y)Z$ for any vectors X, Y, Z tangent to M . We denote $R(X, Y; ZW)$ by

$$R(X, Y; Z, W) = g(R(X, Y)Z, W).$$

Then the sectional curvature of M determined by orthonormal vectors X and Y is given by $K(X, Y) = R(X, Y; Y, X)$. It is easy to see then

$$K(JX, JY) = K(X, Y), \quad K(X, JY) = K(JX, Y)$$

and

$$R(X, Y; Z, W) = R(JX, JY; Z, W) = R(X, Y; JZ, JW).$$

By a plane section we mean a 2-dimensional linear subspace of a tangent space. A plane section π is called holomorphic (respectively antiholomorphic or totally real) if $J\pi = \pi$ (respectively $J\pi$ is perpendicular to π). The sectional curvature for a holomorphic (respectively totally real) plane section is called holomorphic (respectively totally real) sectional curvature. A Kaehler manifold of constant holomorphic sectional curvature is called a *complex space form*. Let X be a unit vector in a holomorphic plane section, then it is clear

Received by the editors December 17, 1974 and, in revised form, March 3, 1975.

AMS (MOS) subject classifications (1970). Primary 53B35, 53B20; Secondary 53C55.

Key words and phrases. Totally real sectional curvature, totally real bisectional curvature, complex space form.

© American Mathematical Society 1976

that the holomorphic sectional curvature of that holomorphic plane section is $K(X, JX)$. We denote $H(X) = K(X, JX)$.

It is easy to see that orthonormal vectors X and Y span a totally real section if and only if X, Y and JY are orthonormal. Hence if we use $\{X, Y\}$ to denote the plane section spanned by orthonormal vectors X and Y , then that $\{X, Y\}$ is totally real implies that each of $\{X, JY\}$ and $\{Y, JX\}$ is also totally real.

We define the *totally real bisectional curvature* $H(X, Y)$ for a totally real section $\{X, Y\}$ by

$$H(X, Y) = -R(X, JX; Y, JY), \quad \{X, Y\} \text{ is totally real.}$$

We say M is of constant totally real bisectional curvature if $H(X, Y) =$ constant for every totally real plane section $\{X, Y\}$.

We shall prove the following:

THEOREM. *A Kaehler manifold M with $\dim M \geq 3$ is a complex space form if and only if M is of constant totally real bisectional curvature.*

2. By the first Bianchi identity we have

$$\begin{aligned} H(X, Y) &= -R(X, JX; Y, JY) = R(X, Y; JY, JX) + R(X, JY; JX, Y) \\ &= R(X, Y; Y, X) + R(X, JY; JY, X) \\ &= K(X, Y) + K(X, JY). \end{aligned}$$

Hence if M is of constant totally real sectional curvature then M is of constant totally real bisectional curvature.

Conversely we assume that $H(X, Y) = -R(X, JX; Y, JY) = C$, C being a constant. Then

$$(2.1) \quad C = K(X, Y) + K(X, JY).$$

Since $\{X, Y\}$ is totally real, the plane section $\{\sqrt{2}^{-1}(X + Y), \sqrt{2}^{-1}(JX - JY)\}$ is also totally real. Thus we have

$$\frac{1}{4}R(X + Y, JX + JY; JX - JY, -X + Y) = -C.$$

After expanding the left side of the above relation and some cancellation we have

$$\begin{aligned} &\frac{1}{4}R(X + Y, JX + JY; JX - JY, -X + Y) \\ &= \frac{1}{4}\{R(X, JX; X, JX) + R(Y, JY; Y, JY) + 2R(X, JX; Y, JY) \\ &\quad - 4R(X, JY; X, JY)\} \\ &= \frac{1}{4}\{-H(X) - H(Y) - 2C + 4K(X, JY)\}. \end{aligned}$$

And hence

$$(2.2) \quad H(X) + H(Y) - 4K(X, JY) = 2C.$$

We can replace Y by JY in (2.2). Noticing that $H(JY) = H(Y)$ we have

$$(2.3) \quad H(X) + H(Y) - 4K(X, Y) = 2C.$$

From (2.1), (2.2) and (2.3) we conclude that

$$H(X) + H(Y) = 4C, \quad K(X, Y) = K(X, JY) = \frac{1}{2}C.$$

Hence M has constant totally real sectional curvature. Thus M is of constant totally real bisectional curvature if and only if M is of constant totally real sectional curvature. On the other hand, Chen and Ogiue in [1] proved that if $\dim M \geq 3$, M has constant totally real sectional curvature if and only if M is a complex space form. The theorem is thus proved.

REFERENCES

1. B-Y. Chen and K. Ogiue, *Some characterizations of complex space forms*, *Duke Math. J.* **40** (1973), 797–799. MR **48** #9623.
2. S. I. Goldberg and S. Kobayashi, *Holomorphic bisectional curvature*, *J. Differential Geometry* **1** (1967), 225–233. MR **37** #3485.

DEPARTMENT OF MATHEMATICS, WAYNE STATE UNIVERSITY, DETROIT, MICHIGAN 48202