

**ON TRANSVERSELY ISOTROPIC FUNCTIONS OF VECTORS,
 SYMMETRIC SECOND-ORDER TENSORS AND SKEW-
 SYMMETRIC SECOND-ORDER TENSORS***

By

G. F. SMITH

Lehigh University

1. Introduction. There are five groups T_1, \dots, T_5 which define the symmetry properties of materials which are referred to as being transversely isotropic. We define these groups by listing the matrices which generate the groups:

$$\begin{aligned}
 T_1: & \mathbf{Q}(\theta), \\
 T_2: & \mathbf{Q}(\theta), \mathbf{R}_1 = \text{diag}(-1, 1, 1), \\
 T_3: & \mathbf{Q}(\theta), \mathbf{R}_3 = \text{diag}(1, 1, -1), \\
 T_4: & \mathbf{Q}(\theta), \mathbf{R}_1 = \text{diag}(-1, 1, 1), \mathbf{R}_3 = \text{diag}(1, 1, -1), \\
 T_5: & \mathbf{Q}(\theta), \mathbf{D}_2 = \text{diag}(-1, 1, -1).
 \end{aligned}
 \tag{1.1}$$

In (1.1), $\mathbf{Q}(\theta)$ denotes the matrix

$$\mathbf{Q}(\theta) = \begin{vmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{vmatrix}.
 \tag{1.2}$$

$\mathbf{Q}(\theta)$ corresponds to a rotation about the x_3 axis. \mathbf{R}_1 and \mathbf{R}_3 correspond to reflections in planes perpendicular to the x_1 axis and the x_3 axis respectively. \mathbf{D}_2 corresponds to a rotation through 180 degrees about the x_2 axis.

In this paper, we determine integrity bases for polynomial functions $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M, \mathbf{W}_1, \dots, \mathbf{W}_P)$ of N three-dimensional second-order symmetric tensors $\mathbf{A}_p = \|A_{ij}^p\|$ ($p = 1, \dots, N$), M three-dimensional vectors $\mathbf{V}_q = V_i^q$ ($q = 1, \dots, M$) and P three-dimensional second-order skew-symmetric tensors $\mathbf{W}_r = \|W_{ij}^r\|$ ($r = 1, \dots, P$) which are invariant under any given group chosen from T_1, \dots, T_5 . Adkins [1, 2] has considered the problem of determining integrity bases for functions $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M)$ which are invariant under the group T_1 and for functions $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M)$ which are invariant under the group T_2 . Long and McIntire [3] have considered the problem of determining an integrity basis for functions $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M, \mathbf{W}_1, \dots, \mathbf{W}_P)$ which are invariant under the group T_4 . The results obtained here for this case differ from those given in [3].

* Received November 6, 1980. This work was supported by a grant from the National Science Foundation to Lehigh University.

2. An integrity basis for functions invariant under T_1 . Let us employ the notation

$$\begin{aligned} B_\alpha^i &= A_{3\alpha}^i & (\alpha = 1, 2; i = 1, \dots, N), \\ B_\alpha^{N+j} &= V_\alpha^j & (\alpha = 1, 2; j = 1, \dots, M), \\ B_\alpha^{N+M+k} &= W_{3\alpha}^k & (\alpha = 1, 2; k = 1, \dots, P). \end{aligned} \tag{2.1}$$

It is readily seen that the problem of determining the form of a polynomial function $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M, \mathbf{W}_1, \dots, \mathbf{W}_P)$ which is invariant under the group T_1 is equivalent to that of determining the form of a polynomial function $G(A_{33}^i, \dots, B_1^r - \iota B_2^r)$ which is subject to the restrictions that

$$\begin{aligned} &G(A_{33}^i, A_{11}^i + A_{22}^i, A_{11}^i - A_{22}^i + 2\iota A_{12}^i, A_{11}^i - A_{22}^i - 2\iota A_{12}^i, V_3^p, \\ &W_{12}^m, B_1^r + \iota B_2^r, B_1^r - \iota B_2^r) = G(A_{33}^i, A_{11}^i + A_{22}^i, (A_{11}^i - A_{22}^i + 2\iota A_{12}^i)e^{-2i\theta}, \tag{2.2} \\ &(A_{11}^i - A_{22}^i - 2\iota A_{12}^i)e^{2i\theta}, V_3^p, W_{12}^m, (B_1^r + \iota B_2^r)e^{-i\theta}, (B_1^r - \iota B_2^r)e^{i\theta}) \end{aligned}$$

shall hold for $0 \leq \theta \leq 2\pi$. In (2.2), $\iota^2 = -1$ and $i = 1, \dots, N; p = 1, \dots, M; m = 1, \dots, P; r = 1, \dots, N + M + P$. It is immediately seen that $G(A_{33}^i, \dots)$ is expressible as a polynomial in the quantities (2.3) listed below which then form an integrity basis for functions $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M, \mathbf{W}_1, \dots, \mathbf{W}_P)$ which are invariant under T_1 .

$$\begin{aligned} &A_{33}^i, A_{11}^i + A_{22}^i, V_3^p, W_{12}^m, \\ &(A_{11}^i - A_{22}^i)(A_{11}^j - A_{22}^j) + 4A_{12}^i A_{12}^j \quad (i \leq j), \\ &(A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i \quad (i < j), \\ &B_1^r B_1^s + B_2^r B_2^s \quad (r \leq s), \quad B_1^r B_2^s - B_2^r B_1^s \quad (r < s), \\ &(A_{11}^i - A_{22}^i)(B_1^r B_1^s - B_2^r B_2^s) + 2A_{12}^i (B_1^r B_2^s + B_2^r B_1^s) \quad (r \leq s), \\ &(A_{11}^i - A_{22}^i)(B_1^r B_2^s + B_2^r B_1^s) - 2A_{12}^i (B_1^r B_1^s - B_2^r B_2^s) \quad (r \leq s). \end{aligned} \tag{2.3}$$

In (2.3), $i, j = 1, \dots, N; p = 1, \dots, M; m = 1, \dots, P; r, s = 1, \dots, N + M + P$ subject to the restrictions indicated. We observe from (2.3) that the integrity basis for functions $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M)$ invariant under T_1 given by Adkins [1, 2] contains redundant terms.

3. An integrity basis for functions invariant under T_2 . The group T_2 is generated by the matrices $\mathbf{Q}(\theta)$ and \mathbf{R}_1 . We have seen in Sec. 2 that any polynomial function $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M, \mathbf{W}_1, \dots, \mathbf{W}_P)$ which is invariant under the group T_1 generated by $\mathbf{Q}(\theta)$ is expressible as a polynomial in the quantities (2.3). In order to determine the general form of the function $F(\mathbf{A}_1, \dots, \mathbf{W}_P)$ which is invariant under T_2 , we need only determine the general form of a polynomial function of the quantities (2.3) which is invariant under \mathbf{R}_1 . The elements of (2.3) either remain invariant under \mathbf{R}_1 or change sign under \mathbf{R}_1 . Let

I_1, \dots, I_a and J_1, \dots, J_b denote the elements of (2.3) which remain invariant under \mathbf{R}_1 and which change sign under \mathbf{R}_1 respectively. With (2.3), we see that the J_1, \dots, J_b are given by

$$W_{12}^m, (A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i \quad (i < j), \quad B_1^r B_2^s - B_2^r B_1^s \quad (r < s),$$

$$(A_{11}^i - A_{22}^i)(B_1^r B_2^s + B_2^r B_1^s) - 2A_{12}^i(B_1^r B_1^s - B_2^r B_2^s) \quad (r \leq s) \quad (3.1)$$

where $i, j = 1, \dots, N$; $m = 1, \dots, P$; $r, s = 1, \dots, N + M + P$ subject to the restrictions indicated. The I_1, \dots, I_a are the elements of (2.3) not listed in (3.1). An integrity basis for functions $F(\mathbf{A}_1, \dots, \mathbf{W}_p)$ which are invariant under T_2 is then given by I_1, \dots, I_a and $J_p J_q$ ($p, q = 1, \dots, b$; $p \leq q$).

After eliminating the redundant elements from the set $J_p J_q$, we obtain the result that an integrity basis for functions $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M, \mathbf{W}_1, \dots, \mathbf{W}_p)$ which are invariant under T_2 is given by

$$A_{33}^i, A_{11}^i + A_{22}^i, V_3^i, (A_{11}^i - A_{22}^i)(A_{11}^j - A_{22}^j) + 4A_{12}^i A_{12}^j \quad (i \leq j),$$

$$B_1^r B_1^s + B_2^r B_2^s \quad (r \leq s),$$

$$(A_{11}^i - A_{22}^i)(B_1^r B_1^s - B_2^r B_2^s) + 2A_{12}^i(B_1^r B_2^s + B_2^r B_1^s) \quad (r \leq s),$$

$$W_{12}^m W_{12}^n \quad (m \leq n), \quad W_{12}^m(A_{11}^i - A_{22}^i)A_{12}^j - W_{12}^n(A_{11}^j - A_{22}^j)A_{12}^i \quad (i < j), \quad (3.2)$$

$$W_{12}^m(B_1^r B_2^s - B_2^r B_1^s) \quad (r < s),$$

$$W_{12}^m(A_{11}^i - A_{22}^i)(B_1^r B_2^s + B_2^r B_1^s) - 2W_{12}^n A_{12}^i(B_1^r B_1^s - B_2^r B_2^s) \quad (r \leq s),$$

$$((A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i)(B_1^r B_2^s - B_2^r B_1^s) \quad (i < j, r < s)$$

where $i, j = 1, \dots, N$; $p = 1, \dots, M$; $m, n = 1, \dots, P$; $r, s = 1, \dots, N + M + P$ subject to the restrictions indicated. The quantities B_α^r ($\alpha = 1, 2$; $r = 1, \dots, N + M + P$) are defined by (2.1). If we set the $W_{12}^m, W_{31}^m, W_{23}^m$ ($m = 1, \dots, P$) appearing in (3.2) equal to zero, we obtain an integrity basis for functions $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M)$ which are invariant under T_2 . The integrity basis so obtained contains fewer terms than that given by Adkins [2], which would indicate the presence of redundant terms in the basis listed in [2].

4. An integrity basis for functions invariant under T_3 . The group T_3 is generated by the matrices $\mathbf{Q}(\theta)$ and \mathbf{R}_3 . We have seen in Sec. 2 that any polynomial function $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M, \mathbf{W}_1, \dots, \mathbf{W}_p)$ which is invariant under the group T_1 generated by $\mathbf{Q}(\theta)$ is expressible as a polynomial in the quantities (2.3). In order to determine the general form of the function $F(\mathbf{A}_1, \dots, \mathbf{W}_p)$ which is invariant under T_3 , we need only determine the general form of a polynomial function of the quantities (2.3) which is invariant under \mathbf{R}_3 . The elements of (2.3) either remain invariant under \mathbf{R}_3 or change sign under \mathbf{R}_3 . Let K_1, \dots, K_c and L_1, \dots, L_d denote the elements of (2.3) which remain invariant under \mathbf{R}_3 and which change sign under \mathbf{R}_3 respectively. Let

$$C_\alpha^i = A_{3\alpha}^i \quad (\alpha = 1, 2; i = 1, \dots, N), \quad C_\alpha^{N+j} = W_{3\alpha}^j \quad (\alpha = 1, 2; j = 1, \dots, P). \quad (4.1)$$

With (2.3) and (4.1), we see that the elements K_1, \dots, K_c of (2.3) which remain invariant under \mathbf{R}_3 are given by

$$\begin{aligned}
 & A_{33}^i, A_{11}^i + A_{22}^i, W_{12}^m, (A_{11}^i - A_{22}^i)(A_{11}^j - A_{22}^j) + 4A_{12}^i A_{12}^j \quad (i \leq j), \\
 & (A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i \quad (i < j), \quad C_1^r C_1^s + C_2^r C_2^s \quad (r \leq s), \\
 & C_1^r C_2^s - C_2^r C_1^s \quad (r < s), \quad V_1^p V_1^q + V_2^p V_2^q \quad (p \leq q), \\
 & V_1^p V_2^q - V_2^p V_1^q \quad (p < q), \\
 & (A_{11}^i - A_{22}^i)(C_1^r C_1^s - C_2^r C_2^s) + 2A_{12}^i(C_1^r C_2^s + C_2^r C_1^s) \quad (r \leq s), \\
 & (A_{11}^i - A_{22}^i)(V_1^p V_1^q - V_2^p V_2^q) + 2A_{12}^i(V_1^p V_2^q + V_2^p V_1^q) \quad (p \leq q), \\
 & (A_{11}^i - A_{22}^i)(C_1^r C_2^s + C_2^r C_1^s) - 2A_{12}^i(C_1^r C_1^s - C_2^r C_2^s) \quad (r \leq s), \\
 & (A_{11}^i - A_{22}^i)(V_1^p V_2^q + V_2^p V_1^q) - 2A_{12}^i(V_1^p V_1^q - V_2^p V_2^q) \quad (p \leq q)
 \end{aligned} \tag{4.2}$$

where $i, j = 1, \dots, N$; $p, q = 1, \dots, M$; $m = 1, \dots, P$; $r, s = 1, \dots, N + P$ subject to the restrictions indicated. With (2.3) and (4.1), we see that the elements L_1, \dots, L_d of (2.3) which change sign under \mathbf{R}_3 are given by

$$\begin{aligned}
 & V_3^p, C_1^r V_1^p + C_2^r V_2^p, C_1^r V_2^p - C_2^r V_1^p, \\
 & (A_{11}^i - A_{22}^i)(C_1^r V_1^p - C_2^r V_2^p) + 2A_{12}^i(C_1^r V_2^p + C_2^r V_1^p), \\
 & (A_{11}^i - A_{22}^i)(C_1^r V_2^p + C_2^r V_1^p) - 2A_{12}^i(C_1^r V_1^p - C_2^r V_2^p)
 \end{aligned} \tag{4.3}$$

where $i = 1, \dots, N$; $p = 1, \dots, M$; $r = 1, \dots, N + P$. An integrity basis for functions $F(\mathbf{A}_1, \dots, \mathbf{W}_p)$ which are invariant under T_3 is then given by the quantities K_1, \dots, K_c and $L_p L_q$ ($p, q = 1, \dots, d$; $p \leq q$). After eliminating the redundant terms from the set of invariants $L_p L_q$, we obtain the result that an integrity basis for functions $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M, \mathbf{W}_1, \dots, \mathbf{W}_p)$ which are invariant under T_3 is given by

$$\begin{aligned}
 & A_{33}^i, A_{11}^i + A_{22}^i, W_{12}^m, (A_{11}^i - A_{22}^i)(A_{11}^j - A_{22}^j) + 4A_{12}^i A_{12}^j \quad (i \leq j), \\
 & (A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i \quad (i < j), \\
 & C_1^r C_1^s + C_2^r C_2^s \quad (r \leq s), \\
 & C_1^r C_2^s - C_2^r C_1^s \quad (r < s), \quad V_1^p V_1^q + V_2^p V_2^q \quad (p \leq q), \\
 & V_1^p V_2^q - V_2^p V_1^q \quad (p < q), \\
 & V_3^p V_3^q \quad (p \leq q), \quad (A_{11}^i - A_{22}^i)(C_1^r C_1^s - C_2^r C_2^s) \\
 & \quad + 2A_{12}^i(C_1^r C_2^s + C_2^r C_1^s) \quad (r \leq s), \\
 & (A_{11}^i - A_{22}^i)(C_1^r C_2^s + C_2^r C_1^s) - 2A_{12}^i(C_1^r C_1^s - C_2^r C_2^s) \quad (r \leq s), \\
 & (A_{11}^i - A_{22}^i)(V_1^p V_1^q - V_2^p V_2^q) + 2A_{12}^i(V_1^p V_2^q + V_2^p V_1^q) \quad (p \leq q), \\
 & (A_{11}^i - A_{22}^i)(V_1^p V_2^q + V_2^p V_1^q) - 2A_{12}^i(V_1^p V_1^q - V_2^p V_2^q) \quad (p \leq q), \\
 & V_3^p(C_1^r V_1^q + C_2^r V_2^q), \quad V_3^p(C_1^r V_2^q - C_2^r V_1^q), \\
 & V_3^p(A_{11}^i - A_{22}^i)(C_1^r V_1^q - C_2^r V_2^q) + 2V_3^p A_{12}^i(C_1^r V_2^q + C_2^r V_1^q),
 \end{aligned} \tag{4.4}$$

$$\begin{aligned}
 &V_3^p(A_{11}^i - A_{22}^i)(C_1^r V_2^q + C_2^r V_1^q) - 2V_3^p A_{12}^i(C_1^r V_1^q - C_2^r V_2^q), \\
 &(C_1^r C_1^s - C_2^r C_2^s)(V_1^p V_1^q - V_2^p V_2^q) \\
 &\quad + (C_1^r C_2^s + C_2^r C_1^s)(V_1^p V_2^q + V_2^p V_1^q) \quad (r \leq s, p \leq q), \\
 &(C_1^r C_1^s - C_2^r C_2^s)(V_1^p V_2^q + V_2^p V_1^q) \\
 &\quad - (C_1^r C_2^s + C_2^r C_1^s)(V_1^p V_1^q - V_2^p V_2^q) \quad (r \leq s, p \leq q)
 \end{aligned}$$

where $i, j = 1, \dots, N$; $p, q = 1, \dots, M$; $m = 1, \dots, P$; $r, s = 1, \dots, N + P$ subject to the restrictions indicated. The quantities C_α^r ($\alpha = 1, 2$; $r = 1, \dots, N + P$) are defined by (4.1).

5. An integrity basis for functions invariant under T_4 . The group T_4 is generated by the matrices $Q(\theta)$, R_1 and R_3 . We have seen in Sec. 3 that any polynomial function $F(A_1, \dots, A_N, V_1, \dots, V_M, W_1, \dots, W_P)$ which is invariant under the group T_2 generated by $Q(\theta)$ and R_1 is expressible as a polynomial in the quantities (3.2). In order to determine the general form of a function $F(A_1, \dots, W_P)$ which is invariant under T_4 , we need only determine the general form of a polynomial function of the quantities (3.2) which is invariant under R_3 . We observe that the elements of (3.2) either remain invariant under R_3 or change sign under R_3 . Let

$$C_\alpha^i = A_{3\alpha}^i \quad (\alpha = 1, 2; i = 1, \dots, N), \quad C_\alpha^{N+j} = W_{3\alpha}^j \quad (\alpha = 1, 2; j = 1, \dots, P). \quad (5.1)$$

With (3.2) and (5.1), we see that the elements M_1, \dots, M_e of (3.2) which remain invariant under R_3 are given by

$$\begin{aligned}
 &A_{33}^i, A_{11}^i + A_{22}^i, (A_{11}^i - A_{22}^i)(A_{11}^i - A_{22}^i) + 4A_{12}^i A_{12}^i \quad (i \leq j), \\
 &C_1^r C_1^s + C_2^r C_2^s \quad (r \leq s), \quad V_1^p V_1^q + V_2^p V_2^q \quad (p \leq q), \\
 &W_{12}^m W_{12}^n \quad (m \leq n), \\
 &(A_{11}^i - A_{22}^i)(C_1^r C_1^s - C_2^r C_2^s) + 2A_{12}^i(C_1^r C_2^s + C_2^r C_1^s) \quad (r \leq s), \\
 &((A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i)(C_1^r C_2^s - C_2^r C_1^s) \quad (i < j, r < s), \\
 &(A_{11}^i - A_{22}^i)(V_1^p V_1^q - V_2^p V_2^q) + 2A_{12}^i(V_1^p V_2^q + V_2^p V_1^q) \quad (p \leq q), \\
 &((A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i)(V_1^p V_2^q - V_2^p V_1^q) \quad (i < j, p < q), \\
 &W_{12}^m(A_{11}^i - A_{22}^i)A_{12}^j - W_{12}^n(A_{11}^j - A_{22}^j)A_{12}^i \quad (i < j), \\
 &W_{12}^m(C_1^r C_2^s - C_2^r C_1^s) \quad (r < s), \\
 &W_{12}^m(V_1^p V_2^q - V_2^p V_1^q) \quad (p < q), \\
 &W_{12}^m(A_{11}^i - A_{22}^i)(C_1^r C_2^s + C_2^r C_1^s) - 2W_{12}^n A_{12}^i(C_1^r C_1^s - C_2^r C_2^s) \quad (r \leq s), \\
 &W_{12}^m(A_{11}^i - A_{22}^i)(V_1^p V_2^q + V_2^p V_1^q) - 2W_{12}^n A_{12}^i(V_1^p V_1^q - V_2^p V_2^q) \quad (p \leq q)
 \end{aligned} \tag{5.2}$$

where $i, j = 1, \dots, N$; $p, q = 1, \dots, M$; $m, n = 1, \dots, P$; $r, s = 1, \dots, N + P$ subject to the restrictions indicated. With (3.2) and (5.1), we see that the elements N_1, \dots, N_f of (3.2)

which change sign under \mathbf{R}_3 are given by

$$\begin{aligned}
 & V_3^p, C_1^r V_1^p + C_2^r V_2^p, W_{12}^m(C_1^r V_2^p - C_2^r V_1^p), \\
 & (A_{11}^i - A_{22}^i)(C_1^r V_1^p - C_2^r V_2^p) + 2A_{12}^i(C_1^r V_2^p + C_2^r V_1^p), \\
 & W_{12}^m(A_{11}^i - A_{22}^i)(C_1^r V_2^p + C_2^r V_1^p) - 2W_{12}^m A_{12}^i(C_1^r V_1^p - C_2^r V_2^p), \\
 & ((A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i)(C_1^r V_2^p - C_2^r V_1^p) \quad (i < j)
 \end{aligned} \tag{5.3}$$

where $i, j = 1, \dots, N$; $p = 1, \dots, M$; $m = 1, \dots, P$; $r = 1, \dots, N + P$ subject to the restrictions indicated. An integrity basis for functions $F(\mathbf{A}_1, \dots, \mathbf{W}_p)$ which are invariant under T_4 is then given by M_1, \dots, M_e and $N_p N_q$ ($p, q = 1, \dots, f$; $p \leq q$). After eliminating the redundant elements from the set of invariants $N_p N_q$, we obtain the result that an integrity basis for functions $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M, \mathbf{W}_1, \dots, \mathbf{W}_P)$ which are invariant under T_4 is given by

$$\begin{aligned}
 & A_{33}^i, A_{11}^i + A_{22}^i, (A_{11}^i - A_{22}^i)(A_{11}^j - A_{22}^j) + 4A_{12}^i A_{12}^j \quad (i \leq j), \\
 & C_1^r C_1^s + C_2^r C_2^s \quad (r \leq s), \quad V_1^p V_1^q + V_2^p V_2^q \quad (p \leq q), \\
 & V_3^p V_3^q \quad (p \leq q), \quad W_{12}^m W_{12}^n \quad (m \leq n), \\
 & (A_{11}^i - A_{22}^i)(C_1^r C_1^s - C_2^r C_2^s) + 2A_{12}^i(C_1^r C_2^s + C_2^r C_1^s) \quad (r \leq s), \\
 & ((A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i)(C_1^r C_2^s - C_2^r C_1^s) \quad (i < j, r < s), \\
 & (A_{11}^i - A_{22}^i)(V_1^p V_1^q - V_2^p V_2^q) + 2A_{12}^i(V_1^p V_2^q + V_2^p V_1^q) \quad (p \leq q), \\
 & ((A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i)(V_1^p V_2^q - V_2^p V_1^q) \quad (i < j, p < q), \\
 & W_{12}^m(A_{11}^i - A_{22}^i)A_{12}^j - W_{12}^n(A_{11}^j - A_{22}^j)A_{12}^i \quad (i < j), \\
 & (C_1^r C_1^s - C_2^r C_2^s)(V_1^p V_1^q - V_2^p V_2^q) \\
 & \quad + (C_1^r C_2^s + C_2^r C_1^s)(V_1^p V_2^q + V_2^p V_1^q) \quad (r \leq s, p \leq q), \\
 & (C_1^r C_2^s - C_2^r C_1^s)(V_1^p V_2^q - V_2^p V_1^q) \quad (r < s, p < q), \\
 & V_3^p(C_1^r V_1^q + C_2^r V_2^q), W_{12}^m(C_1^r C_2^s - C_2^r C_1^s) \quad (r < s), \\
 & W_{12}^m(V_1^p V_2^q - V_2^p V_1^q) \quad (p < q), \\
 & ((A_{11}^i - A_{22}^i)(C_1^r C_2^s + C_2^r C_1^s) \\
 & \quad - 2A_{12}^i(C_1^r C_1^s - C_2^r C_2^s))(V_1^p V_2^q - V_2^p V_1^q) \quad (r \leq s, p < q), \\
 & ((A_{11}^i - A_{22}^i)(V_1^p V_2^q + V_2^p V_1^q) \\
 & \quad - 2A_{12}^i(V_1^p V_1^q - V_2^p V_2^q))(C_1^r C_2^s - C_2^r C_1^s) \quad (p \leq q, r < s), \\
 & W_{12}^m(A_{11}^i - A_{22}^i)(C_1^r C_2^s + C_2^r C_1^s) - 2W_{12}^m A_{12}^i(C_1^r C_1^s - C_2^r C_2^s) \quad (r \leq s), \\
 & W_{12}^m(A_{11}^i - A_{22}^i)(V_1^p V_2^q + V_2^p V_1^q) - 2W_{12}^m A_{12}^i(V_1^p V_1^q - V_2^p V_2^q) \quad (p \leq q),
 \end{aligned} \tag{5.4}$$

$$\begin{aligned}
 &W_{12}^m(C_1^r C_1^s - C_2^r C_2^s)(V_1^p V_2^q + V_2^p V_1^q) \\
 &\quad - W_{12}^m(C_1^r C_2^s + C_2^r C_1^s)(V_1^p V_1^q - V_2^p V_2^q) \quad (r \leq s, p \leq q), \\
 &V_3^p(A_{11}^i - A_{22}^i)(C_1^r V_1^q - C_2^r V_2^q) + 2V_3^p A_{12}^i(C_1^r V_2^q + C_2^r V_1^q), \\
 &V_3^p((A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i)(C_1^r V_2^q - C_2^r V_1^q) \quad (i < j), \\
 &V_3^p W_{12}^m(C_1^r V_2^q - C_2^r V_1^q), \\
 &V_3^p W_{12}^m(A_{11}^i - A_{22}^i)(C_1^r V_2^q + C_2^r V_1^q) - 2V_3^p W_{12}^m A_{12}^i(C_1^r V_1^q - C_2^r V_2^q)
 \end{aligned}$$

where $i, j = 1, \dots, N$; $p, q = 1, \dots, M$; $m, n = 1, \dots, P$; $r, s = 1, \dots, N + P$ subject to the restrictions indicated. The quantities C_α^r ($\alpha = 1, 2$; $r = 1, \dots, N + P$) are defined by (5.1).

6. An integrity basis for functions invariant under T_5 . The group T_5 is generated by the matrices $Q(\theta)$ and D_2 . We have seen in Sec. 2 that any polynomial function $F(A_1, \dots, A_N, V_1, \dots, V_M, W_1, \dots, W_P)$ which is invariant under the group T_1 generated by $Q(\theta)$ is expressible as a polynomial in the quantities (2.3). In order to determine the general form of the function $F(A_1, \dots, W_P)$ which is invariant under T_5 , we need only determine the general form of a polynomial function of the quantities (2.3) which is invariant under D_2 . The elements of (2.3) either remain invariant under D_2 or change sign under D_2 . Let

$$C_\alpha^i = A_{3\alpha}^i \quad (\alpha = 1, 2; i = 1, \dots, N), \quad C_\alpha^{N+j} = W_{3\alpha}^j \quad (\alpha = 1, 2; j = 1, \dots, P). \quad (6.1)$$

With (2.3) and (6.1), we see that the elements P_1, \dots, P_g of (2.3) which remain invariant under D_2 are given by

$$\begin{aligned}
 &A_{33}^i, A_{11}^i + A_{22}^i, (A_{11}^i - A_{22}^i)(A_{11}^j - A_{22}^j) + 4A_{12}^i A_{12}^j \quad (i \leq j), \\
 &C_1^s C_1^t + C_2^s C_2^t \quad (s \leq t), \quad V_1^p V_1^q + V_2^p V_2^q \quad (p \leq q), \quad C_1^s V_2^p - C_2^s V_1^p, \\
 &(A_{11}^i - A_{22}^i)(C_1^s C_1^t - C_2^s C_2^t) + 2A_{12}^i(C_1^s C_2^t + C_2^s C_1^t) \quad (s \leq t), \\
 &(A_{11}^i - A_{22}^i)(V_1^p V_1^q - V_2^p V_2^q) + 2A_{12}^i(V_1^p V_2^q + V_2^p V_1^q) \quad (p \leq q), \\
 &(A_{11}^i - A_{22}^i)(C_1^s V_2^p + C_2^s V_1^p) - 2A_{12}^i(C_1^s V_1^p - C_2^s V_2^p)
 \end{aligned} \quad (6.2)$$

where $i, j = 1, \dots, N$; $p, q = 1, \dots, M$; $s, t = 1, \dots, N + P$ subject to the restrictions indicated. With (2.3) and (6.1), we see that the elements Q_1, \dots, Q_h of (2.3) which change sign under D_2 are given by

$$\begin{aligned}
 &V_3^p, W_{12}^m, (A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i \quad (i < j), \\
 &C_1^s V_1^p + C_2^s V_2^p, C_1^s C_2^t - C_2^s C_1^t \quad (s < t), \quad V_1^p V_2^q - V_2^p V_1^q \quad (p < q), \\
 &(A_{11}^i - A_{22}^i)(C_1^s V_1^p - C_2^s V_2^p) + 2A_{12}^i(C_1^s V_2^p + C_2^s V_1^p), \\
 &(A_{11}^i - A_{22}^i)(C_1^s C_2^t + C_2^s C_1^t) - 2A_{12}^i(C_1^s C_1^t - C_2^s C_2^t) \quad (s \leq t), \\
 &(A_{11}^i - A_{22}^i)(V_1^p V_2^q + V_2^p V_1^q) - 2A_{12}^i(V_1^p V_1^q - V_2^p V_2^q) \quad (p \leq q)
 \end{aligned} \quad (6.3)$$

where $i, j = 1, \dots, N$; $p, q = 1, \dots, M$; $m = 1, \dots, P$; $s, t = 1, \dots, N + P$ subject to the restrictions indicated. An integrity basis for functions $F(A_1, \dots, W_p)$ which are invariant under T_5 is then given by P_1, \dots, P_g and Q_r, Q_s ($r, s = 1, \dots, h$; $r \leq s$). After eliminating the redundant elements from the set of invariants Q_r, Q_s , we obtain the result that an integrity

basis for functions $F(\mathbf{A}_1, \dots, \mathbf{A}_N, \mathbf{V}_1, \dots, \mathbf{V}_M, \mathbf{W}_1, \dots, \mathbf{W}_P)$ which are invariant under T_5 is given by

$$\begin{aligned}
 & A_{33}^i, A_{11}^i + A_{22}^i, (A_{11}^i - A_{22}^i)(A_{11}^j - A_{22}^j) + 4A_{12}^i A_{12}^j \quad (i \leq j), \\
 & C_1^s C_1^t + C_2^s C_2^t \quad (s \leq t), \quad C_1^s V_1^p - C_2^s V_1^p, V_1^p V_1^q + V_2^p V_2^q \quad (p \leq q), \\
 & (A_{11}^i - A_{22}^i)(C_1^s C_1^t - C_2^s C_2^t) + 2A_{12}^i(C_1^s C_2^t + C_2^s C_1^t) \quad (s \leq t), \\
 & (A_{11}^i - A_{22}^i)(C_1^s V_2^p + C_2^s V_1^p) - 2A_{12}^i(C_1^s V_1^p - C_2^s V_2^p), \\
 & (A_{11}^i - A_{22}^i)(V_1^p V_1^q - V_2^p V_2^q) + 2A_{12}^i(V_1^p V_2^q + V_2^p V_1^q) \quad (p \leq q), \\
 & ((A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i)(C_1^s C_2^t - C_2^s C_1^t) \quad (i < j, s < t), \\
 & ((A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i)(C_1^s V_1^p + C_2^s V_2^p) \quad (i < j), \\
 & ((A_{11}^i - A_{22}^i)A_{12}^j - (A_{11}^j - A_{22}^j)A_{12}^i)(V_1^p V_2^q - V_2^p V_1^q) \quad (i < j, p < q), \\
 & W_{12}^m W_{12}^n \quad (m \leq n), \quad W_{12}^m(A_{11}^i - A_{22}^i)A_{12}^j - W_{12}^n(A_{11}^i - A_{22}^i)A_{12}^j \quad (i < j), \\
 & W_{12}^m(C_1^s C_2^t - C_2^s C_1^t) \quad (s < t), \quad W_{12}^m(C_1^s V_1^p + C_2^s V_2^p), \\
 & W_{12}^m(V_1^p V_2^q - V_2^p V_1^q) \quad (p < q), \tag{6.4} \\
 & W_{12}^m(A_{11}^i - A_{22}^i)(C_1^s C_2^t + C_2^s C_1^t) - 2W_{12}^n A_{12}^i(C_1^s C_1^t - C_2^s C_2^t) \quad (s \leq t), \\
 & W_{12}^m(A_{11}^i - A_{22}^i)(C_1^s V_1^p - C_2^s V_2^p) + 2W_{12}^n A_{12}^i(C_1^s V_2^p + C_2^s V_1^p), \\
 & W_{12}^m(A_{11}^i - A_{22}^i)(V_1^p V_2^q + V_2^p V_1^q) - 2W_{12}^n A_{12}^i(V_1^p V_1^q - V_2^p V_2^q) \quad (p \leq q), \\
 & W_{12}^m V_3^p, V_3^p V_3^q \quad (p \leq q), \\
 & V_3^p(A_{11}^i - A_{22}^i)A_{12}^j - V_3^q(A_{11}^i - A_{22}^i)A_{12}^j \quad (i < j), \\
 & V_3^p(C_1^s C_2^t - C_2^s C_1^t) \quad (s < t), \quad V_3^p(C_1^s V_1^q + C_2^s V_2^q), \\
 & V_3^p(V_1^q V_2^r - V_2^q V_1^r) \quad (q < r), \\
 & V_3^p(A_{11}^i - A_{22}^i)(C_1^s V_1^q - C_2^s V_2^q) + 2V_3^q A_{12}^i(C_1^s V_2^q + C_2^s V_1^q), \\
 & V_3^p(A_{11}^i - A_{22}^i)(C_1^s C_2^t + C_2^s C_1^t) - 2V_3^q A_{12}^i(C_1^s C_1^t - C_2^s C_2^t) \quad (s \leq t), \\
 & V_3^p(A_{11}^i - A_{22}^i)(V_1^q V_2^r + V_2^q V_1^r) - 2V_3^q A_{12}^i(V_1^q V_1^r - V_2^q V_2^r) \quad (q \leq r)
 \end{aligned}$$

where $i, j = 1, \dots, N$; $p, q, r = 1, \dots, M$; $m, n = 1, \dots, P$; $s, t = 1, \dots, N + P$ subject to the restrictions indicated. The quantities C_α^s ($\alpha = 1, 2$; $s = 1, \dots, N + P$) are defined by (6.1).

REFERENCES

- [1] J. E. Adkins, Arch. Rat. Mech. Anal. **4**, 193 (1960)
- [2] J. E. Adkins, Arch. Rat. Mech. Anal. **5**, 263 (1960)
- [3] R. L. Long and L. V. McIntire, private communication (1973)