

ON UNEQUALLY SPACED TIME POINTS IN TIME SERIES

BY WILLIAM CLINGER¹ AND JOHN W. VAN NESS²

The University of Texas at Dallas

This article discusses the sampling of stationary discrete-time stochastic processes at fixed but unequally spaced time points. The pattern of the sampling times is periodic with a cycle of p time units. One of the major problems is to determine given p the minimum number of sampling points required per cycle in order to estimate the covariances at all lags. The second problem is to find a pattern of distribution for the sampling points within the cycle which will allow the estimation of all covariances. A discussion of the references which describe the statistical properties of the estimates of covariances and spectra in this sampling situation is given.

1. Introduction. One of the authors (Van Ness) was approached by oceanographers who wanted to study the spectra of certain time-varying features of the ocean (salinity, temperature, etc.). They wished to measure these features at several different depths and for a length of time of the order of magnitude of the cruise time. However, they had only one sensing device on board and could not keep the ship out long enough to hold the device at each depth for a suitable length of time. They proposed, therefore, to lower and raise the sensor in a cyclic fashion keeping its velocity magnitude constant, thus giving a depth versus time plot as shown in Figure 1.

At a fixed depth, one has a time series $\{X(t_1), X(t_2), \dots\}$ observed at unequally spaced time points. If $t_3 - t_2$ and $t_2 - t_1$ are both integral multiples of some number, the problem involves observations at unequally but cyclically spaced time points of a discrete time series.

We generalize this problem as follows. Given an integer $p > 2$, to be called the sampling period, let the sampling pattern, $S = \{s_1, \dots, s_k\}$, be a subset of the integers from 0 to $p - 1$. Sample $\{X(t)\}$ only at those values of t for which $0 \leq t \leq T$ and $t \pmod{p} \in S$. For example, if $\{Z(t)\}$ is the underlying continuous time process in the example of Figure 1 and $t_2 - t_1$ is an integral multiple of $t_3 - t_2$, one could define $X(t) = Z(t(t_3 - t_2) + t_1)$, $p = (t_3 - t_1)/(t_3 - t_2)$ and $S = \{0, (t_2 - t_1)/(t_3 - t_2)\}$. General sampling patterns are treated in this paper.

Jones (1962 and 1971) and Parzen (1963) discuss special cases of this problem. The latter gives other examples where such problems arise. Loynes (1970) reviews the state of the literature on random processes observed at unequally spaced time points. As Loynes (1969) points out, very little additional work

Received March 1975; revised January 1976.

¹ Supported by funds from the Clark Foundation.

² Supported by NSF Grant GP 42659.

AMS 1970 subject classifications. 62M15, 62M10.

Key words and phrases. Time series, spectral analysis, missing observations, autocorrelation, autocovariance, covariance sequence, unequally spaced samples.




FIG. 1. Oceanographer's suggested sampling plan.

has been done on the case of missing observations at fixed (nonrandom) time points.

The major concern in this paper is to find the minimum number of observations per cycle which will allow estimation of the complete covariance sequence and spectrum in the discrete time case.

2. Cyclic sampling. Let $\{X_t, t = 0, \pm 1, \pm 2, \dots\}$ be a stationary process with mean zero and finite second-order moments. Let p and $S = \{s_1, \dots, s_k\}$ be as defined above, only for convenience assume S is ordered so that $0 \leq s_1 < s_2 < \dots < s_k \leq p - 1$. Denote the covariance of lag v by $R(v) = EX(t)X(t + v)$; $v, t = 0, \pm 1, \pm 2, \dots$.

Assume $\{X(t)\}$ has been sampled according to S over a time period $[0, T]$ giving observations $\{X_t : t \pmod p \in S, 0 \leq t \leq T\}$. It is convenient to use the following picture: divide a circle into p equal segments and number the division points from 0 to $p - 1$. Put a dot at those points whose numbers are in S , e.g. if $p = 8$ and $S = \{0, 1, 4\}$, we have Figure 2.

Define Q to be a "translation operator" on S such that

$$Q(S) \equiv Q(s_1, s_2, \dots, s_k) \\ \equiv (s_1 + 1 \pmod p, s_2 + 1 \pmod p, \dots, s_k + 1 \pmod p).$$




FIG. 2. Graphical representation of S .

Q shifts the dots in our picture one unit clockwise. Define $Q^2(S) \equiv Q(Q(S))$. It will also be convenient to use the notation of Parzen (1963). Let

$$(2.1) \quad \begin{aligned} g(t) &= 0 && \text{if } X(t) \text{ is not observed} \\ &= 1 && \text{if } X(t) \text{ is observed} \end{aligned}$$

i.e. $g(t) = 1$ if and only if $t \pmod p \in S$. Define

$$(2.2) \quad Y(t) = g(t)X(t),$$

then $\{Y(t)\}$ is a particular amplitude modulated version of $\{X(t)\}$. Note that if $g(t) = 1$ then $g(t + jp) = 1$ for all integers j .

3. Estimation of covariances. The natural estimates $R^*(v)$ of the covariances $R(\pm v)$, $v \geq 0$ are

$$(3.1) \quad \begin{aligned} R^*(v) = R^*(-v) &\equiv \frac{1}{M_T(v)} \sum_{t=0}^{T-v} Y(t)Y(t+v) \\ &= \frac{1}{M_T(v)} \sum_{t=0}^{T-v} g(t)X(t)g(t+v)X(t+v) \end{aligned}$$

when $M_T(v) = \sum_{t=0}^{T-v} g(t)g(t+v) > 0$. If $M_T(v) = 0$ for all T we cannot use (3.1), no matter how large the sample.

First we determine those v for which $M_T(v) > 0$ for some T . Due to symmetry we need henceforth only consider $v \geq 0$. First note that $M_T(v) > 0$ for some T if and only if $Q^v(S) \cap S \neq \emptyset$.

A simple way of determining which $R^*(v)$ exist can be given using the pictures corresponding to Figure 2. Let $\text{card } A$ denote the number of points in the set A . Then $k = \text{card } S$ is the number of observations per p time units. Take the circle with p segment division points and K dots distributed as described in Section 2. Draw lines connecting each dot to each other dot. With each of these $\binom{k}{2}$ lines associate two numbers giving the number of segments between its two dots counting in the two possible directions. Define V to be the set containing 0 and all the different numbers so obtained. Then one can form $R^*(v)$ if and only if $|v| \pmod p \in V$. Note that $\text{card } V \leq 2\binom{k}{2} + 1 = k^2 - k + 1$.

Secondly we would like to know the size of $M_T(v)$. Let $[\cdot]$ denote the greatest integer function.

PROPOSITION 1.

$$\left(\left[\frac{T-v}{p} \right] \right) \text{card } (Q^v(S) \cap S) \leq M_T(v) \leq \left(\left[\frac{T-v}{p} \right] + 1 \right) \text{card } (Q^v(S) \cap S).$$

COROLLARY 2.

$$\lim_{T \rightarrow \infty} M_T(v)/T = \frac{1}{p} \text{card } (Q^v(S) \cap S) \equiv m(v).$$

Following Parzen (1963) we define

$$(3.2) \quad R_T(v) = R_T(-v) = \frac{1}{Tm(v)} \sum_{t=0}^{T-v} Y(t)Y(t+v),$$

when $m(v) > 0$, which is a simple estimate useful in spectral estimation (see Section 7).

4. Number theory results. In number theory, if $\{0, 1, 2, \dots, p-1\} \subset V$ then S is called a difference cover for p . We wish to find how few observations we can get away with and still estimate all covariances using (3.1) or (3.2). Thus we are interested in the minimum cardinality that a difference cover for p can have. Call this number $f(p)$. As we have shown, $p \leq f^2(p) - f(p) + 1$.

If equality holds in the previous equation then every integer $0, 1, \dots, p-1$ is uniquely expressible as a difference. Such difference coverings are called simple difference coverings. The existence of known simple difference coverings is given by

THEOREM 3 (Singer (1938)). *A sufficient condition for the existence of $k+1$ integers v_1, \dots, v_{k+1} , having the property that their k^2+k differences $v_j - v_i$; $i \neq j$; $i, j = 1, \dots, k+1$, are congruent, modulo $p = k^2 + k + 1$, to $1, 2, \dots, k^2 + k$ in some order is that k be a power of a prime.*

It has been conjectured that no simple difference set exists for $p = m^2 + m + 1$ when m is not a prime power. This conjecture has been verified for the 1321 cases in which $m \leq 1600$, but has not been proved for all m (Mann (1965), page 86). A number of criteria for the existence of simple difference sets have been established however (see Mann (1965) or Ryser (1963) for discussion and bibliography).

A trivial covering gives the following result.

PROPOSITION 4. *If $p \leq 2m(n+1) + 1$ when m, n positive integers, $n \geq 2$, then $f(p) \leq m + n$.*

COROLLARY 5. $1 < f^2(p)/p \leq 2$ for $p \geq 12$.

PROOF. Suppose $f^2(p) \leq p$, then $f^2(p) \leq f^2(p) - f(p) + 1$, giving $f(p) = 1$ which yields $p = 1$. Thus $f^2(p) > p$.

Let x be the least integer such that $x \geq (2p+1)^{\frac{1}{2}} - 1$. Then $2p \leq x^2 + 2x$. If x is even, $p \leq \frac{1}{2}x^2 + x = 2(\frac{1}{2}x)(\frac{1}{2}x + 1)$ so by Proposition 4 $f(p) \leq \frac{1}{2}x + \frac{1}{2}x = x$. If x is odd, so is $x^2 + 2x$, so $2p \leq x^2 + 2x - 1$, from which $p \leq (x^2 - 1)/2 + x < 2((x+1)/2)((x-1)/2 + 1)$, and again by Proposition 4, $f(p) \leq (x+1)/2 + (x-1)/2 = x$ provided $(x-1)/2 \geq 2$, which is the case for odd x corresponding to $p \geq 12$. Therefore, $f(p) < (2p+1)^{\frac{1}{2}}$. Suppose $f^2(p) > 2p$. Then $2p < f^2(p) < 2p + 1$, which is impossible; hence $f^2(p) \leq 2p$, for $p \geq 12$. (By inspection of cases, $f^2(p) \leq 2p$ holds for all p except $p = 4$.) \square

An asymptotic upper bound for $f^2(p)/p$ follows from the work of Redei and Rényi (1949) on difference bases. A difference basis with respect to n is a set of integers a_1, \dots, a_n such that every positive integer $0 < v \leq n$ can be written in the form $v = a_j - a_i$ for some $i \neq j$. Let $k(n)$ denote the minimum of k for given n . Redei and Rényi showed that $\lim_{n \rightarrow \infty} k^2(n)/n$ exists, and gave numerical

bounds which were later improved by Leech (1956) and then Golay (1972). Since a difference basis for n is immediately a difference cover for $p \leq 2n + 1$, it follows that

$$\lim_{p_0 \rightarrow \infty} \sup_{p > p_0} \frac{f^2(p)}{p} \leq \frac{1}{2} \lim_{n \rightarrow \infty} \frac{k^2(n)}{n} < 1.3286 .$$

Actual calculation of $f(p)$ appears very difficult. The congruence conditions stated in the next theorem are often useful, especially for p having small divisors, but no reasonable algorithm has been developed for arbitrary p values. The following corollaries will be used later to get some further practical partial results about f .

THEOREM 6. *Let n be a divisor of p . Let v_1, \dots, v_b be a difference cover for p and let x_i be the number of elements in the cover congruent to i modulo n . Then*

$$\begin{aligned} \sum_{i=0}^{n-1} x_i &= b ; \\ \sum_{i=0}^{n-1} x_i(x_i - 1) &\geq \frac{p}{n} - 1 ; \\ \sum_{i=0}^{n-1} x_i x_{\text{rem}(i+j) \bmod n} &\geq \frac{p}{n}, \quad j = 1, \dots, n - 1, \end{aligned}$$

where $\text{rem}(i) \bmod n$ denotes the reduced residue representative congruent to i modulo n .

PROOF. Consider the integers $1, \dots, p - 1$. $p/n - 1$ of them are congruent to $0 \bmod n$, and p/n are congruent to $j \bmod n$ for $j = 1, \dots, n - 1$. The inequalities merely state that sufficient differences $v_j - v_i$ are congruent to $0, 1, \dots, n - 1$ so that p may be covered by the difference cover. \square

COROLLARY 7. *Let the conditions of Theorem 6 hold. Then $\sum_{i=0}^{n-1} x_i^2 \leq b^2 - ((n - 1)/n)p$.*

PROOF.

$$\begin{aligned} b^2 &= \left(\sum_{i=0}^{n-1} x_i\right)^2 \\ &= \sum_{i=0}^{n-1} x_i^2 + \sum_{j=1}^{n-1} \sum_{i=0}^{n-1} x_i x_{\text{rem}(i+j) \bmod n} \\ &\geq \sum_{i=0}^{n-1} x_i^2 + \sum_{j=1}^{n-1} \frac{p}{n}. \end{aligned} \quad \square$$

Furthermore, equality holds if $b = m + 1$ and $p = b^2 - b + 1 = m^2 + m + 1$.

COROLLARY 8. *Let the conditions of Theorem 6 hold with $b = m + 1$ and $p = m^2 + m + 1$, then*

$$\sum_{i=0}^{n-1} x_i^2 = (m + 1)^2 - \frac{n - 1}{n} (m^2 + m + 1) = \frac{1}{n} [m^2 + (n + 1)m + 1] .$$

COROLLARY 9. *Suppose $f^2(p)/p \rightarrow 1$. Let $n \geq 2$. For each $p = kn$, $k = 1, 2, 3, \dots$, let $V_k = \{v_1, \dots, v_{f(p)}\}$ be a difference cover for p , and let x_{ki} be the number of elements of V_k congruent to i modulo n , $i = 0, \dots, n - 1$. Then*

$$\lim_{k \rightarrow \infty} \frac{1}{k} \sum_{i=0}^{n-1} x_{ki}^2 = 1 .$$

PROOF. For each $k = 1, 2, \dots, f(kn) = \sum_{i=0}^{n-1} x_{ki}$. By a well-known isoperimetric inequality, then,

$$\begin{aligned} \frac{1}{k} \sum_{i=0}^{n-1} x_{ki}^2 &\geq \frac{1}{k} \sum_{i=0}^{n-1} \left[\frac{f(kn)}{n} \right]^2 \\ &> \frac{1}{k} \sum_{i=0}^{n-1} \left[\frac{(kn)^{\frac{1}{2}}}{n} \right]^2. \end{aligned}$$

Let $\varepsilon > 0$. Let p_0 be such that $p > p_0$ implies $f^2(p)/p < 1 + \varepsilon/n$. Let k_0 be such that $k_0 n > p_0$. For $k > k_0$, then,

$$\begin{aligned} 1 &< \frac{1}{k} \sum_{i=0}^{n-1} x_{ki}^2 \\ &\leq \left(\frac{f^2(kn)}{kn} - \frac{n-1}{n} \right) n \\ &< \left(1 + \frac{\varepsilon}{n} - \frac{n-1}{n} \right) n. \end{aligned} \quad \square$$

The next few corollaries apply the congruence conditions modulo 2.

COROLLARY 10. Let m and k be integers with $m \geq 1, k \geq 0$. If m is odd (even) and $f(m^2 + m - 2k) = m + 1$, then there exists an even (odd) integer y such that

$$m - (2k + 1) \leq y^2 \leq m + (2k + 1).$$

PROOF. Given a difference cover of $m + 1$ elements for $p = m^2 + m - 2k$, let x be the number of elements congruent to 0 mod 2; then $(m + 1) - x$ elements are congruent to 1 mod 2. We may assume $x \leq (m + 1)/2$ (for if that is not so, adding 1 to each element of the cover produces a new cover in which it is true). Since $m^2 + m - 2k$ is even, Theorem 6 gives

$$x(x - 1) + (m + 1 - x)(m - x) \geq \frac{m^2 + m - 2k}{2} - 1$$

and

$$2x(m + 1 - x) \geq \frac{m^2 + m - 2k}{2}.$$

Thus

$$x \leq \frac{m + 1 - (m - (2k + 1))^{\frac{1}{2}}}{2}$$

and

$$\frac{m + 1 - (m + 2k + 1)^{\frac{1}{2}}}{2} \leq x \leq \frac{m + 1 + (m + 2k + 1)^{\frac{1}{2}}}{2}$$

giving

$$m - (2k + 1) \leq [(m + 1) - 2x]^2 \leq m + (2k + 1)$$

and set $y = (m + 1) - 2x$. \square

COROLLARY 11. For every positive integer k_0 , there exists an integer m such that

$$f(m^2 + m - 2k) > m + 1 \quad \text{for } k = 0, 1, \dots, k_0 - 1.$$

PROOF. Given k_0 , let $m = (k_0 + 1)^2$. Then

$$m - (2k + 1) = k_0^2 + 2k_0 - 2k > k_0^2$$

and

$$\begin{aligned} m + (2k + 1) &= k_0^2 + 2k_0 + 2k + 2 \\ &\leq k_0^2 + 2k_0 + 2k_0 < (k_0 + 2)^2, \quad k = 0, 1, \dots, k_0 - 1, \end{aligned}$$

so by the preceding corollary $f(m^2 + m - 2k) > m + 1$ for $k = 0, 1, \dots, k_0 - 1$. \square

The congruence conditions for p divisible by three are a little more complicated.

COROLLARY 12. Let $p \equiv 0$ modulo 3. Consider a difference cover for p of b elements and let x_i ($i = 0, 1, 2$) denote the number of elements in the cover congruent to i modulo 3. If $A_j = -3x_j^2 + 2bx_j + b^2 - 4p/3$, and $B_j = -3x_j^2 + 2bx_j - b^2 + 2b - 2 + 2p/3$, then

- 1) $x_0 + x_1 + x_2 = b$
- 2) $(b - 2(b^2 - p)^{\frac{1}{2}})/3 \leq x_i \leq (b + 2(b^2 - p)^{\frac{1}{2}})/3, i = 0, 1, 2$
- 3) $(b - x_j - A_j^{\frac{1}{2}})/2 \leq x_i \leq (b - x_j + A_j^{\frac{1}{2}})/2, i \neq j$
- 4) $x_i \leq (b - x_j - B_j^{\frac{1}{2}})/2$ or $x_i > (b - x_j + B_j^{\frac{1}{2}})/2, i \neq j, \text{ if } B_j > 0.$

PROOF. From Theorem 6 (sums are from 0 to 2):

$$\begin{aligned} \sum x_i &= b, \\ \sum x_i(x_i - 1) &\geq \frac{P}{3} - 1, \\ x_0x_1 + x_1x_2 + x_2x_0 &\geq \frac{P}{3}. \end{aligned}$$

Thus

$$\begin{aligned} x_2 &= b - x_0 - x_1, \\ x_0(x_0 - 1) + x_1(x_1 - 1) + (b - x_0 - x_1)(b - x_0 - x_1 - 1) &\geq \frac{P}{3} - 1, \\ x_0x_1 + x_1(b - x_0 - x_1) + (b - x_0 - x_1)x_0 &\geq \frac{P}{3}, \end{aligned}$$

or

$$(4.1) \quad 2x_0^2 - 2bx_0 + 2x_1^2 - 2bx_1 + 2x_0x_1 + b^2 - b + 1 - \frac{P}{3} \geq 0,$$

$$(4.2) \quad -x_0^2 - x_1^2 - x_0x_1 + bx_0 + bx_1 - \frac{P}{3} \geq 0.$$

$A_1 < 0$ gives no solution to (4.2), so

$$\frac{b - 2(b^2 - p)^{\frac{1}{2}}}{3} \leq x_1 \leq \frac{b + 2(b^2 - p)^{\frac{1}{2}}}{3}.$$

Then, from (4.2)

$$\frac{(b - x_1) - A_1^{\frac{1}{2}}}{2} \leq x_0 \leq \frac{(b - x_1) + A_1^{\frac{1}{2}}}{2}.$$

From (4.1), if $B_1 > 0$, then

$$x_0 \leq \frac{(b - x_1) - B_1^{\frac{1}{2}}}{2} \quad \text{or} \quad x_0 \geq \frac{(b - x_1) + B_1^{\frac{1}{2}}}{2}.$$

The symmetry of the congruence conditions permits interchanging the subscripts 0, 1, 2. \square

COROLLARY 13. *Let $m \equiv 1$ modulo 3. Set $p = m^2 + m + 1$ and let x_0, x_1 and x_2 be defined as in the preceding corollary. If $b = m + 1$*

- 1) $(m + 1 - 2m^{\frac{1}{2}})/3 \leq x_0, x_1, x_2 \leq (m + 1 + 2m^{\frac{1}{2}})/3$; and
- 2) if $D = -3x_0^2 + 2(m + 1)x_0 - \frac{1}{3}(m - 1)^2$, then $x_1 = (m + 1 - x_0 \pm D^{\frac{1}{2}})/2$ and $x_2 = (m + 1 - x_0 \mp D^{\frac{1}{2}})/2$; and
- 3) two of the x_i are congruent to 1 modulo 3, and the other is congruent to 0 modulo 3.

PROOF. Part 1 is just the preceding corollary. In the preceding corollary, $A_0 = B_0 = D$ and this together with the fact that $x_0 + x_1 + x_2 = m + 1$, gives part 2.

$D \equiv x_0 \pmod{3}$. From part 2, $D \equiv (2x_1 + x_0 - m - 1)^2 \pmod{3}$, from which $x_0 \equiv 0$ or $x_0 \equiv 1 \pmod{3}$ since 2 is a quadratic nonresidue mod 3. If $x_0 \equiv 0$, then $2x_i \equiv m + 1$ gives $x_i \equiv 1$. If $x_0 \equiv 1$, then $2x_i \equiv 2$ or $2x_i \equiv 0$, giving $x_i \equiv 0$ or $x_i \equiv 1$; since $x_0 + x_1 + x_2 = m + 1 \equiv 2$, one of x_1, x_2 , is congruent to 0 and the other to 1 mod 3. \square

5. Results. The previous corollaries provide partial information about $f(p)$. For example, Corollary 10 shows that $f(20) \geq 6$ and $f(42) \geq 8$ while Corollary 12 shows that $f(30) \geq 7$ and Corollary 13 shows $f(111) \geq 12$.

Using these corollaries, published results and calculations of difference covers in special cases we can construct Table 1. Obviously $f(p)$ is still difficult to compute.

6. Convergence. In view of Corollary 5 and the result on difference bases, it is natural to ask if the ratio $f^2(p)/p$ converges. If it does, then by Singer's theorem it must converge to 1. As of now it is not even known that f can take on a value as much as two above the bound given by $p \leq f^2(p) - f(p) + 1$.

An interesting necessary condition for convergence is given by Corollary 9. If n divides p , then a smallest difference cover must be fairly evenly distributed among the residues mod n in order to satisfy Theorem 6, but Corollary 9 states that, if $f^2(p)/p \rightarrow 1$, then a smallest difference cover for large multiples of n must be almost perfectly distributed.

7. Conclusions. While we have made progress in determining optimal sampling

TABLE 1
f(p) for small *p*

<i>p</i>	<i>f(p)</i>	<i>a difference cover</i>									
1	1	0									
2	2	0	1								
3	2	0	1								
4	3	0	1	2							
5	3	0	1	4							
6	3	0	1	3							
7	3	0	1	3							
8	4	0	1	3	7						
9	4	0	1	3	7						
10	4	0	1	3	6						
11	4	0	1	4	6						
12	4	0	1	4	6						
13	4	0	1	3	9						
14	5	0	1	3	7	9					
15	5	0	1	3	7	10					
16	5	0	1	3	7	11					
17	5	0	1	3	7	12					
18	5	0	1	3	10	15					
19	5	0	1	3	9	15					
20	6	0	1	3	9	13	15				
21	5	0	1	4	14	16					
22	6	0	1	4	9	11	15				
23	6	0	1	4	11	13	19				
24	6	0	1	3	7	12	17				
25	6	0	1	3	7	12	15				
26	6	0	1	4	14	19	21				
27	6	0	1	4	10	12	17				
28	6	0	1	4	15	20	22				
29		?									
30	7	0	1	2	6	10	13	16			
31	6	0	1	3	8	12	18				
32	7	0	1	4	9	11	17	23			
33	7	0	1	2	3	8	13	17			
34	7	0	1	2	3	8	13	17			
35	7	0	1	2	3	8	13	17			
36	7	0	1	11	16	19	23	25			
37	7	0	1	6	10	17	23	35			
38-41		?									
42	8	0	1	6	10	19	26	37	40		
43	8	0	8	18	19	22	24	31	39		
44	8	0	8	18	19	22	24	31	39		
45	8	0	8	18	19	22	24	31	39		
46	8	0	8	18	19	22	24	31	39		
47	8	0	8	18	19	22	24	31	39		
48	8	0	8	18	19	22	24	31	39		
49	8	0	8	18	19	22	24	31	39		
50-55		≤ 9									
56	9	0	1	10	19	21	37	43	49	52	
57	8	0	1	3	13	32	36	43	52		

(Recall Mann (1965) has results for $p = m^2 + m + 1$, $m \leq 1600$.)

patterns, the general answer seems extremely difficult. If one has a sample which does allow the estimation of all covariances, questions concerning the estimates and their properties are discussed in Parzen (1963).

Note the strange behavior of $f(p)$ shown by Table 1. For example, 6 points per cycle are required for $p = 20$ but only 5 for $p = 21$. This disturbance in the order of $f(p)$ tends to occur around the points $p = m^2 + m + 1$, $m = 4, 5, 6, \dots$.

Interesting open questions besides the complete determination of $f(p)$ remain. One, mentioned above, is whether $f(p)$ is ever more than one above the bound given by $p \leq f^2(p) - f(p) + 1$. A second is how many different difference covers for p exist. Of course, one could try to add information to Table 1. We have, for example, written an APL program which conducts a trial and error search for difference covers. It was this program that found the smallest difference covers for $p = 36, 37, 42, 56$ and found difference covers of size 9 for $p = 50, \dots, 55$.

8. Acknowledgment. The authors would like to thank Dr. Richard Duke for his helpful discussions on this problem.

REFERENCES

- [1] GOLAY, M. (1972). Notes on the representation of $1, 2, \dots, n$ by differences. *J. London Math. Soc.* (2) **4** 729-734.
- [2] JONES, R. H. (1962). Spectral analysis with regularly missed observations. *Ann. Math. Statist.* **33** 455-461.
- [3] JONES, R. H. (1971). Spectrum estimation with missing observations. *Ann. Inst. Statist. Math.* **23** 387-398.
- [4] LEECH, J. (1956). On the representation of $1, 2, \dots, n$ by differences. *J. London Math. Soc.* **31** 160-169.
- [5] LOYNES, R. M. (1969). Aliasing and related questions in stationary processes. *Proc. Twelfth Biennial Sem. Canad. Math. Cong. on Time Series and Stochastic Processes; Convexity and Combinatorics* (Vancouver, B.C.), 89-98.
- [6] MANN, H. (1965). *Addition Theorems: The Addition Theorems of Group Theory and Number Theory*, Tracts in Mathematics Number 18, Interscience, New York.
- [7] PARZEN, E. (1963). On spectral analysis with missing observations and amplitude modulation. *Sankhyā Ser. A* **25** 383-392.
- [8] REDEI, L. and RÉNYI, A. (1949). On the representation of $1, 2, \dots, n$ by differences. *Mat. Sb.* **66** (NS 24) 385-389 (in Russian).
- [9] RYSER, H. J. (1963). *Combinatorial Mathematics*. Carus Math. Monograph No. 14, Wiley, New York.
- [10] SINGER, J. (1938). A theorem in finite projective geometry and some applications to number theory. *Trans. Amer. Math. Soc.* **43** 377-385.

PROGRAMS IN MATHEMATICAL SCIENCES
THE UNIVERSITY OF TEXAS AT DALLAS
BOX 688
RICHARDSON, TEXAS 75080