

On upper and lower bounds of the numerical radius and an equality condition

by

TAKEAKI YAMAZAKI (Yokohama)

Abstract. We give an inequality relating the operator norm of T and the numerical radii of T and its Aluthge transform. It is a more precise estimate of the numerical radius than Kittaneh's result [Studia Math. 158 (2003)]. Then we obtain an equivalent condition for the numerical radius to be equal to half the operator norm.

1. Introduction. For a bounded linear operator T on a complex Hilbert space \mathcal{H} , we denote the operator norm and the numerical radius of T by $\|T\|$ and $w(T)$, respectively. It is well known that $w(T)$ is an equivalent norm of T , since (see [5, Theorem 1.3-1])

$$(1.1) \quad \frac{1}{2} \|T\| \leq w(T) \leq \|T\|.$$

Concerning the second inequality, Kittaneh [8] has shown the following precise estimate of $w(T)$ by using several norm inequalities and ingenious techniques:

$$(1.2) \quad w(T) \leq \frac{1}{2} \|T\| + \frac{1}{2} \|T^2\|^{1/2}.$$

Obviously, (1.2) is sharper than the right inequality of (1.1). We remark that we cannot compare $w(T)$ with $\|T^2\|^{1/2}$, generally. In fact, if $T = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, then $0 = \|T^2\|^{1/2} < w(T) = 1/2$; but if

$$T = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix},$$

then $1/\sqrt{2} = w(T) < \|T^2\|^{1/2} = 1$.

We obtain a sufficient condition for $w(T) = \frac{1}{2}\|T\|$ to hold from (1.1), (1.2) and [8]: if $T^2 = 0$, then $w(T) = \frac{1}{2}\|T\|$. But this condition is not

2000 *Mathematics Subject Classification*: Primary 47A12; Secondary 47A30, 47A20.

Key words and phrases: numerical radius, Aluthge transform, normaloid operators.

necessary: if $T = 1 \oplus \begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix}$, then $w(T) = \frac{1}{2}\|T\| = 1$, but $T^2 \neq 0$. We remark that some (necessary or sufficient) conditions for $w(T) = \frac{1}{2}\|T\|$ to hold are given in [5, Theorems 1.3-4 and 1.3-5], but no equivalent condition has been known yet.

Let $T = U|T|$ be the polar decomposition of T . The *Aluthge transform* \tilde{T} of T is defined by $\tilde{T} = |T|^{1/2}U|T|^{1/2}$ (see [1]). The following properties of \tilde{T} are well known:

- (i) $\|\tilde{T}\| \leq \|T\|$,
- (ii) $w(\tilde{T}) \leq w(T)$,
- (iii) $r(\tilde{T}) = r(T)$.

The first and last properties are easy by the definition of \tilde{T} , and the second one is shown in [7], [9] and [11]. Moreover for a non-negative integer n , we denote the n th Aluthge transform by \tilde{T}_n , i.e.,

$$\tilde{T}_n = \tilde{\tilde{T}}_{n-1} \quad \text{and} \quad \tilde{T}_0 = T.$$

This was first considered in [7] and [10], independently.

In this paper, first, we obtain a more precise estimate than (1.2). In the inequality, we use a bigger term, $\|T\|$, and a smaller one, $w(\tilde{T})$, than $w(T)$. Moreover the proof is very simple and needs only a generalized polarization identity. Next, we give a condition equivalent to $w(T) = \frac{1}{2}\|T\|$.

2. An inequality sharper than Kittaneh's. In this section, we prove a sharper estimate of $w(T)$ than Kittaneh's [8], as follows:

THEOREM 2.1. *For any $T \in B(\mathcal{H})$, $w(T) \leq \frac{1}{2}\|T\| + \frac{1}{2}w(\tilde{T})$.*

We remark that by the Heinz inequality [6], $\|A^r X B^r\| \leq \|A X B\|^r \|X\|^{1-r}$ for $A, B \geq 0$ and $r \in [0, 1]$, we have

$$(2.1) \quad w(\tilde{T}) \leq \|\tilde{T}\| = \||T|^{1/2}U|T|^{1/2}\| \leq \||T|U|T|\|^{1/2}\|U\|^{1/2} = \|T^2\|^{1/2},$$

i.e., Theorem 2.1 is sharper than (1.2).

To prove Theorem 2.1, we use the following famous formula which is called the generalized polarization identity:

THEOREM A (Generalized Polarization Identity). *For each $T \in B(\mathcal{H})$ and $x, y \in \mathcal{H}$,*

$$(2.2) \quad \begin{aligned} \langle Tx, y \rangle &= \frac{1}{4}(\langle T(x+y), x+y \rangle - \langle T(x-y), x-y \rangle) \\ &\quad + \frac{1}{4}i(\langle T(x+iy), x+iy \rangle - \langle T(x-iy), x-iy \rangle). \end{aligned}$$

Proof of Theorem 2.1. First of all, we note that

$$(2.3) \quad w(T) = \sup_{\theta \in \mathbb{R}} \|\operatorname{Re}(e^{i\theta}T)\|,$$

since

$$\sup_{\theta \in \mathbb{R}} \operatorname{Re}\{e^{i\theta} \langle Tx, x \rangle\} = |\langle Tx, x \rangle|$$

and

$$\sup_{\theta \in \mathbb{R}} \|\operatorname{Re}(e^{i\theta} T)\| = \sup_{\theta \in \mathbb{R}} w(\operatorname{Re}(e^{i\theta} T)) = w(T).$$

Let $T = U|T|$ be the polar decomposition. Then by (2.2), we have

$$\begin{aligned} \langle e^{i\theta} Tx, x \rangle &= \langle e^{i\theta} |T|x, U^*x \rangle \\ &= \frac{1}{4}(\langle |T|(e^{i\theta} + U^*)x, (e^{i\theta} + U^*)x \rangle - \langle |T|(e^{i\theta} - U^*)x, (e^{i\theta} - U^*)x \rangle) \\ &\quad + \frac{1}{4}i(\langle |T|(e^{i\theta} + iU^*)x, (e^{i\theta} + iU^*)x \rangle - \langle |T|(e^{i\theta} - iU^*)x, (e^{i\theta} - iU^*)x \rangle). \end{aligned}$$

Note that the inner products on the right hand side are all positive since $|T|$ is positive. Hence we have

$$\begin{aligned} \operatorname{Re}\langle e^{i\theta} Tx, x \rangle &= \frac{1}{4}(\langle (e^{-i\theta} + U)|T|(e^{i\theta} + U^*)x, x \rangle - \langle (e^{-i\theta} - U)|T|(e^{i\theta} - U^*)x, x \rangle) \\ &\leq \frac{1}{4}\langle (e^{-i\theta} + U)|T|(e^{i\theta} + U^*)x, x \rangle \\ &\leq \frac{1}{4}\|(e^{-i\theta} + U)|T|(e^{i\theta} + U^*)\| \\ &= \frac{1}{4}\||T|^{1/2}(e^{i\theta} + U^*)(e^{-i\theta} + U)|T|^{1/2}\| \quad (\text{since } \|X^*X\| = \|XX^*\|) \\ &= \frac{1}{4}\|2|T| + e^{i\theta}\tilde{T} + e^{-i\theta}(\tilde{T})^*\| \\ &= \frac{1}{2}\| |T| + \operatorname{Re}(e^{i\theta}\tilde{T}) \| \\ &\leq \frac{1}{2}\|T\| + \frac{1}{2}\|\operatorname{Re}(e^{i\theta}\tilde{T})\| \\ &\leq \frac{1}{2}\|T\| + \frac{1}{2}w(\tilde{T}) \quad (\text{by (2.3)}). \end{aligned}$$

Hence we have the desired inequality. ■

COROLLARY 2.2. *If $\tilde{T} = 0$, then $w(T) = \frac{1}{2}\|T\|$.*

Proof. The proof is easy by Theorem 2.1 and (1.1). ■

REMARK.

- (i) In Corollary 2.2, the conditions $\tilde{T} = 0$ and $w(T) = \frac{1}{2}\|T\|$ are not equivalent: if $T = 1 \oplus \begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix}$, then $w(T) = \frac{1}{2}\|T\| = 1$, but $\tilde{T} = 1 \oplus 0 \neq 0$.
- (ii) The conditions $\tilde{T} = 0$ and $T^2 = 0$ are equivalent. Indeed, let $T = U|T|$ be the polar decomposition. If $\tilde{T} = 0$, then

$$T^2 = U|T|U|T| = U|T|^{1/2}\tilde{T}|T|^{1/2} = 0.$$

Conversely, if $T^2 = 0$, then by (2.1) we have $\|\tilde{T}\| \leq \|T^2\|^{1/2} = 0$.

COROLLARY 2.3. For $T \in B(\mathcal{H})$,

$$w(T) \leq \sum_{n=1}^{\infty} \frac{1}{2^n} \|\tilde{T}_{n-1}\|.$$

Proof. By using Theorem 2.1 several times, we have

$$\begin{aligned} w(T) &\leq \frac{1}{2}\|T\| + \frac{1}{2}w(\tilde{T}) \leq \frac{1}{2}\|T\| + \frac{1}{2}\left\{\frac{1}{2}\|\tilde{T}\| + \frac{1}{2}w(\tilde{T}_2)\right\} \\ &= \frac{1}{2}\|T\| + \frac{1}{4}\|\tilde{T}\| + \frac{1}{4}w(\tilde{T}_2) \\ &\leq \frac{1}{2}\|T\| + \frac{1}{4}\|\tilde{T}\| + \frac{1}{8}\|\tilde{T}_2\| + \frac{1}{8}w(\tilde{T}_3) \leq \cdots \leq \sum_{n=1}^{\infty} \frac{1}{2^n} \|\tilde{T}_{n-1}\|. \quad \blacksquare \end{aligned}$$

Let

$$s(T) = \sum_{n=1}^{\infty} \frac{1}{2^n} \|\tilde{T}_{n-1}\|.$$

By (2.1), $\|\tilde{A}\| \leq \|A^2\|^{1/2} \leq \|A\|$ for any $A \in B(\mathcal{H})$, and we obtain

$$(2.4) \quad r(T) \leq w(T) \leq \frac{1}{2}\|T\| + \frac{1}{2}w(\tilde{T}) \leq s(T) \leq \frac{1}{2}\|T\| + \frac{1}{2}\|T^2\|^{1/2} \leq \|T\|,$$

where $r(T)$ means the spectral radius of T .

It is well known that T is *normaloid* (i.e., $\|T\| = r(T)$) if and only if $\|T\| = w(T)$. Here we give other conditions of normaloidity of T :

COROLLARY 2.4. *The following conditions are equivalent:*

- (i) T is *normaloid*,
- (ii) $\|T\| = s(T)$,
- (iii) $r(T) = \frac{1}{2}\|T\| + \frac{1}{2}w(\tilde{T})$,
- (iv) $s(T) = s(\tilde{T})$.

REMARK.

- (i) In Corollary 2.4, condition (ii) cannot be replaced by the weaker condition $\|T\| = \frac{1}{2}\|T\| + \frac{1}{2}\|T^2\|^{1/2}$. For example, let

$$T = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}.$$

Then $\|T\| = \frac{1}{2}\|T\| + \frac{1}{2}\|T^2\|^{1/2} = 1$ but $0 = r(T) < \|T\|$.

- (ii) In Corollary 2.4, condition (iii) cannot be replaced by the weaker condition $r(T) = w(T)$ either. In fact, let $T = 1 \oplus \begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix}$. Then $1 = r(T) = w(T) < \|T\| = 2$. (We call an operator satisfying $r(T) = w(T)$ *spectraloid*.)

To prove Corollary 2.4, the following formula will be used.

THEOREM B ([10]). For any $T \in B(\mathcal{H})$, $\lim_{n \rightarrow \infty} \|\tilde{T}_n\| = r(T)$.

Proof of Corollary 2.4. (i) \Rightarrow (ii), (iii) and (iv) are obvious by (2.4) and $r(T) = r(\tilde{T}) \leq s(\tilde{T}) \leq s(T) \leq \|T\|$.

(ii) \Rightarrow (i). By the definition of $s(T)$,

$$(2.5) \quad s(T) = \frac{1}{2}\|T\| + \frac{1}{2}s(\tilde{T}).$$

Hence (ii) yields $s(\tilde{T}) = \|T\|$. Since $\|\tilde{T}\| \leq \|T\|$, this gives

$$s(\tilde{T}) \leq \|\tilde{T}\| \leq \|T\| = s(\tilde{T}),$$

and so $s(\tilde{T}) = \|\tilde{T}\| = \|T\|$. By using the same technique, we have $\|T\| = \|\tilde{T}_n\|$ for all $n \in \mathbb{N}$. Hence by Theorem B, we have

$$\|T\| = \lim_{n \rightarrow \infty} \|\tilde{T}_n\| = r(T),$$

that is, T is normaloid.

(iii) \Rightarrow (i). Since $r(\tilde{T}) = r(T)$, by (iii) we have

$$r(T) = \frac{1}{2}\|T\| + \frac{1}{2}w(\tilde{T}) \geq \frac{1}{2}\|T\| + \frac{1}{2}r(\tilde{T}) = \frac{1}{2}\|T\| + \frac{1}{2}r(T),$$

that is, $r(T) \geq \|T\|$ and so $r(T) = \|T\|$.

(iv) \Rightarrow (ii). Evident by (2.5). ■

In [2], Ando shows that the equality $W(T) = W(\tilde{T})$ of numerical ranges is equivalent to $\text{co } \sigma(T) = W(T)$ (i.e., T is convexoid) for any matrix T , where $\text{co } \sigma(T)$ means the convex hull of the spectrum of T . We think that this result is parallel to the equivalence between (i) and (iv). So we expect that $s(T)$ has some interesting properties.

3. Condition equivalent to $w(T) = \frac{1}{2}\|T\|$. In Corollary 2.2, we have obtained a sufficient condition for $w(T) = \frac{1}{2}\|T\|$ to hold. Some (necessary or sufficient) conditions for $w(T) = \frac{1}{2}\|T\|$ to hold are given in [5, Theorems 1.3-4 and 1.3-5]. But no condition equivalent to $w(T) = \frac{1}{2}\|T\|$ has been known. In this section, we give such a condition:

THEOREM 3.1. *Let $T \in B(\mathcal{H})$. The following conditions are equivalent:*

- (i) $w(T) = \frac{1}{2}\|T\|$,
- (ii) $\|T\| = \|\text{Re}(e^{i\theta}T)\| + \|\text{Im}(e^{i\theta}T)\|$ for all $\theta \in \mathbb{R}$.

We remark that (ii) cannot be replaced by “ $\|T\| = \|\text{Re}(e^{i\theta}T)\| + \|\text{Im}(e^{i\theta}T)\|$ for some $\theta \in \mathbb{R}$,” because if T is a non-zero self-adjoint operator, then $\|T\| = \|\text{Re } T\| + \|\text{Im } T\| = \|\text{Re } T\|$, but $w(T) = \|T\| > \frac{1}{2}\|T\|$.

To prove Theorem 3.1, we need the following theorem:

THEOREM C ([3]). *Let $A, B \in B(\mathcal{H})$ be non-zero. Then the equality $\|A + B\| = \|A\| + \|B\|$ holds if and only if $\|A\| \|B\| \in \overline{W(A^*B)}$.*

Proof of Theorem 3.1. Let $e^{i\theta}T = H_\theta + iK_\theta$ be the Cartesian decomposition of $e^{i\theta}T$. We remark that

$$(3.1) \quad K_\theta = H_{\theta-\pi/2},$$

because $e^{i(\theta-\pi/2)}T = -ie^{i\theta}T = K_\theta - iH_\theta$.

(i) \Rightarrow (ii). Since $w(T) = \sup_{\theta \in \mathbb{R}} \|H_\theta\| = \sup_{\theta \in \mathbb{R}} \|K_\theta\|$ by (2.3) and (3.1), we have

$$\|T\| = \|e^{i\theta}T\| = \|H_\theta + iK_\theta\| \leq \|H_\theta\| + \|K_\theta\| \leq w(T) + w(T) = \|T\|,$$

proving (ii).

(ii) \Rightarrow (i). For any $\theta \in \mathbb{R}$, (ii) ensures $\|H_\theta\| \|K_\theta\| \in \overline{W(H_\theta^*(iK_\theta))}$ by Theorem C, i.e., $-i\|H_\theta\| \|K_\theta\| \in \overline{W(H_\theta K_\theta)}$. Since $-i\|H_\theta\| \|K_\theta\|$ is a purely imaginary number and $\text{Im}(H_\theta K_\theta) = \text{Im}(H_0 K_0)$ for all $\theta \in \mathbb{R}$, we have

$$\|H_\theta\| \|K_\theta\| = w(H_\theta K_\theta) = \|\text{Im}(H_\theta K_\theta)\| = \|\text{Im}(H_0 K_0)\|.$$

Thus for all $\theta \in \mathbb{R}$,

$$\|H_\theta\| + \|K_\theta\| = \|T\|, \quad \|H_\theta\| \|K_\theta\| = \|\text{Im}(H_0 K_0)\|,$$

that is,

$$\|H_\theta\| = \frac{\|T\| + \sqrt{\|T\|^2 - 4\|\text{Im}(H_0 K_0)\|}}{2},$$

$$\|K_\theta\| = \frac{\|T\| - \sqrt{\|T\|^2 - 4\|\text{Im}(H_0 K_0)\|}}{2},$$

or the other way round. We remark that these values do not depend on $\theta \in \mathbb{R}$. So the function $\|H_\theta\|$ of $\theta \in \mathbb{R}$ takes only two values by (3.1). By an easy calculation, we have

$$H_\theta = H_0 \cos \theta - K_0 \sin \theta.$$

Hence by the continuity of the operator norm, the function $\|H_\theta\|$ is continuous on $\theta \in \mathbb{R}$. Therefore it must be constant, i.e.,

$$\|H_\theta\| = \|K_\theta\| = \frac{1}{2}\|T\|.$$

Hence we have (i). ■

References

- [1] A. Aluthge, *Some generalized theorems on p -hyponormal operators*, Integral Equations Operator Theory 24 (1996), 497–501.
- [2] T. Ando, *Aluthge transforms and the convex hull of the eigenvalues of a matrix*, Linear Multilinear Algebra 52 (2004), 281–292.
- [3] M. Barraa and M. Boumazgour, *Inner derivations and norm equality*, Proc. Amer. Math. Soc. 130 (2002), 471–476.
- [4] T. Furuta, *Invitation to Linear Operators*, Taylor and Francis, London, 2001.
- [5] K. E. Gustafson and D. K. M. Rao, *Numerical Range*, Springer, New York, 1997.

- [6] E. Heinz, *Beiträge zur Störungstheorie der Spektralzerlegung*, Math. Ann. 123 (1951), 415–438.
- [7] I. B. Jung, E. Ko and C. Pearcy, *Aluthge transforms of operators*, Integral Equations Operator Theory 37 (2000), 437–448.
- [8] F. Kittaneh, *A numerical radius inequality and an estimate for the numerical radius of the Frobenius companion matrix*, Studia Math. 158 (2003), 11–17.
- [9] P. Y. Wu, *Numerical range of Aluthge transform of operator*, Linear Algebra Appl. 357 (2002), 295–298.
- [10] T. Yamazaki, *An expression of spectral radius via Aluthge transformation*, Proc. Amer. Math. Soc. 130 (2002), 1131–1137.
- [11] —, *On numerical range of the Aluthge transformation*, Linear Algebra Appl. 341 (2002), 111–117.

Department of Mathematics
Kanagawa University
Yokohama 221-8686, Japan
E-mail: yamazt26@kanagawa-u.ac.jp

Received April 7, 2006

Revised version September 25, 2006

(5894)