



COMPOSITIO MATHEMATICA

On volumes of arithmetic line bundles

Xinyi Yuan

Compositio Math. **145** (2009), 1447–1464.

[doi:10.1112/S0010437X0900428X](https://doi.org/10.1112/S0010437X0900428X)



FOUNDATION
COMPOSITIO
MATHEMATICA

*The London
Mathematical
Society*





On volumes of arithmetic line bundles

Xinyi Yuan

ABSTRACT

We show an arithmetic generalization of the recent work of Lazarsfeld–Mustață which uses Okounkov bodies to study linear series of line bundles. As applications, we derive a log-concavity inequality on volumes of arithmetic line bundles and an arithmetic Fujita approximation theorem for big line bundles.

1. Introduction

In their recent paper [LM08], Lazarsfeld and Mustață explored a systematic way of using Okounkov bodies, originated in [Oko96, Oko03], to study the volumes of line bundles over algebraic varieties. They easily recovered many positivity results in algebraic geometry (cf. [Laz04]). A similar construction with a different viewpoint was also taken by Kaveh–Khovanskii [KK08]. Our paper is the expected arithmetic analogue of [LM08]. Our main results are as follows.

- Introduce arithmetic Okounkov bodies associated to an arithmetic line bundle, and prove that the volumes of the former approximate the volume of the later.
- Show some log-concavity inequalities on the volumes and top intersection numbers, which can be viewed as a high-dimensional generalization of the Hodge index theorem on arithmetic surfaces of Faltings [Fal84].
- Prove an arithmetic analogue of Fujita’s approximation theorem, which is proved independently by Huayi Chen [Che08b] during the preparation of this paper.
- As by-products, we recover the convergence of $\hat{h}^0(X, m\bar{\mathcal{L}})/(m^d/d!)$ proved by Chen [Che08a] and the arithmetic Hodge index theorem in codimension one proved by Moriwaki [Mor96].

1.1 Volume of an arithmetic line bundle

Let X be an arithmetic variety of dimension d . That is, X is a d -dimensional integral scheme, projective and flat over $\text{Spec}(\mathbb{Z})$. For any Hermitian line bundle $\bar{\mathcal{L}} = (\mathcal{L}, \|\cdot\|)$ over X , denote

$$\hat{H}^0(X, \bar{\mathcal{L}}) = \{s \in H^0(X, \mathcal{L}) : \|s\|_{\text{sup}} \leq 1\}$$

and

$$\hat{h}^0(X, \bar{\mathcal{L}}) = \log \# \hat{H}^0(X, \bar{\mathcal{L}}).$$

Define the volume to be

$$\text{vol}(\bar{\mathcal{L}}) = \limsup_{m \rightarrow \infty} \frac{\hat{h}^0(X, m\bar{\mathcal{L}})}{m^d/d!}.$$

Received 12 November 2008, accepted in final form 5 March 2009.

2000 Mathematics Subject Classification 14G40 (primary), 11G35, 11G50 (secondary).

Keywords: arithmetic varieties, Hermitian line bundles, Arakelov theory, volumes of line bundles, big line bundles, Fujita approximation, Hodge index theorem.

The author is fully supported by a research fellowship of the Clay Mathematics Institute.

This journal is © Foundation Compositio Mathematica 2009.

Note that in this paper we write line bundles additively, so $m\bar{\mathcal{L}}$ means $\bar{\mathcal{L}}^{\otimes m}$.

A line bundle $\bar{\mathcal{L}}$ is said to be big if $\text{vol}(\bar{\mathcal{L}}) > 0$; it is effective if $\hat{h}^0(X, \bar{\mathcal{L}}) > 0$. If $\bar{\mathcal{L}}$ is ample in the sense of Zhang [Zha95] (cf. § 2.1), then it is big by the formula

$$\text{vol}(\bar{\mathcal{L}}) = \bar{\mathcal{L}}^d > 0.$$

This is a result combining the works of Gillet–Soulé [GS91, GS92], Bismut–Vasserot [BV89] and Zhang [Zha95]. See [Yua08, Corollary 2.7] for example.

As pointed out above, Huayi Chen proved that ‘limsup = lim’ in the definition of $\text{vol}(\bar{\mathcal{L}})$ in his recent work [Che08a] using Harder–Narasimhan filtrations. We will derive this result by means of Okounkov bodies in Theorem 2.7.

Some basic properties of big line bundles are proved in [Mor00, Mor09, Yua08]. They use different definitions of bigness, but [Yua08, Corollary 2.4] shows that all of these definitions are equivalent. The result ‘big = ample + effective’ of [Yua08] rephrased in Theorem 2.2 will be widely used in this paper. It is also worth noting that the main result of Moriawaki [Mor09] asserts that the volume function is continuous at all Hermitian line bundles.

1.2 Okounkov body

Assume that X is normal with smooth generic fibre $X_{\mathbb{Q}}$. Let

$$X \supset Y_1 \supset \cdots \supset Y_d$$

be a flag on X , where each Y_i is a regular irreducible closed subscheme of codimension i in X . We require that Y_1 is a vertical divisor lying over some finite prime p , and write $\text{char}(Y_i) = p$ in this case. We further require that the residue field of Y_d is isomorphic to \mathbb{F}_p . There is a positive density of such prime p for which Y_i exists.

Define a valuation map

$$\nu_Y = (\nu_1, \dots, \nu_d) : H^0(X, \mathcal{L}) - \{0\} \rightarrow \mathbb{Z}^d$$

with respect to the flag Y as in [LM08]. We explain it here. For any non-zero $s \in H^0(X, \mathcal{L})$, we first set $\nu_1(s) = \text{ord}_{Y_1}(s)$. Let s_{Y_1} be a section of the line bundle $\mathcal{O}(Y_1)$ with zero locus Y_1 . Then $s_{Y_1}^{\otimes(-\nu_1(s))} s$ is non-zero on Y_1 , and let $s_1 = (s_{Y_1}^{\otimes(-\nu_1(s))} s)|_{Y_1}$ be the restriction. Set $\nu_2(s) = \text{ord}_{Y_2}(s_1)$. Continue this process on the section s_1 on Y_2 , we can define $\nu_3(s)$ and thus $\nu_4(s), \dots, \nu_d(s)$.

For any Hermitian line bundle $\bar{\mathcal{L}}$ on X , denote

$$v_Y(\bar{\mathcal{L}}) = \nu_Y(\hat{H}^0(X, \bar{\mathcal{L}}) - \{0\})$$

to be the image in \mathbb{Z}^d . Note that we only pick up the image of the finite set $\hat{H}^0(X, \bar{\mathcal{L}}) - \{0\}$.

Let $\Delta_Y(\bar{\mathcal{L}})$ be the closure of $\Lambda_Y(\bar{\mathcal{L}}) = \bigcup_{m \geq 1} (1/m^d)v_Y(m\bar{\mathcal{L}})$ in \mathbb{R}^d . It turns out that $\Delta_Y(\bar{\mathcal{L}})$ is a bounded convex subset of \mathbb{R}^d if non-empty. It has a finite volume under the Lebesgue measure of \mathbb{R}^d . See Lemma 2.4. We have the following counterpart of [LM08, Theorem A].

THEOREM A. *If $\bar{\mathcal{L}}$ is big, then*

$$\lim_{p=\text{char}(Y_i) \rightarrow \infty} \text{vol}(\Delta_Y(\bar{\mathcal{L}})) \log p = \frac{1}{d!} \text{vol}(\bar{\mathcal{L}}).$$

In the geometric case of [LM08], exact equality between two volumes are easily obtained without taking limit on p . However, it seems hard to be true in the arithmetic case if $\bar{\mathcal{L}}$ is not ample. Nevertheless, the above result is apparently sufficient for applications of $\text{vol}(\bar{\mathcal{L}})$.

1.3 Log-concavity and Hodge index theorem

As what Lazarsfeld and Mustață do, we also show the log-concavity of volume functions by the classical Brunn–Minkowski theorem in Euclidean geometry.

THEOREM B. *For any two effective line bundles $\overline{\mathcal{L}}_1, \overline{\mathcal{L}}_2$, we have*

$$\text{vol}(\overline{\mathcal{L}}_1 + \overline{\mathcal{L}}_2)^{1/d} \geq \text{vol}(\overline{\mathcal{L}}_1)^{1/d} + \text{vol}(\overline{\mathcal{L}}_2)^{1/d}.$$

When $\overline{\mathcal{L}}_1$ and $\overline{\mathcal{L}}_2$ are ample, the above volumes are exactly equal to arithmetic intersection numbers. Even in this case, the inequality is not as transparent as the geometric case. The result can be viewed as a generalization of the Hodge index theorem on arithmetic surfaces. See [Laz04] for many related inequalities in the geometric case.

An easy consequence of the above result is the relation

$$(\overline{\mathcal{L}}_1^{d-1} \cdot \overline{\mathcal{L}}_2)^2 \geq (\overline{\mathcal{L}}_1^d)(\overline{\mathcal{L}}_1^{d-2} \cdot \overline{\mathcal{L}}_2^2)$$

on intersection numbers for any two ample line bundles $\overline{\mathcal{L}}_1, \overline{\mathcal{L}}_2$. This simple-looking relation is equivalent to the Hodge index theorem for divisors on arithmetic varieties, which was proved by Moriwaki [Mor96]. See Corollaries 3.1 and 3.2.

1.4 Arithmetic Fujita approximation

In the geometric case, one of the most important properties of big line bundles is Fujita’s approximation in [Fuj94]. It asserts that a big line bundle can be arbitrarily closely approximated by ample line bundles. In this way, many properties of ample line bundles can be carried to big line bundles. Our arithmetic analogue is as follows.

THEOREM C. *Let $\overline{\mathcal{L}}$ be a big line bundle over X . Then for any $\epsilon > 0$, there exist an integer $n > 0$, a birational morphism $\pi : X' \rightarrow X$ from another arithmetic variety X' to X , and an isomorphism*

$$n\pi^*\overline{\mathcal{L}} = \overline{\mathcal{A}} + \overline{\mathcal{E}}$$

for an effective line bundle $\overline{\mathcal{E}}$ on X' and an ample line bundle $\overline{\mathcal{A}}$ on X' satisfying

$$\frac{1}{n^d} \text{vol}(\overline{\mathcal{A}}) > \text{vol}(\overline{\mathcal{L}}) - \epsilon.$$

Our proof of this theorem consists of a finite part and an infinite part. The finite part is an analogue in [LM08, Theorem 3.3], the infinite part is solved by taking pluri-subharmonic envelope which is well-known in complex analysis.

As pointed out at the beginning, Chen [Che08b] also proves this result. His proof also relies on [LM08]. The difference between our approaches is that, he really applies the original [LM08, Theorem 3.5] combining with Bost’s slope theory, while here we prove an arithmetic analogue of the theorem. Furthermore, he does not obtain our Theorems A and B.

1.5 Notations

We use $\widehat{\text{Pic}}(X), \widehat{\text{Amp}}(X), \widehat{\text{Big}}(X), \widehat{\text{Eff}}(X)$ to denote respectively the **isometry classes** of Hermitian line bundles, ample Hermitian line bundles, big Hermitian line bundles, effective Hermitian line bundles.

When we treat the valuation ν , we always ignore the fact that the section 0 has no image. For example, $\nu(S)$ is understood as $\nu(S - \{0\})$ for any subset $S \subset H^0(X, \mathcal{L})$.

For any smooth function $f : X(\mathbb{C}) \rightarrow \mathbb{R}$, denote by $\overline{\mathcal{O}}(f) = (\mathcal{O}, e^{-f})$ the trivial bundle \mathcal{O} endowed with the metric given by $\|1\| = e^{-f}$. In particular, it makes sense if $f = \alpha \in \mathbb{R}$ is a constant function. For any vertical Cartier divisor V of X , denote by $\overline{\mathcal{O}}(V) = (\mathcal{O}(V), \|\cdot\|)$ the line bundle $\mathcal{O}(V)$ associated to V with a metric given by $\|s_V\| = 1$. Here s_V denotes a fixed section defining V . We further denote $\overline{\mathcal{L}}(f + V) = \overline{\mathcal{L}} + \overline{\mathcal{O}}(f) + \overline{\mathcal{O}}(V)$ for any Hermitian line bundle $\overline{\mathcal{L}}$.

2. The arithmetic Okounkov body

The main goal of this section is to prove Theorem A. Section 2.1 recalls some results on ample line bundles and big line bundles. After considering some easy properties followed from [LM08], the proof of Theorem A is reduced to Theorem 2.6 in §2.2. Then we prove Theorem 2.6 in the next two subsections.

2.1 Basics on arithmetic ampleness and bigness

We follow the arithmetic intersection theory of Gillet–Soulé [GS90] and the notion of arithmetic ampleness by Zhang [Zha95].

Recall that an arithmetic variety is an integral scheme, projective and flat over $\text{Spec}(\mathbb{Z})$. The dimension means the absolute dimension. Let X be an arithmetic variety of dimension d . The notion of Hermitian line bundles needs more words if the complex space $X(\mathbb{C})$ is not smooth.

A metrized line bundle $\overline{\mathcal{L}} = (\mathcal{L}, \|\cdot\|)$ over X is an invertible sheaf \mathcal{L} over X together with a Hermitian metric $\|\cdot\|$ on each fibre of $\mathcal{L}(\mathbb{C})$ over $X(\mathbb{C})$. We say this metric is smooth if the pull-back metric over $f^*\mathcal{L}$ under any analytic map $f : B^{d-1} \rightarrow X(\mathbb{C})$ is smooth in the usual sense. Here B^{d-1} denotes the unit ball in \mathbb{C}^{d-1} . We call $\overline{\mathcal{L}}$ a Hermitian line bundle if its metric is smooth and invariant under complex conjugation. For a Hermitian line bundle $\overline{\mathcal{L}}$, we say the metric or the curvature of $\overline{\mathcal{L}}$ is semipositive if the curvature of $f^*\mathcal{L}$ with the pull-back metric under any analytic map $f : B^{d-1} \rightarrow X(\mathbb{C})$ is semipositive definite.

A Hermitian line bundle $\overline{\mathcal{L}}$ over X is called ample if the following three conditions are satisfied.

- (a) The generic fibre $\mathcal{L}_{\mathbb{Q}}$ is ample.
- (b) The Hermitian line bundle $\overline{\mathcal{L}}$ is relatively semipositive: the curvature of $\overline{\mathcal{L}}$ is semipositive and $\deg(\mathcal{L}|_C) \geq 0$ for any closed curve C on any special fibre of X over $\text{Spec}(\mathbb{Z})$.
- (c) The Hermitian line bundle $\overline{\mathcal{L}}$ is horizontally positive: the intersection number $(\overline{\mathcal{L}}|_Y)^{\dim Y}$ is greater than zero for any horizontal irreducible closed subvariety Y .

Zhang proved an arithmetic Nakai–Moishezon theorem which includes a special case as follows.

THEOREM 2.1 [Zha95, Corollary 4.8]. *Let $\overline{\mathcal{L}}$ be an ample Hermitian line bundle on an arithmetic variety X such that $X_{\mathbb{Q}}$ is smooth. Then for any Hermitian line bundle $\overline{\mathcal{E}}$ over X , the \mathbb{Z} -module $H^0(X, \mathcal{E} + N\mathcal{L})$ has a basis consisting of strictly effective sections for N large enough.*

Here an effective section is a non-zero section with supremum norm less than or equal to 1. If the supremum norm of the section is less than 1, the section and the line bundle are said to be strictly effective.

As for big line bundles, we need the following result.

THEOREM 2.2 [Yua08, Theorem 2.1]. *A Hermitian line bundle $\overline{\mathcal{L}}$ on X is big if and only if $N\overline{\mathcal{L}} = \overline{\mathcal{A}} + \overline{\mathcal{E}}$ for some integer $N > 0$, some $\overline{\mathcal{A}} \in \widehat{\text{Amp}}(X)$ and some $\overline{\mathcal{E}} \in \widehat{\text{Eff}}(X)$.*

Fujita's approximation roughly says that we can make the ample part $(1/N)\overline{\mathcal{A}}$ arbitrarily close to $\overline{\mathcal{L}}$.

In the end, we quote a theorem of Moriwaki which says that the volume function is invariant under birational morphisms. We need it when we use generic resolution of singularities.

THEOREM 2.3 [Mor09, Theorem 4.2]. *Let $\pi : \widetilde{X} \rightarrow X$ be a birational morphism of arithmetic varieties. Then for any Hermitian line bundle $\overline{\mathcal{L}} \in \widehat{\text{Pic}}(X)$, we have $\text{vol}(\pi^*\overline{\mathcal{L}}) = \text{vol}(\overline{\mathcal{L}})$.*

2.2 Volumes of Okounkov bodies

We always assume X to be normal with smooth generic fibre when we consider Okounkov bodies. Recall that for the flag

$$X \supset Y_1 \supset Y_2 \supset \cdots \supset Y_d,$$

we require that Y_1 is vertical over some prime p and the residue field of Y_d is isomorphic to \mathbb{F}_p . This is not essential, but we will only stick on this case for simplicity. We will first explain why such p has a positive density.

Let K be the largest algebraic number field contained in the fraction field of X . Then the structure morphism $X \rightarrow \text{Spec}(\mathbb{Z})$ factors through $X \rightarrow \text{Spec}(O_K)$ where O_K is the ring of integers of K , and X is geometrically connected over O_K . It follows that the fiber X_φ over any prime ideal φ of O_K is connected. It is smooth for almost all φ . We must have $Y_1 = X_\varphi$ for some φ lying over p .

We require that the residue field $\mathbb{F}_\varphi = O_K/\varphi$ is isomorphic to \mathbb{F}_p . For example, it is true if p splits completely in O_K , and it happens with a positive density by Chebotarev's density theorem. Once this is true, it is easy to choose Y_2, Y_3, \dots, Y_{d-1} . The existence of a point Y_d follows from Weil's conjecture for curves over finite fields.

We start with some basic properties of the Okounkov body we defined. Recall that

$$\nu = \nu_Y = (\nu_1, \dots, \nu_d) : H^0(X, \mathcal{L}) - \{0\} \rightarrow \mathbb{Z}^d$$

is the corresponding valuation map,

$$v(\overline{\mathcal{L}}) = \nu_Y(\overline{\mathcal{L}}) = \nu_Y(\widehat{H}^0(X, \overline{\mathcal{L}}) - \{0\})$$

is the image in \mathbb{Z}^d , and $\Delta = \Delta_Y(\overline{\mathcal{L}})$ is the closure of $\Lambda = \Lambda_Y(\overline{\mathcal{L}}) = \bigcup_{m \geq 1} (1/m^d)v(m\overline{\mathcal{L}})$ in \mathbb{R}^d .

LEMMA 2.4. *The Okounkov body $\Delta_Y(\overline{\mathcal{L}})$ is convex and bounded for any $\overline{\mathcal{L}} \in \widehat{\text{Pic}}(X)$.*

Proof. We first show convexity. By taking limit, it suffices to show that $\sum_{i=0}^k a_i x_i \in \Lambda$ for all $x_i \in \Lambda$ and $a_i \in \mathbb{Q}_{>0}$ satisfying $\sum_{i=0}^k a_i = 1$. Assume that x_i comes from the section $s_i \in \widehat{H}^0(X, m_i \overline{\mathcal{L}})$. Let N be a positive common denominator of a_i/m_i , and write $a_i/m_i = b_i/N$. Then the section

$$\bigotimes_{i=0}^k s_i^{\otimes b_i} \in \widehat{H}^0(X, N\overline{\mathcal{L}})$$

gives the point $\sum_{i=0}^k a_i x_i \in \Lambda$.

Next we show boundedness. Similar to [LM08], we will show that there exists an integer $b > 0$ such that

$$\nu_i(s) \leq mb \quad \forall s \in \widehat{H}^0(X, m\overline{\mathcal{L}}) - \{0\}, i = 1, \dots, d. \quad (1)$$

The valuation (ν_2, \dots, ν_d) is exactly the valuation of dimension $d - 1$ with respect to the flag $Y_1 \supset Y_2 \supset \cdots \supset Y_d$ on the ambient variety Y_1 . Hence, the bound of ν_i for $i > 1$ follows from [LM08, Proposition 2.1]. It remains to bound ν_1 .

Fix an ample line bundle $\overline{\mathcal{A}}$ on X . For any $s \in \widehat{H}^0(X, m\overline{\mathcal{L}}) - \{0\}$, we have

$$\overline{\mathcal{L}} \cdot \overline{\mathcal{A}}^{d-1} = \frac{1}{m}(\overline{\mathcal{A}}|_{\text{div}(s)})^{d-1} - \frac{1}{m} \int_{X(\mathbb{C})} \log \|s\|_{c_1(\overline{\mathcal{M}})}^{d-1} \geq \frac{\nu_1(s)}{m}(\overline{\mathcal{A}}|_{Y_1})^{d-1}.$$

Thus we get a bound

$$\nu_1(s) \leq \frac{\overline{\mathcal{L}} \cdot \overline{\mathcal{A}}^{d-1}}{(\overline{\mathcal{A}}|_{Y_1})^{d-1}} m. \quad \square$$

It is natural to describe the volume of the Okounkov body in terms of the order of the images of the valuations. The proof is actually an argument of Okounkov [Oko03] using some results of Khovanskii [Kho92] in convex geometry, but we will only refer to the setting of [LM08] below.

PROPOSITION 2.5. *If $\overline{\mathcal{L}}$ is big, then*

$$\lim_{m \rightarrow \infty} \frac{\#v_Y(m\overline{\mathcal{L}})}{m^d} = \text{vol}(\Delta_Y).$$

Proof. Note that in our arithmetic case,

$$\Gamma = \bigcup_{m \geq 0} (v(m\overline{\mathcal{L}}), m) \subset \mathbb{Z}^{d+1}$$

is also a semigroup. We will apply [LM08, Proposition 2.1] on Γ . We only need to check that Γ satisfies conditions (2.3)–(2.5) required by the proposition. The proof is similar to Lemma 2.2 of the paper.

Condition (2.3) is trivial. Condition (2.4) follows from (1) in the proof of Lemma 2.4. In fact, Γ is contained in the the semigroup generated by $\{(x_1, x_2, \dots, x_d, 1) : x_i = 0, 1, \dots, b\}$. It remains to check (2.5).

We first look at the case that $\overline{\mathcal{L}}$ is ample. By the arithmetic Nakai–Moishezon theorem proved by Zhang [Zha95] (cf. Theorem 2.1), when m is sufficiently large, $H^0(m\overline{\mathcal{L}})$ has a \mathbb{Z} -basis consisting of effective sections. By this it is easy to find an $s \in \widehat{H}^0(m\overline{\mathcal{L}})$ which is non-zero on Y_d , or equivalently $\nu(s) = 0$. It follows that $(0, \dots, 0, m) \in \Gamma$. We also have $(0, \dots, 0, m+1) \in \Gamma$. Then we see that $(0, \dots, 0, 1)$ is generated by two elements of Γ . It remains to show that $\bigcup_{m \geq 0} v(m\overline{\mathcal{L}})$ generates \mathbb{Z}^d . We will show that one $v(m\overline{\mathcal{L}})$ is enough if m is sufficiently large.

For any $i = 1, 2, \dots, d$, we can find a line bundle \mathcal{M}_i on X with a section $t_i \in H^0(X, \mathcal{M}_i)$ such that t_i does not vanish on Y_{i-1} , vanishes on Y_i , and vanishes on Y_d with order one. Then $\{\nu(t_i)\}$ is exactly the standard basis of \mathbb{Z}^d . Choose and fix one metric on \mathcal{M}_i such that t_i is effective. Denote the Hermitian line bundle so obtained by $\overline{\mathcal{M}}_i$. Consider the line bundle $m\overline{\mathcal{L}} - \overline{\mathcal{M}}_i$. We can find a section $s_i \in \widehat{H}^0(m\overline{\mathcal{L}} - \overline{\mathcal{M}}_i)$ with $\nu(s_i) = 0$. The existence is still a simple consequence of Zhang’s theorem, which works on $m\overline{\mathcal{L}} - \overline{\mathcal{M}}_i$ when m is large enough. The section $s_i \otimes t_i \in \widehat{H}^0(m\overline{\mathcal{L}})$, and $\{\nu(s_i \otimes t_i)\}$ form the standard basis of \mathbb{Z}^d .

Now we assume that $\overline{\mathcal{L}}$ is any big line bundle. By Theorem 2.2, we get $N\overline{\mathcal{L}} = \overline{\mathcal{L}}' + \overline{\mathcal{E}}$ for some integer $N > 0$, some ample line bundle $\overline{\mathcal{L}}'$ and some effective line bundle $\overline{\mathcal{E}}$. Following the line above, we first show that $(0, \dots, 0, 1) \in \mathbb{Z}^{d+1}$ is generated by Γ . Fix an effective non-zero section $e \in \overline{\mathcal{E}}$. For m large enough, by the above argument we have non-zero sections $s \in \widehat{H}^0(m\overline{\mathcal{L}}')$ and $s' \in \widehat{H}^0(m\overline{\mathcal{L}}' + \overline{\mathcal{L}})$ with valuation $\nu(s) = \nu(s') = 0$. Now the sections $s \otimes e^{\otimes m} \in \widehat{H}^0(mN\overline{\mathcal{L}})$ and $s' \otimes e^{\otimes m} \in \widehat{H}^0((mN+1)\overline{\mathcal{L}})$ give

$$(\nu(s' \otimes e^{\otimes m}), mN+1) - (\nu(s \otimes e^{\otimes m}), mN) = (0, \dots, 0, 1).$$

It remains to show that $\bigcup_{m \geq 0} v(m\bar{\mathcal{L}})$ generates \mathbb{Z}^d . Let s, e be as above. For any non-zero section $u \in \widehat{H}^0(m\bar{\mathcal{L}}')$, we have $\nu(e^{\otimes m} \otimes u) - \nu(e^{\otimes m} \otimes s) = \nu(u)$ is the difference of two elements in $v(mN\bar{\mathcal{L}})$. If m is sufficiently large, $v(mN\bar{\mathcal{L}})$ generates \mathbb{Z}^d since $v(m\bar{\mathcal{L}}')$ does. \square

In the end, we state a theorem whose proof will take up the rest of this section. Combined with Proposition 2.5, it simply implies Theorem A and the convergence result of Chen [Che08a].

THEOREM 2.6. *For any $\bar{\mathcal{L}} \in \widehat{\text{Pic}}(X)$, there exists a constant $c = c(\bar{\mathcal{L}})$ depending only on $(X, \bar{\mathcal{L}})$, such that*

$$\limsup_{m \rightarrow \infty} \left| \frac{\#v_Y(m\bar{\mathcal{L}})}{m^d} \log p - \frac{\hat{h}^0(X, m\bar{\mathcal{L}})}{m^d} \right| \leq \frac{c}{\log p}.$$

Furthermore, we can take

$$c(\bar{\mathcal{L}}) = 2e_0 \frac{\text{vol}(\mathcal{L}_{\mathbb{Q}}) \bar{\mathcal{L}} \cdot \bar{\mathcal{A}}^{d-1}}{\text{vol}(\mathcal{A}_{\mathbb{Q}}) (d-1)!}.$$

Here e_0 is the number of connected components of $X_{\bar{\mathbb{Q}}}$, and $\bar{\mathcal{A}}$ is any ample line bundle on X .

We first see how to induce the following result of Chen [Che08a].

THEOREM 2.7. *The limit $\lim_{m \rightarrow \infty} \hat{h}^0(X, m\bar{\mathcal{L}})/(m^d/d!)$ exists for any $\bar{\mathcal{L}} \in \widehat{\text{Pic}}(X)$.*

Proof. The case that $\bar{\mathcal{L}}$ is not big is easy. Assume that $\bar{\mathcal{L}}$ is big. The key is that $\#v_Y(m\bar{\mathcal{L}})/m^d$ is convergent by Proposition 2.5. Then Theorem 2.6 implies

$$\limsup_{m \rightarrow \infty} \frac{\hat{h}^0(X, m\bar{\mathcal{L}})}{m^d} - \liminf_{m \rightarrow \infty} \frac{\hat{h}^0(X, m\bar{\mathcal{L}})}{m^d} \leq \frac{2c}{\log p}.$$

Let $p \rightarrow \infty$, we get $\limsup = \liminf$ and thus the convergence. \square

It is also immediate to show Theorem A. In fact, since both limits exist, the result in Theorem 2.6 simplifies as

$$\left| \text{vol}(\Delta_Y(\bar{\mathcal{L}})) \log p - \frac{1}{d!} \text{vol}(\bar{\mathcal{L}}) \right| \leq \frac{c}{\log p}.$$

Hence Theorem A is true under the assumption of Theorem 2.6.

2.3 Some preliminary results

We show some simple results which will be needed in the proof of Theorem 2.6 in next subsection.

Let K be a number field and O_K be the ring of integers of K . Let X be an arithmetic variety over O_K . In another word, the structure morphism $X \rightarrow \text{Spec}(\mathbb{Z})$ factors through $X \rightarrow \text{Spec}(O_K)$. For any non-zero ideal I of O_K , consider the reduction modulo I map

$$r_I : H^0(X, \mathcal{L}) \rightarrow H^0(X_{O_K/I}, \mathcal{L}_{O_K/I}).$$

We want to bound the order of $r_I(\widehat{H}^0(X, \bar{\mathcal{L}}))$. Denote by Z_I the zero locus of I in X . Recall that the notation $\bar{\mathcal{L}}(f + V)$ is explained at the end of the introduction. The following result is a bridge from the arithmetic case to the geometric case.

PROPOSITION 2.8. *For any $\bar{\mathcal{L}} \in \widehat{\text{Pic}}(X)$,*

$$\begin{aligned} \log \#r_I(\widehat{H}^0(X, \bar{\mathcal{L}})) &\leq \hat{h}^0(X, \bar{\mathcal{L}}(\log 2)) - \hat{h}^0(X, \bar{\mathcal{L}}(-Z_I)) \\ \log \#r_I(\widehat{H}^0(X, \bar{\mathcal{L}})) &\geq \hat{h}^0(X, \bar{\mathcal{L}}) - \hat{h}^0(X, \bar{\mathcal{L}}(\log 2 - Z_I)). \end{aligned}$$

Proof. For each $t \in H^0(X_{O_K/I}, \mathcal{L}_{O_K/I})$, fix one lifting $s_0 \in r_I^{-1}(t) \cap \widehat{H}^0(X, \overline{\mathcal{L}})$ if it exists. For any other $s \in r_I^{-1}(t) \cap \widehat{H}^0(X, \overline{\mathcal{L}})$, we have $s_{Z_I}^{-1} \otimes (s - s_0)$ regular everywhere and $\|s - s_0\|_{\text{sup}} \leq 2$. Thus we have an element

$$s_{Z_I}^{-1} \otimes (s - s_0) \in \widehat{H}^0(X, \overline{\mathcal{L}}(\log 2 - Z_I)).$$

It follows that

$$\#(r_I^{-1}(t) \cap \widehat{H}^0(X, \overline{\mathcal{L}})) \leq \#\widehat{H}^0(X, \overline{\mathcal{L}}(\log 2 - Z_I)).$$

It induces the inequality

$$\#r_I(\widehat{H}^0(X, \overline{\mathcal{L}})) \geq \frac{\#\widehat{H}^0(X, \overline{\mathcal{L}})}{\#\widehat{H}^0(X, \overline{\mathcal{L}}(\log 2 - Z_I))}.$$

Now we seek the upper bound of $\#r_I(\widehat{H}^0(X, \overline{\mathcal{L}}))$. Consider the set

$$S = \widehat{H}^0(X, \overline{\mathcal{L}}) + s_{Z_I} \otimes \widehat{H}^0(X, \overline{\mathcal{L}}(-Z_I)).$$

Apparently $r_I(S) = r_I(\widehat{H}^0(X, \overline{\mathcal{L}}))$. We further have $S \subset \widehat{H}^0(X, \overline{\mathcal{L}}(\log 2))$ since any $s \in S$ satisfies $\|s\|_{\text{sup}} \leq 1 + 1 = 2$.

For each $t \in r_I(S)$, there is a lifting s_0 of t in $\widehat{H}^0(X, \overline{\mathcal{L}})$, so

$$s_0 + s_{Z_I} \otimes \widehat{H}^0(X, \overline{\mathcal{L}}(-Z_I)) \subset r_I^{-1}(t) \cap S.$$

Hence, $\#r_I^{-1}(t) \cap S \geq \#\widehat{H}^0(X, \overline{\mathcal{L}}(-Z_I))$. It follows that

$$\#r_I(\widehat{H}^0(X, \overline{\mathcal{L}})) = \#r_I(S) \leq \frac{\#S}{\#\widehat{H}^0(X, \overline{\mathcal{L}}(-Z_I))} \leq \frac{\#\widehat{H}^0(X, \overline{\mathcal{L}}(\log 2))}{\#\widehat{H}^0(X, \overline{\mathcal{L}}(-Z_I))}. \quad \square$$

LEMMA 2.9. For any $\overline{\mathcal{L}} \in \widehat{\text{Pic}}(X)$,

$$0 \leq \hat{h}^0(X, \overline{\mathcal{L}}) - \hat{h}^0(X, \overline{\mathcal{L}}(-\alpha)) \leq (\alpha + \log 3) \text{rank}_{\mathbb{Z}} H^0(X, \mathcal{L})$$

for any $\alpha \in \mathbb{R}_+$.

Proof. The first inequality is trivial, and we only need to show the second one. Take $I = (n)$ for an integer $n \geq 2$ in Proposition 2.8, we get

$$\hat{h}^0(X, \overline{\mathcal{L}}) - \hat{h}^0(X, \overline{\mathcal{L}}(\log 2 - Z_n)) \leq \log \#r_n(\widehat{H}^0(X, \overline{\mathcal{L}})).$$

The right-hand side has an easy bound

$$\log \#r_n(\widehat{H}^0(X, \overline{\mathcal{L}})) \leq \log \#(H^0(X, \mathcal{L})/nH^0(X, \mathcal{L})) = \text{rank}_{\mathbb{Z}} H^0(X, \mathcal{L}) \log n.$$

It is easy to see that $\overline{\mathcal{O}}(Z_n) \cong \overline{\mathcal{O}}(\log n)$, so the above gives

$$\hat{h}^0(X, \overline{\mathcal{L}}) - \hat{h}^0\left(X, \overline{\mathcal{L}}\left(-\log \frac{n}{2}\right)\right) \leq \text{rank}_{\mathbb{Z}} H^0(X, \mathcal{L}) \log n.$$

For general $\alpha > 0$, taking $n = [2e^\alpha] + 1$, then the above gives

$$\hat{h}^0(X, \overline{\mathcal{L}}) - \hat{h}^0(X, \overline{\mathcal{L}}(-\alpha)) \leq \hat{h}^0(X, \overline{\mathcal{L}}) - \hat{h}^0\left(X, \overline{\mathcal{L}}\left(-\log \frac{n}{2}\right)\right) \leq \text{rank}_{\mathbb{Z}} H^0(X, \mathcal{L}) \log n.$$

It proves the result since

$$\log n \leq \log(2e^\alpha + 1) \leq \alpha + \log 3. \quad \square$$

2.4 Comparison of the volumes

In this subsection, we will prove Theorem 2.6. Resume the notation in § 2.2. That is, K is the number field such that $X \rightarrow \text{Spec}(O_K)$ is geometrically connected. And $Y_1 = X_\wp$ for some prime \wp of O_K lying over a prime number p .

Recall that $\nu = (\nu_1, \dots, \nu_d)$ is the valuation on X with respect to the flag

$$X \supset Y_1 \supset Y_2 \supset \dots \supset Y_d.$$

Then the flag

$$Y_1 \supset Y_2 \supset \dots \supset Y_d$$

on the ambient variety Y_1 induces a valuation map $\nu^\circ = (\nu_2, \dots, \nu_d)$ of dimension $d - 1$ in the geometric case. They are compatible in the sense that

$$\nu(s) = (\nu_1(s), \nu^\circ((s_{Y_1}^{\otimes(-\nu_1(s))} s)|_{Y_1})),$$

where s_{Y_1} is the section of $\mathcal{O}(Y_1)$ defining Y_1 . The notation such as $\overline{\mathcal{L}}(\log \beta - Y_1)$ in the proposition below is explained at the end of the introduction.

PROPOSITION 2.10. *For any $\mathcal{L} \in \widehat{\text{Pic}}(X)$:*

- (1) $\#\nu^\circ(\widehat{H}^0(X, \overline{\mathcal{L}})|_{Y_1}) \log p \leq \hat{h}^0(X, \overline{\mathcal{L}}(\log(2\beta))) - \hat{h}^0(X, \overline{\mathcal{L}}(\log \beta - Y_1));$
- (2) $\#\nu^\circ(\widehat{H}^0(X, \overline{\mathcal{L}})|_{Y_1}) \log p \geq \hat{h}^0(X, \overline{\mathcal{L}}(-\log \beta)) - \hat{h}^0(X, \overline{\mathcal{L}}(-\log(\beta/2) - Y_1)).$

Here we denote $\beta = p \dim_{\mathbb{F}_\wp} H^0(X_{\mathbb{F}_\wp}, \mathcal{L}_{\mathbb{F}_\wp})$.

Proof. We first prove part (1). The key point is to pass to the \mathbb{F}_\wp -subspace $\langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle$ of $H^0(Y_1, \mathcal{L}|_{Y_1})$ generated by $\widehat{H}^0(\overline{\mathcal{L}})|_{Y_1}$. For vector spaces, we can apply the last result of [LM08, Lemma 1.3] to have the order of its valuation image. It is easy to see it also works for non-algebraically closed field as long as the residue field of Y_d agrees with the field of definition of the ambient variety.

We first use the trivial bound

$$\#\nu^\circ(\widehat{H}^0(X, \overline{\mathcal{L}})|_{Y_1}) \leq \#\nu^\circ \langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle.$$

Then [LM08, Lemma 1.3] implies

$$\#\nu^\circ \langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle = \dim_{\mathbb{F}_\wp} \langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle.$$

Thus

$$\#\nu^\circ(\widehat{H}^0(X, \overline{\mathcal{L}})|_{Y_1}) \leq \dim_{\mathbb{F}_\wp} \langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle = \log_p \# \langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle.$$

Now we seek an upper bound on the order of $\langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle$. The key is to put this space into $\widehat{H}^0(\overline{\mathcal{L}}')|_{Y_1}$ for some ‘bigger’ Hermitian line bundle $\overline{\mathcal{L}}'$.

Choose a basis $\{t\}$ of $\langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle$ lying in $\widehat{H}^0(\overline{\mathcal{L}})|_{Y_1}$, and fix a lifting $\tilde{t} \in \widehat{H}^0(\overline{\mathcal{L}})$ for each $t \in \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1}$. Since the residue field \mathbb{F}_\wp is equal to \mathbb{F}_p , the set

$$S = \left\{ \sum_t a_t \tilde{t} : a_t = 0, 1, \dots, p - 1 \right\}$$

maps surjectively to $\langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle$ under the reduction r_\wp . For any such element $\sum_t a_t \tilde{t}$, the norm

$$\left\| \sum_t a_t \tilde{t} \right\|_{\text{sup}} \leq p \sum_t 1 = p \dim_{\mathbb{F}_\wp} \langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle \leq p h^0(\mathcal{L}|_{Y_1}) = \beta.$$

It follows that $S \subset \widehat{H}^0(\overline{\mathcal{L}}(\log \beta))$, and thus their reductions have the relation

$$\langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle \subset \widehat{H}^0(\overline{\mathcal{L}}(\log \beta))|_{Y_1}.$$

By this we get a bound

$$\#\langle \widehat{H}^0(\overline{\mathcal{L}})|_{Y_1} \rangle \leq \#\widehat{H}^0(\overline{\mathcal{L}}(\log \beta))|_{Y_1}.$$

By Proposition 2.8, we obtain

$$\log \#\widehat{H}^0(\overline{\mathcal{L}}(\log \beta))|_{Y_1} \leq \hat{h}^0(\overline{\mathcal{L}}(\log(2\beta))) - \hat{h}^0(\overline{\mathcal{L}}(\log \beta - Y_1)).$$

Putting the inequalities together, we achieve part (1) of Proposition 2.10.

Now we prove part (2) of Proposition 2.10. Similar to the above, we construct a set

$$T = \left\{ \sum_t a_t \tilde{t} : a_t = 0, 1, \dots, p-1 \right\}.$$

Here $\{\tilde{t}\}$ is a basis of $\langle \widehat{H}^0(\overline{\mathcal{L}}(-\log \beta))|_{Y_1} \rangle$ lying in $\widehat{H}^0(\overline{\mathcal{L}}(-\log \beta))|_{Y_1}$, and $\tilde{t} \in \widehat{H}^0(\overline{\mathcal{L}}(-\log \beta))$ is a fixed lifting for each $t \in \widehat{H}^0(\overline{\mathcal{L}}(-\log \beta))|_{Y_1}$. By the same reason, we see that

$$T \subset \widehat{H}^0(X, \overline{\mathcal{L}})$$

and

$$\widehat{H}^0(X, \overline{\mathcal{L}})|_{Y_1} \supset T|_{Y_1} = \langle \widehat{H}^0(\overline{\mathcal{L}}(-\log \beta))|_{Y_1} \rangle.$$

Thus we have

$$\#\nu^\circ(\widehat{H}^0(X, \overline{\mathcal{L}})|_{Y_1}) \geq \#\nu^\circ\langle \widehat{H}^0(\overline{\mathcal{L}}(-\log \beta))|_{Y_1} \rangle.$$

By [LM08, Lemma 1.3] again, we get

$$\begin{aligned} \#\nu^\circ\langle \widehat{H}^0(\overline{\mathcal{L}}(-\log \beta))|_{Y_1} \rangle &= \dim_{\mathbb{F}_p}\langle \widehat{H}^0(\overline{\mathcal{L}}(-\log \beta))|_{Y_1} \rangle \\ &= \log_p \#\langle \widehat{H}^0(\overline{\mathcal{L}}(-\log \beta))|_{Y_1} \rangle \geq \log_p \#\widehat{H}^0(\overline{\mathcal{L}}(-\log \beta))|_{Y_1}. \end{aligned}$$

Apply Proposition 2.8 again. We have

$$\begin{aligned} \#\nu^\circ(\widehat{H}^0(X, \overline{\mathcal{L}})|_{Y_1}) \log p &\geq \log \#\widehat{H}^0(\overline{\mathcal{L}}(-\log \beta))|_{Y_1} \\ &\geq \hat{h}^0(X, \overline{\mathcal{L}}(-\log \beta)) - \hat{h}^0\left(X, \overline{\mathcal{L}}\left(-\log \frac{\beta}{2} - Y_1\right)\right). \end{aligned}$$

This proves part (2) of Proposition 2.10. □

COROLLARY 2.11.

$$\begin{aligned} \#v(\overline{\mathcal{L}}) \log p &\leq \hat{h}^0(X, \overline{\mathcal{L}}(\log(2\beta))) + \sum_{k \geq 1} (\hat{h}^0(X, \overline{\mathcal{L}}(\log(2\beta) - kY_1)) - \hat{h}^0(X, \overline{\mathcal{L}}(\log \beta - kY_1))), \\ \#v(\overline{\mathcal{L}}) \log p &\geq \hat{h}^0(X, \overline{\mathcal{L}}(-\log \beta)) - \sum_{k \geq 1} \left(\hat{h}^0\left(X, \overline{\mathcal{L}}\left(-\log \frac{\beta}{2} - kY_1\right)\right) - \hat{h}^0(X, \overline{\mathcal{L}}(-\log \beta - kY_1)) \right). \end{aligned}$$

Proof. Denote

$$M_k = \{s_{Y_1}^{-k} \otimes s : s \in \widehat{H}^0(X, \overline{\mathcal{L}}), \nu_1(s) \geq k\}.$$

Then the compatibility between $\nu^\circ = (\nu_2, \dots, \nu_d)$ and $\nu = (\nu_1, \nu_2, \dots, \nu_d)$ gives

$$\#v(\overline{\mathcal{L}}) = \sum_{k \geq 0} \#\nu^\circ(M_k|_{Y_1}).$$

Since the metric of s_{Y_1} is identically 1 in $\overline{\mathcal{O}}(Y_1)$, we have an interpretation

$$M_k = \widehat{H}^0(X, \overline{\mathcal{L}}(-kY_1)).$$

Therefore

$$\#v(\overline{\mathcal{L}}) = \sum_{k \geq 0} \#v^\circ(\widehat{H}^0(X, \overline{\mathcal{L}}(-kY_1))|_{Y_1}).$$

Apply the above result to each $\overline{\mathcal{L}}(-kY_1)$ and rearrange the summations. □

Remark. The summations in both inequalities in the proposition have only finitely many non-zero terms, as we will see below.

Now we can prove Theorem 2.6 which asserts

$$\limsup_{m \rightarrow \infty} \left| \frac{\#v_{Y_1}(m\overline{\mathcal{L}})}{m^d} \log p - \frac{\widehat{h}^0(X, m\overline{\mathcal{L}})}{m^d} \right| \leq \frac{c}{\log p}.$$

Proof of Theorem 2.6. The above corollary gives

$$\begin{aligned} \#v(m\overline{\mathcal{L}}) \log p &\leq \widehat{h}^0(m\overline{\mathcal{L}} + \overline{\mathcal{O}}(\log(2\beta_m))) \\ &+ \sum_{k \geq 1} (\widehat{h}^0(m\overline{\mathcal{L}} + \overline{\mathcal{O}}(\log(2\beta_m) - kY_1)) - \widehat{h}^0(m\overline{\mathcal{L}} + \overline{\mathcal{O}}(\log \beta_m - kY_1))). \end{aligned}$$

Here $\beta_m = ph^0(m\mathcal{L}|_{Y_1})$. By Lemma 2.9, we get

$$\widehat{h}^0(m\overline{\mathcal{L}} + \overline{\mathcal{O}}(\log(2\beta_m))) \leq \widehat{h}^0(m\overline{\mathcal{L}}) + \log(6\beta_m) h^0(m\mathcal{L}_\mathbb{Q})$$

and

$$\widehat{h}^0(m\overline{\mathcal{L}} + \overline{\mathcal{O}}(\log(2\beta_m) - kY_1)) - \widehat{h}^0(m\overline{\mathcal{L}} + \overline{\mathcal{O}}(\log \beta_m - kY_1)) \leq (\log 6) h^0(m\mathcal{L}_\mathbb{Q}).$$

Let S be the set of $k \geq 1$ such that

$$\widehat{h}^0(m\overline{\mathcal{L}} + \overline{\mathcal{O}}(\log(2\beta_m) - kY_1)) \neq 0.$$

Then we have

$$\#v(m\overline{\mathcal{L}}) \log p \leq \widehat{h}^0(m\overline{\mathcal{L}}) + h^0(m\mathcal{L}_\mathbb{Q}) \log(6\beta_m) + (\log 6) h^0(m\mathcal{L}_\mathbb{Q})(\#S).$$

Next we bound $\#S$ which gives the main error term.

Fix an ample line bundle $\overline{\mathcal{A}}$. We are going to give an upper bound of S in terms of intersection numbers with $\overline{\mathcal{A}}$. Assume that $k \in S$, so $m\overline{\mathcal{L}} + \overline{\mathcal{O}}(\log(2\beta_m) - kY_1)$ is effective. We must have

$$(m\overline{\mathcal{L}} + \overline{\mathcal{O}}(\log(2\beta_m) - kY_1)) \cdot \overline{\mathcal{A}}^{d-1} \geq 0.$$

Equivalently,

$$m\overline{\mathcal{L}} \cdot \overline{\mathcal{A}}^{d-1} + \log(2\beta_m) \deg(\mathcal{A}_\mathbb{Q}) - kY_1 \cdot \overline{\mathcal{A}}^{d-1} \geq 0.$$

Note that

$$Y_1 \cdot \overline{\mathcal{A}}^{d-1} = \frac{1}{[K : \mathbb{Q}]} \deg(\mathcal{A}_\mathbb{Q}) \log p.$$

We get

$$k \leq [K : \mathbb{Q}] \frac{\overline{\mathcal{L}} \cdot \overline{\mathcal{A}}^{d-1}}{\deg(\mathcal{A}_\mathbb{Q})} \frac{m}{\log p} + [K : \mathbb{Q}] \frac{\log(2\beta_m)}{\log p}.$$

The order of S has the same bound.

Therefore,

$$\begin{aligned} & \#v(m\bar{\mathcal{L}}) \log p - \hat{h}^0(m\bar{\mathcal{L}}) \\ & \leq h^0(m\mathcal{L}_{\mathbb{Q}}) \log(6\beta_m) + (\log 6) h^0(m\mathcal{L}_{\mathbb{Q}})[K : \mathbb{Q}] \left(\frac{\bar{\mathcal{L}} \cdot \bar{\mathcal{A}}^{d-1}}{\deg(\mathcal{A}_{\mathbb{Q}})} \frac{m}{\log p} + \frac{\log(2\beta_m)}{\log p} \right) \\ & = (\log 6)[K : \mathbb{Q}] \frac{\text{vol}(\mathcal{L}_{\mathbb{Q}}) \bar{\mathcal{L}} \cdot \bar{\mathcal{A}}^{d-1}}{(d-1)! \deg(\mathcal{A}_{\mathbb{Q}}) \log p} m^d + O(m^{d-1} \log m). \end{aligned}$$

Here we have used the fact that $\beta_m = ph^0(m\mathcal{L}|_{Y_1})$ is at most a Hilbert polynomial of degree $d - 1$. It gives one direction of what we need to prove. Similarly, we can obtain the other direction. \square

3. Consequences

In this section, X is any arithmetic variety. To have good flags to apply Theorem A, we take the normalization \tilde{X} of the generic resolution of X . Consider the pull-back of Hermitian line bundles. The volume does not change by Moriwaki's result quoted in Theorem 2.3.

3.1 Log-concavity

We also show the log-concavity of volume functions. The key is still the Brunn–Minkowski theorem which asserts that

$$\text{vol}(S_1 + S_2)^{1/d} \geq \text{vol}(S_1)^{1/d} + \text{vol}(S_2)^{1/d},$$

for any two compact subsets S_1 and S_2 of \mathbb{R}^d . Unlike in [LM08], we do not explore any universal Okounkov body since the space of numerical classes in our setting is too big.

THEOREM B. *For any two effective line bundles $\bar{\mathcal{L}}_1, \bar{\mathcal{L}}_2$, we have*

$$\text{vol}(\bar{\mathcal{L}}_1 + \bar{\mathcal{L}}_2)^{1/d} \geq \text{vol}(\bar{\mathcal{L}}_1)^{1/d} + \text{vol}(\bar{\mathcal{L}}_2)^{1/d}.$$

Proof. It is easy to see that the inequality is true if one of $\text{vol}(\bar{\mathcal{L}}_1)$ and $\text{vol}(\bar{\mathcal{L}}_2)$ is zero by the effectivity property. So we can assume that $\bar{\mathcal{L}}_1$ and $\bar{\mathcal{L}}_2$ are big. We can further assume that X is normal with smooth generic fibre by Moriwaki's theorem of pull-back quoted in Theorem 2.3.

Take a flag Y . on X . It is easy to have

$$\Lambda_Y(\bar{\mathcal{L}}_1) + \Lambda_Y(\bar{\mathcal{L}}_2) \subset \Lambda_Y(\bar{\mathcal{L}}_1 + \bar{\mathcal{L}}_2).$$

Taking closures in \mathbb{R}^d , we get

$$\Delta_Y(\bar{\mathcal{L}}_1) + \Delta_Y(\bar{\mathcal{L}}_2) \subset \Delta_Y(\bar{\mathcal{L}}_1 + \bar{\mathcal{L}}_2).$$

The Brunn–Minkowski theorem gives

$$\text{vol}(\Delta_Y(\bar{\mathcal{L}}_1 + \bar{\mathcal{L}}_2))^{1/n} \geq \text{vol}(\Delta_Y(\bar{\mathcal{L}}_1))^{1/n} + \text{vol}(\Delta_Y(\bar{\mathcal{L}}_2))^{1/n}.$$

It implies the result by taking $\text{char}(Y.) \rightarrow \infty$. \square

COROLLARY 3.1. *For any two ample line bundles $\bar{\mathcal{L}}_1, \bar{\mathcal{L}}_2$, we have*

$$\begin{aligned} \bar{\mathcal{L}}_1^{d-1} \cdot \bar{\mathcal{L}}_2 & \geq (\bar{\mathcal{L}}_1^d)^{(d-1)/d} (\bar{\mathcal{L}}_2^d)^{1/d}, \\ (\bar{\mathcal{L}}_1^{d-1} \cdot \bar{\mathcal{L}}_2)^2 & \geq (\bar{\mathcal{L}}_1^d) (\bar{\mathcal{L}}_1^{d-2} \cdot \bar{\mathcal{L}}_2^2). \end{aligned}$$

Proof. Note that volumes for ample line bundles are equal to top self-intersection numbers. We first show the first inequality. Let t be any positive rational number. Then the above theorem gives

$$((\bar{\mathcal{L}}_1 + t\bar{\mathcal{L}}_2)^d)^{1/d} \geq (\bar{\mathcal{L}}_1^d)^{1/d} + t(\bar{\mathcal{L}}_2^d)^{1/d}.$$

In other words,

$$(\bar{\mathcal{L}}_1 + t\bar{\mathcal{L}}_2)^d - ((\bar{\mathcal{L}}_1^d)^{1/d} + t(\bar{\mathcal{L}}_2^d)^{1/d})^d \geq 0.$$

The left-hand side is a polynomial in t whose constant term is zero. The coefficient of degree one must be non-negative by considering $t \rightarrow 0$. It gives the result exactly.

Now we prove the second relation. Use the same trick on the first inequality. We get

$$\bar{\mathcal{L}}_1^{d-1} \cdot (\bar{\mathcal{L}}_1 + t\bar{\mathcal{L}}_2) \geq (\bar{\mathcal{L}}_1^d)^{(d-1)/d} ((\bar{\mathcal{L}}_1 + t\bar{\mathcal{L}}_2)^d)^{1/d}.$$

It becomes

$$(\bar{\mathcal{L}}_1^d + t\bar{\mathcal{L}}_1^{d-1} \cdot \bar{\mathcal{L}}_2)^d \geq (\bar{\mathcal{L}}_1^d)^{d-1} (\bar{\mathcal{L}}_1 + t\bar{\mathcal{L}}_2)^d.$$

Let $t \rightarrow 0$. The terms of degree less than or equal to 2 give

$$\begin{aligned} & (\bar{\mathcal{L}}_1^d)^d + dt(\bar{\mathcal{L}}_1^d)^{d-1}(\bar{\mathcal{L}}_1^{d-1} \cdot \bar{\mathcal{L}}_2) + \frac{d(d-1)}{2}t^2(\bar{\mathcal{L}}_1^d)^{d-2}(\bar{\mathcal{L}}_1^{d-1} \cdot \bar{\mathcal{L}}_2)^2 \\ & \geq (\bar{\mathcal{L}}_1^d)^{d-1} \left(\bar{\mathcal{L}}_1^d + dt\bar{\mathcal{L}}_1^{d-1} \cdot \bar{\mathcal{L}}_2 + \frac{d(d-1)}{2}t^2\bar{\mathcal{L}}_1^{d-2} \cdot \bar{\mathcal{L}}_2^2 \right). \end{aligned}$$

It turns out the terms of degree less than or equal to 1 are canceled, and the degree two terms give

$$(\bar{\mathcal{L}}_1^d)^{d-2}(\bar{\mathcal{L}}_1^{d-1} \cdot \bar{\mathcal{L}}_2)^2 \geq (\bar{\mathcal{L}}_1^d)^{d-1}(\bar{\mathcal{L}}_1^{d-2} \cdot \bar{\mathcal{L}}_2^2).$$

This gives the second inequality. □

Remark. The above method can also induce some inequalities on big line bundles. For example, if we allow $\bar{\mathcal{L}}_2$ to be big in the proof of the first inequality, then $\bar{\mathcal{L}}_1 + t\bar{\mathcal{L}}_2$ is still ample for t small enough. Then we have

$$\text{vol}(\bar{\mathcal{L}}_2) \leq \frac{(\bar{\mathcal{L}}_1^{d-1} \cdot \bar{\mathcal{L}}_2)^d}{(\bar{\mathcal{L}}_1^d)^{d-1}}$$

for any ample line bundle $\bar{\mathcal{L}}_1$.

It turns out that the second inequality above implies the arithmetic Hodge index theorem in codimension one proved by Moriwaki [Mor96].

COROLLARY 3.2. *Let $\bar{\mathcal{A}} \in \widehat{\text{Amp}}(X)$, $\bar{\mathcal{L}} \in \widehat{\text{Pic}}(X)$. If $\bar{\mathcal{A}}^{d-1} \cdot \bar{\mathcal{L}} = 0$, then $\bar{\mathcal{A}}^{d-2} \cdot \bar{\mathcal{L}}^2 \leq 0$.*

Proof. By replacing $\bar{\mathcal{A}}$ by its positive multiples, we can assume that $\bar{\mathcal{B}} = \bar{\mathcal{A}} + \bar{\mathcal{L}}$ is ample. Then $\bar{\mathcal{L}} = \bar{\mathcal{B}} - \bar{\mathcal{A}}$. We need to show that

$$\bar{\mathcal{A}}^{d-2} \cdot \bar{\mathcal{L}}^2 = \bar{\mathcal{A}}^{d-2} \cdot \bar{\mathcal{B}}^2 - 2\bar{\mathcal{A}}^{d-1}\bar{\mathcal{B}} + \bar{\mathcal{A}}^d \leq 0. \tag{2}$$

The above lemma gives

$$(\bar{\mathcal{A}}^{d-1} \cdot \bar{\mathcal{B}})^2 \geq (\bar{\mathcal{A}}^d)(\bar{\mathcal{A}}^{d-2} \cdot \bar{\mathcal{B}}^2). \tag{3}$$

The condition $\bar{\mathcal{A}}^{d-1} \cdot \bar{\mathcal{L}} = 0$ becomes $\bar{\mathcal{A}}^{d-1} \cdot \bar{\mathcal{B}} = \bar{\mathcal{A}}^d$. However, it is easy to see that (2) and (3) are equivalent under this condition. □

3.2 Fujita approximation

For any $\overline{\mathcal{L}} \in \widehat{\text{Pic}}(X)$, denote

$$V_{k,n}(\overline{\mathcal{L}}) = \{s_1 \otimes s_2 \otimes \cdots \otimes s_k : s_1, s_2, \dots, s_k \in \widehat{H}^0(X, n\overline{\mathcal{L}})\}.$$

It is a subset of $\widehat{H}^0(X, kn\overline{\mathcal{L}})$. Then a consequence of [LM08, Proposition 3.1] is the following result.

THEOREM 3.3. *Assume that $\overline{\mathcal{L}}$ is big and X is normal with smooth generic fibre, and let Y be a flag on X . Then for any $\epsilon > 0$, there exists $n_0 > 0$ such that*

$$\lim_{k \rightarrow \infty} \frac{\#v_Y(V_{k,n}(\overline{\mathcal{L}}))}{(nk)^d} \geq \text{vol}(\Delta_Y(\overline{\mathcal{L}})) - \epsilon \quad \forall n > n_0.$$

This theorem serves as an analogue of [LM08, Theorem 3.3] in the proof of the arithmetic Fujita approximation. It is not as neat as the direct arithmetic analogue:

$$\lim_{k \rightarrow \infty} \frac{\log \#V_{k,n}(\overline{\mathcal{L}})}{(nk)^d/d!} \geq \text{vol}(\overline{\mathcal{L}}) - \epsilon \quad \forall n > n_0.$$

However, we do not know whether this analogue is true.

For the arithmetic Fujita approximation theorem, we need an extra argument to take care of the archimedean part.

THEOREM 3.4. *Let L be an ample line bundle on a projective complex manifold M , and let $\|\cdot\|$ be any smooth Hermitian metric on L . Then the following results hold.*

- (a) *There is a canonical positive continuous metric $\|\cdot\|'$ on L such that $\|\cdot\|' \geq \|\cdot\|$ pointwise and $\|\cdot\|_{\text{sup}} = \|\cdot\|'_{\text{sup}}$ as norms on $H^0(M, mL)$ for any positive integer m .*
- (b) *The above metric $\|\cdot\|'$ can be uniformly approximated by a sequence $\{\|\cdot\|_j\}_j$ of positive smooth Hermitian metrics $\|\cdot\|_j \geq \|\cdot\|'$ on L in the sense that $\lim_{j \rightarrow \infty} \|\cdot\|_j/\|\cdot\|' = 1$ uniformly on M .*

Proof. The results are standard in complex analysis. We will take the metric $\|\cdot\|'$ to be the equilibrium metric of $\|\cdot\|$ defined as the following envelope:

$$\|s(z)\|' := \inf\{\|s(z)\|_+ : \|\cdot\|_+ \text{ positive singular metric on } L, \|\cdot\|_+ \geq \|\cdot\|\}.$$

By definition, it is positive and satisfies those two norm relations with $\|\cdot\|$ in Theorem 3.4(a). As for the continuity, see [Ber07, Theorem 2.3 (1)], where the metric is proved to be $C^{1,1}$.

Now we consider Theorem 3.4(b). In fact, any positive continuous metric can be approximated uniformly by positive smooth ones. It is essentially due to the technique of Demailly [Dem92]. See also [BK07, Theorem 1] for a shorter proof. Note that monotone convergence to a continuous function is uniform over any compact space. Once the convergence is uniform, it is easy to obtain $\|\cdot\|_j \geq \|\cdot\|'$ by scalar perturbations on the metrics. For more general results, we refer to [BB08], especially [BB08, Proposition 1.13]. \square

Now we are prepared to prove our arithmetic Fujita approximation theorem.

THEOREM C. *Let $\overline{\mathcal{L}}$ be a big line bundle over X . Then for any $\epsilon > 0$, there exist an integer $n > 0$, a birational morphism $\pi : X' \rightarrow X$ from another arithmetic variety X' to X , and an isomorphism*

$$n \pi^* \overline{\mathcal{L}} = \overline{\mathcal{A}} + \overline{\mathcal{E}}$$

for an effective line bundle $\bar{\mathcal{E}}$ on X' and an ample line bundle $\bar{\mathcal{A}}$ on X' satisfying

$$\frac{1}{n^d} \text{vol}(\bar{\mathcal{A}}) > \text{vol}(\bar{\mathcal{L}}) - \epsilon.$$

Proof. We can assume that X is normal with smooth generic fibre by Moriwaki's pull-back result.

For any positive integer n , let $\pi_n : X_n \rightarrow X$ be the blowing-up of the base ideal generated by $\widehat{H}^0(X, n\bar{\mathcal{L}})$. Denote by \mathcal{E}_n the line bundle associated to the exceptional divisor, with $e \in H^0(X_n, \mathcal{E}_n)$ a section defining the exceptional divisor. Denote by $\mathcal{A}_n = n \pi^* \mathcal{L} - \mathcal{E}_n$, which has regular sections $\lambda(s) = \pi^* s \otimes e^{\otimes(-1)}$ for any $s \in \widehat{H}^0(X, n\bar{\mathcal{L}})$. Furthermore, $\lambda(\widehat{H}^0(X, n\bar{\mathcal{L}}))$ is base-point free.

For any $s \in \widehat{H}^0(X, n\bar{\mathcal{L}})$, we have $\|\pi^* s\|_{\text{sup}} = \|s\|_{\text{sup}} \leq 1$ under the pull-back metric on $\pi^* \mathcal{L}$. We claim that there exist metrics on \mathcal{E}_n and \mathcal{A}_n such that under these metrics $n \pi^* \mathcal{L} = \mathcal{A}_n + \mathcal{E}_n$ is isometric, e is effective, and $\lambda(s)$ is effective for all $s \in \widehat{H}^0(X, n\bar{\mathcal{L}})$.

To get such metrics on \mathcal{E}_n and \mathcal{A}_n , start from any metric $\|\cdot\|_0$ on \mathcal{E}_n . It suffices to find a smooth function $f : X(\mathbb{C}) \rightarrow \mathbb{R}_+$ such that $f\|e\|_0 \leq 1$ and $\|s\|/(f\|e\|_0) \leq 1$ for all $s \in \widehat{H}^0(X, n\bar{\mathcal{L}})$. Equivalently, we need

$$\sup_s \frac{\|s\|}{\|e\|_0} \leq f \leq \frac{1}{\|e\|_0}.$$

This is possible because the left-hand side is always less than or equal to the right-hand side, and the left-hand side is actually bounded everywhere.

Endowed with the above metrics, we get line Hermitian bundles $\bar{\mathcal{E}}_n$ and $\bar{\mathcal{A}}_n$ such that

$$n \pi^* \bar{\mathcal{L}} = \bar{\mathcal{A}}_n + \bar{\mathcal{E}}_n.$$

Here $\bar{\mathcal{E}}_n$ is effective and $\bar{\mathcal{A}}_n$ is base-point free since $\lambda(\widehat{H}^0(X, n\bar{\mathcal{L}})) \subset \widehat{H}^0(X, \bar{\mathcal{A}}_n)$. Furthermore, each element of $V_{k,n}(\bar{\mathcal{L}})$ gives an effective section in $\widehat{H}^0(X, k\bar{\mathcal{A}}_n)$ by the same way. We claim that these sections give

$$\lim_{n \rightarrow \infty} \frac{1}{n^d} \text{vol}(\bar{\mathcal{A}}_n) = \text{vol}(\bar{\mathcal{L}}). \quad (4)$$

Once it is true, we have obtained decompositions satisfying all requirements of the theorem except that $\bar{\mathcal{A}}_n$ may not be ample, though it is base-point free. We will make some adjustment to get ampleness, but let us first verify (4). It is a consequence of Theorems 2.6 and 3.3.

In fact, it suffices to show that

$$\liminf_{n \rightarrow \infty} \frac{1}{n^d} \text{vol}(\bar{\mathcal{A}}_n) \geq \text{vol}(\bar{\mathcal{L}}).$$

Replacing $\bar{\mathcal{L}}$ by its tensor power if necessary, we can assume that $\widehat{H}^0(X, \bar{\mathcal{L}})$ is non-zero. Let p_0 be an integer such that the base locus of $\widehat{H}^0(X, \bar{\mathcal{L}})$ has no vertical irreducible components lying above any prime $p > p_0$. Let Y be a flag on X with $\text{char}(Y) = p > p_0$ such that Y_d is not contained in the base locus of $\widehat{H}^0(X, \bar{\mathcal{L}})$. Then the same property is true for all $\widehat{H}^0(X, n\bar{\mathcal{L}})$ since its base locus is contained in that of $\widehat{H}^0(X, \bar{\mathcal{L}})$. The strict transform $\pi_n^* Y$ gives a flag on X_n . It follows that

$$\# \nu_{\pi_n^* Y}(\widehat{H}^0(X, k\bar{\mathcal{A}}_n)) \geq \# \nu_{\pi_n^* Y}(\lambda(V_{k,n}(\bar{\mathcal{L}}))) = \# \nu_Y(V_{k,n}(\bar{\mathcal{L}})).$$

Divide both sides by $(nk)^d$, and set $k \rightarrow \infty$. We get

$$\frac{1}{n^d} \text{vol}(\Delta_{\pi_n^* Y}(\bar{\mathcal{A}}_n)) \geq \lim_{k \rightarrow \infty} \frac{\# \nu_Y(V_{k,n}(\bar{\mathcal{L}}))}{(nk)^d}.$$

By Theorem 3.3,

$$\liminf_{n \rightarrow \infty} \frac{1}{n^d} \text{vol}(\Delta_{\pi_n^* Y}(\overline{\mathcal{A}}_n)) \geq \text{vol}(\Delta_Y(\overline{\mathcal{L}})). \quad (5)$$

Now we want to take the limit as $p \rightarrow \infty$ to get volumes of the line bundles. It can be done by interchanging the orders of limits, which is possible since Theorem A is somehow uniform on $\overline{\mathcal{A}}_n$. In fact, Theorem 2.6 gives

$$\begin{aligned} \left| \text{vol}(\Delta_Y(\overline{\mathcal{L}})) \log p - \frac{1}{d!} \text{vol}(\overline{\mathcal{L}}) \right| &\leq \frac{c(\overline{\mathcal{L}})}{\log p}, \\ \left| \text{vol}(\Delta_{\pi_n^* Y}(\overline{\mathcal{A}}_n)) \log p - \frac{1}{d!} \text{vol}(\overline{\mathcal{A}}_n) \right| &\leq \frac{c(\overline{\mathcal{A}}_n)}{\log p}. \end{aligned}$$

The constants $c(\overline{\mathcal{L}}), c(\overline{\mathcal{A}}_n)$ can be taken as follows. By choosing $\overline{\mathcal{B}} \in \widehat{\text{Amp}}(X)$, we can set

$$c(\overline{\mathcal{L}}) = 2e_0 \frac{\text{vol}(\mathcal{L}_{\mathbb{Q}}) \overline{\mathcal{L}} \cdot \overline{\mathcal{B}}^{d-1}}{\text{vol}(\mathcal{B}_{\mathbb{Q}}) (d-1)!}.$$

The line bundle $\pi_n^* \overline{\mathcal{B}}$ is not ample on X_n , but it is nef in the sense that $\pi_n^* \overline{\mathcal{B}}^{d-1} \cdot \overline{\mathcal{T}} \geq 0$ for any $\overline{\mathcal{T}} \in \widehat{\text{Eff}}(X_n)$. It follows that $\pi_n^* \overline{\mathcal{B}}$ can also be a reference line bundle in the proof of Theorem 2.6, and thus we can take

$$c(\overline{\mathcal{A}}_n) = 2e_0 \frac{\text{vol}(\mathcal{A}_{n, \mathbb{Q}}) \overline{\mathcal{A}}_n \cdot \pi_n^* \overline{\mathcal{B}}^{d-1}}{\text{vol}(\pi_n^* \mathcal{B}_{\mathbb{Q}}) (d-1)!} \leq 2e_0 \frac{\text{vol}(n\mathcal{L}_{\mathbb{Q}}) n\overline{\mathcal{L}} \cdot \overline{\mathcal{B}}^{d-1}}{\text{vol}(\mathcal{B}_{\mathbb{Q}}) (d-1)!} = n^d c(\overline{\mathcal{L}}).$$

It follows that

$$\left| \frac{1}{n^d} \text{vol}(\Delta_{\pi_n^* Y}(\overline{\mathcal{A}}_n)) \log p - \frac{1}{d!} \frac{\text{vol}(\overline{\mathcal{A}}_n)}{n^d} \right| \leq \frac{c(\overline{\mathcal{L}})}{\log p}.$$

Therefore, taking $p \rightarrow \infty$ on (5) yields

$$\liminf_{n \rightarrow \infty} \frac{1}{n^d} \text{vol}(\overline{\mathcal{A}}_n) \geq \text{vol}(\overline{\mathcal{L}}).$$

This finishes proving (5).

As mentioned above, the line bundle $\overline{\mathcal{A}}_n$ in the decomposition above is base-point free but not ample in general. In order to make it ample, we are going to do a lot of modifications; for simplicity, we write $(\pi_n, X_n, \overline{\mathcal{A}}_n)$ as $(\pi, X', \overline{\mathcal{A}})$.

- We can further assume that X' is normal with smooth generic fibre by pass to a generic resolution of singularity.
- It suffices to treat the case that $\mathcal{A}_{\mathbb{Q}}$ is ample. In fact, we have $n_0 \pi^* \overline{\mathcal{L}} = \overline{\mathcal{A}}_0 + \overline{\mathcal{E}}_0$ for some $n_0 > 0$, $\overline{\mathcal{A}}_0 \in \text{amp}(X')$ and $\overline{\mathcal{E}}_0 \in \text{eff}(X')$. Then consider

$$(nN + n_0) \pi^* \overline{\mathcal{L}} = (N\overline{\mathcal{A}} + \overline{\mathcal{A}}_0) + (N\overline{\mathcal{E}} + \overline{\mathcal{E}}_0).$$

Make $N \gg n_0$.

- By the same trick above, we can further assume that $\overline{\mathcal{L}}(-c) = (\mathcal{L}, e^c \|\cdot\|)$ is base-point free for some $c > 0$.
- We can assume that $\overline{\mathcal{A}}$ has a positive smooth metric. By Theorem 3.4, we can find a positive metric $\|\cdot\|_j$ of $\overline{\mathcal{A}}$ which is pointwise greater than $\|\cdot\|$ and satisfies

$$1 \leq \frac{\|\cdot\|_{j, \text{sup}}}{\|\cdot\|_{\text{sup}}} < e^c.$$

Then the decomposition

$$n\pi^*\bar{\mathcal{L}} = (\mathcal{A}, \|\cdot\|_j) + \bar{\mathcal{E}}\left(\log \frac{\|\cdot\|_j}{\|\cdot\|}\right)$$

gives what we want.

- The above $\bar{\mathcal{A}}$ is already ample. By the condition that the metric of $\bar{\mathcal{A}}$ is positive and $\widehat{H}^0(X', \bar{\mathcal{A}})$ is base-point free, we need to check that $\bar{\mathcal{A}}$ is ample. Since \mathcal{A} is base-point free, it is automatically nef on any vertical subvarieties. We only need to check that $\hat{c}_1(\bar{\mathcal{L}}|_Z)^{\dim Z} > 0$ for any horizontal irreducible closed subvariety Z of X' . We can find an $s \in \widehat{H}^0(X', \bar{\mathcal{A}})$ such that $\text{div}(s)$ does not contain Z by this base-point-free property. Use this section to compute intersection. We get

$$\begin{aligned} (\bar{\mathcal{A}}|_Z)^{\dim Z} &= (\bar{\mathcal{A}}|_{\text{div}(s).Z})^{\dim Z-1} - \int_{Z(\mathbb{C})} \log \|s\| c_1(\bar{\mathcal{L}})^{\dim Z-1} \\ &> (\bar{\mathcal{A}}|_{\text{div}(s).Z})^{\dim Z-1}. \end{aligned}$$

By writing $\text{div}(s).Z$ as a positive linear combination of irreducible cycles, we reduce the problem to a smaller dimension. It means that the proof can be finished by induction on $\dim Z$. \square

ACKNOWLEDGEMENTS

I am indebted so much to Robert Lazarsfeld for introducing his joint work with Mircea Mustață to me, and for his hospitality during my visit at the University of Michigan at Ann Arbor. Almost all results of this paper are based on their work. I would also like to thank Shou-wu Zhang for many illustrating communications. Thanks also go to Sébastien Boucksom for pointing out the continuity property of the envelope, to Atsushi Moriawaki for pointing out a gap in an early version of the paper, and to Yuan Yuan for clarifying many concepts in complex analysis.

REFERENCES

- Ber07 R. Berman, *Bergman kernels and equilibrium measures for ample line bundles*, Preprint (2007), arXiv: 0704.1640v1 [math.CV].
- BB08 R. Berman and S. Boucksom, *Growth of balls of holomorphic sections and energy at equilibrium*, Preprint (2008), arXiv: 0803.1950v2 [math.CV].
- BV89 J.-M. Bismut and E. Vasserot, *The asymptotics of the Ray–Singer analytic torsion associated with high powers of a positive line bundle*, Comm. Math. Phys. **125** (1989), 355–367.
- BK07 Z. Blocki and S. Kolodziej, *On regularization of plurisubharmonic functions on manifolds*, Proc. Amer. Math. Soc. **135** (2007), 2089–2093.
- Che08a H. Chen, *Positive degree and arithmetic bigness*, Preprint (2008), arXiv: 0803.2583v3 [math.AG].
- Che08b H. Chen, *Arithmetic Fujita approximation*, Preprint (2008), arXiv: 0810.5479v2 [math.AG].
- Dem92 J.-P. Demailly, *Regularization of closed positive currents and intersection theory*, J. Algebraic Geom. **1** (1992), 361–409.
- Fal84 G. Faltings, *Calculus on arithmetic surfaces*, Ann. of Math. (2) **119** (1984), 387–424.
- Fuj94 T. Fujita, *Approximating Zariski decomposition of big line bundles*, Kodai Math. J. **17** (1994), 1–3.
- GS90 H. Gillet and C. Soulé, *Arithmetic intersection theory*, Publ. Math. Inst. Hautes Études Sci. **72** (1990), 93–174.

- GS91 H. Gillet and C. Soulé, *On the number of lattice points in convex symmetric bodies and their duals*, Israel J. Math. **74** (1991), 347–357.
- GS92 H. Gillet and C. Soulé, *An arithmetic Riemann–Roch theorem*, Invent. Math. **110** (1992), 473–543.
- KK08 K. Kaveh and A. Khovanskii, *Convex bodies and algebraic equations on affine varieties*, Preprint (2008), arXiv: 0804.4095v1 [math.AG].
- Kho92 A. Khovanskii, *The Newton polytope, the Hilbert polynomial and sums of finite sets*, Funct. Anal. Appl. **26** (1992), 276–281.
- Laz04 R. Lazarsfeld, *Positivity in algebraic geometry. I. Classical setting: line bundles and linear series*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3, vol. 48 (Springer, Berlin, 2004).
- LM08 R. Lazarsfeld and M. Mustață, *Convex bodies associated to linear series*, Preprint (2008), arXiv: 0805.4559v1 [math.AG], Ann. Sci. École Norm. Sup., to appear.
- Mor96 A. Moriwaki, *Hodge index theorem for arithmetic cycles of codimension one*, Math. Res. Lett. **3** (1996), 173–183.
- Mor00 A. Moriwaki, *Arithmetic height functions over finitely generated fields*, Invent. Math. **140** (2000), 101–142.
- Mor09 A. Moriwaki, *Continuity of volumes on arithmetic varieties*, J. Algebraic Geom. **18** (2009), 407–457.
- Oko96 A. Okounkov, *Brunn–Minkowski inequality for multiplicities*, Invent. Math. **125** (1996), 405–411.
- Oko03 A. Okounkov, *Why would multiplicities be log-concave?* in *The orbit method in geometry and physics*, Progress in Mathematics, vol. 213 (Birkhäuser, Boston, MA, 2003), 329–347.
- Yua08 X. Yuan, *Big line bundle over arithmetic varieties*, Invent. Math. **173** (2008), 603–649.
- Zha95 S. Zhang, *Positive line bundles on arithmetic varieties*, J. Amer. Math. Soc. **8** (1995), 187–221.

Xinyi Yuan xyy@ias.edu

School of Mathematics, Institute for Advanced Study, Einstein Drive, Princeton, NJ 08540, USA