

This article was downloaded by: [Matematicky Ustav Av Cr]

On: 19 December 2011, At: 11:44

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Stochastic Analysis and Applications

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Isaa20>

On Weak Solutions of Stochastic Differential Equations

Martina Hofmanová^{a b} & Jan Seidler^b

^a Department of Mathematical Analysis, Faculty of Mathematics and Physics, Charles University, Praha, Czech Republic

^b Institute of Information Theory and Automation of the ASCR, Praha, Czech Republic

Available online: 19 Dec 2011

To cite this article: Martina Hofmanová & Jan Seidler (2012): On Weak Solutions of Stochastic Differential Equations, *Stochastic Analysis and Applications*, 30:1, 100-121

To link to this article: <http://dx.doi.org/10.1080/07362994.2012.628916>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

On Weak Solutions of Stochastic Differential Equations

MARTINA HOFMANOVÁ^{1,2} AND JAN SEIDLER²

¹Department of Mathematical Analysis, Faculty of Mathematics
and Physics, Charles University, Praha, Czech Republic

²Institute of Information Theory and Automation of the ASCR,
Praha, Czech Republic

A new proof of existence of weak solutions to stochastic differential equations with continuous coefficients based on ideas from infinite-dimensional stochastic analysis is presented. The proof is fairly elementary, in particular, neither theorems on representation of martingales by stochastic integrals nor results on almost sure representation for tight sequences of random variables are needed.

Keywords Fractional integrals; Stochastic differential equations; Weak solutions.

Mathematics Subject Classification 60H10.

0. Introduction

In this article, we provide a modified proof of Skorokhod's classical theorem on existence of (weak) solutions to a stochastic differential equation

$$dX = b(t, X)dt + \sigma(t, X)dW, \quad X(0) = \varphi,$$

where $b : [0, T] \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ and $\sigma : [0, T] \times \mathbb{R}^m \rightarrow \mathbb{M}_{m \times n}$ are Borel functions of at most linear growth continuous in the second variable. (Henceforth, by $\mathbb{M}_{m \times n}$ we shall denote the space of all m -by- n matrices over \mathbb{R} endowed with the Hilbert–Schmidt norm $\|A\| = (\text{Tr } AA^*)^{1/2}$.) Our proof combines tools that were proposed for handling weak solutions of stochastic evolution equations in infinite-dimensional spaces, where traditional methods cease to work, with results on preservation of the local martingale property under convergence in law. In a finite-dimensional situation, the “infinite-dimensional” methods simplify considerably and, in our opinion, the alternative proof based on them is more lucid and elementary than the

Received December 22, 2010; Accepted December 22, 2010

This research was supported by the GAČR Grant No. P201/10/0752.

The authors are indebted to Martin Ondřeját for many useful discussions.

Address correspondence to Jan Seidler, ÚTIA AV ČR, Pod vodárenskou věží 4, 182 08 Praha 8, Czech Republic; E-mail: seidler@utia.cas.cz

standard one. A positive teaching experience of the second author was, in fact, the main motivation for writing this article. Moreover, we believe that the reader may find the comparison with other available approaches illuminating.

To explain our argument more precisely, let us recall the structure of the usual proof; for notational simplicity, we shall consider (in the informal introduction only) autonomous equations. Kiyosi Itô showed in his seminal articles (see, e.g., [9, 10]) that a stochastic differential equation

$$dX = b(X)dt + \sigma(X)dW \quad (0.1)$$

$$X(0) = \varphi \quad (0.2)$$

driven by an n -dimensional Wiener process W has a unique solution provided that $b : \mathbb{R}^m \rightarrow \mathbb{R}^m$, $\sigma : \mathbb{R}^m \rightarrow \mathbb{M}_{m \times n}$ are Lipschitz continuous functions. A next important step was taken by Skorokhod [16, 17] in 1961, who proved that there exists a solution to (0.1), (0.2) if b and σ are continuous functions of at most linear growth, that is,

$$\sup_{x \in \mathbb{R}^m} \frac{\|b(x)\| + \|\sigma(x)\|}{1 + \|x\|} < \infty.$$

It was realized only later that two different concepts of a solution are involved: For Lipschitzian coefficients, there exists an (\mathcal{F}_t) -progressively measurable process in \mathbb{R}^m solving (0.1) and such that $X(0) = \varphi$, whenever $(\Omega, \mathcal{F}, (\mathcal{F}_t), \mathbf{P})$ is a stochastic basis carrying an n -dimensional (\mathcal{F}_t) -Wiener process and φ is an \mathcal{F}_0 -measurable function. (We say that (0.1), (0.2) has a strong solution.) On the other hand, for continuous coefficients, a stochastic basis $(\Omega, \mathcal{F}, (\mathcal{F}_t), \mathbf{P})$, an n -dimensional (\mathcal{F}_t) -Wiener process W and an (\mathcal{F}_t) -progressively measurable process X may be found such that X solves (0.1) and $X(0)$ and φ have the same law. (We speak about existence of a weak solution to (0.1), (0.2) in such a case.) It is well known that this difference is substantial in general: under assumptions of the Skorokhod theorem strong solutions need not exist (see [1]).

Skorokhod's existence theorem is remarkable not only by itself, but also because of the method of its proof. To present it, we need some notation: If M and N are continuous real local martingales, then by $\langle M \rangle$ we denote the quadratic variation of M and by $\langle M, N \rangle$ the cross-variation of M and N . Let $M = (M^i)_{i=1}^m$ and $N = (N^j)_{j=1}^n$ be continuous local martingales with values in \mathbb{R}^m and \mathbb{R}^n , respectively. By $\langle\langle M \rangle\rangle$ we denote the tensor quadratic variation of M , $\langle\langle M \rangle\rangle = (\langle M^i, M^k \rangle)_{i,k=1}^m$, and we set $\langle M \rangle = \text{Tr} \langle\langle M \rangle\rangle$. Analogously, we define

$$M \otimes N = (M^i N^j)_{i=1, j=1}^{m, n}, \quad \langle\langle M, N \rangle\rangle = (\langle M^i, N^j \rangle)_{i=1, j=1}^{m, n}.$$

Let X and Y be random variables with values in the same measurable space (E, \mathcal{E}) , we write $X \stackrel{\mathcal{L}}{\sim} Y$ if X and Y have the same law on \mathcal{E} . Similarly, $X \stackrel{\mathcal{L}}{\sim} \nu$ means that the law of X is a probability measure ν on \mathcal{E} .

Let

$$dX_r = b_r(X_r)dt + \sigma_r(X_r)dW, \quad X_r(0) = \varphi$$

be a sequence of equations that have strong solutions and approximate (0.1) in a suitable sense. (We shall approximate b and σ by Lipschitz continuous functions having the same growth as b and σ , but likewise it is possible to use, for example, finite difference approximations.) The linear growth hypothesis makes it possible to prove that

$$\text{the laws of } \{X_r; r \geq 1\} \text{ are tight,} \quad (0.3)$$

that is, form a relatively weakly compact set of measures on the space of continuous trajectories. Then Skorokhod's theorem on almost surely converging realizations of converging laws (see, e.g., [5, Theorem 11.7.2]) may be invoked, which yields a subsequence $\{X_{r_k}\}$ of $\{X_r\}$, a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbf{P}})$ and sequences $\{\tilde{X}_k; k \geq 0\}$, $\{\tilde{W}_k; k \geq 0\}$ such that

$$(X_{r_k}, W) \stackrel{\mathcal{D}}{\sim} (\tilde{X}_k, \tilde{W}_k), \quad k \geq 1; \quad (\tilde{X}_k, \tilde{W}_k) \xrightarrow[k \rightarrow \infty]{\tilde{\mathbf{P}}\text{-a.s.}} (\tilde{X}_0, \tilde{W}_0). \quad (0.4)$$

It is claimed that \tilde{X}_0 is the (weak) solution looked for. Skorokhod's articles [16, 17] are written in a very concise way and details of proofs are not offered; nowadays a standard version of Skorokhod's proof is as follows (see [18, Theorem 6.1.6], [8, Theorem IV.2.2], [12, Theorem 5.4.22]): under a suitable integrability assumption upon the initial condition,

$$M_k = X_{r_k} - X_{r_k}(0) - \int_0^{\cdot} b_{r_k}(X_{r_k}(s)) ds$$

is a martingale with a (tensor) quadratic variation

$$\langle\langle M_k \rangle\rangle = \int_0^{\cdot} \sigma_{r_k}(X_{r_k}(s)) \sigma_{r_k}^*(X_{r_k}(s)) ds,$$

for all $k \geq 1$. Equality in law (0.4) implies that also

$$\tilde{M}_k = \tilde{X}_k - \tilde{X}_k(0) - \int_0^{\cdot} b_{r_k}(\tilde{X}_k(s)) ds$$

are martingales for $k \geq 1$, with quadratic variations

$$\langle\langle \tilde{M}_k \rangle\rangle = \int_0^{\cdot} \sigma_{r_k}(\tilde{X}_k(s)) \sigma_{r_k}^*(\tilde{X}_k(s)) ds.$$

Using convergence $\tilde{\mathbf{P}}$ -almost everywhere, it is possible to show that

$$\tilde{M}_0 = \tilde{X}_0 - \tilde{X}_0(0) - \int_0^{\cdot} b(\tilde{X}_0(s)) ds$$

is a martingale with a quadratic variation

$$\langle\langle \tilde{M}_0 \rangle\rangle = \int_0^{\cdot} \sigma(\tilde{X}_0(s)) \sigma^*(\tilde{X}_0(s)) ds.$$

By the integral representation theorem for martingales with an absolutely continuous quadratic variation (see, e.g., [12, Theorem 3.4.2] or [8, Theorem II.7.1']), there exists a Wiener process \widehat{W} (on an extended probability space) satisfying

$$\widetilde{M}_0 = \int_0^\cdot \sigma(\widetilde{X}_0(s)) d\widehat{W}(s).$$

Therefore, $(\widehat{W}, \widetilde{X}_0)$ is a weak solution to (0.1), (0.2). (In the cited books, martingale problems are used instead of weak solutions. Then the integral representation theorem is hidden in the construction of a weak solution from a solution to the martingale problem, so a complete proof is essentially the one sketched above.)

This procedure has two rather nontrivial inputs: the Skorokhod representation theorem, and the integral representation theorem whose proof, albeit based on a simple and beautiful idea, becomes quite technical if the space dimension is greater than one. An alternative approach to identification of the limit was discovered recently (see [3, 14]) in the course of study of stochastic wave maps between manifolds, where integral representation theorems for martingales are no longer available. The new method, which refers only to basic properties of martingales and stochastic integrals, may be described in the case of the problem (0.1), (0.2) in the following way: One starts again with a sequence $\{(\widetilde{X}_k, \widetilde{W}_k)\}$ such that (0.4) holds true. If the initial condition is p -integrable for some $p > 2$, it can be shown in a straightforward manner, using the almost sure convergence, that

$$\widetilde{M}_0, \quad \|\widetilde{M}_0\|^2 - \int_0^\cdot \|\sigma(\widetilde{X}_0(s))\|^2 ds, \quad \widetilde{M}_0 \otimes \widetilde{W}_0 - \int_0^\cdot \sigma(\widetilde{X}_0(s)) ds$$

are martingales, in other words,

$$\left\langle \widetilde{M}_0 - \int_0^\cdot \sigma(\widetilde{X}_0(s)) d\widetilde{W}_0(s) \right\rangle = 0,$$

whence one concludes that $(\widetilde{W}_0, \widetilde{X}_0)$ is a weak solution. If the additional integrability hypothesis on φ is not satisfied, the proof remains almost the same, only a suitable cut-off procedure must be amended.

We take a step further and eliminate also the Skorokhod representation theorem. Let \widetilde{P}_k be the laws of (X_{r_k}, W) on the space $U = \mathcal{C}([0, T]; \mathbb{R}^m) \times \mathcal{C}([0, T]; \mathbb{R}^n)$; we know that the sequence $\{\widetilde{P}_k\}$ converges weakly to some measure \widetilde{P}_0 . Denote by (Y, B) the canonical process on U and set

$$\overline{M}_k = Y - Y(0) - \int_0^\cdot b_{r_k}(Y(s)) ds, \quad k \geq 0$$

(with $b_{r_0} = b$, $\sigma_{r_0} = \sigma$). Then

$$\overline{M}_k, \quad \|\overline{M}_k\|^2 - \int_0^\cdot \|\sigma_{r_k}(Y(s))\|^2 ds, \quad \overline{M}_k \otimes B - \int_0^\cdot \sigma_{r_k}(Y(s)) ds, \quad (0.5)$$

are local martingales under the measure \widetilde{P}_k for every $k \geq 1$, as can be inferred quite easily from the definition of the measure \widetilde{P}_k . Now one may try to use Theorem IX.1.17 from Jacod and Shiryaev [11] stating, roughly speaking, that a limit in law

of a sequence of continuous local martingales is a local martingale. We do not use this theorem explicitly, since to establish convergence in law of the processes (0.5) as $k \rightarrow \infty$ is not simpler than to check the local martingale property for $k = 0$ directly, but our argument is inspired by the proofs in the book [11]. The proof we propose is not difficult and it is almost self-contained, it requires only two auxiliary lemmas (with simple proofs) from Jacod and Shiryaev [11] on continuity properties of certain first entrance times, which we recall in the Appendix. Once we know that the processes (0.5) are local martingales for $k = 0$ as well, the trick from Brzeźniak and Ondreját [3] and Ondreját [14] may be used yielding that (B, Y) is a weak solution to (0.1), (0.2). It is worth mentioning that this procedure is independent of any integrability hypothesis on φ .

The proof of (0.3) not being our main concern notwithstanding, we decided to include a less standard proof of tightness inspired also by the theory of stochastic partial differential equations. We adopt an argument proposed by Gątarek and Goldys [6] (cf. also [4, chapter 8]), who introduced it when studying weak solutions to stochastic evolution equations in Hilbert spaces, and which relies on the factorization method of Da Prato et al. (see [4, chapters 5 and 7] for a thorough exposition) and on compactness properties of fractional integral operators. The fractional calculus has become popular amongst probabilists recently because of its applications to fractional Brownian motion driven stochastic integrals and a proof of tightness using it may suit some readers more than the traditional one based on estimates of moduli of continuity.

Let us close this Introduction by stating the result to be proved precisely.

Theorem 0.1. *Let $b : [0, T] \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ and $\sigma : [0, T] \times \mathbb{R}^m \rightarrow \mathbb{M}_{m \times n}$ be Borel functions such that $b(t, \cdot)$ and $\sigma(t, \cdot)$ are continuous on \mathbb{R}^m for any $t \in [0, T]$ and the linear growth hypothesis is satisfied, that is*

$$\exists K_* < \infty \forall t \in [0, T] \forall x \in \mathbb{R}^m \quad \|b(t, x)\| \vee \|\sigma(t, x)\| \leq K_* (1 + \|x\|). \quad (0.6)$$

Let ν be a Borel probability measure on \mathbb{R}^m . Then there exists a weak solution to the problem

$$dX = b(t, X)dt + \sigma(t, X)dW, \quad X(0) \stackrel{\mathcal{D}}{\sim} \nu. \quad (0.7)$$

We recall that a weak solution to (0.7) is a triple $((G, \mathcal{G}, (\mathcal{G}_t), \mathbf{Q}), W, X)$, where $(G, \mathcal{G}, (\mathcal{G}_t), \mathbf{Q})$ is a stochastic basis with a filtration (\mathcal{G}_t) that satisfies the usual conditions, W is an n -dimensional (\mathcal{G}_t) -Wiener process and X is an \mathbb{R}^m -valued (\mathcal{G}_t) -progressively measurable process such that $\mathbf{Q} \circ X(0)^{-1} = \nu$ and

$$X(t) = X(0) + \int_0^t b(r, X(r))dr + \int_0^t \sigma(r, X(r))dW(r)$$

for all $t \in [0, T]$ \mathbf{Q} -almost surely.

The rest of this article is devoted to the proof of Theorem 0.1. In Section 1, a sequence of equations with Lipschitzian coefficients approximation (0.7) is constructed, tightness of the set of their solutions being shown in Section 2. In Section 3, cluster points of the set of approximating solutions are identified as weak solutions to (0.7).

1. Approximations

In this section, we introduce a sequence of equations that have strong solutions and approximate the problem (0.7). If E and F are metric spaces, we denote by $\mathcal{C}(E; F)$ the space of all continuous mappings from E to F . For brevity, we shall sometimes write \mathcal{C}_V instead of $\mathcal{C}([0, T]; \mathbb{R}^V)$ if $V \in \mathbb{N}$. If $f \in \mathcal{C}([0, T]; F)$ and $s \in [0, T]$ then the restriction of f to the interval $[0, s]$ will be denoted by $q_s f$. Plainly, $q_s : \mathcal{C}([0, T]; F) \rightarrow \mathcal{C}([0, s]; F)$ is a continuous mapping. Finally, $L^q(G; \mathbb{R}^V)$ stands for the space of q -integrable functions on G with values in \mathbb{R}^V .

Our construction is based on the following proposition.

Proposition 1.1. *Suppose that $F : \mathbb{R}_+ \times \mathbb{R}^N \rightarrow \mathbb{R}^V$ is a Borel function of at most linear growth, that is,*

$$\exists L < \infty \forall t \geq 0 \forall x \in \mathbb{R}^N \quad \|F(t, x)\| \leq L(1 + \|x\|),$$

such that $F(t, \cdot) \in \mathcal{C}(\mathbb{R}^N; \mathbb{R}^V)$ for any $t \in \mathbb{R}_+$. Then there exists a sequence of Borel functions $F_k : \mathbb{R}_+ \times \mathbb{R}^N \rightarrow \mathbb{R}^V$, $k \geq 1$, which have at most linear growth uniformly in k , namely

$$\forall k \geq 1 \forall t \geq 0 \forall x \in \mathbb{R}^N \quad \|F_k(t, x)\| \leq L(2 + \|x\|),$$

which are Lipschitz continuous in the second variable uniformly in the first one,

$$\forall k \geq 1 \exists L_k < \infty \forall t \geq 0 \forall x, y \in \mathbb{R}^N \quad \|F_k(t, x) - F_k(t, y)\| \leq L_k \|x - y\|,$$

and that satisfy

$$\lim_{k \rightarrow \infty} F_k(t, \cdot) = F(t, \cdot) \quad \text{locally uniformly on } \mathbb{R}^N$$

for all $t \geq 0$.

The proof is rather standard so it is not necessary to dwell on its details: one takes a smooth function $\zeta \in \mathcal{C}^\infty(\mathbb{R}^N)$ such that $\zeta \geq 0$, $\text{supp } \zeta \subseteq \{x \in \mathbb{R}^N; \|x\| \leq 1\}$ and $\int_{\mathbb{R}^N} \zeta dx = 1$ and sets

$$G_k(t, x) = k^N \int_{\mathbb{R}^N} F(t, y) \zeta(k(x - y)) dy$$

for $k \geq 1$, $t \geq 0$ and $x \in \mathbb{R}^N$. The functions G_k have all desired properties except for being only locally Lipschitz, but it is possible to modify them outside a sufficiently large ball in an obvious manner.

Let the coefficients b and σ satisfy the assumptions of Theorem 0.1. Using Proposition 1.1, we find Borel functions $b_k : [0, T] \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ and $\sigma_k : [0, T] \times \mathbb{R}^m \rightarrow \mathbb{M}_{m \times n}$, $k \geq 1$, such that

$$\sup_{k \geq 1} \sup_{t \in [0, T]} \{ \|b_k(t, x)\| \vee \|\sigma_k(t, x)\| \} \leq K_* (2 + \|x\|), \quad x \in \mathbb{R}^m, \quad (1.1)$$

$b_k(t, \cdot)$ and $\sigma_k(t, \cdot)$ are Lipschitz continuous uniformly in $t \in [0, T]$ and converge locally uniformly on \mathbb{R}^m as $k \rightarrow \infty$ to $b(t, \cdot)$ and $\sigma(t, \cdot)$, respectively, for all $t \in [0, T]$.

Fix an arbitrary stochastic basis $(\Omega, \mathcal{F}, (\mathcal{F}_t), \mathbf{P})$, on which an n -dimensional (\mathcal{F}_t) -Wiener process W and an \mathcal{F}_0 -measurable random variable $\varphi : \Omega \rightarrow \mathbb{R}^m$ with $\varphi \stackrel{\mathcal{D}}{\sim} \nu$ are defined. It is well known that for any $k \geq 1$ there exists a unique (\mathcal{F}_t) -progressively measurable \mathbb{R}^m -valued stochastic process X_k solving the equation

$$dX_k = b_k(t, X_k)dt + \sigma_k(t, X_k)dW, \quad X_k(0) = \varphi. \quad (1.2)$$

Moreover, for any $p \in [2, \infty[$ there exists a constant $C_* < \infty$, depending only on p , T and K_* , such that

$$\sup_{k \geq 1} \mathbf{E} \sup_{0 \leq t \leq T} \|X_k(t)\|^p \leq C_* (1 + \mathbf{E}\|\varphi\|^p), \quad (1.3)$$

provided that

$$\int_{\mathbb{R}^m} \|x\|^p d\nu(x) = \mathbf{E}\|\varphi\|^p < \infty.$$

2. Tightness

Let $\{X_k; k \geq 1\}$ be the sequence of solutions to (1.2). Plainly, the processes X_k may be viewed as random variables $X_k : \Omega \rightarrow \mathcal{C}_m$ (where the Polish metric space \mathcal{C}_m is endowed with its Borel σ -algebra). In this section, we aim at establishing the following proposition.

Proposition 2.1. *The set $\{\mathbf{P} \circ X_k^{-1}; k \geq 1\}$ of Borel probability measures on $\mathcal{C}([0, T]; \mathbb{R}^m)$ is tight.*

To this end, let us recall the definition of the Riemann–Liouville (or fractional integral) operator: if $q \in]1, \infty]$, $\alpha \in]\frac{1}{q}, 1]$ and $f \in L^q([0, T]; \mathbb{R}^m)$, we define a function $R_\alpha f : [0, T] \rightarrow \mathbb{R}^m$ by

$$(R_\alpha f)(t) = \int_0^t (t-s)^{\alpha-1} f(s) ds, \quad 0 \leq t \leq T.$$

The definition is correct, as an easy application of the Hölder inequality shows. Note that, in particular, $R_1 f = \int_0^\cdot f(t) dt$. It is well known (and may be checked by very straightforward calculations) that R_α is a bounded linear operator from $L^q([0, T]; \mathbb{R}^m)$ to the space $\mathcal{C}^{0, \alpha-1/q}([0, T]; \mathbb{R}^m)$ of $(\alpha - \frac{1}{q})$ -Hölder continuous functions (see, e.g., [15, Theorem 3.6]). Balls in $\mathcal{C}^{0, \alpha-1/q}([0, T]; \mathbb{R}^m)$ are relatively compact in $\mathcal{C}([0, T]; \mathbb{R}^m)$ by the Arzelà–Ascoli theorem, hence, we arrive at

Lemma 2.2. *If $q \in]1, \infty]$ and $\alpha \in]\frac{1}{q}, 1]$, then R_α is a compact linear operator from $L^q([0, T]; \mathbb{R}^m)$ to $\mathcal{C}([0, T]; \mathbb{R}^m)$.*

We shall need also a Fubini-type theorem for stochastic integrals in the following form (see [4, Theorem 4.18] for a more general result):

Lemma 2.3. *Let (X, Σ, μ) be a finite measure space, $(G, \mathcal{G}, (\mathcal{G}_t), \mathbf{Q})$ a stochastic basis, and B an n -dimensional (\mathcal{G}_t) -Wiener process. Denote by \mathcal{M} the σ -algebra of*

(\mathcal{G}_t) -progressively measurable sets and assume that $\psi : [0, T] \times G \times X \rightarrow \mathbb{M}_{m \times n}$ is an $\mathcal{M} \otimes \Sigma$ -measurable mapping such that

$$\int_X \left(\int_0^T \int_G \|\psi(s, x)\|^2 d\mathbf{Q} ds \right)^{1/2} d\mu(x) < \infty. \quad (2.1)$$

Then

$$\int_X \left[\int_0^T \psi(s, x) dB(s) \right] d\mu(x) = \int_0^T \left[\int_X \psi(s, x) d\mu(x) \right] dB(s)$$

\mathbf{Q} -almost surely.

The last auxiliary result to be recalled is the Young inequality for convolutions (see, e.g., [13, Theorem 4.2]).

Lemma 2.4. Let $p, r, s \in [1, \infty]$ satisfy

$$\frac{1}{p} + \frac{1}{q} = 1 + \frac{1}{s}.$$

If $f \in L^p(\mathbb{R}^d)$ and $g \in L^q(\mathbb{R}^d)$, then the integral

$$(f * g)(x) \equiv \int_{\mathbb{R}^d} f(x - y)g(y)dy$$

converges for almost all $x \in \mathbb{R}^d$, $f * g \in L^s(\mathbb{R}^d)$ and

$$\|f * g\|_{L^s} \leq \|f\|_{L^p} \|g\|_{L^q}.$$

In fact, we shall need only a particular one-dimensional case of Lemma 2.4: if $f \in L^p(0, T)$, $g \in L^q(0, T)$, $\frac{1}{p} + \frac{1}{q} = 1 + \frac{1}{s}$, then

$$\int_0^T \left| \int_0^t f(t-r)g(r)dr \right|^s dt \leq \|f\|_{L^p(0,T)}^s \|g\|_{L^q(0,T)}^s. \quad (2.2)$$

Now we derive a representation formula that plays a key role in our proof of Proposition 2.1.

Lemma 2.5. Let ψ be an $\mathbb{M}_{m \times n}$ -valued progressively measurable process such that

$$\mathbf{E} \int_0^T \|\psi(s)\|^q ds < \infty$$

for some $q > 2$. Choose $\alpha \in]\frac{1}{q}, \frac{1}{2}[$ and set

$$Z(t) = \int_0^t (t-u)^{-\alpha} \psi(u) dW(u), \quad 0 \leq t \leq T.$$

Then

$$\int_0^t \psi(s) dW(s) = \frac{\sin \pi \alpha}{\pi} (R_x Z)(t)$$

for all $t \in [0, T]$ \mathbf{P} -almost surely.

Proof. The result is well known and widely used for infinite-dimensional systems (see, e.g., [4, §5.3]). For finite-dimensional equations, the proof is slightly simpler and, thus, it is repeated here for the reader's convenience.

Since $s^{-2\alpha} \in L^1(0, T)$, $\mathbf{E}\|\psi(\cdot)\|^2 \in L^1(0, T)$, their convolution

$$t \mapsto \int_0^t (t-s)^{-2\alpha} \mathbf{E}\|\psi(s)\|^2 ds = \mathbf{E} \int_0^t |(t-s)^{-\alpha} \psi(s)|^2 ds$$

belongs to $L^1(0, T)$ as well and so is finite almost everywhere in $[0, T]$, which implies that $Z(t)$ is well defined for almost all $t \in [0, T]$. By the Burkholder-Davis-Gundy inequality,

$$\begin{aligned} \mathbf{E} \int_0^T \|Z(t)\|^q dt &= \int_0^T \mathbf{E} \left\| \int_0^s (s-u)^{-\alpha} \psi(u) dW(u) \right\|^q ds \\ &\leq C_q \mathbf{E} \int_0^T \left(\int_0^s (s-u)^{-2\alpha} \|\psi(u)\|^2 du \right)^{q/2} ds \\ &\leq C_q \left(\int_0^T s^{-2\alpha} ds \right)^{q/2} \left(\int_0^T \mathbf{E} \|\psi(u)\|^q du \right); \end{aligned}$$

the last estimate being a consequence of (2.2) and the fact that $\mathbf{E}\|\psi(\cdot)\|^2 \in L^{q/2}(0, T)$. Hence $Z(\cdot, \omega) \in L^q(0, T; \mathbb{R}^m)$ for \mathbf{P} -almost all $\omega \in \Omega$ and $R_\alpha Z$ is well defined \mathbf{P} -almost surely.

Further,

$$\begin{aligned} &\int_0^t \left(\mathbf{E} \int_0^t \|(t-s)^{\alpha-1} \mathbf{1}_{[0,s]}(u) (s-u)^{-\alpha} \psi(u)\|^2 du \right)^{1/2} ds \\ &= \int_0^t (t-s)^{\alpha-1} \left(\int_0^s (s-u)^{-2\alpha} \mathbf{E} \|\psi(u)\|^2 du \right)^{1/2} ds \\ &\leq \left(\int_0^t s^{(\alpha-1)q^*} ds \right)^{1/q^*} \left(\int_0^t \left(\int_0^s (s-u)^{-2\alpha} \mathbf{E} \|\psi(u)\|^2 du \right)^{q/2} ds \right)^{1/q} \\ &\leq \left(\int_0^t s^{(\alpha-1)q^*} ds \right)^{1/q^*} \left(\int_0^t s^{-2\alpha} ds \right)^{1/2} \left(\int_0^t \mathbf{E} \|\psi(u)\|^q du \right)^{1/q} < \infty, \end{aligned}$$

where $\frac{1}{q^*} + \frac{1}{q} = 1$ and the Hölder and Young inequalities were used consecutively. This means that the hypothesis (2.1) of Lemma 2.3 is satisfied and this lemma may be used to obtain

$$\begin{aligned} (R_\alpha Z)(t) &= \int_0^t (t-s)^{\alpha-1} \left(\int_0^s (s-u)^{-\alpha} \psi(u) dW(u) \right) ds \\ &= \int_0^t \int_0^t (t-s)^{\alpha-1} \mathbf{1}_{[0,s]}(u) (s-u)^{-\alpha} \psi(u) dW(u) ds \\ &= \int_0^t \left[\int_0^t (t-s)^{\alpha-1} \mathbf{1}_{[0,s]}(u) (s-u)^{-\alpha} ds \right] \psi(u) dW(u) \end{aligned}$$

$$\begin{aligned}
&= \int_0^t \left[\int_u^t (t-s)^{\alpha-1} (s-u)^{-\alpha} ds \right] \psi(u) dW(u) \\
&= \int_0^t \underbrace{\left[\int_0^1 (1-v)^{\alpha-1} v^{-\alpha} dv \right]}_{\frac{\pi}{\sin \pi \alpha}} \psi(u) dW(u).
\end{aligned}$$

□

Proof of Proposition 2.1. Let an arbitrary $\varepsilon > 0$ be given, we have to find a relatively compact set $K \subseteq \mathcal{C}_m$ such that

$$\inf_{k \geq 1} \mathbf{P}\{X_k \in K\} \geq 1 - \varepsilon.$$

In what follows, we shall denote by D_i constants independent of k and by $|\cdot|_q$ the norm of $L^q(0, T; \mathbb{R}^m)$.

First, we prove our claim under an additional assumption that there exists $p > 2$ such that

$$\mathbf{E}\|\varphi\|^p < \infty. \quad (2.3)$$

Plainly, a compact set $\Gamma \subseteq \mathbb{R}^m$ may be found satisfying

$$v(\Gamma) = \mathbf{P}\{\varphi \in \Gamma\} \geq 1 - \frac{\varepsilon}{3}.$$

Take an $\alpha \in]\frac{1}{p}, \frac{1}{2}[$. By Lemma 2.5,

$$\begin{aligned}
X_k(t) &= \varphi + \int_0^t b_k(s, X_k(s)) ds + \int_0^t \sigma_k(s, X_k(s)) dW(s) \\
&= \varphi + [R_1 b(\cdot, X_k(\cdot))](t) + \frac{\sin \pi \alpha}{\pi} (R_\alpha Z_k)(t), \quad 0 \leq t \leq T,
\end{aligned}$$

\mathbf{P} -almost surely, where

$$Z_k(s) = \int_0^s (s-u)^{-\alpha} \sigma_k(u, X_k(u)) dW(u), \quad 0 \leq s \leq T.$$

Applying the Chebyshev inequality, (1.1) and (1.3) we get

$$\begin{aligned}
\mathbf{P}\{|b_k(\cdot, X_k(\cdot))|_p \geq \Lambda\} &\leq \frac{1}{\Lambda^p} \mathbf{E} \int_0^T \|b_k(t, X_k(t))\|^p dt \\
&\leq \frac{1}{\Lambda^p} K_*^p \mathbf{E} \int_0^T (2 + \|X_k(t)\|)^p dt \\
&\leq \frac{D_1}{\Lambda^p} (1 + \mathbf{E}\|\varphi\|^p).
\end{aligned}$$

Similarly, invoking in addition the Burkholder–Davis–Gundy and Young inequalities,

$$\mathbf{P}\{|Z_k|_p \geq \Lambda\} \leq \frac{1}{\Lambda^p} \mathbf{E} \int_0^T \|Z_k(t)\|^p dt$$

$$\begin{aligned} &\leq \frac{D_2}{\Lambda^p} \mathbf{E} \int_0^T \left(\int_0^t (t-s)^{-2\alpha} \|\sigma_k(s, X_k(s))\|^2 ds \right)^{p/2} dt \\ &\leq \frac{D_2}{\Lambda^p} \left(\int_0^T s^{-2\alpha} ds \right)^{p/2} \left(\int_0^T \mathbf{E} \|\sigma_k(s, X_k(s))\|^p ds \right) \\ &\leq \frac{D_3}{\Lambda^p} (1 + \mathbf{E} \|\varphi\|^p). \end{aligned}$$

Let us choose $\Lambda_0 < \infty$ so that

$$\frac{D_1 + D_3}{\Lambda_0^p} (1 + \mathbf{E} \|\varphi\|^p) < \frac{\varepsilon}{3}$$

and set

$$K = \left\{ f \in \mathcal{C}([0, T]; \mathbb{R}^m); f = x + R_1 r + \frac{\sin \pi\alpha}{\pi} R_\alpha v, x \in \Gamma, \right. \\ \left. r, v \in L^p(0, T; \mathbb{R}^m), |r|_p \vee |v|_p \leq \Lambda_0 \right\}.$$

Since the operators R_1 and R_α are compact, the set K is relatively compact and

$$\begin{aligned} \mathbf{P}\{X_k \notin K\} &\leq \mathbf{P}\{\varphi \notin \Gamma\} + \mathbf{P}\{|b_k(\cdot, X_k(\cdot))|_p > \Lambda_0\} + \mathbf{P}\{|Z_k|_p > \Lambda_0\} \\ &\leq \frac{2}{3} \varepsilon < \varepsilon \end{aligned}$$

for any $k \geq 1$, which completes the proof of tightness under the additional assumption (2.3).

Finally, let φ be arbitrary. Let $\varepsilon > 0$ be fixed, we may find $\Pi \geq 0$ such that $\mathbf{P}\{\|\varphi\| > \Pi\} < \frac{\varepsilon}{2}$. Let $\widehat{X}_k, k \geq 1$, be the solutions to

$$d\widehat{X}_k = b_k(t, \widehat{X}_k)dt + \sigma_k(t, \widehat{X}_k)dW, \quad \widehat{X}_k(0) = \mathbf{1}_{\{\|\varphi\| \leq \Pi\}} \varphi. \tag{2.4}$$

The initial condition in (2.4) satisfies (2.3), so by the first part of the proof we know that the set $\{\mathbf{P} \circ \widehat{X}_k^{-1}; k \geq 1\}$ is tight and there exists a compact set $K \subseteq \mathcal{C}_m$ such that

$$\inf_{k \geq 1} \mathbf{P}\{\widehat{X}_k \notin K\} \leq \frac{\varepsilon}{2}.$$

Since the coefficients b_k, σ_k are Lipschitz continuous in space variables,

$$\mathbf{1}_{\{\|\varphi\| \leq \Pi\}} \widehat{X}_k = \mathbf{1}_{\{\|\varphi\| \leq \Pi\}} X_k \quad \mathbf{P}\text{-almost surely}$$

for all $k \geq 1$, this implies

$$\mathbf{P}\{X_k \notin K\} \leq \mathbf{P}\{\widehat{X}_k \notin K\} + \mathbf{P}\{\|\varphi\| > \Pi\} < \varepsilon$$

for any $k \geq 1$ and tightness of the set $\{\mathbf{P} \circ X_k^{-1}; k \geq 1\}$ follows. □

Corollary 2.6. *The set $\{\mathbf{P} \circ (X_k, W)^{-1}; k \geq 1\}$ is a tight set of probability measures on $\mathcal{C}([0, T]; \mathbb{R}^m) \times \mathcal{C}([0, T]; \mathbb{R}^n)$.*

By the Prokhorov theorem, the set $\{\mathbf{P} \circ (X_k, W)^{-1}; k \geq 1\}$ is relatively (sequentially) compact in the weak topology of probability measures, so it contains a weakly convergent subsequence. Without loss of generality we may (and shall) assume that the sequence $\{\mathbf{P} \circ (X_k, W)^{-1}\}_{k=1}^\infty$ itself is weakly convergent. Let us set for brevity $\tilde{\mathbf{P}}_k = \mathbf{P} \circ (X_k, W)^{-1}, k \geq 1$, and denote the weak limit of $\{\tilde{\mathbf{P}}_k\}_{k=1}^\infty$ by $\tilde{\mathbf{P}}_0$. Set further

$$U = \mathcal{C}_m \times \mathcal{C}_n, \quad \mathcal{U} = \text{Borel}(\mathcal{C}_m) \otimes \text{Borel}(\mathcal{C}_n),$$

and let (Y, B) be the process of projections on U , that is

$$(Y_t, B_t) : \mathcal{C}_m \times \mathcal{C}_n \longrightarrow \mathbb{R}^m \times \mathbb{R}^n, (h, g) \longmapsto (h(t), g(t)), \quad 0 \leq t \leq T.$$

Finally, let (\mathcal{U}_t) be the $\tilde{\mathbf{P}}_0$ -augmented canonical filtration of the process (Y, B) , that is,

$$\mathcal{U}_t = \sigma(\varrho_s Y, \varrho_s B) \cup \{N \in \mathcal{U}; \tilde{\mathbf{P}}_0(N) = 0\}, \quad 0 \leq t \leq T.$$

3. Identification of the Limit

In this section, we shall show that $((U, \mathcal{U}, (\mathcal{U}_t), \tilde{\mathbf{P}}_0), B, Y)$ is a weak solution to the problem (0.7). Toward this end, define

$$M_k = Y - Y(0) - \int_0^\cdot b_k(r, Y(r)) dr, \quad k \geq 0,$$

where we set $b_0 = b, \sigma_0 = \sigma$. The proof is an immediate consequence of the following four lemmas.

Lemma 3.1. *The process M_0 is an m -dimensional local (\mathcal{U}_t) -martingale on $(U, \mathcal{U}, \tilde{\mathbf{P}}_0)$.*

Lemma 3.2. *The process B is an n -dimensional (\mathcal{U}_t) -Wiener process on $(U, \mathcal{U}, \tilde{\mathbf{P}}_0)$.*

Lemma 3.3. *The process*

$$\|M_0\|^2 - \int_0^\cdot \|\sigma(r, Y(r))\|^2 dr$$

is a local (\mathcal{U}_t) -martingale on $(U, \mathcal{U}, \tilde{\mathbf{P}}_0)$.

Lemma 3.4. *The process*

$$M_0 \otimes B - \int_0^\cdot \sigma(r, Y(r)) dr$$

is an $\mathbb{M}_{m \times n}$ -valued local (\mathcal{U}_t) -martingale on $(U, \mathcal{U}, \tilde{\mathbf{P}}_0)$.

Proofs of these lemmas have an identical structure, so we prove only the first of them in detail, the other ones being treated only in a concise manner. In the course

of the proof, we shall need two easy results on continuity properties of the first entrance times as functionals of paths. Let $V \geq 1$, for any $L \in \mathbb{R}_+$ define

$$\tau_L : \mathcal{C}_V \longrightarrow [0, T], f \longmapsto \inf\{t \geq 0; \|f(t)\| \geq L\}$$

(with a convention $\inf \emptyset = T$).

Lemma 3.5.

- (a) For any $f \in \mathcal{C}_V$, the function $L \mapsto \tau_L(f)$ is nondecreasing and left-continuous on \mathbb{R}_+ .
 (b) For each $L \in \mathbb{R}_+$, the mapping τ_L is lower semicontinuous. Moreover, τ_L is continuous at every point $f \in \mathcal{C}_V$ for which $\tau_\bullet(f)$ is continuous at L .

If $(Z_t)_{t \in [0, T]}$ is a continuous \mathbb{R}^V -valued stochastic process defined on a probability space $(G, \mathcal{G}, \mathbf{q})$, then $(\tau_L(Z))_{L \geq 0}$ is a stochastic process with nondecreasing left-continuous trajectories, whence we get

Lemma 3.6. *The set*

$$\{L \in \mathbb{R}_+; \mathbf{q}\{\tau_\bullet(Z) \text{ is not continuous at } L\} > 0\}$$

is at most countable.

Lemma 3.5 is proved (but not stated exactly in this form) in Jacod and Shiryaev [11], see Lemma VI.2.10 and Proposition VI.2.11 there. For Lemma 3.6, see [11, Lemma VI.3.12]. In Jacod and Shiryaev [11], τ_L is considered as a function on the Skorokhod space \mathbb{D} , in our case the proofs simplify further; they are recalled in the Appendix to keep the article self-contained.

Further, let us quote an useful result on weak convergence of measures (cf., e.g., [2, Proposition IX.5.7]).

Lemma 3.7. *Let $\{\nu_r\}_{r \geq 1}$ be a sequence of Borel probability measures on a metric space Θ converging weakly to a Borel probability measure ν_0 . Let $f : \Theta \longrightarrow \mathbb{R}$ be a bounded real function continuous at ν_0 -almost all points of Θ . Then*

$$\lim_{r \rightarrow \infty} \int_{\Theta} f \, d\nu_r = \int_{\Theta} f \, d\nu_0.$$

Proof of Lemma 3.1. The idea of the proof is simple: define processes

$$\mu_k = X_k - X_k(0) - \int_0^\cdot b_k(r, X_k(r))dr, \quad k \geq 1,$$

in analogy with the definition of M_k but using the solutions X_k to the problem (1.2) instead of the process Y . We shall prove: i) μ_k , $k \geq 1$, are local martingales, ii) M_k , $k \geq 1$, are local martingales with respect to the measure $\tilde{\mathbf{P}}_k$ due to the equality of laws $\tilde{\mathbf{P}}_k \circ (Y, B)^{-1} = \mathbf{P} \circ (X_k, W)^{-1}$, iii) M_0 is a local martingale as a limit of local martingales M_k .

First, as X_k solves (1.2),

$$\mu_k(t) = \int_0^t \sigma_k(r, X_k(r)) dW_r, \quad 0 \leq t \leq T,$$

and so μ_k is a local (\mathcal{F}_t) -martingale. Take an $L \in \mathbb{R}_+$, for the time being arbitrary. Obviously, $\tau_L(X_k)$ is a stopping time and $\mu_k(\cdot \wedge \tau_L(X_k))$ is a bounded process by (1.1) and the definition of τ_L , hence, $\mu_k(\cdot \wedge \tau_L(X_k))$ is a martingale.

Hereafter, times $s, t \in [0, T]$, $s \leq t$, and a continuous function

$$\gamma : \mathcal{C}([0, s]; \mathbb{R}^m) \times \mathcal{C}([0, s]; \mathbb{R}^n) \longrightarrow [0, 1]$$

will be fixed but otherwise arbitrary. Obviously, $\gamma(\varrho_s X_k, \varrho_s W)$ is a bounded \mathcal{F}_s -measurable function, hence,

$$\mathbf{E}\gamma(\varrho_s X_k, \varrho_s W)\mu_k(t \wedge \tau_L(X_k)) = \mathbf{E}\gamma(\varrho_s X_k, \varrho_s W)\mu_k(s \wedge \tau_L(X_k)) \quad (3.1)$$

by the martingale property of $\mu_k(\cdot \wedge \tau_L(X_k))$.

Note that the mapping

$$[0, T] \times \mathcal{C}_m \longrightarrow \mathbb{R}^m, \quad (u, h) \longmapsto h(u) - h(0) - \int_0^u b_k(r, h(r))dr$$

is continuous for any $k \geq 0$ due to the continuity of $b_k(r, \cdot)$, and the mapping

$$\mathcal{C}_m \longrightarrow [0, T] \times \mathcal{C}_m, \quad h \longmapsto (\zeta \wedge \tau_L(h), h)$$

is Borel for any $\zeta \in [0, T]$ fixed by Lemma 3.5(b), thus, also their superposition

$$H_k(\zeta, \cdot) : \mathcal{C}_m \longrightarrow \mathbb{R}^m, \quad h \longmapsto h(\zeta \wedge \tau_L(h)) - h(0) - \int_0^{\zeta \wedge \tau_L(h)} b_k(r, h(r))dr$$

is Borel. Consequently, the mapping

$$\mathcal{C}_m \times \mathcal{C}_n \longrightarrow \mathbb{R}^m, \quad (h, g) \longmapsto \gamma(\varrho_s h, \varrho_s g)H_k(\zeta, h)$$

is Borel. Since $\mu_k(\zeta \wedge \tau_L(X_k)) = H_k(\zeta, X_k)$, $k \geq 1$, and $M_k(\zeta \wedge \tau_L(Y)) = H_k(\zeta, Y)$, $k \geq 0$, we get

$$\mathbf{P} \circ [\gamma(\varrho_s X_k, \varrho_s W)\mu_k(\zeta \wedge \tau_L(X_k))]^{-1} = \tilde{\mathbf{P}}_k \circ [\gamma(\varrho_s Y, \varrho_s B)M_k(\zeta \wedge \tau_L(Y))]^{-1}$$

for all $k \geq 1$ by the definition of $\tilde{\mathbf{P}}_k$, which together with (3.1) implies

$$\tilde{\mathbf{E}}_k \gamma(\varrho_s Y, \varrho_s B)M_k(t \wedge \tau_L(Y)) = \tilde{\mathbf{E}}_k \gamma(\varrho_s Y, \varrho_s B)M_k(s \wedge \tau_L(Y)), \quad k \geq 1. \quad (3.2)$$

Now, suppose in addition that L is chosen so that

$$\tilde{\mathbf{P}}_0\{\tau_\bullet(Y) \text{ is continuous at } L\} = 1. \quad (3.3)$$

(Lemma 3.6 shows that such a choice is possible.) Then

$$\tilde{\mathbf{P}}_0\{(f, g) \in U; \tau_L(\cdot) \text{ is continuous at } f\} = 1$$

by Lemma 3.5(b) and the fact that Y is a canonical projection from U onto \mathcal{E}_m , so also

$$\tilde{\mathbf{P}}_0\{(f, g) \in U; H_0(\xi, \cdot) \text{ is continuous at } f\} = 1.$$

This implies that $\gamma(\varrho_s Y, \varrho_s B)H_0(\xi, Y)$ is a bounded function continuous $\tilde{\mathbf{P}}_0$ -almost everywhere on U for any ξ fixed. We may estimate

$$\begin{aligned} & \|\tilde{\mathbf{E}}_k \gamma(\varrho_s Y, \varrho_s B)H_k(\xi, Y) - \tilde{\mathbf{E}}_0 \gamma(\varrho_s Y, \varrho_s B)H_0(\xi, Y)\| \\ & \leq \|\tilde{\mathbf{E}}_k \gamma(\varrho_s Y, \varrho_s B)[H_k(\xi, Y) - H_0(\xi, Y)]\| \\ & \quad + \|\tilde{\mathbf{E}}_k \gamma(\varrho_s Y, \varrho_s B)H_0(\xi, Y) - \tilde{\mathbf{E}}_0 \gamma(\varrho_s Y, \varrho_s B)H_0(\xi, Y)\|. \end{aligned}$$

From Lemma 3.7 we obtain that

$$\lim_{k \rightarrow \infty} \tilde{\mathbf{E}}_k \gamma(\varrho_s Y, \varrho_s B)H_0(\xi, Y) = \tilde{\mathbf{E}}_0 \gamma(\varrho_s Y, \varrho_s B)H_0(\xi, Y).$$

Further,

$$\begin{aligned} & \|\tilde{\mathbf{E}}_k \gamma(\varrho_s Y, \varrho_s B)[H_k(\xi, Y) - H_0(\xi, Y)]\| \\ & \leq \tilde{\mathbf{E}}_k \|H_k(\xi, Y) - H_0(\xi, Y)\| \\ & = \tilde{\mathbf{E}}_k \left\| \int_0^{\xi \wedge \tau_L(Y)} [b_k(r, Y(r)) - b_0(r, Y(r))] dr \right\| \\ & = \tilde{\mathbf{E}}_k \mathbf{1}_{\{\tau_L(Y) > 0\}} \left\| \int_0^{\xi \wedge \tau_L(Y)} [b_k(r, Y(r)) - b_0(r, Y(r))] dr \right\| \\ & \leq \tilde{\mathbf{E}}_k \mathbf{1}_{\{\tau_L(Y) > 0\}} \int_0^{\xi \wedge \tau_L(Y)} \|b_k(r, Y(r)) - b_0(r, Y(r))\| dr \\ & \leq \tilde{\mathbf{E}}_k \mathbf{1}_{\{\tau_L(Y) > 0\}} \int_0^T \|b_k(r, Y(r \wedge \tau_L(Y))) - b_0(r, Y(r \wedge \tau_L(Y)))\| dr \\ & \leq \tilde{\mathbf{E}}_k \mathbf{1}_{\{\tau_L(Y) > 0\}} \int_0^T \sup_{\|z\| \leq L} \|b_k(r, z) - b_0(r, z)\| dr \\ & \leq \int_0^T \sup_{\|z\| \leq L} \|b_k(r, z) - b_0(r, z)\| dr, \end{aligned}$$

as $\|Y(r \wedge \tau_L(Y))\| \leq L$ on the set $\{\tau_L(Y) > 0\}$. Since $b_k(r, \cdot) \rightarrow b_0(r, \cdot)$ locally uniformly on \mathbb{R}^m for every $r \in [0, T]$ and

$$\sup_{\|z\| \leq L} \|b_k(r, z) - b_0(r, z)\| \leq 2K_*(2 + L)$$

by (0.6) and (1.1), we have

$$\lim_{k \rightarrow \infty} \int_0^T \sup_{\|z\| \leq L} \|b_k(r, z) - b_0(r, z)\| dr = 0$$

by the dominated convergence theorem, hence,

$$\lim_{k \rightarrow \infty} \tilde{E}_k \gamma(\varrho_s Y, \varrho_s B) H_k(\zeta, Y) = \tilde{E}_0 \gamma(\varrho_s Y, \varrho_s B) H_0(\zeta, Y)$$

for any $\zeta \in [0, T]$. Therefore,

$$\tilde{E}_0 \gamma(\varrho_s Y, \varrho_s B) M_0(t \wedge \tau_L(Y)) = \tilde{E}_0 \gamma(\varrho_s Y, \varrho_s B) M_0(s \wedge \tau_L(Y)) \quad (3.4)$$

follows from (3.2). If $G \subseteq \mathcal{C}([0, s]; \mathbb{R}^m \times \mathbb{R}^n)$ is an arbitrary open set, then there exist continuous functions $g_l : \mathcal{C}([0, s]; \mathbb{R}^m \times \mathbb{R}^n) \rightarrow [0, 1]$ such that $g_l \nearrow \mathbf{1}_G$ on $\mathcal{C}([0, s]; \mathbb{R}^m \times \mathbb{R}^n)$ as $l \rightarrow \infty$. Therefore, using the Levi monotone convergence theorem we derive from (3.4) that

$$\tilde{E}_0 \mathbf{1}_G(\varrho_s Y, \varrho_s B) M_0(t \wedge \tau_L(Y)) = \tilde{E}_0 \mathbf{1}_G(\varrho_s Y, \varrho_s B) M_0(s \wedge \tau_L(Y)). \quad (3.5)$$

Further,

$$\{G \subseteq \mathcal{C}([0, s]; \mathbb{R}^m \times \mathbb{R}^n); G \text{ Borel and (3.5) holds for } \mathbf{1}_G\}$$

is a λ -system containing, as we have just shown, the system of all open sets in $\mathcal{C}([0, s]; \mathbb{R}^m \times \mathbb{R}^n)$ closed under finite intersections. Consequently, (3.5) holds for all Borel sets $G \subseteq \mathcal{C}([0, s]; \mathbb{R}^m \times \mathbb{R}^n)$, that is,

$$\tilde{E}_0 \mathbf{1}_A M_0(t \wedge \tau_L(Y)) = \tilde{E}_0 \mathbf{1}_A M_0(s \wedge \tau_L(Y))$$

holds for all $A \in \sigma(\varrho_s Y, \varrho_s B)$, thus, for all $A \in \mathcal{U}_s$. We see that $M_0(\cdot \wedge \tau_L(Y))$ is a (\mathcal{U}_t) -martingale, whenever $L \in \mathbb{R}_+$ satisfies (3.3). It remains to note that by Lemma 3.6 there exists a sequence $L_r \nearrow \infty$ such that

$$\tilde{P}_0 \{\tau_\bullet(Y) \text{ is continuous at } L_r \text{ for every } r \geq 1\} = 1.$$

As $\{\tau_{L_r}(Y)\}$ is plainly a localizing sequence of stopping times, we conclude that M_0 is a local (\mathcal{U}_t) -martingale on $(U, \mathcal{U}, \tilde{P}_0)$, as claimed. \square

Proof of Lemma 3.2. By our construction, $\mathbf{P} \circ W^{-1} = \tilde{\mathbf{P}}_k \circ B^{-1}$ for each $k \geq 1$, so also $\mathbf{P} \circ W^{-1} = \tilde{\mathbf{P}}_0 \circ B^{-1}$ and B is an n -dimensional Wiener process (with respect to its canonical filtration) on $(U, \mathcal{U}, \tilde{\mathbf{P}}_0)$. In particular, its tensor quadratic variation satisfies $\langle\langle B \rangle\rangle_t = tI$. Mimicking the procedure from the previous proof we may check easily that B is a local (\mathcal{U}_t) -martingale, hence an (\mathcal{U}_t) -Wiener process by the Lévy theorem. \square

Proof of Lemma 3.3. We know that $\mu_k, k \geq 1$, are local martingales and

$$\langle \mu_k \rangle = \left\langle \int_0^\cdot \sigma_k(r, X_k(r)) dW_r \right\rangle = \int_0^\cdot \|\sigma_k(r, X_k(r))\|^2 dr,$$

thus,

$$\|\mu_k\|^2 = \int_0^\cdot \|\sigma_k(r, X_k(r))\|^2 dr, \quad k \geq 1,$$

are continuous local martingales. For times $s \leq t$ and a function γ introduced in the proof of Lemma 3.1 we get

$$\begin{aligned} & \mathbf{E}\gamma(\varrho_s X_k, \varrho_s W) \left[\|\mu_k(t \wedge \tau_L(X_k))\|^2 - \int_0^{t \wedge \tau_L(X_k)} \|\sigma_k(r, X_k(r))\|^2 dr \right] \\ &= \mathbf{E}\gamma(\varrho_s X_k, \varrho_s W) \left[\|\mu_k(s \wedge \tau_L(X_k))\|^2 - \int_0^{s \wedge \tau_L(X_k)} \|\sigma_k(r, X_k(r))\|^2 dr \right]. \end{aligned} \quad (3.6)$$

Note that

$$\mathcal{E}_m \longrightarrow \mathbb{R}, h \longmapsto \|H_k(\xi, h)\|^2 - \int_0^{\xi \wedge \tau_L(h)} \|\sigma_k(r, h(r))\|^2 dr$$

is a Borel mapping for all $k \geq 0$ and $\xi \in [0, T]$. It can be seen easily that it suffices to check that

$$\mathcal{E}_m \longrightarrow \mathbb{R}, h \longmapsto \int_0^u \|\sigma_k(r, h(r))\|^2 dr$$

is a continuous mapping for any $u \in [0, T]$; this follows from the estimate

$$\begin{aligned} & \left| \int_0^u \|\sigma_k(r, h_1(r))\|^2 dr - \int_0^u \|\sigma_k(r, h_2(r))\|^2 dr \right| \\ & \leq \int_0^u \left\{ \|\sigma_k(r, h_1(r))\| + \|\sigma_k(r, h_2(r))\| \right\} \left| \|\sigma_k(r, h_1(r))\| - \|\sigma_k(r, h_2(r))\| \right| dr \\ & \leq K_* \left(4 + \|h_1\|_{\mathcal{E}_m} + \|h_2\|_{\mathcal{E}_m} \right) \int_0^u \|\sigma_k(r, h_1(r)) - \sigma_k(r, h_2(r))\| dr \end{aligned}$$

for $h_1, h_2 \in \mathcal{E}_m$, continuity of functions $\sigma_k(r, \cdot)$ and the dominated convergence theorem.

Hence, (3.6) yields

$$\begin{aligned} & \tilde{\mathbf{E}}_k \gamma(\varrho_s Y, \varrho_s B) \left[\|M_k(t \wedge \tau_L(Y))\|^2 - \int_0^{t \wedge \tau_L(Y)} \|\sigma_k(r, Y(r))\|^2 dr \right] \\ &= \tilde{\mathbf{E}}_k \gamma(\varrho_s Y, \varrho_s B) \left[\|M_k(s \wedge \tau_L(Y))\|^2 - \int_0^{s \wedge \tau_L(Y)} \|\sigma_k(r, Y(r))\|^2 dr \right]. \end{aligned}$$

Passing to the limit exactly in the same way as in the proof of Lemma 3.1 we obtain

$$\begin{aligned} & \tilde{\mathbf{E}}_0 \gamma(\varrho_s Y, \varrho_s B) \left[\|M_0(t \wedge \tau_L(Y))\|^2 - \int_0^{t \wedge \tau_L(Y)} \|\sigma_0(r, Y(r))\|^2 dr \right] \\ &= \tilde{\mathbf{E}}_0 \gamma(\varrho_s Y, \varrho_s B) \left[\|M_0(s \wedge \tau_L(Y))\|^2 - \int_0^{s \wedge \tau_L(Y)} \|\sigma_0(r, Y(r))\|^2 dr \right] \end{aligned}$$

provided that $L \in \mathbb{R}_+$ satisfies (3.3), and the proof may be completed easily. \square

Proof of Lemma 3.4. Since μ_k and W are continuous local martingales, the process $\mu_k \otimes W - \langle \mu_k, W \rangle$ is an $\mathbb{M}_{m \times n}$ -valued local martingale. Let us denote $\mu_k = (\mu_k^i)_{i=1}^m$,

$W = (W^j)_{j=1}^n$ and $\sigma_k = (\sigma_k^{ij})_{i=1, j=1}^m, n$. Then

$$\begin{aligned} \langle \mu_k^i, W^j \rangle &= \left\langle \sum_{l=1}^n \int_0^\cdot \sigma_k^{il}(r, X_k(r)) dW^l(r), W^j \right\rangle \\ &= \sum_{l=1}^n \int_0^\cdot \sigma_k^{il}(r, X_k(r)) d\langle W^l, W^j \rangle_r \\ &= \int_0^\cdot \sigma_k^{ij}(r, X_k(r)) dr, \end{aligned}$$

therefore,

$$\mu_k \otimes W - \int_0^\cdot \sigma_k(r, X_k(r)) dr \tag{3.7}$$

is an $\mathbb{M}_{m \times n}$ -valued local martingale. The process (3.7) stopped at $\tau_L(X_k, W)$ is bounded, hence it is a martingale and so

$$\begin{aligned} E\gamma(\varrho_s X_k, \varrho_s W) &\left[(\mu_k \otimes W)(t \wedge \tau_L(X_k, W)) - \int_0^{t \wedge \tau_L(X_k, W)} \sigma_k(r, X_k(r)) dr \right] \\ &= E\gamma(\varrho_s X_k, \varrho_s W) \left[(\mu_k \otimes W)(s \wedge \tau_L(X_k, W)) - \int_0^{s \wedge \tau_L(X_k, W)} \sigma_k(r, X_k(r)) dr \right], \end{aligned}$$

whenever $0 \leq s \leq t \leq T$ and γ is a continuous function as above. (Since $\mathcal{C}_m \times \mathcal{C}_n \cong \mathcal{C}_{m+n}$, it is clear how $\tau_L(f, g)$ is defined for $(f, g) \in \mathcal{C}_m \times \mathcal{C}_n$.) Now we may proceed as in the proof of Lemma 3.1. \square

Proof of Theorem 0.1. Lemmas 3.1–3.4 having been established, it is straightforward to prove that $((U, \mathcal{U}, (\mathcal{U}_t), \tilde{\mathbf{P}}_0), B, Y)$ is a weak solution of (0.7). Since $\tilde{\mathbf{P}}_0 \circ Y(0)^{-1} = \tilde{\mathbf{P}}_k \circ Y(0)^{-1} = \mathbf{P} \circ \varphi^{-1} = \nu$ by our construction, it remains only to show that

$$Y(t) = Y(0) + \int_0^t b(r, Y(r)) dr + \int_0^t \sigma(r, Y(r)) dB(r)$$

for any $t \in [0, T]$ $\tilde{\mathbf{P}}_0$ -almost surely, that is

$$M_0(t) = \int_0^t \sigma(r, Y(r)) dB(r) \quad \text{for all } t \in [0, T] \quad \tilde{\mathbf{P}}_0\text{-almost surely.} \tag{3.8}$$

Obviously, (3.8) is equivalent to

$$\left\langle M_0 - \int_0^\cdot \sigma(r, Y(r)) dB(r) \right\rangle_T = 0 \quad \tilde{\mathbf{P}}_0\text{-almost surely.} \tag{3.9}$$

We have

$$\begin{aligned} \left\langle M_0 - \int_0^\cdot \sigma(r, Y(r)) dB(r) \right\rangle_T &= \langle M_0 \rangle_T + \left\langle \int_0^\cdot \sigma(r, Y(r)) dB(r) \right\rangle_T \\ &\quad - 2 \sum_{i=1}^m \left\langle M_0^i, \sum_{j=1}^n \int_0^\cdot \sigma^{ij}(r, Y(r)) dB^j(r) \right\rangle_T \end{aligned}$$

$$\begin{aligned}
 &= \langle M_0 \rangle_T + \int_0^T \|\sigma(r, Y(r))\|^2 dr \\
 &\quad - 2 \sum_{i=1}^m \left\langle M_0^i, \sum_{j=1}^n \int_0^{\cdot} \sigma^{ij}(r, Y(r)) dB^j(r) \right\rangle_T.
 \end{aligned}$$

By Lemma 3.3,

$$\langle M_0 \rangle_T = \int_0^T \|\sigma(r, X(r))\|^2 dr,$$

and by Lemma 3.4 we obtain

$$\begin{aligned}
 \sum_{i=1}^m \sum_{j=1}^n \left\langle M_0^i, \int_0^{\cdot} \sigma^{ij}(r, Y(r)) dB^j(r) \right\rangle_T &= \sum_{i=1}^m \sum_{j=1}^n \int_0^T \sigma^{ij}(r, Y(r)) d\langle M_0^i, B^j \rangle_r \\
 &= \sum_{i=1}^m \sum_{j=1}^n \int_0^T (\sigma^{ij}(r, Y(r)))^2 dr \\
 &= \int_0^T \|\sigma(r, Y(r))\|^2 dr,
 \end{aligned}$$

hence (3.9) holds true. □

Remark 3.1. If the coefficients b and σ of the Equation (0.7) are defined on $\mathbb{R}_+ \times \mathbb{R}^m$ and satisfy the assumptions of Theorem 0.1 there, then there exists a weak solution to (0.7) defined for all times $t \geq 0$. The proof remains almost the same, only its part concerning tightness requires small modifications. However, it suffices to realize that the space $\mathcal{C}(\mathbb{R}_+; \mathbb{R}^V)$ equipped with the topology of locally uniform convergence is a Polish space whose Borel σ -algebra is generated by the projections $f \mapsto f(t)$, $t \geq 0$ and whose closed subset K is compact if and only if $\{q_T f; f \in K\}$ is a compact subset of $\mathcal{C}([0, T]; \mathbb{R}^V)$ for all $T \geq 0$.

Remark 3.2. Tracing the proofs in Section 3, we can check easily that, unlike the proof of tightness in Section 2, they depend only on the following properties of the coefficients $b = b_0$, $\sigma = \sigma_0$ and their approximations b_k, σ_k :

1. the functions $b_k(r, \cdot), \sigma_k(r, \cdot)$ are continuous on \mathbb{R}^m for any $r \in [0, T]$ and $k \geq 0$,
2. $b_k(r, \cdot) \rightarrow b(r, \cdot), \sigma_k(r, \cdot) \rightarrow \sigma(r, \cdot)$ locally uniformly on \mathbb{R}^m as $k \rightarrow \infty$ for any $r \in [0, T]$,
3. the functions b_k, σ_k are locally bounded uniformly in $k \geq 0$, i.e.

$$\sup_{k \geq 0} \sup_{r \in [0, T]} \sup_{\|z\| \leq L} \{ \|b_k(r, z)\| \vee \|\sigma_k(r, z)\| \} < \infty$$

for each $L \geq 0$.

As a consequence, Theorem 0.1 remains valid if existence of a suitable Lyapunov function is supposed instead of the linear growth hypothesis. One proceeds as in the proof of Theorem 0.1, approximating the coefficients b and σ by bounded continuous functions that satisfy the same Lyapunov estimate as b and σ . However, the proof of tightness is more technical, although no fundamentally new ideas are needed; details may be found in a companion article [7].

References

1. Barlow, M.T. 1982. One dimensional stochastic differential equation with no strong solution. *Journal of the London Mathematical Society* 2 26(2):335–347.
2. Bourbaki, N. 2004. *Integration II*. Springer, Berlin.
3. Brzeźniak, Z., and Ondreját, M. 2007. Strong solutions to stochastic wave equations with values in Riemannian manifolds. *Journal of Functional Analysis* 253(2):449–481.
4. Da Prato, G., and Zabczyk, J. 1992. *Stochastic Equations in Infinite Dimensions*. Cambridge University Press, Cambridge, UK.
5. Dudley, R.M. 2002. *Real Analysis and Probability*. Cambridge University Press, Cambridge, UK.
6. Gątarek, D., and Gołdys, B. 1994. On weak solutions of stochastic equations in Hilbert spaces. *Stochastics and Stochastics Reports* 46(1):41–51.
7. Hofmanová, M., and Seidler, J. to appear. On weak solutions of stochastic differential equations II. Dissipative drifts.
8. Ikeda, N., and Watanabe, S. 1989. *Stochastic Differential Equations and Diffusion Processes*, 2nd ed. North-Holland, Amsterdam.
9. Itô, K. 1946. On a stochastic integral equation. *Proceedings of the Japan Academy* 22(1–4):32–35.
10. Ito, K. 1951. On stochastic differential equations. *Memoirs of the American Mathematical Society* 4:1–51.
11. Jacod, J., and Shiryaev, A.N. 2003. *Limit Theorems for Stochastic Processes*, 2nd ed. Springer, Berlin.
12. Karatzas, I., and Shreve, S. 1988. *Brownian Motion and Stochastic Calculus*. Springer, New York.
13. Lieb, E.H., and Loss, M. 2001. *Analysis*, 2nd ed. American Mathematical Society, Providence, RI.
14. Ondreját, M. 2010. Stochastic nonlinear wave equations in local Sobolev spaces. *Electronic Journal of Probability* 15(33):1041–1091.
15. Samko, S.S., Kilbas, A.A., and Marichev, A.A. 1993. *Fractional Integrals and Derivatives*, Gordon and Breach, Yverdon.
16. Skorokhod, A.V. 1961. On existence and uniqueness of solutions to stochastic diffusion equations. *Sibirskii Matematicheskii Zhurnal* 2(1):129–137. (in Russian)
17. Skorokhod, A.V. 1962. On stochastic differential equations. In: *Proceedings of the 6th All-Union Conference on Probability Theory and Mathematical Statistics*, GIPNL Litovskoi SSR, Vil'nyus, pp. 159–168. (in Russian)
18. Stroock, D.W., and Varadhan, S.R.S. 1979. *Multidimensional Diffusion Processes*. Springer, Berlin.

Appendix

To keep this article self-contained as much as possible, we provide here proofs of Lemmas 3.5 and 3.6.

Proof of Lemma 3.5. Choose $f \in \mathcal{C}_V$ and $L > 0$ arbitrarily. The function $K \mapsto \tau_K(f)$ is obviously nondecreasing, hence it has a left-hand limit at the point L and

$$\lim_{K \rightarrow L^-} \tau_K(f) \leq \tau_L(f). \quad (\text{A.1})$$

If $\|f\|_{\mathcal{C}_V} < L$ then $\|f\|_{\mathcal{C}_V} < L - \delta$ for some $\delta > 0$ and, thus, $\tau_L(f) = T = \tau_K(f)$ for all $K \in [L - \delta, L]$, so we may assume that $\|f\|_{\mathcal{C}_V} \geq L$. Then $\|f(\tau_K(f))\| \geq K$ for all

$K \in [0, L]$ and continuity of f yields

$$\|f\left(\lim_{K \rightarrow L^-} \tau_K(f)\right)\| = \lim_{K \rightarrow L^-} \|f(\tau_K(f))\| \geq \lim_{K \rightarrow L^-} K = L,$$

whence

$$\tau_L(f) \leq \lim_{K \rightarrow L^-} \tau_K(f),$$

which together with (A.1) proves the statement (a).

To prove (b), take an arbitrary sequence $\{f_r\}$ in \mathcal{C}_V such that $f_r \rightarrow f$ uniformly on $[0, T]$ as $r \rightarrow \infty$. Let $\varepsilon > 0$, then

$$\max_{[0, \tau_L(f) - \varepsilon]} \|f\| < L,$$

so there exists $r_0 \in \mathbb{N}$ such that

$$\max_{[0, \tau_L(f) - \varepsilon]} \|f_r\| < L$$

for all $r \geq r_0$, thus, $\tau_L(f_r) \geq \tau_L(f) - \varepsilon$ for all $r \geq r_0$. Since ε was arbitrary,

$$\liminf_{r \rightarrow \infty} \tau_L(f_r) \geq \tau_L(f),$$

that is, τ_L is lower semicontinuous at the point f .

Finally, assume in addition that $\tau_\bullet(f)$ is continuous at the point L . If $\tau_L(f) = T$ then

$$T = \tau_L(f) \leq \liminf_{r \rightarrow \infty} \tau_L(f_r) \leq \limsup_{r \rightarrow \infty} \tau_L(f_r) \leq T$$

(note that τ_L is $[0, T]$ -valued) and we are done. So assume that $\tau_L(f) < T$ and take an arbitrary $\varepsilon > 0$ satisfying $\tau_L(f) + \varepsilon < T$. By continuity, a $K > L$ may be found such that $\tau_K(f) < \tau_L(f) + \varepsilon$. Consequently,

$$\max_{[0, \tau_L(f) + \varepsilon]} \|f\| \geq K > L,$$

thus,

$$\max_{[0, \tau_L(f) + \varepsilon]} \|f_r\| \geq L$$

for all r sufficiently large, that is, $\tau_L(f_r) \leq \tau_L(f) + \varepsilon$ for all r sufficiently large, which implies

$$\limsup_{r \rightarrow \infty} \tau_L(f_r) \leq \tau_L(f)$$

and τ_L is upper semicontinuous at f . □

Proof of Lemma 3.6. Here we follow Jacod and Shiryaev [11] closely. First, note that for any given $u > 0$ \mathbf{q} -almost any trajectory of $\tau_{\bullet}(Z)$ has only finitely many jumps of size greater than u . For brevity, set

$$\Delta\tau_L(Z) = \lim_{M \rightarrow L^+} \tau_M(Z) - \tau_L(Z)$$

and define recursively random times

$$\Sigma_0(u) = 0, \Sigma_p(u) = \inf\{L > \Sigma_p(u); \Delta\tau_L(Z) > u\}, \quad u > 0, p \in \mathbb{N}.$$

Plainly, the set

$$\{L \geq 0; \mathbf{q}\{\Sigma_p(u) = L\} > 0\}$$

is at most countable for any $p \in \mathbb{N}$ and $u > 0$, hence it only remains to note that

$$\{L \geq 0; \mathbf{q}\{\Delta\tau_L(Z) > 0\} > 0\} = \bigcup_{p=0}^{\infty} \bigcup_{r=1}^{\infty} \{L \geq 0; \mathbf{q}\{\Sigma_p(r^{-1}) = L\} > 0\}. \quad \square$$