

## ON WEIGHTED SOBOLEV SPACES

SENG-KEE CHUA

**ABSTRACT.** We study density and extension problems for weighted Sobolev spaces on bounded  $(\epsilon, \delta)$  domains  $\mathcal{D}$  when a doubling weight  $w$  satisfies the weighted Poincaré inequality on cubes near the boundary of  $\mathcal{D}$  and when it is in the Muckenhoupt  $A_p$  class locally in  $\mathcal{D}$ . Moreover, when the weights  $w_i(x)$  are of the form  $\text{dist}(x, M_i)^{\alpha_i}$ ,  $\alpha_i \in \mathbb{R}$ ,  $M_i \subset \mathcal{D}$  that are doubling, we are able to obtain some extension theorems on  $(\epsilon, \infty)$  domains.

**1. Introduction.** Recently there has been quite a number of works related to weighted Sobolev spaces. For example, Kufner [23] studied various properties of weighted Sobolev spaces on certain domains  $\mathcal{D}$  for weights arising from  $\text{dist}(\cdot, M)$  with  $M \subset \partial \mathcal{D}$ . Also, Brown and Hinton [2], [3], [4] and Gutierrez and Wheeden [20] obtained weighted Sobolev interpolation inequalities. Meanwhile, the author [9], [11], [13] has studied the extension and restriction problems on weighted Sobolev spaces. In this paper, we would like to improve some results in [9]. Namely, we will study density problems and extension problems on weighted Sobolev spaces. Note that some of our results overlap some of those in [23] and [17].

By a weight  $w$ , we mean a non-negative locally integrable function on  $\mathbb{R}^n$ . By abusing notation, we will also write  $w$  for the measure induced by  $w$ . Sometimes we write  $dw$  to denote  $w dx$ . We always assume  $w$  is doubling, by which we mean  $w(2Q) \leq Cw(Q)$  for every cube  $Q$ , where  $2Q$  denotes the cube with the same center as  $Q$  and twice its edglength. All cubes in this paper are assumed to be closed and with edges parallel to the axes. By  $w \in A_p$ , we mean  $w$  satisfies the Muckenhoupt  $A_p$  condition, *i.e.*,

$$\frac{1}{|Q|} \left( \int_Q w dx \right)^{1/p} \left( \int_Q w^{-1/(p-1)} dx \right)^{1/p'} \leq C \quad \text{when } 1 < p < \infty, \text{ and}$$

$$\frac{1}{|Q|} \int_Q w(x) dx \leq C \operatorname{ess\,inf}_{x \in Q} w(x) \quad \text{when } p = 1,$$

for all cubes  $Q$  in  $\mathbb{R}^n$ . Note that  $w$  is doubling when it is in  $A_p$ . Moreover, when  $\mathcal{D}$  is an open set, we will write  $w \in A_p^{\text{loc}}(\mathcal{D})$  if for any cube  $Q_0 \subset \mathcal{D}$ , there exists  $C_{Q_0} > 0$  such that

$$\frac{1}{|Q|} w(Q \cap Q_0)^{1/p} \left( \int_{Q \cap Q_0} w^{\frac{-1}{p-1}}(x) dx \right)^{1/p'} \leq C_{Q_0} \quad \text{when } 1 < p < \infty, \text{ and}$$

---

Received by the editors March 14, 1993; revised May 1995.

AMS subject classification: 46E35.

Key words and phrases: Poincaré inequalities,  $A_p$  weights, power weights, doubling, locally  $A_p$  weights,  $(\epsilon, \delta)$  and  $(\epsilon, \infty)$  domains.

© Canadian Mathematical Society, 1996.

$$\frac{w(Q \cap Q_0)}{|Q|} \leq C_{Q_0} \operatorname{ess\,inf}_{x \in Q \cap Q_0} w(x) \quad \text{when } p = 1,$$

for all cubes  $Q$  in  $\mathbb{R}^n$ .<sup>1</sup>

Let  $\mathcal{D}$  be an open set in  $\mathbb{R}^n$ . If  $\alpha$  is a multi-index,  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in \mathbb{Z}_+^n$ , we will denote  $\sum_{j=1}^n \alpha_j$  by  $|\alpha|$  and  $D^\alpha = (\frac{\partial}{\partial x_1})^{\alpha_1} \dots (\frac{\partial}{\partial x_n})^{\alpha_n}$ . By  $\alpha \geq \beta$ , we mean  $\alpha_j \geq \beta_j$  for all  $1 \leq j \leq n$ . Moreover we write  $\alpha > \beta$  if  $\alpha \geq \beta$  and  $\alpha \neq \beta$ . We denote by  $\nabla$  the vector  $(\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \dots, \frac{\partial}{\partial x_n})$  and by  $\nabla^m$  the vector of all possible  $m^{\text{th}}$  order derivatives for  $m \in \mathbb{N}$ . A locally integrable function  $f$  on  $\mathcal{D}$  (we will write  $f \in L^1_{\text{loc}}(\mathcal{D})$ ) has a weak derivative of order  $\alpha$  if there is a locally integrable function (denoted by  $D^\alpha f$ ) such that

$$\int_{\mathcal{D}} f(D^\alpha \varphi) \, dx = (-1)^{|\alpha|} \int_{\mathcal{D}} (D^\alpha f) \varphi \, dx$$

for all  $C^\infty$  functions  $\varphi$  with compact support in  $\mathcal{D}$  (we will write  $\varphi \in C^\infty_0(\mathcal{D})$ ).

If  $1 < p < \infty$ ,  $p'$  is always equal to  $p/(p - 1)$  and  $p' = \infty$  when  $p = 1$ .  $Q$  will always be a cube and  $l(Q)$  will be its edglength. Following [22], we say that two cubes touch if a face of one cube is contained in a face of the other. For  $1 \leq p < \infty, k \in \mathbb{N}$ , and any weight  $w, L^p_{w,k}(\mathcal{D})$  and  $E^p_{w,k}(\mathcal{D})$  are the spaces of functions having weak derivatives of all orders  $\alpha, |\alpha| \leq k$ , and satisfying

$$\|f\|_{L^p_{w,k}(\mathcal{D})} = \sum_{0 \leq |\alpha| \leq k} \|D^\alpha f\|_{L^p_w(\mathcal{D})} = \sum_{0 \leq |\alpha| \leq k} \left( \int_{\mathcal{D}} |D^\alpha f|^p \, dw \right)^{1/p} < \infty,$$

and

$$\|f\|_{E^p_{w,k}(\mathcal{D})} = \sum_{|\alpha|=k} \|D^\alpha f\|_{L^p_w(\mathcal{D})} < \infty$$

respectively. Moreover, in the case when  $w \equiv 1$ , we will denote  $L^p_{w,k}(\mathcal{D})$  and  $E^p_{w,k}(\mathcal{D})$  by  $L^p_k(\mathcal{D})$  and  $E^p_k(\mathcal{D})$  respectively. Also, let  $\hat{E}^p_{w,k}(\mathcal{D})$  be the factor space  $E^p_{w,k}(\mathcal{D})/\mathcal{P}_{k-1}$  where  $\mathcal{P}_l$  is the subspace of polynomials of degree not greater than  $l$ . By  $f \in L^p_{w,1,\text{loc}}(\mathcal{D})$ , we mean  $f \in L^p_{w,1}(K^\circ)$  for all compact sets  $K$  in  $\mathcal{D}$ .

Let  $\mathcal{D}$  be an open connected set. It is easy to see that  $L^p_{w,k}(\mathcal{D})$  is a Banach space when  $w^{-1/p} \in L^p_{\text{loc}}(\mathcal{D})$  [17]. Moreover, the author [9] prove that  $\hat{E}^p_{w,k}(\mathcal{D})$  is a Banach space when  $w \in A_p$ . Note that it is just a weighted version of Theorem 1.1.13.1 in [26]. We will show that indeed the following is true.

**THEOREM 1.1.** *Let  $1 \leq p < \infty$  and let  $w$  be a doubling weight. If  $w^{-1/p} \in L^p_{\text{loc}}(\mathcal{D})$  then  $\hat{E}^p_{w,k}(\mathcal{D})$  is a Banach space for any connected open set  $\mathcal{D}$ .*

**DEFINITION 1.2.** An open set  $\mathcal{D}$  is an  $(\varepsilon, \delta)$  domain if for all  $x, y \in \mathcal{D}, |x - y| < \delta$ , there exists a rectifiable curve  $\gamma$  connecting  $x, y$  such that  $\gamma$  lies in  $\mathcal{D}$  and

$$(1.1) \quad l(\gamma) < \frac{|x - y|}{\varepsilon}$$

<sup>1</sup> Note that  $w \in A^{\text{loc}}_p(\mathcal{D}) \Rightarrow w \in A^K_p$  for all compact sets  $K \subset \mathcal{D}$  in the notation of Wolff [35].

$$(1.2) \quad d(z, \partial \mathcal{D}) > \frac{\varepsilon|x-z||y-z|}{|x-y|} \quad \forall z \in \gamma.$$

Here  $l(\gamma)$  is the length of  $\gamma$  and  $d(z, \partial \mathcal{D})$  is the distance between  $z$  and the boundary of  $\mathcal{D}$ . Moreover, we will write  $d(Q, S) = \inf_{x \in Q, y \in S} |x - y|$ ,  $d(Q) = d(Q, \partial \mathcal{D})$  and  $d(z) = d(\{z\}, \partial \mathcal{D})$ .

In 1981, P. Jones [22] extended a famous extension theorem on Lipschitz domains to  $(\varepsilon, \delta)$  domains.

**THEOREM 1.3.** *If  $\mathcal{D}$  is a connected  $(\varepsilon, \delta)$  domain and  $1 \leq p \leq \infty$ , then  $C^\infty(\mathbb{R}^n) \cap L^p_k(\mathcal{D})$  is dense in  $L^p_k(\mathcal{D})$  and  $L^p_k(\mathcal{D})$  has a bounded extension operator. Moreover the norm of the extension operator depends only on  $\varepsilon, \delta, k, p, \text{rad}(\mathcal{D})$ , and the dimension  $n$ .*

Furthermore he proved that

**THEOREM 1.4.** *If  $\mathcal{D}$  is an  $(\varepsilon, \infty)$  domain in  $\mathbb{R}^n$ , then  $E^1_1(\mathcal{D})$  has a bounded extension operator, i.e., there exists  $\Lambda: E^1_1(\mathcal{D}) \rightarrow E^1_1(\mathbb{R}^n)$  such that  $\Lambda|_{\mathcal{D}} = f$  a.e. and  $\|\Lambda\|$  is bounded.*

Recently, the author extended Theorems 1.3 and 1.4 to weighted Sobolev spaces when the weight is in  $A_p$  [9]. In this paper, we will extend these results further by relaxing the  $A_p$  assumption on the weight  $w$  to the following conditions on a bounded  $(\varepsilon, \delta)$  domain  $\mathcal{D}$ :

- $w$  is doubling on  $\mathbb{R}^n$ ,  $w \in A^{\text{loc}}_p(\mathcal{D})$
- $w$  satisfies a local Poincaré inequality on  $\mathcal{D}$ .

Indeed, we prove that

**THEOREM 1.5.** *Let  $\mathcal{D}$  be a bounded  $(\varepsilon, \delta)$  domain. Let  $1 \leq p < \infty$  and let  $w$  be a doubling weight such that  $w \in A^{\text{loc}}_p(\mathcal{D})$ . Suppose further that*

$$(1.3) \quad \|f - f_{Q,w}\|_{L^p_w(Q)} \leq C(A)l(Q)\|\nabla f\|_{L^p_w(Q)} \quad \forall f \in L^p_{w,1,\text{loc}}(\mathcal{D})$$

for all cubes  $Q \subset \mathcal{D}$  near  $\partial \mathcal{D}$  such that  $Ad(Q) \leq l(Q) \leq d(Q)/A$ ,  $A > 0$  where  $f_{Q,w} = \int_Q f dw / w(Q)$ . Then given any  $f \in L^p_{w,k}(\mathcal{D})$  (resp.  $E^p_{w,k}(\mathcal{D})$ ) and  $\eta > 0$ , there exists  $f_\eta \in C^\infty(\mathbb{R}^n)$  such that

$$\|f - f_\eta\|_{L^p_{w,k}(\mathcal{D})} < \eta \quad (\text{resp. } \|\nabla^k(f - f_\eta)\|_{L^p_w(\mathcal{D})} < \eta).$$

Moreover, with the help of [11, Theorems 1.1 and 1.2] and the previous theorem, we show that:

**THEOREM 1.6.** *Let  $\mathcal{D}$  be a bounded  $(\varepsilon, \delta)$  domain. Let  $1 \leq p < \infty$  and  $w$  a doubling weight. If  $w \in A^{\text{loc}}_p(\mathcal{D})$ ,  $w^{-1/p} \in L^p_{\text{loc}}(\mathbb{R}^n)$  and (3.3) holds, then there exists an extension operator  $\Lambda$  on  $L^p_{w,k}(\mathcal{D})$  such that*

$$\|\Lambda f\|_{L^p_{w,k}(\mathbb{R}^n)} \leq C\|f\|_{L^p_{w,k}(\mathcal{D})}.$$

Moreover, if in addition that  $\mathcal{D}$  is a bounded  $(\varepsilon, \infty)$  domain, then there exists an extension operator  $\Lambda'$  on  $E_{w,k}^p(\mathcal{D})$  such that

$$\|\nabla^k \Lambda' f\|_{L_w^p(\mathbb{R}^n)} \leq C \|\nabla^k f\|_{L_w^p(\mathcal{D})}.$$

REMARK 1.7. (a) Let  $M \subset \partial \mathcal{D}$  and  $1 \leq p < \infty$ . It is easy to see that if  $w(x) = \text{dist}(x, M)^\alpha$ ,  $\alpha \in \mathbb{R}$ , then it follows from the non-weighted Poincaré inequality that

$$(1.4) \quad \|f - f_Q\|_{L_w^p(Q)} \leq C l(Q) \|\nabla f\|_{L_w^p(Q)} \quad \forall f \in L_{w,1,\text{loc}}^p(\mathcal{D})$$

for all cubes  $Q$  with  $l(Q)$  comparable to  $d(Q)$ . Moreover, it is clear that  $w \in A_p^{\text{loc}}(\mathcal{D})$ . Hence it follows from Theorem 1.5 that  $C^\infty(\mathbb{R}^n) \cap L_{w,k}^p(\mathcal{D})$  is dense in  $L_{w,k}^p(\mathcal{D})$  when  $w(x) = \text{dist}(x, M)^\alpha$  is doubling (note that (1.4) implies (1.3)). Thus when  $w$  is doubling and  $\mathcal{D}$  is a bounded  $(\varepsilon, \delta)$  domain, we obtain those density theorems in [23].

(b) Furthermore, if  $w(x) = s(\text{dist}(x, M))$  where  $s$  is a positive and continuous function on positive real numbers that satisfies certain properties described in Kufner [23] or [17], similar conclusion can be obtained by Theorem 1.5 if we know that  $w$  is doubling.

(c) We do not know exactly when will the weights  $w$  defined as above will be doubling. However, in the case that  $M$  is just a finite subset of  $\partial \mathcal{D}$ , it is easy to see that  $\text{dist}(x, M)^\alpha$  is doubling if and only if  $\alpha > -n$ . For more details, refer to [15].

REMARK 1.8. (a) Let  $w$  be as in Remark 1.7. If in addition that  $w^{-1/p} \in L_{\text{loc}}^{p'}(\mathbb{R}^n)$ , then we can apply Theorem 1.6 to get extension operator for  $L_{w,k}^p(\mathcal{D})$  or  $E_{w,k}^p(\mathcal{D})$ . This overlaps some results in [17].

(b) The assumption that  $w^{-1/p} \in L_{\text{loc}}^{p'}(\mathbb{R}^n)$  in Theorem 1.6 is somewhat too strong. Indeed, we need only to assume that  $w^{-1/p} \in L^{p'}(\mathcal{D})$ . For the details, see [10]. Note that when  $\mathcal{D}$  is a bounded  $(\varepsilon, \infty)$  domain,  $w \in A_p^{\text{loc}}(\mathcal{D})$  and (3.3) holds, it follows from [14, Corollary 1.5] that  $f \in E_{w,k}^p(\mathcal{D})$  if and only if  $f \in L_{w,k}^p(\mathcal{D})$ .

Finally, when the weights are of the form as in Remark 1.7(a), we are able to obtain extension theorems similar to Theorems 1.4 and 1.5 in [9]; see Remark 4.3.

ACKNOWLEDGEMENT. The author is grateful to the reviewer for his suggestions to improve the presentation of the paper and the proof of Lemma 2.5.

**2. Preliminaries.** In what follows,  $C$  denotes various positive constants, they may differ even in a same string of estimates. Moreover, sometimes, we will use  $C(\alpha, \beta, \dots)$  instead of  $C$  to emphasize that the constant is depending on  $\alpha, \beta, \dots$ . Following [22], we say that two cubes touch if a face of one cube is contained in a face of the other. In particular, the union of two touching cubes of equal size is a rectangle.

First, let us state a theorem on polynomials.

THEOREM 2.1 ([9, LEMMA 2.3]). *Let  $F, Q$  be cubes such that  $F \subset Q$  and  $|F| > \gamma|Q|$ . If  $w$  is a doubling weight,  $1 \leq q < \infty$ , and  $p$  is a polynomial of degree  $m$ , then*

$$\|p\|_{L_w^q(E)} \leq C(\gamma, m, n, w) \left( \frac{w(E)}{w(F)} \right)^{1/q} \|p\|_{L_w^q(F)}$$

for all measurable sets  $E \subset Q$ .

Next, the following lemma is indeed a special case of a result in [12].

LEMMA 2.2 ([12, THEOREM 2.1]). *Let  $f$  be a measurable function on  $\mathbb{R}^n$  and let  $w$  be a doubling weight. Also, let  $1 \leq p \leq \infty$ ,  $k \in \mathbb{N}$  and  $L > 0$ . For each cube  $Q$  in  $\mathbb{R}^n$ , let  $a(f, Q)$  be a polynomial of degree  $k$  associated to  $f$  on  $Q$  for each cube  $Q$ . Suppose that  $\{Q_i\}_{i=0}^l$  is a sequence of cubes such that  $Q_i \cap Q_{i+1}$  contains a cube  $Q^i$  with  $|Q^i| \geq L \max\{|Q_i|, |Q_{i+1}|\}$  for each  $i = 0, 1, \dots, l - 1$ . Then*

$$(2.1) \quad \|f - a(f, Q_0)\|_{L^p_w(Q_0)} \leq C \sum_i \|f - a(f, Q_i)\|_{L^p_w(Q_i)}$$

where  $C$  depends only on  $L, l, w, k, p$  and the dimension  $n$ .

PROOF OF THEOREM 1.1. We will modify the proof of [26, Theorem 1.1.13.1] and [9, Theorem 4.9].

Let  $Q_0$  be a Whitney cube in  $\mathcal{D}$  and let  $\{\Omega_i\}$  be a sequence of open connected sets which are the interiors of finite unions of touching Whitney cubes of  $\mathcal{D}$  (when  $\mathcal{D} = \mathbb{R}^n$ , just take  $\{\Omega_i\}$  be a sequence of nested cubes) such that  $Q_0 \subset \Omega_i, \bar{\Omega}_i \subset \Omega_{i+1}, \cup_i \Omega_i = \mathcal{D}$ .

Given any Cauchy sequence  $\{u_j\} \subset E^p_{w,k}(\mathcal{D})$ , and any cube  $Q$  in  $\mathcal{D}$ , let  $P(Q, u_j)$  be the unique polynomial of degree  $< k$  such that  $\int_Q D^\beta(u_j - P(Q, u_j)) dx = 0$  for all  $|\beta| < k$ . Since

$$\begin{aligned} \|D^\beta(u_j - u_l - P(Q, u_j - u_l))\|_{L^1(Q)} &= \|D^\beta(u_j - u_l - (P(Q, u_j) - P(Q, u_l)))\|_{L^1(Q)} \\ &\leq C|Q|^{k-|\beta|} \|\nabla^k(u_j - u_l)\|_{L^1(Q)} \end{aligned}$$

for all cubes  $Q$  in  $\mathcal{D}$  by the unweighted Poincaré inequality, we have if  $P_j = P(Q_0, u_j)$ ,

$$\begin{aligned} \|D^\beta(u_j - u_l - (P_j - P_l))\|_{L^1(\Omega_i)} &\leq C(\Omega_i) \|\nabla^k(u_j - u_l)\|_{L^1(\Omega_i)} \\ &\leq C(\Omega_i) \|\nabla^k(u_j - u_l)\|_{L^p_w(\Omega_i)} \|w^{-1/p}\|_{L^{p'}(\Omega_i)} \\ &\leq C(\Omega_i) \|\nabla^k(u_j - u_l)\|_{L^p_w(\Omega_i)}, \end{aligned}$$

by the previous lemma, the Hölder inequality and the assumption on  $w$ . Hence if  $v_j = u_j - P_j$ , then  $\{D^\beta v_j\}$  is a Cauchy sequence in  $L^1(\Omega_i)$  for any  $i$  and  $|\beta| \leq k$ . Thus it follows that for each  $i$  and  $\beta$  with  $|\beta| < k$ , there exists  $h_{i,\beta} \in L^1(\Omega_i)$  such that  $\|D^\beta v_j - h_{i,\beta}\|_{L^1(\Omega_i)} \rightarrow 0$  as  $j \rightarrow \infty$ . (When  $|\beta| = k$ , clearly there exists  $h_\beta \in L^p_w(\mathcal{D})$  such that  $\|D^\beta v_j - h_\beta\|_{L^p_w(\mathcal{D})} \rightarrow 0$  as  $L^p_w(\mathcal{D})$  is complete.) Using subsequences, it is clear that  $h_{i+1,\beta} = h_{i,\beta}$  a.e. on  $\Omega_i$ . If we define  $h_\beta$  on  $\mathcal{D}$  by setting  $h_\beta = h_{i,\beta}$  on  $\Omega_i$ , it follows that for each compact set  $K \subset \mathcal{D}$  we have  $h_\beta \in L^1(K)$  and  $D^\beta v_j \rightarrow h_\beta$  in  $L^1(K)$  for all  $|\beta| \leq k$  (for  $|\beta| = k$ , just use the Hölder inequality and the fact that  $w^{-1/p} \in L^{p'}_{loc}(\mathcal{D})$ ). Thus if  $\varphi \in C^\infty_0(\mathcal{D})$ , then (let us write  $h_\beta$  as  $h$  when  $\beta = 0$ )

$$\int_{\mathcal{D}} h D^\beta \varphi dx = \lim_{j \rightarrow \infty} \int_{\mathcal{D}} v_j D^\beta \varphi dx = \lim_{j \rightarrow \infty} (-1)^{|\beta|} \int_{\mathcal{D}} (D^\beta v_j) \varphi dx = (-1)^{|\beta|} \int_{\mathcal{D}} h_\beta \varphi dx.$$

Hence  $D^\beta h = h_\beta$  exists. Moreover  $D^\alpha h = \lim D^\alpha u_j$  when  $|\alpha| = k$  since  $D^\alpha u_j = D^\alpha v_j$ . This completes the proof of the theorem.

COROLLARY 2.3. Let  $\mathcal{D}$  be an open connected set, let  $\{u_j\}$  be a Cauchy sequence in  $E_{w,k}^p(\mathcal{D})$  and let  $u$  be a function in  $E_{w,k}^p(\mathcal{D})$  such that

$$\|\nabla^k(u_j - u)\|_{L_w^p(\mathcal{D})} \rightarrow 0.$$

Then there exists a sequence of polynomials  $\{P_j\}$  of degree  $< k$  with  $u_j - P_j \rightarrow u$  in  $L^1(K)$  for all compact sets  $K$  in  $\mathcal{D}$ .

PROOF. By the previous proof, we know  $v_j = u_j - P_j \rightarrow h$  in  $L^1(K)$  for each compact set  $K$  in  $\mathcal{D}$ , and  $\nabla^k u_j \rightarrow \nabla^k h$  in  $L_w^p(\mathcal{D})$ . Since also  $\nabla^k u_j \rightarrow \nabla^k u$  in  $L_w^p(\mathcal{D})$ , we see that  $\nabla^k(u - h) = 0$ , so  $u - h = P$  for some polynomial  $P$  of degree  $< k$ . Thus  $u_j - P_j + P \rightarrow h + P = u$  in  $L^1(K)$ .

Now we will state a well-known lemma; see for example, Theorem III.2 in [31].

LEMMA 2.4. Let  $k(x)$  be nonnegative and integrable on  $\mathbb{R}^n$  and suppose  $k(x)$  depends only on  $|x|$  and decreases as  $|x|$  increases. Then for all non-negative measurable functions  $f$ ,

$$\sup_{t>0} |f * k_t(x)| \leq C \|k\|_{L^1(\mathbb{R}^n)} Mf(x)$$

with  $C$  independent of  $x, f$  and  $k$ . Here  $k_t(y) = t^{-n}k(y/t)$  and  $Mf$  is the Hardy-Littlewood maximal function of  $f$ .

Similar to  $A_p$  weights [27], [18], we have the following results.

LEMMA 2.5. Let  $1 < p < \infty$ , and  $w \in A_p^{\text{loc}}(\mathcal{D})$ . Then

$$(2.2) \quad \|M(f\chi_K)\|_{L_w^p(K)} \leq C_K \|f\|_{L_w^p(K)}$$

for all compact sets  $K$  in  $\mathcal{D}$ .

PROOF. We will only prove it for the case when  $w$  is doubling.<sup>2</sup> It suffices to show that (2.2) holds for  $K = Q_0$  for all cubes  $Q_0$  in  $\mathcal{D}$  such that  $3Q_0 \subset \mathcal{D}$ .

Let  $\mu = \chi_{3Q_0}$ ,  $\nu = \chi_{3Q_0}w$  and  $\tilde{w} = \chi_{Q_0}w$ . Note that  $(\frac{d\mu}{d\nu})^{p'-1} = \chi_{3Q_0}w^{1-p'}$ . Let  $M_\mu h(x) = \sup \int_F h(y) d\mu / \mu(F)$  where the supremum is taken over all cubes  $F$  containing  $x$ . Let  $Q$  be any cube. We will now show that  $\nu, \tilde{w}$  and  $M_\mu$  satisfies the  $S_p$  condition [29]. Let  $x \in Q_0 \cap Q$ , we now consider two cases:

CASE (i)  $Q \subset 3Q_0$ . Then there exists a cube  $F \subset Q$  and  $x \in F$  such that  $M_\mu \chi_{Q \cap 3Q_0} w^{1-p'}(x) \leq C \int_F w^{1-p'} dy / |F|$ . Thus

$$(2.3) \quad \begin{aligned} M_\mu(\chi_{Q \cap 3Q_0} w^{1-p'})(x) &\leq C \left( \frac{1}{|F|} \int_F w dy \right)^{1-p'} \quad \text{since } w \in A_p^{\text{loc}}(\mathcal{D}) \\ &= C \left( \frac{1}{w(F)} \int_F w^{-1} w dy \right)^{p'-1} \leq C (M_w(\chi_{Q \cap 3Q_0} w^{-1})(x))^{p'-1}. \end{aligned}$$

<sup>2</sup> The idea of this proof was provided by the referee.

Hence

$$\begin{aligned}
 \int_Q [M_\mu(\chi_{Q \cap 3Q_0} w^{1-p'})(x)]^p d\tilde{w}(x) &= \int_{Q \cap Q_0} [M_\mu(\chi_{Q \cap 3Q_0} w^{1-p'})(x)]^p w(x) dx \\
 &\leq C \int_{Q \cap 3Q_0} [M_w(\chi_{Q \cap 3Q_0} w^{-1})(x)]^{p'} w(x) dx \\
 &\leq \int_{Q \cap 3Q_0} (w^{-1})^{p'} w(x) dx \\
 (2.4) \qquad \qquad \qquad &= \int \chi_Q \left( \frac{d\mu}{dv} \right)^{p'-1} v(x) dx
 \end{aligned}$$

since  $w$  is doubling<sup>3</sup> on  $\mathbb{R}^n$ ; see for example [21].

CASE (ii).  $Q$  is not contained in  $3Q_0$ . Since there is nothing to prove when  $Q \cap Q_0 = \emptyset$ , we may assume  $3^n |Q \cap 3Q_0| \geq |3Q_0|$ . Thus

$$\begin{aligned}
 \int_Q [M_\mu(\chi_{Q \cap 3Q_0} w^{1-p'})(x)]^p d\tilde{w}(x) &\leq \int_{Q_0} [M_\mu(\chi_{3Q_0} w^{1-p'})(x)]^p w(x) dx \\
 &\leq C \int_{3Q_0} w^{1-p'}(x) dx \leq \int_{Q \cap 3Q_0} w^{1-p'}(x) dx
 \end{aligned}$$

since  $w \in A_p^{\text{loc}}(\mathcal{D})$ . Hence by Theorem A of [29], we have

$$\begin{aligned}
 \|M(\chi_{Q_0} f)\|_{L_w^p(Q_0)} &= \|M_\mu(\chi_{Q_0} f)\|_{L_w^p(Q_0)} = \|M_\mu(\chi_{Q_0} f)\|_{L_w^p(\mathbb{R}^n)} \\
 &\leq \|\chi_{Q_0} f\|_{L_w^p(\mathbb{R}^n)} = C \|f\|_{L_w^p(Q_0)}
 \end{aligned}$$

and hence (2.2) holds for  $K = Q_0$ .

LEMMA 2.6. Let  $1 \leq p < \infty$ ,  $w \in A_p^{\text{loc}}(\mathcal{D})$  and let  $\xi \in C_0^\infty$  be a non-negative decreasing radial function with support in  $\{x \in \mathbb{R}^n : |x| \leq 1\}$  and  $\int \xi(x) dx = 1$ . Then for  $f \in L_w^p(\mathcal{D})$ ,  $f * \xi_t \rightarrow f$  in  $L_w^p(K)$  as  $t \rightarrow 0$  for all compact sets  $K$  in  $\mathcal{D}$ . Moreover, if  $f \in L_{w,k}^p(\mathcal{D})$  then  $f * \xi_t \rightarrow f$  in  $L_{w,k}^p(K)$  for all compact sets  $K$  in  $\mathcal{D}$ .

PROOF. When  $1 < p < \infty$ , it follows from Lemmas 2.4 and 2.5 and the Lebesgue dominated convergence theorem. Now if  $p = 1$ , given any compact set  $K \subset \mathcal{D}$ , let us first choose a continuous function  $g$  such that

$$(2.5) \qquad \qquad \qquad \|f - g\|_{L_w^1(K^s)} \leq \eta$$

where  $K^s = \{x + y : |y| \leq s, x \in K\}$ , and  $s$  is chosen so that  $K^s \subset \mathcal{D}$ . Next since  $g$  is continuous, there exists  $L > 0$  such that  $|g(x) - g(y)| < \eta$  for  $x, y \in K^s$  and  $|x - y| \leq L$ . Next if  $sB = \{x \in \mathbb{R}^n : |x| \leq s\}$  and  $0 < t < s$ ,

$$\begin{aligned}
 \|f * \xi_t - f\|_{L_w^1(K)} &\leq \int_K \int_{sB} |f(x - y) - f(x)| \xi_t(y) dy w(x) dx \\
 &\leq \int_K \int_{sB} |f(x - y) - g(x - y)| \xi_t(y) dy w(x) dx \\
 &\quad + \int_K \int_{sB} |g(x - y) - g(x)| \xi_t(y) dy w(x) dx \\
 &\quad + \int_K \int_{sB} |g(x) - f(x)| \xi_t(y) dy w(x) dx \\
 &= I + II + III.
 \end{aligned}$$

<sup>3</sup> However, the theorem can be proved without assuming  $w$  is doubling i.e., assuming only  $w \in A_p^{\text{loc}}(\mathcal{D})$ .

However,  $II \leq w(K)\eta$  when  $0 < t < s \leq L$  and

$$III = \int_K |g(x) - f(x)|w(x) dx \leq \eta$$

by (2.5). Finally, note that

$$\begin{aligned} I &\leq \int_K \int_{K^s} |f(y) - g(y)|\xi_t(x - y) dy w(x) dx \\ &\leq \int_{K^s} \int_K \xi_t(x - y)w(x) dx |f(y) - g(y)| dy \\ &\leq C \int_{K^s} M(w\chi_K)(y) |f(y) - g(y)| dy \\ &\leq C \|f - g\|_{L^1_w(K^s)} \leq C(K)\eta. \end{aligned}$$

Lemma 2.6 now follows from the fact that  $D^\alpha(f * \xi_t) = (D^\alpha f) * \xi_t$ .<sup>4</sup>

**THEOREM 2.7.** *Let  $1 \leq p < \infty$  and  $w \in A_p^{loc}(\mathcal{D})$ . Then for all compact sets  $K$  in  $\mathcal{D}$ ,*

$$(2.6) \quad \|f - a(f, Q)\|_{L^p_w(Q)} \leq C(K)l(Q)\|\nabla f\|_{L^p_w(Q)}$$

for all  $f \in L^p_{w,1,loc}(\mathcal{D})$  and cube  $Q \subset K$  where  $a(f, Q) = \int_Q f dx / |Q|$  or  $\int_Q f dw / w(Q)$ .

**PROOF.** Let  $K$  be any compact set in  $\mathcal{D}$ . First, note that it suffices to show that (2.6) holds with  $a(f, Q) = f_Q = \int_Q f dx / |Q|$ . However,

$$|f(x) - f_Q| \leq \frac{1}{|Q|} \int_Q |f(x) - f(y)| dy \leq C \int_Q \frac{|\nabla f(y)|}{|x - y|^{n-1}} dy$$

for  $x \in Q, f \in C^\infty(\mathbb{R}^n)$  (see [33, Proposition 4.2]). Hence if  $f \in C^\infty(\mathbb{R}^n)$  it suffices to show that

$$(2.7) \quad \left\| \int_Q \frac{g(y)}{|\cdot - y|^{n-1}} dy \right\|_{L^p_w(Q)} \leq C(K)l(Q)\|g\|_{L^p_w(Q)}$$

for all cubes  $Q \subset K$ . However, in the case  $1 < p < \infty$ , (2.7) is just a consequence of Lemma 2.5. Moreover, the case  $p = 1$  follows immediately from the fact that  $w \in A_1^{loc}(\mathcal{D})$ . Finally, with the help of Lemma 2.6, by similar argument as the proof of Theorem 4.3 in [9], our assertion follows.

Next we will state a theorem which is similar to [26, Theorem 1.1.2.1] and [9, Theorem 4.2]. Since it can be proved by very similar method as the proof of [9, Theorem 4.2] with the help of Lemma 2.6 and Theorem 2.7, we will omit the proof.

**THEOREM 2.8.** *Let  $\mathcal{D}$  be any open set in  $\mathbb{R}^n$  and let  $1 \leq p < \infty, w \in A_p^{loc}(\mathcal{D})$ . If  $f \in E^p_{w,k}(\mathcal{D})$ , then*

$$\int_K |D^\gamma f|^p dw < \infty \quad \text{for all compact sets } K \subset \mathcal{D}, \forall 0 \leq |\gamma| \leq k.$$

<sup>4</sup> For the case  $p = 1$ , indeed we just modify the proof of Lemma 8 in [28].



3. **Density theorems.** Let  $\mathcal{D}$  be an  $(\varepsilon, \delta)$  domain, we will decompose  $\mathcal{D} = \cup \mathcal{D}_\alpha$  into connected components and define

$$r = \text{rad}(\mathcal{D}) = \inf_{\alpha} \inf_{x \in \mathcal{D}_\alpha} \sup_{y \in \mathcal{D}_\alpha} |x - y|.$$

We will assume  $r > 0$  in most cases. Then for any  $x \in \mathcal{D}$ , there is a point  $y$  in the same component with  $|x - y| \geq \frac{3r}{4}$ . Note that we always have  $r > 0$  when  $\mathcal{D}$  is an  $(\varepsilon, \infty)$  domain since  $\mathcal{D}$  is then connected.

Let us recall that two cubes touch if a face of one cube is contained in a face of the other. In particular, the union of two touching cubes of equal size is a rectangle. A collection of cubes  $\{S_i\}_{i=0}^m$  is called a *chain* if  $S_i$  touches  $S_{i+1}$  for all  $i$ .

Next let us recall some properties of the cubes in the Whitney decomposition of an open set  $\mathcal{D}$  [31]. Since these properties are well-known, we will often make use of them without explicitly mentioning them.

$$\begin{aligned} l(Q) &= 2^{-k} \quad \text{for some } k \in \mathbb{Z}, \\ Q_1 \cap Q_2 &= \emptyset \quad \text{if } Q_1 \neq Q_2, \\ 1/4 \leq \frac{l(Q_1)}{l(Q_2)} &\leq 4 \quad \text{if } Q_1 \cap Q_2 \neq \emptyset, \\ 1 \leq \frac{d(Q)}{l(Q)} &\leq 4\sqrt{n}. \end{aligned}$$

The purpose of this section is to prove the density theorem.

PROOF OF THEOREM 1.5. Our proof is similar to that of [22] and [9]. Let  $\rho = 2^{-m}$ ,  $m \in \mathbb{Z}_+$ . Let  $\mathcal{W}_1$  be the Whitney decomposition of  $\mathcal{D}$ . Define

$$\begin{aligned} \mathfrak{R}' &= \{\text{dyadic cubes } R \text{ with edgelenh } \rho, R \subset \mathcal{D}\} \text{ and} \\ \mathfrak{R} &= \{R \in \mathfrak{R}' : R \subset S \text{ for some } S \in \mathcal{W}_1, l(S) \geq 32n^3\rho/\varepsilon\}. \end{aligned}$$

Moreover, for each  $R \in \mathfrak{R}$  let  $\tilde{R}, \check{R}$  be cubes concentric with  $R$  with sides parallel to the axes and  $l(\tilde{R}) = 1281n^4\rho/\varepsilon^2$  and  $l(\check{R}) = 2562n^4\rho/\varepsilon^2$ . For  $s > 0$ , let  $\mathcal{D}_s = \{x \in \mathcal{D} : d(x) \geq s\}$ . First, let us make the following two observations.

- (I)  $\mathcal{D} \subset \cup_{R \in \mathfrak{R}} \tilde{R}$  provided  $\text{rad}(\mathcal{D}) > 0$  and  $\rho$  is small enough.
- (II) Let  $\mathcal{D}$  be an  $(\varepsilon, \delta)$  domain with  $\text{rad}(\mathcal{D}) > 0$  and let  $s = 3203n^5\rho/\varepsilon^3 < \delta$ .

Then for all  $R_0, R_j \in \mathfrak{R}$  with  $\tilde{R}_0 \cap \tilde{R}_j \neq \emptyset$  and  $\check{R}_0 \cap (\mathcal{D} \setminus \mathcal{D}_{2s}) \neq \emptyset$ , there exists a chain  $G_{0,j} = \{R_0 = S_1, S_2, \dots, S_m = R_j\}$  in  $\mathfrak{R}'$  connecting  $R_0, R_j$  with  $m \leq C$  that depends only on  $\varepsilon, \delta$  and  $n$ , and  $\cup G_{0,j} \subset \mathcal{D} \setminus \mathcal{D}_{3s}$ ,  $d(\cup G_{0,j}) \geq 20n^2\rho$ .

(I) is first stated in [22] without proof. Nevertheless, the reader can refer to the proof of Theorem 6.1 in [9]. A similar conclusion as (II) can indeed be found in [22, Lemma 4.1] or [9]. However, since (II) is slightly stronger than the conclusion in [22] or [9], we will prove it.

First note that since  $d(R_0, R_j) \leq \sqrt{n}(2561n^4\rho/\varepsilon^2) < \delta$ , there exists  $\gamma$  connecting  $R_0, R_j$  which satisfies (1.1) and (1.2). Next if  $z \in \gamma$ , we will show that  $d(z, \mathcal{D}_{3s}) > \sqrt{n}\rho$ .

First, we have

$$d(z, R_0) \leq l(\gamma) < d(R_0, R_j)/\varepsilon \leq 2561n^5\rho/\varepsilon^3, \\ d(R_0, (\mathcal{D}_{2s})^c) \leq \sqrt{n}(640n^4\rho/\varepsilon^2) \leq 640n^5\rho/\varepsilon^2$$

as  $\tilde{R}_0 \cap (\mathcal{D}_{2s})^c \neq \emptyset$ . Moreover,

$$d(R_0, \mathcal{D}_{3s}) \geq d((\mathcal{D}_{2s})^c, \mathcal{D}_{3s}) - d(R_0, (\mathcal{D}_{2s})^c) - \sqrt{nl}(R_0) \\ \geq 3203n^5\rho/\varepsilon^3 - 640n^5\rho/\varepsilon^2 - \sqrt{n}\rho \\ \geq 2562n^5\rho/\varepsilon^3.$$

Next, without loss of generality, we may assume that  $d(z, R_0) \leq d(z, R_j)$ . We now consider two cases:

CASE (i).  $d(z, R_0) \leq 42n^2\rho/\varepsilon$ . Then  $d(z) \geq 32n^3\rho/\varepsilon - 42n^2\rho/\varepsilon \geq 22n^2\rho/\varepsilon$ . (Note that we may restrict ourself to the case  $n \geq 2$ .)

CASE (ii).  $d(z, R_0) > 42n^2\rho/\varepsilon$ . Then by (1.2),

$$d(z) \geq \frac{\varepsilon d(z, R_0)d(z, R_j)}{d(R_0, R_j)} \geq 21n^2\rho.$$

Finally let us note that an appropriate subcollection of  $\{R \in \mathfrak{R}' : R \cap \gamma \neq \emptyset\}$  will provide us the required chain. Moreover,  $m \leq C$  as  $l(\gamma) \leq d(R_0, R_j)/\varepsilon$ .

Now, given  $f \in L^p_{w,k}(\mathcal{D})$ , we will let  $P_j = P(R_j)$  be the unique polynomial of degree  $k - 1$  such that

$$\int_{R_j} D^\alpha (f - P(R_j)) dw = 0, \quad 0 \leq |\alpha| \leq k - 1.$$

Next let  $R_0, R_j \in \mathfrak{R}$ ,  $R_0, R_j$  be as in (II). Suppose that  $G_{0,j}$  is the chain connecting  $R_0, R_j$  guaranteed by (II). If  $P_0 = P(R_0)$  and  $P_j = P(R_j)$ , similar to the proof of [9, Lemma 6.3], by the triangle inequality, (1.3), Lemma 2.2 and the fact that  $\varepsilon^3 d(R)/10000n^5 \leq l(R) \leq 20n^2 d(R)$  for all  $R \in \cup G_{0,j}$ , we can show that

$$(3.1) \quad \|D^\alpha(P_0 - P_j)\|_{L^p_{w,k}(R_0)} \leq C\rho^{k-|\alpha|} \|\nabla^k f\|_{L^p_{w,k}(\cup G_{0,j})} \quad \forall 0 \leq |\alpha| \leq k$$

where  $C$  is independent of  $f, R_0, R_j$  and  $\rho$ .

Next given  $\eta > 0$ , let us choose  $s > 0$  such that  $\|f\|_{L^p_{w,k}(\mathcal{D} \setminus \mathcal{D}_{3s})} \leq \eta$ . We then choose  $\psi \in C^\infty$  such that  $\chi_{\mathcal{D}_{2s}} \leq \psi \leq \chi_{\mathcal{D}_s}$  and  $|D^\alpha \psi| \leq C(\alpha)s^{-|\alpha|}$ .

Recall that by Lemma 2.6, there exists  $\xi \in C_0^\infty$  such that  $\int \xi dx = 1$  and

$$\|f - f * \xi_t\|_{L^p_{w,k}(\mathcal{D}_s)} \rightarrow 0 \quad \text{as } t \rightarrow 0 \text{ for } f \in L^p_{w,k}(\mathcal{D}), \text{ where } \xi_t(x) = t^{-n} \xi\left(\frac{x}{t}\right).$$

Thus we can choose  $0 < t < s/2$  such that

$$(3.2) \quad \|D^\alpha(f - f * \xi_t)\|_{L^p_{w,k}(\mathcal{D}_s)} = \|D^\alpha f - (D^\alpha f) * \xi_t\|_{L^p_{w,k}(\mathcal{D}_s)} \leq \eta s^{k-|\alpha|}, \quad 0 \leq |\alpha| \leq k.$$

For each  $R_j \in \mathfrak{R}$ , let us choose  $\varphi_j \in C^\infty$  with  $0 \leq \varphi_j \leq \chi_{\tilde{R}_j}$  such that  $\sum_{R_j \in \mathfrak{R}} \varphi_j \equiv 1$  on  $\bigcup_{R_j \in \mathfrak{R}} \tilde{R}_j$ ,  $0 \leq \sum_{R_j \in \mathfrak{R}} \varphi_j \leq 1$  and  $|D^\alpha \varphi_j| \leq C \varrho^{-|\alpha|}$ .

Fixing  $t$  and  $s$ , let  $g_0 = \sum_{R_j \in \mathfrak{R}} P_j \varphi_j$ ,  $g_1 = g_0(1 - \psi)$  and  $g_2 = (f * \xi_t)\psi$ . Then clearly  $g_0, g_1, g_2 \in C^\infty(\mathbb{R}^n)$ . We now show that  $\|f - (g_1 + g_2)\|_{L^p_{w,k}(\mathcal{D})} \leq C\eta$ . First, we will show that  $\|f - (g_1 + g_2)\|_{L^p_{w,k}(\mathcal{D}_{2s})} \leq C\eta$ . Let us note that since  $g_1 \equiv 0$  on  $\mathcal{D}_{2s}$  and  $g_2 = f * \xi_t$  on  $\mathcal{D}_{2s}$ , for  $|\alpha| \leq k$  we have

$$\|D^\alpha(f - (g_1 + g_2))\|_{L^p_w(\mathcal{D}_{2s})} = \|D^\alpha(f - f * \xi_t)\|_{L^p_w(\mathcal{D}_{2s})} \leq C\eta \quad \text{by (3.2)}.$$

Next write

$$\begin{aligned} D^\alpha(f - (g_1 + g_2)) &= D^\alpha(\psi(f - f * \xi_t)) + D^\alpha((1 - \psi)(f - g_0)) \\ &= \sum_{\beta \leq \alpha} C_{\alpha,\beta} D^{\alpha-\beta} \psi D^\beta(f - f * \xi_t) + \sum_{\beta \leq \alpha} C_{\alpha,\beta} D^{\alpha-\beta} (1 - \psi) D^\beta(f - g_0) \\ &= A + B. \end{aligned}$$

Since  $|D^{\alpha-\beta} \psi| \leq C s^{-|\alpha-\beta|}$ ,  $0 \leq \beta \leq \alpha$  and  $\psi \equiv 0$  on  $(\mathcal{D}_s)^c$ , we have  $\|A\|_{L^p_w(\mathcal{D} \setminus \mathcal{D}_{2s})} \leq C\eta$  by (3.2).

To complete the proof, we need only to prove that  $\|B\|_{L^p_w(\mathcal{D} \setminus \mathcal{D}_{2s})} \leq C\|\nabla^k f\|_{L^p_w(\mathcal{D} \setminus \mathcal{D}_{3s})}$ . To this end, first note that if  $\tilde{R}_0 \cap (\mathcal{D} \setminus \mathcal{D}_{2s}) \neq \emptyset$ ,  $\tilde{R}_0 \cap \tilde{R}_j \neq \emptyset$  then by the triangle inequality and (3.1),

$$\begin{aligned} \sum_{R_j \in \mathfrak{R}} \|D^\beta((P_0 - P_j)\varphi_j)\|_{L^p_w(R_0)} &\leq C \sum_{\tilde{R}_0 \cap \tilde{R}_j \neq \emptyset} \sum_{\gamma \leq \beta} l(R_0)^{-|\gamma|} \|D^{\beta-\gamma}(P_0 - P_j)\|_{L^p_w(R_0)} \\ (3.3) \qquad \qquad \qquad &\leq C \sum_{\tilde{R}_0 \cap \tilde{R}_j \neq \emptyset} \varrho^{k-|\beta|} \|\nabla^k f\|_{L^p_w(\cup G_{0j})}. \end{aligned}$$

Also, note that

$$(3.4) \quad |D^\beta(f - g_0)| = \left| D^\beta\left(f - \sum_{R_j \in \mathfrak{R}} P_j \varphi_j\right) \right| \leq |D^\beta(f - P_0)| + \left| D^\beta \sum_{R_j \in \mathfrak{R}} (P_0 - P_j)\varphi_j \right|.$$

We now consider two cases:

CASE (i).  $\beta < \alpha$ . Then  $D^{\alpha-\beta}(1 - \psi) = 0$  on  $\mathcal{D} \setminus \mathcal{D}_s$  and hence

$$\begin{aligned} &\|D^{\alpha-\beta}(1 - \psi)D^\beta(f - g_0)\|_{L^p_w(\mathcal{D} \setminus \mathcal{D}_{2s})} \\ &\leq C s^{-|\alpha-\beta|p} \sum_{R_0 \in \mathfrak{R}, R_0 \cap (\mathcal{D}_s \setminus \mathcal{D}_{2s}) \neq \emptyset} [\varrho^{k-|\beta|} \|\nabla^k f\|_{L^p_w(R_0)}]^p \\ &\quad + C s^{-|\alpha-\beta|p} \sum_{R_0 \in \mathfrak{R}, R_0 \cap (\mathcal{D}_s \setminus \mathcal{D}_{2s}) \neq \emptyset} \sum_{\tilde{R}_0 \cap \tilde{R}_j \neq \emptyset} [\varrho^{k-|\beta|} \|\nabla^k f\|_{L^p_w(\cup G_{0j})}]^p \end{aligned}$$

by (3.4) and (3.3) since  $\mathcal{D}_s \setminus \mathcal{D}_{2s} \subset \bigcup_{R_0 \in \mathfrak{R}} R_0$ . Next note that  $\|\sum_{R_0 \in \mathfrak{R}} \sum_{\tilde{R}_i \cap \tilde{R}_0 \neq \emptyset} \chi_{\cup G_{0j}}\|_{L^\infty} \leq C$  where  $C$  is independent of  $\varrho$ . Moreover by (II), if  $R_0 \cap (\mathcal{D}_s \setminus \mathcal{D}_{2s}) \neq \emptyset$ ,  $\tilde{R}_i \cap \tilde{R}_0 \neq \emptyset$ , then  $\cup G_{0j} \subset \mathcal{D} \setminus \mathcal{D}_{3s}$ , and in particular  $R_0 \subset \mathcal{D} \setminus \mathcal{D}_{3s}$ . Hence if  $\alpha > \beta$  (then  $|\beta| < k$ ),

$$\|D^{\alpha-\beta}(1 - \psi)D^\beta(f - g_0)\|_{L^p_w(\mathcal{D} \setminus \mathcal{D}_{2s})} \leq C s^{-|\alpha-\beta|} \varrho^{k-|\beta|} \|\nabla^k f\|_{L^p_w(\mathcal{D} \setminus \mathcal{D}_{3s})} \leq C\eta.$$

CASE (ii).  $\beta = \alpha$ . First observe that for each  $R_0 \in \mathfrak{R}$ ,  $\tilde{R}_0 \cap (\mathcal{D} \setminus \mathcal{D}_{2s}) \neq \emptyset$ , similar to (3.3) we have

$$\sum_{R_j \in \mathfrak{R}} \|D^\alpha((P_0 - P_j)\varphi_j)\|_{L_w^p(\tilde{R}_0)} \leq C \sum_{R_j \in \mathfrak{R}, \tilde{R}_0 \cap \tilde{R}_j \neq \emptyset} \varrho^{k-|\alpha|} \|\nabla^k f\|_{L_w^p(\cup G_{0j})}$$

by Lemma 2.1. Thus

$$\begin{aligned} \|D^\alpha \sum P_j \varphi_j\|_{L_w^p(\tilde{R}_0)} &\leq \|D^\alpha P_0\|_{L_w^p(\tilde{R}_0)} + \|D^\alpha \sum (P_j - P_0)\varphi_j\|_{L_w^p(\tilde{R}_0)} \\ &\leq C \|D^\alpha P_0\|_{L_w^p(R_0)} + C \sum_{R_j \in \mathfrak{R}, \tilde{R}_0 \cap \tilde{R}_j \neq \emptyset} \varrho^{k-|\alpha|} \|\nabla^k f\|_{L_w^p(\cup G_{0j})} \\ &\leq C \|D^\alpha f\|_{L_w^p(R_0)} + C \varrho^{k-|\alpha|} \|\nabla^k f\|_{L_w^p(R_0)} \\ &\quad + C \varrho^{k-|\alpha|} \sum_{R_j \in \mathfrak{R}, \tilde{R}_0 \cap \tilde{R}_j \neq \emptyset} \|\nabla^k f\|_{L_w^p(\cup G_{0j})}. \end{aligned}$$

Note that again by (II), if  $\tilde{R}_0 \cap (\mathcal{D} \setminus \mathcal{D}_{2s}) \neq \emptyset$  and  $\tilde{R}_0 \cap \tilde{R}_j \neq \emptyset$  then  $\cup G_{0j} \subset \mathcal{D} \setminus \mathcal{D}_{3s}$ , and in particular  $R_0 \subset \mathcal{D} \setminus \mathcal{D}_{3s}$ . Hence by the previous estimate,

$$\begin{aligned} \|D^\alpha(f - g_0)\|_{L_w^p(\mathcal{D} \setminus \mathcal{D}_{2s})}^p &\leq C \|D^\alpha f\|_{L_w^p(\mathcal{D} \setminus \mathcal{D}_{2s})}^p \\ &\quad + \sum_{R_0 \in \mathfrak{R}, \tilde{R}_0 \cap (\mathcal{D} \setminus \mathcal{D}_{2s}) \neq \emptyset} C \|D^\alpha \sum_{R_j \in \mathfrak{R}} P_j \varphi_j\|_{L_w^p(\tilde{R}_0)}^p \\ &\leq C \|D^\alpha f\|_{L_w^p(\mathcal{D} \setminus \mathcal{D}_{2s})}^p + C \|D^\alpha f\|_{L_w^p(\mathcal{D} \setminus \mathcal{D}_{3s})}^p \\ &\quad + C \varrho^{(k-|\alpha|)p} \|\nabla^k f\|_{L_w^p(\mathcal{D} \setminus \mathcal{D}_{3s})}^p \leq C \eta^p \end{aligned}$$

since  $\|\sum_{R_0 \in \mathfrak{R}} \sum_{\tilde{R}_j \cap \tilde{R}_0 \neq \emptyset} \chi_{\cup G_{0j}}\|_{L^\infty} < C$ . Thus  $\|D^\alpha(f - (g_1 + g_2))\|_{L_w^p(\mathcal{D} \setminus \mathcal{D}_{2s})} \leq C \eta$ .

Finally, if  $f \in E_{w,k}^p(\mathcal{D})$ , let us note that by Theorem 2.8, we have  $f \in L_{w,k}^p(\mathcal{D}_s)$ . We can then construct  $g_1 + g_2$  as before since (3.2) still hold. One can just check through the proof and see that  $g_1 + g_2$  satisfies our assertion.

**4. Extension theorems.** First, let us state an extension theorem from [11].

**THEOREM 4.1** ([11, THEOREMS 1.1 AND 1.2]). *Let  $\mathcal{D}$  be an  $(\varepsilon, \delta)$  domain. Let  $1 \leq p < \infty$  and let  $w$  be a doubling weight such that*

$$(4.1) \quad \|f - f_{Q,w}\|_{L_w^p(Q)} \leq C_0 l(Q) \|\nabla f\|_{L_w^p(Q)} \quad \forall f \in \text{Lip}_{\text{loc}}(\mathbb{R}^n)$$

for all cubes  $Q$  in  $\mathcal{D}$  where  $f_{Q,w} = \int_Q f dw / w(Q)$ . Then there exists an extension operator  $\Lambda$  on  $\mathcal{D}$  (i.e.,  $\Lambda f = f$  on  $\mathcal{D}$  a.e.) such that

$$\|\Lambda f\|_{L_{w,k}^p(\mathbb{R}^n)} \leq C \|f\|_{L_{w,k}^p(\mathcal{D})}$$

for all  $f \in \text{Lip}_{\text{loc}}^{k-1}(\mathbb{R}^n)$  ( $= \{f : D^\alpha f \in \text{Lip}_{\text{loc}}(\mathbb{R}^n)$  for all  $|\alpha| < k$ ) where  $C$  depends only on  $\varepsilon, \delta, \text{rad}(\mathcal{D}), p, w, k, C_0$  and  $n$ . Moreover, if  $\mathcal{D}$  is an  $(\varepsilon, \infty)$  domain, then there exists another extension operator  $\Lambda'$  on  $\mathcal{D}$  such that

$$\|\nabla^k \Lambda' f\|_{L_w^p(\mathbb{R}^n)} \leq C \|\nabla^k f\|_{L_w^p(\mathcal{D})}$$

for all  $f \in \text{Lip}_{\text{loc}}^{k-1}(\mathbb{R}^n)$  where  $C$  depends only on  $\varepsilon, p, w, k, C_0$  and  $n$ .

REMARK 4.2. Checking through the proof of Theorem 1.1 in [11], let us note that indeed we need only to assume (4.1) holds for all cubes  $Q$  near  $\partial \mathcal{D}$  such that  $l(Q)$  is comparable to  $d(Q)$  for the first part. However, for the second part, we need to assume in addition that  $\mathcal{D}$  is bounded.

With the help of the preceding theorem and the density theorem in the previous section, we can now prove our extension theorem.

PROOF OF THEOREM 1.6. First given  $f \in L^p_{w,k}(\mathcal{D})$ , by Theorem 1.5, there exists a sequence  $\{f_j\} \subset C^\infty(\mathbb{R}^n)$  such that  $f_j \rightarrow f$  in  $L^p_{w,k}(\mathcal{D})$ . Next since  $L^p_{w,k}(\mathbb{R}^n)$  is a Banach space, the first part of the theorem now follows from the preceding theorem (see Remark 4.2). Now let  $f \in E^p_{w,k}(\mathcal{D})$ . By Theorem 1.5 there exists  $\{f_j\} \subset C^\infty(\mathbb{R}^n)$  such that  $\|\nabla^k f_j - \nabla^k f\|_{L^p_w(\mathcal{D})} \rightarrow 0$ . Then  $\{\Lambda' f_j\}$  is a Cauchy sequence in  $E^p_{w,k}(\mathbb{R}^n)$  by the preceding theorem. Since  $E^p_{w,k}(\mathbb{R}^n)$  is complete by Theorem 1.1, there exists  $g \in E^p_{w,k}(\mathbb{R}^n)$  such that  $\nabla^k \Lambda' f_j \rightarrow \nabla^k g$  in  $L^p_w(\mathbb{R}^n)$ . Since  $\Lambda' f_j = f_j$  on  $\mathcal{D}$ , we obtain  $\|\nabla^k g - \nabla^k f\|_{L^p_w(\mathcal{D})} = 0$ . Hence there exists a polynomial  $P$  of degree  $< k$  such that  $g = f + P$  a.e. on  $\mathcal{D}$ . Define  $\Lambda' f = g - P$ . Then  $\Lambda' f = f$  a.e. on  $\mathcal{D}$ . Also,  $\nabla^k \Lambda' f = \nabla^k g$  and consequently  $\nabla^k \Lambda' f_j \rightarrow \nabla^k \Lambda' f$  in  $L^p_w(\mathbb{R}^n)$ . The proof of the theorem is now complete by passing to the limit.

REMARK 4.3. (a) Let  $\mathcal{D}$  be a bounded  $(\varepsilon, \infty)$  domain with  $r = \text{rad}(\mathcal{D})$  and let  $\Omega$  be a bounded open set containing  $\mathcal{D}$ . Let  $W_2$  be the collection of cubes in the Whitney decomposition of  $(\mathcal{D}^c)^o$  and define

$$W_3 = \left\{ Q \in W_2 : l(Q) \leq \frac{\varepsilon r}{16nL} \right\}, \quad L = 2^{-m}, \quad m \in \mathbb{Z}_+,$$

where  $L$  is chosen so that  $\Omega \subset (\cup_{Q \in W_3} Q) \cup \mathcal{D}$ . Finally, when the weights are of the form as in Remark 1.7(a), we have better extension theorems.

THEOREM 4.4. Let  $1 \leq p_i < \infty$ ,  $w_i = \text{dist}(x, M_i)^{\alpha_i}$ ,  $\alpha_i \in \mathbb{R}$ ,  $M_i \subset \partial \mathcal{D}$  such that  $w_i$  is doubling for  $i = 0, 1, \dots, N$ . Let  $\Omega$  be a bounded open set containing an  $(\varepsilon, \infty)$  domain  $\mathcal{D}$  and let  $L$  and  $r$  be defined as above. Suppose that  $k_i = 0$  for  $0 \leq i \leq N_1$ ,  $k_i = k > 0$  for  $N_2 < i \leq N$  and  $0 < k_i < k$  otherwise. Then there exist extension operators  $\Lambda$  and  $\Lambda'$  on  $\mathcal{D}$  such that

$$\begin{aligned} \|\Lambda f\|_{L^{p_i}_{w_i}(\mathbb{R}^n)} &\leq C_i \|f\|_{L^{p_i}_{w_i}(\mathcal{D})} \quad \text{for } 0 \leq i \leq N_1 \\ \|\nabla^{k_i} \Lambda f\|_{L^{p_i}_{w_i}(\Omega)} &\leq C_i \|\nabla^{k_i} f\|_{L^{p_i}_{w_i}(\mathcal{D})} \quad \text{for } N_1 < i \leq N \\ \|\nabla^{k_i} \Lambda' f\|_{L^{p_i}_{w_i}(\Omega)} &\leq C_i \|\nabla^{k_i} f\|_{L^{p_i}_{w_i}(\mathcal{D})} \quad \text{for } 0 \leq i \leq N_2 \\ \|\nabla^k \Lambda' f\|_{L^{p_i}_{w_i}(\mathbb{R}^n)} &\leq C_i \|\nabla^k f\|_{L^{p_i}_{w_i}(\mathcal{D})} \quad \text{for } N_2 < i \leq N \end{aligned}$$

for all  $f \in \text{Lip}^{k-1}_{\text{loc}}(\mathbb{R}^n)$ . Here  $C_i$  depends only on  $\varepsilon$ ,  $p_i$ ,  $w_i$ ,  $k_i$ ,  $n$ ,  $L$  and  $\max_i k_i$ . (Unfortunately  $L$  usually depends on  $r$ , but there are cases where  $L$  is independent of  $r$  and consequently  $C_i$  is independent of  $r$ .)

**THEOREM 4.5.** *Let  $1 \leq p_i < \infty$ ,  $w_i = \text{dist}(x, M_i)^{\alpha_i}$ ,  $\alpha_i \in \mathbb{R}$ ,  $M_i \subset \partial \mathcal{D}$  such that  $w_i$  is doubling for  $i = 0, 1, \dots, N$ . If  $\mathcal{D}$  is an unbounded  $(\varepsilon, \infty)$  domain, then there exists an extension operator on  $\mathcal{D}$  such that*

$$\|\nabla^{k_i} \Lambda f\|_{L_{w_i}^{p_i}(\mathbb{R}^n)} \leq C_i \|\nabla^{k_i} f\|_{L_{w_i}^{p_i}(\mathcal{D})}$$

for all  $i$  and  $f \in \text{Lip}_{\text{loc}}^{k-1}(\mathbb{R}^n)$ . Here  $C_i$  depends only on  $\varepsilon$ ,  $w_i$ ,  $p_i$ ,  $k_i$  and  $\max_i k_i$ .

**PROOF OF THEOREMS 4.4 AND 4.5.** If  $w(x) = \text{dist}(x, M)^\alpha$  for  $M \subset \mathcal{D}$ ,  $\alpha \in \mathbb{R}$ , let us make the following two observations:

$$(4.2) \quad \|f' - f_Q\|_{L_w^p(Q)} \leq C(A)l(Q)\|\nabla f\|_{L_w^p(Q)}$$

$$(4.3) \quad \frac{1}{|Q|} \|f\|_{L^1(Q)} \leq C(A)w(Q)^{-1/p} \|f\|_{L_w^p(Q)}$$

for all cubes  $Q$  in  $\mathcal{D}$  such that  $Al(Q) \leq d(Q) \leq l(Q)/A$  for  $A > 0$ . We can now check through the proof of Theorems 1.4 and 1.5 in [9] using (4.2) and (4.3) as the substitute of the condition that  $w \in A_p$  to obtain Theorems 4.4 and 4.5.

(b) In Theorem 4.4, if we assume in addition that  $w^{-1/p} \in L_{\text{loc}}^{p'}(\mathbb{R}^n)$ , we can indeed replace  $\text{Lip}_{\text{loc}}^{k-1}(\mathbb{R}^n)$  by  $\cap E_{w_i, k_i}^{p_i}(\mathcal{D})$  as  $C^\infty(\mathbb{R}^n) \cap (\cap E_{w_i, k_i}^{p_i}(\mathcal{D}))$  is dense in  $\cap E_{w_i, k_i}^{p_i}(\mathcal{D})$ . For the details, check through the proof of Theorem 6.1 in [9].

#### REFERENCES

1. Richard Adams, *Sobolev Spaces*, Academic Press, New York, 1975.
2. R. C. Brown and D. B. Hinton, *Sufficient conditions for weighted inequalities of sum form*, J. Math. Anal. Appl. **112**(1985), 563–578.
3. ———, *Weighted interpolation inequalities of sum and product form in  $\mathbb{R}^n$* , Proc. London Math. Soc. (3) **56**(1988), 261–280.
4. ———, *Weighted interpolation inequalities and embeddings in  $\mathbb{R}^n$* , Canad. J. Math. (6) **62**(1990), 959–980.
5. Alberto P. Calderón, *Lebesgue spaces of differentiable functions and distributions*, Proc. Symp. Pure Math. **IV**(1961), 33–49.
6. Sagun Chanillo and Richard L. Wheeden, *Poincaré inequalities for a class of non- $A_p$  weights*, Indiana Math. J. (3) **41**(1992), 605–623.
7. Filippo Chiarenza and Michele Frasca, *A note on a weighted Sobolev inequality*, Proc. Amer. Math. Soc. (4) **93**(1985), 703–704.
8. Michael Christ, *The extension problem for certain function spaces involving fractional orders of differentiability*, Ark. Mat. (1) **22**(1984), 63–81.
9. Seng-Kee Chua, *Extension theorems on weighted Sobolev spaces*, Indiana Math. J. **41**(1992), 1027–1076.
10. ———, *Extension and restriction theorems on weighted Sobolev spaces*, Ph.D. thesis, Rutgers University, 1991.
11. ———, *Some Remarks on extension theorems for weighted Sobolev spaces*, Illinois J. Math., **38**(1994), 95–126.
12. ———, *Weighted Sobolev's inequalities on domains satisfying the chain condition*, Proc. Amer. Math. Soc. **117**(1993), 449–457.
13. ———, *Restriction theorems on weighted Sobolev spaces of mixed norm*, Real Anal. Exchange (2) **17**(1991/92), 633–651.
14. ———, *On weighted Sobolev interpolation inequalities*, Proc. Amer. Math. Soc. **121**(1994), 441–449.
15. ———, *Weighted Sobolev interpolation inequalities on certain domains*, J. London Math. Soc. (2) **51**(1995), 532–544.

16. Ronald R. Coifman and Charles Fefferman, *Weighted norm inequalities for maximal functions and singular integrals*, *Studia Math.* **51**(1974), 241–250.
17. D. E. Edmunds, Alois Kufner and Jiong Sun, *Extension of functions in weighted Sobolev spaces*, *Rend. Accad. Naz. Sci. XL Mem. Mat. (5)*, 108<sup>o</sup> (17) **16**(1990), 327–339.
18. Eugene B. Fabes, Carlos E. Kenig and Raul P. Serapioni, *The local regularity of solutions of degenerate elliptic equations*, *Comm. Partial Differential Equations* **7**(1982), 77–116.
19. V. M. Gol'dshtein and Yu. G. Reshetnyak, *Quasiconformal Mappings and Sobolev Spaces*, Kluwer Academic Publishers, 1990.
20. Cristian E. Gutierrez and Richard L. Wheeden, *Sobolev interpolation inequalities with weights*, *Trans. Amer. Math. Soc.* **323**(1991), 263–281.
21. R. A. Hunt, D. S. Kurtz and C. I. Neugebauer, *A note on the equivalence of  $A_p$  and Sawyer's condition for equal weights*, *Conf. on Harm. Anal. in Honor of A. Zygmund*, Wadsworth, 1983, 156–158.
22. Peter Jones, *Quasiconformal mappings and extendability of functions in Sobolev spaces*, *Acta Math.* (1–2) **147**(1981), 71–88.
23. Alois Kufner, *Weighted Sobolev Spaces*, John Wiley & Sons Ltd, 1985.
24. ———, *How to define reasonable Sobolev spaces*, *Comment. Math. Univ. Carolin.* (3) **25**(1984), 537–554.
25. Olli Martio and J. Sarvas, *Injectivity theorems in plane and space*, *Ann. Acad. Sci. Fenn. Series A I Math.* (2) **4**(1979), 383–401.
26. Vladimir G. Maz'ja, *Sobolev Spaces*, Springer-Verlag, New York, 1985.
27. Benjamin Muckenhoupt, *Weighted norm inequalities for the Hardy maximal function*, *Trans. Amer. Math. Soc.* **165**(1972), 207–226.
28. Benjamin Muckenhoupt and Richard Wheeden, *On the dual of weighted  $H^1$  of the half-space*, *Studia Math.* (1) **63**(1978), 57–79.
29. Eric Sawyer, *A characterization of a two-weighted norm inequality for maximal operators*, *Studia Math.* **75**(1982), 1–11.
30. Eric Sawyer and Richard L. Wheeden, *Weighted inequalities for fractional integrals on Euclidean and homogeneous spaces*, *Amer. J. Math.* (4) **114**(1992), 813–874.
31. Elias M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, 1970.
32. Jan-Olov Stromberg and Alberto Torchinsky, *Weighted Hardy Spaces*, Lecture Notes in Math. **1381**, Springer, New York, 1989.
33. Alberto Torchinsky, *Real-Variable Methods in Harmonic Analysis*, Academic Press Inc., New York, 1986.
34. Richard L. Wheeden and Antoni Zygmund, *Measure and Integral*, Marcel Dekker Inc., New York, 1977.
35. Thomas H. Wolff, *Restriction of  $A_p$  weights*, preprint.

National University of Singapore

Department of Mathematics

10, Kent Ridge Crescent

Singapore 0511

e-mail: matcsk@math.nus.sg