

On Wigner's theorem

DANIEL S FREED

Wigner's theorem asserts that any symmetry of a quantum system is unitary or antiunitary. In this short note we give two proofs based on the geometry of the Fubini–Study metric.

81R99; 81Q70, 82B03, 82D25, 19L50

For Mike Freedman, on the occasion of his 60th birthday

The space of pure states of a quantum mechanical system is the projective space $\mathbb{P}\mathcal{H}$ of lines in a separable complex Hilbert space $(\mathcal{H}, \langle -, - \rangle)$, which may be finite or infinite dimensional. It carries a symmetric function $p: \mathbb{P}\mathcal{H} \times \mathbb{P}\mathcal{H} \rightarrow [0, 1]$ whose value $p(L_1, L_2)$ on states $L_1, L_2 \in \mathbb{P}\mathcal{H}$ is the *transition probability*: if $\psi_i \in L_i$ is a unit norm vector in the line L_i , then

$$p(L_1, L_2) = |\langle \psi_1, \psi_2 \rangle|^2.$$

Let $\text{Aut}_{\text{qtm}}(\mathbb{P}\mathcal{H})$ denote the group of symmetries of $(\mathbb{P}\mathcal{H}, p)$, the group of quantum symmetries. A fundamental theorem of Wigner¹ [12, Sections 20A and 26] (see also Bargmann [2] and Weinberg [11, Section 2A]) expresses $\text{Aut}_{\text{qtm}}(\mathbb{P}\mathcal{H})$ as a quotient of linear and antilinear symmetries of \mathcal{H} . This note began with the rediscovery of a formula which relates the quantum geometry of $(\mathbb{P}\mathcal{H}, p)$ to a more familiar structure in differential geometry: the Fubini–Study Kähler metric on $\mathbb{P}\mathcal{H}$. It leads to two proofs of Wigner's theorem, **Theorem 8** of this note, based on the differential geometry of projective space.

The proofs here use more geometry than the elementary proofs [2], [11, Section 2A]. We take this opportunity to draw attention to Wigner's theorem and to the connection between quantum mechanics and projective geometry. It is a fitting link for a small tribute to Mike Freedman, whose dual careers in topology and condensed matter physics continue to inspire.

Let $d: \mathbb{P}\mathcal{H} \times \mathbb{P}\mathcal{H} \rightarrow \mathbb{R}^{\geq 0}$ be the distance function associated to the Fubini–Study metric.

¹As I learned in Bonolis [3, page 74], this theorem was first asserted in a 1928 joint paper [10, page 207] of von Neumann and Wigner, though with only a brief justification. A more complete account appeared in Wigner's book (in the original German) in 1931.

1 Theorem *The functions p and d are related by*

$$(2) \quad \cos(d) = 2p - 1.$$

As a gateway into the literature on ‘geometric quantum mechanics’, where (2) can be found,² see Brody and Hughston [4] and the references therein.

3 Corollary *$\text{Aut}_{\text{qtm}}(\mathbb{P}\mathcal{H})$ is the group of isometries of $\mathbb{P}\mathcal{H}$ with the Fubini–Study distance function.*

4 Remark If \mathcal{H} is infinite dimensional, then $\mathbb{P}\mathcal{H}$ is an infinite dimensional smooth manifold modeled on a Hilbert space. Basic notions of calculus and differential geometry carry over to Hilbert manifolds (Lang [8]). The Myers–Steenrod theorem asserts that a distance-preserving map between two Riemannian manifolds is smooth and preserves the Riemannian metric. That theorem is also true on Riemannian manifolds modeled on Hilbert manifolds (Garrido, Jaramillo and Rangel [6]).³ So in the sequel we use that a distance-preserving map $\phi: \mathbb{P}\mathcal{H} \rightarrow \mathbb{P}\mathcal{H}$ is smooth and is an isometry in the sense of Riemannian geometry.

The tangent space to $\mathbb{P}\mathcal{H}$ at a line $L \subset \mathcal{H}$ is canonically $T_L \mathbb{P}\mathcal{H} \cong \text{Hom}_{\mathbb{C}}(L, L^{\perp})$, where $L^{\perp} \subset \mathcal{H}$ is the orthogonal complement to L , a closed subspace and therefore itself a Hilbert space. If $f_1, f_2: L \rightarrow L^{\perp}$, then the Fubini–Study hermitian metric is defined by

$$(5) \quad \langle f_1, f_2 \rangle = \text{Tr}(f_1^* f_2).$$

The adjoint f_1^* is computed using the inner products on L and L^{\perp} . The composition $f_1^* f_2$ is an endomorphism of L , hence multiplication by a complex number which we identify as the trace of the endomorphism. If $\ell \in L$ has unit norm, then the map

$$(6) \quad \begin{aligned} \text{Hom}_{\mathbb{C}}(L, L^{\perp}) &\longrightarrow L^{\perp} \\ f &\longmapsto f(\ell) \end{aligned}$$

is a linear isometry for the induced metric on $L^{\perp} \subset \mathcal{H}$. The underlying Riemannian metric is the real part of the hermitian metric (5); it only depends on the real part of the inner product on \mathcal{H} .

²Notice that (2) is equivalent to $p = \cos^2(d/2)$.

³The proof depends on the existence of geodesic convex neighborhoods, proved in [8, Section VIII.5]. For the Fubini–Study metric on $\mathbb{P}\mathcal{H}$ such neighborhoods may easily be constructed explicitly. I thank Karl-Hermann Neeb for his inquiry about the Myers–Steenrod theorem in infinite dimensions.

Proof of Theorem 1 Equation (2) is obvious on the diagonal in $\mathbb{P}\mathcal{H} \times \mathbb{P}\mathcal{H}$, as well as if $\dim \mathcal{H} = 1$. Henceforth we rule out both possibilities. Fix $L_1 \neq L_2 \in \mathbb{P}\mathcal{H}$ and let V be the 2-dimensional space $L_1 + L_2 \subset \mathcal{H}$. The unitary automorphism of $\mathcal{H} = V \oplus V^\perp$ which is $+1$ on V and -1 on V^\perp induces an isometry of $\mathbb{P}\mathcal{H}$ which has $\mathbb{P}V$ as a component of its fixed point set. It follows that $\mathbb{P}V$ is totally geodesic. Therefore, to compute $d(L_1, L_2)$ we are reduced to the case of the complex projective line with its Fubini–Study metric: the round 2-sphere.

Let $e_1 \in L_1$ have unit norm and choose $e_2 \in V$ to fill out a unitary basis $\{e_1, e_2\}$. Then $\lambda e_1 + e_2 \in L_2$ for a unique $\lambda \in \mathbb{C}$. If $\lambda = 0$ then it is easy to check that $d = \pi$ and $p = 0$, consistent with (2), so we now assume $\lambda \neq 0$. Identify $\mathbb{P}V \setminus \{\mathbb{C} \cdot e_2\} \approx \mathbb{C}$ by $\mathbb{C} \cdot (e_1 + \mu e_2) \leftrightarrow \mu$. Use stereographic projection from the north pole $(1, 0)$ in Euclidean 3-space $\mathbb{R} \times \mathbb{C}$ to identify $\{0\} \times \mathbb{C} \approx S^2 \setminus \{(1, 0)\}$, where $S^2 \subset \mathbb{R} \times \mathbb{C}$ is the unit sphere. Under these identifications we have

$$L_1 \longleftrightarrow (-1, 0)$$

$$L_2 \longleftrightarrow \left(-\frac{|\lambda|^2 - 1}{|\lambda|^2 + 1}, \frac{2|\lambda|^2}{|\lambda|^2 + 1} \frac{1}{\lambda} \right)$$

from which $\cos(d) = (|\lambda|^2 - 1)/(|\lambda|^2 + 1)$ can be computed as the inner product of vectors in the 3-dimensional vector space $\mathbb{R} \oplus \mathbb{C}$. Since $p = |\lambda|^2/(|\lambda|^2 + 1)$, equation (2) is satisfied. \square

A real linear map $S: \mathcal{H} \rightarrow \mathcal{H}$ is *antiunitary* if it is conjugate linear and

$$\langle S\psi_1, S\psi_2 \rangle = \overline{\langle \psi_1, \psi_2 \rangle} \quad \text{for all } \psi_1, \psi_2 \in \mathcal{H}.$$

Let $G(\mathcal{H})$ denote the group consisting of all unitary and antiunitary operators on \mathcal{H} . In the norm topology it is a Banach Lie group (Milnor [9]) with two contractible components; the same is true in the compact–open topology (Freed and Moore [5, Appendix D]). The identity component is the group $U(\mathcal{H})$ of unitary transformations. Any $S \in G(\mathcal{H})$ maps complex lines to complex lines, so induces a diffeomorphism of $\mathbb{P}\mathcal{H}$, and since S preserves the real part of $\langle -, - \rangle$ the induced diffeomorphism is an isometry. The unit norm scalars $\mathbb{T} \subset G(\mathcal{H})$ act trivially on $\mathbb{P}\mathcal{H}$, so there is an exact⁴ sequence of Lie groups

$$(7) \quad 1 \longrightarrow \mathbb{T} \longrightarrow G(\mathcal{H}) \longrightarrow \text{Aut}_{\text{qtm}}(\mathbb{P}\mathcal{H}).$$

Note that \mathbb{T} is not central since antiunitary maps conjugate scalars.

⁴We assume $\dim \mathcal{H} > 1$.

8 Theorem (Wigner [12]) *The homomorphism $G(\mathcal{H}) \rightarrow \text{Aut}_{\text{qtm}}(\mathbb{P}\mathcal{H})$ is surjective: every quantum symmetry of $\mathbb{P}\mathcal{H}$ lifts to a unitary or antiunitary operator on \mathcal{H} .*

By [Corollary 3](#) the same is true for isometries of the Fubini–Study metric, and indeed we prove Wigner’s Theorem by computing the group of isometries.

9 Remark If $\rho: G \rightarrow \text{Aut}_{\text{qtm}}(\mathbb{P}\mathcal{H})$ is any group of quantum symmetries, then the surjectivity of $G(\mathcal{H}) \rightarrow \text{Aut}_{\text{qtm}}(\mathbb{P}\mathcal{H})$ implies the extension (7) pulls back to a twisted central extension of G . The twist is the homomorphism $G \rightarrow \mathbb{Z}/2\mathbb{Z}$ which tells whether a symmetry lifts to be unitary or antiunitary. The isomorphism class of this twisted central extension is then an invariant of ρ . This is the starting point for joint work with Greg Moore [5] about quantum symmetry classes and topological phases in condensed matter physics.

10 Example $\mathbb{P}(\mathbb{C}^2) = \mathbb{C}\mathbb{P}^1$ with the Fubini–Study metric is the round 2–sphere of unit radius. Its isometry group is the group $O(3)$ of orthogonal transformations of $\text{SO}(3)$. The identity component $\text{SO}(3)$ is the image of the group $U(2)$ of unitary transformations of \mathbb{C}^2 . The other component of $O(3)$ consists of orientation-reversing orthogonal transformations, such as reflections, and they lift to antiunitary symmetries of \mathbb{C}^2 . In this case the group $G(\mathcal{H})$ is also known as $\text{Pin}^c(3)$; see Atiyah, Bott and Shapiro [1].

We present two proofs of [Theorem 8](#). The first is based on the following standard fact in Riemannian geometry.

11 Lemma *Let M be a Riemannian manifold, $p \in M$, and $\phi: M \rightarrow M$ an isometry with $\phi(p) = p$. Suppose $B_r \subset T_p M$ is the open ball of radius r centered at the origin and assume the Riemannian exponential map \exp_p maps B_r diffeomorphically into M . Then in exponential coordinates $\phi|_{B_r}$ equals the restriction of the linear isometry $d\phi_p$ to B_r .*

Proof If $\xi \in B_r$, then $\exp_p(\xi) = \gamma_\xi(1)$, where $\gamma_\xi: [0, 1] \rightarrow M$ is the unique geodesic which satisfies $\gamma_\xi(0) = p$, $\dot{\gamma}_\xi(0) = \xi$. Since ϕ maps geodesics to geodesics, $\phi \circ \exp_p = \exp_p \circ d\phi_p$ on B_r , as desired. \square

If $\rho: [0, r') \rightarrow [0, r)$ is a diffeomorphism for some $r' > 0$, then

$$(12) \quad \xi \longmapsto \exp_p(\rho(|\xi|)\xi)$$

maps $B_{r'}$ diffeomorphically into M , and ϕ in this coordinate system is also linear.

First Proof of Theorem 8 Let $\phi: \mathbb{P}\mathcal{H} \rightarrow \mathbb{P}\mathcal{H}$ be an isometry. Composing with an isometry in $G(\mathcal{H})$ we may assume $\phi(L) = L$ for some $L \in \mathbb{P}\mathcal{H}$. The tangent space $T_L\mathbb{P}\mathcal{H}$ is canonically $\text{Hom}_{\mathbb{C}}(L, L^\perp)$, and also $f \in \text{Hom}_{\mathbb{C}}(L, L^\perp)$ determines $\Gamma_f \in \mathbb{P}\mathcal{H}$ by $\Gamma_f \subset \mathcal{H} = L \oplus L^\perp$ is the graph of f . We claim $f \mapsto \Gamma_f$ has the form (12) for some $\rho: [0, \infty) \rightarrow [0, \pi)$. It suffices to show that for any $f \in \text{Hom}_{\mathbb{C}}(L, L^\perp)$ of unit norm, the map $t \mapsto \Gamma_{tf}$ traces out a (reparametrized) geodesic in a parametrization independent of f . As in the proof of Theorem 1 this reduces to $\dim \mathcal{H} = 2$ and so to an obvious statement about the round 2–sphere. It follows from Lemma 11 that ϕ is a real isometry $S \in \text{End}_{\mathbb{R}}(\text{Hom}_{\mathbb{C}}(L, L^\perp))$. It remains to prove that S is complex linear or antilinear; then we extend S by the identity on L to obtain a unitary or antiunitary operator on $\mathcal{H} = L \oplus L^\perp$.

If $\dim \mathcal{H} = 2$ then Theorem 8 can be verified (see Example 10), so assume $\dim \mathcal{H} > 2$. Identify $\text{Hom}_{\mathbb{C}}(L, L^\perp) \approx L^\perp$ as in (6). Since $S \in \text{End}_{\mathbb{R}}(L^\perp)$ maps complex lines in L^\perp to complex lines, there is a function $\alpha: L^\perp \setminus \{0\} \rightarrow \mathbb{C}$ such that $S(i\xi) = \alpha(\xi)S(\xi)$ for all nonzero $\xi \in L^\perp$. Fix $\xi \neq 0$ and choose $\eta \in L^\perp$ which is linearly independent. Then

$$\begin{aligned} S(i(\xi + \eta)) &= \alpha(\xi + \eta)[S(\xi) + S(\eta)] \\ &= \alpha(\xi)S(\xi) + \alpha(\eta)S(\eta) \end{aligned}$$

from which $\alpha(\xi) = \alpha(\eta)$. Applied to $i\xi, \eta$ we learn $\alpha(\xi) = \alpha(i\xi)$. On the other hand,

$$-S(\xi) = S(-\xi) = \alpha(i\xi)S(i\xi) = \alpha(i\xi)\alpha(\xi)S(\xi),$$

whence $\alpha(\xi)^2 = -1$. By continuity either $\alpha \equiv i$ or $\alpha \equiv -i$, which proves that S is linear or S is antilinear. \square

The second proof leans on complex geometry.

13 Lemma *An isometry $\phi: \mathbb{P}\mathcal{H} \rightarrow \mathbb{P}\mathcal{H}$ is either holomorphic or antiholomorphic.*

Proof Let $I: T\mathbb{P}\mathcal{H} \rightarrow T\mathbb{P}\mathcal{H}$ be the (almost) complex structure. Then I is parallel with respect to the Levi–Civita covariant derivative, since $\mathbb{P}\mathcal{H}$ is Kähler, and so therefore is ϕ^*I . We claim any parallel almost complex structure J equals $\pm I$; the lemma follows immediately.

If J is parallel, then it commutes with the Riemann curvature tensor R . Compute at $L \in \mathbb{P}\mathcal{H}$ and identify $T_L\mathbb{P}\mathcal{H} \approx L^\perp$, as in (6). Then if $\xi, \eta \in L^\perp$ and $\langle \xi, \eta \rangle = 0$, since $\mathbb{P}(L \oplus \mathbb{C} \cdot \xi \oplus \mathbb{C} \cdot \eta) \subset \mathbb{P}\mathcal{H}$ is totally geodesic and has constant holomorphic

sectional curvature one (Kobayashi and Nomizu [7, Section IX.7]), we compute

$$\begin{aligned} R(\xi, I\xi)\xi &= -|\xi|^2 I\xi, \\ R(\xi, I\xi)\eta &= -\frac{1}{2}|\xi|^2 I\eta. \end{aligned}$$

It follows that J preserves every complex line $K = \mathbb{C} \cdot \xi \subset L^\perp$ and commutes with I on K . Therefore, $J = \pm I$ on K . By continuity, the sign is independent of K and L . \square

Second Proof of Theorem 8 First, recall that if U is finite dimensional, then every holomorphic symmetry of $\mathbb{P}U$ is linear. The proof is as follows. Let $\mathcal{L} \rightarrow \mathbb{P}U$ be the canonical holomorphic line bundle whose fiber at $L \in \mathbb{P}U$ is L . A holomorphic line bundle on $\mathbb{P}U$ is determined by its Chern class, so $\phi^* \mathcal{L} \cong \mathcal{L}$. Fix an isomorphism; it is unique up to scale. There is an induced linear map on the space $H^0(\mathbb{P}U; \mathcal{L}^*) \cong U^*$ of global holomorphic sections:

$$(14) \quad \phi^*: H^0(\mathbb{P}U; \mathcal{L}^*) \longrightarrow H^0(\mathbb{P}U; \phi^* \mathcal{L}^*) \cong H^0(\mathbb{P}U; \mathcal{L}^*).$$

The transpose $\hat{\phi}$ of (14) is the desired linear lift of ϕ .

Let $\phi: \mathbb{P}\mathcal{H} \rightarrow \mathbb{P}\mathcal{H}$ be an isometry. After composition with an element of $G(\mathcal{H})$ we may, by Lemma 13, assume ϕ is holomorphic and fixes some $L \in \mathbb{P}\mathcal{H}$. Let $U \subset \mathcal{H}$ be a finite dimensional subspace containing L . Then the pullback of $\mathcal{L}_{\mathcal{H}} \rightarrow \mathbb{P}\mathcal{H}$ to $\phi^* \mathcal{L}_{\mathcal{H}}|_{\mathbb{P}U} \rightarrow \mathbb{P}U$ has degree one, so is isomorphic to $\mathcal{L}_U \rightarrow \mathbb{P}U$, and there is a unique isomorphism which is the identity on the fiber over L . A functional $\alpha \in \mathcal{H}^*$ restricts to a holomorphic section of $\phi^* \mathcal{L}_{\mathcal{H}}^*|_{\mathbb{P}U} \rightarrow \mathbb{P}U$, so by composition with the isomorphism $\phi^* \mathcal{L}_{\mathcal{H}}^*|_{\mathbb{P}U} \cong \mathcal{L}_U^*$ to an element of U^* . The resulting map $\mathcal{H}^* \rightarrow U^*$ is linear, and its transpose $\hat{\phi}: U \rightarrow \mathcal{H}$ is the identity on L . Let U run over all finite dimensional subspaces of \mathcal{H} to define $\hat{\phi}: \mathcal{H} \rightarrow \mathcal{H}$. The uniqueness of the isomorphism $\phi^* \mathcal{L}_{\mathcal{H}}|_{\mathbb{P}U} \cong \mathcal{L}_U$ implies that $\hat{\phi}$ is well-defined and a linear lift of ϕ . It is unitary since ϕ is an isometry. \square

Acknowledgement The author is supported by the National Science Foundation under grant DMS-0603964

References

- [1] MF Atiyah, R Bott, A Shapiro, *Clifford modules*, Topology 3 (1964) 3–38 [MR0167985](#)
- [2] V Bargmann, *Note on Wigner’s theorem on symmetry operations*, J. Mathematical Phys. 5 (1964) 862–868 [MR0164620](#)

- [3] **L Bonolis**, *From the rise of the group concept to the stormy onset of group theory in the new quantum mechanics: A saga of the invariant characterization of physical objects, events and theories*, Nuovo Cimento Rivista Serie 27 (2004)
- [4] **D C Brody, L P Hughston**, *Geometric quantum mechanics*, J. Geom. Phys. 38 (2001) 19–53 [MR1817511](#)
- [5] **D S Freed, G W Moore**, *Twisted equivariant matter*, in preparation
- [6] **I Garrido, J A Jaramillo, Y C Rangel**, *Algebras of differentiable functions on Riemannian manifolds*, Bull. Lond. Math. Soc. 41 (2009) 993–1001 [MR2575330](#)
- [7] **S Kobayashi, K Nomizu**, *Foundations of differential geometry*, volume II, John Wiley & Sons, New York (1996) [MR1393941](#)
- [8] **S Lang**, *Introduction to differentiable manifolds*, 2nd edition, Springer, New York (2002) [MR1931083](#)
- [9] **J Milnor**, *Remarks on infinite-dimensional Lie groups*, from: “Relativity, groups and topology, II (Les Houches, 1983)”, North-Holland, Amsterdam (1984) 1007–1057 [MR830252](#)
- [10] **J von Neumann, E P Wigner**, *Zur Erklärung einiger Eigenschaften der Spektren aus der Quantenmechanik des Drehelektrons*, Zeitschrift für Physik A Hadrons and Nuclei 49 (1928) 73–94
- [11] **S Weinberg**, *The quantum theory of fields*, volume I, Cambridge Univ. Press (2005) [MR2148466](#)
- [12] **E P Wigner**, *Group theory: And its application to the quantum mechanics of atomic spectra*, Pure and Applied Physics 5, Academic Press, New York (1959) [MR0106711](#)

Department of Mathematics, University of Texas
1 University Station C1200, Austin, TX 78712-0257, USA

dafr@math.utexas.edu

www.ma.utexas.edu/users/dafr

Received: 9 December 2011 Revised: 2 March 2012