

One-Pot Synthesis of Simple Alkaloids: 2,3-Polymethylene-4(3*H*)-quinazolinones, Luotonin A, Tryptanthrin, and Rutaecarpine

Katherine Chae JAHNG, Seung Ill KIM, Dong Hyeon KIM, Chang Seob SEO, Jong-Keun SON, Seung Ho LEE, Eung Seok LEE, and Yurngdong JAHNG*

College of Pharmacy, Yeungnam University; Gyeongsan 712–749, Korea.

Received November 27, 2007; accepted January 8, 2008; published online January 25, 2008

One-pot synthesis of various 2,3-polymethylene-4(3*H*)-quinazolinones from anthranilic acid, corresponding lactam and SOCl₂ is described, which can be applicable to the synthesis of simple 4(3*H*)-quinazolinone-derived alkaloids, such as luotonin A, tryptanthrin, and rutaecarpine.

Key words deoxyvasicinone; mackinazolinone; luotonin A; tryptanthrin; rutaecarpine; anthranilic acid

2,3-Trimethylene-4(3*H*)-quinazolinone (deoxyvasicinone, **1a**)¹ and 2,3-tetramethylene-4(3*H*)-quinazolinone (mackinazolinone, **1b**)^{2,3} are not only the alkaloids isolated from *Peganum* and *Mackinlaya* species, respectively, but are also prepared chemically before the isolation from natural sources.^{4–6} The establishment of a simple and efficient synthetic method for these alkaloids has long been a challenging subject.^{7–20} In addition, these substances are a part of the family of intriguing alkaloids that include the bronchodilator vasicinone (**2a**) from *Adhatoda vasica*,²¹ antiendotoxic isaindigotone (**2b**) from *Isatis indigotica*,²² topoisomerase I-dependent cytotoxic luotonin A (**3**) from *Peganum nigellastrum*,²³ antibiotic tryptanthrin (**4**) from yeast (*Candida lipolytica*)^{24,25} and later from cannon ball tree (*Cououputa guianensis*),²⁶ anti-inflammatory rutaecarpine (**5**)²⁷ from *Evodia rutaecarpa*, and related alkaloids (Fig. 1).²⁸

Since Niementowski's preparation of 4(3*H*)-quinazolinone by fusing anthranilic acid with formamide,²⁹ several methods attempting the synthesis of modified quinazolinones have been reported.^{1,30–33} Although most of these are applicable to acyclic amides, methods for cyclic amides, especially without an aryl group, are somewhat limited.^{34–37} These limitations have led to the continuous development of new procedures for cyclic amides, such as cyclocondensation of isoatoic anhydride with lactams,¹² metal catalyzed reductive *N*-heterocyclization of *N*-(2-nitrobenzoyl)lactams in the presence of CO,^{13,14} reaction of 2-aminobenzamide with succinic anhydride,¹⁵ intramolecular aza-Wittig reaction of *N*-(2-azidobenzoyl)lactams,^{9,16} condensation of anthranilic acid with *O*-alkyllactams^{6,18–20} or *S*-alkyllactams,⁸ reaction of an-

thranilic acid with SOCl₂ followed by cyclization with lactams,¹⁷ cyclization of iminochlorides generated from lactam and methyl anthranilate,¹¹ solid-phase synthesis,^{7,9} and microwave assisted synthesis.^{10,12} Nevertheless, the biological importance of 4(3*H*)-quinazolinone and related compounds still spurred the development of a new and simple synthetic method for these compounds.^{38–43}

As a part of our research on biologically important natural products, we herein described a simple one-pot procedure for the preparation of **1** as well as related alkaloids **3**, **4** and **5** in multi-gram quantity.

Although an earlier study revealed that iminoketene (**8**), generated from the reaction of anthranilic acid (**6**) and SOCl₂, underwent cycloaddition to cyclic iminols, which are tautomers of the corresponding lactams (**9**), to yield **1** in 70–82% yield,¹⁹ we sometimes have been unable to repeat the reactions.⁴⁴ Whenever we failed to generate the desired products, imine compounds (**10**) were isolated as major products (Chart 1). Although we could convert these imines (**10**) to the desired 2,3-tri- and 2,3-tetramethylene-4(3*H*)-quinazolinones by employing polyphosphoric acid-catalyzed dehydration (see Experimental section in details), we examined and adjusted the reaction condition to find that such reaction could be carried out in a single step. One-pot reaction of anthranilic acid, lactams (**9**) and SOCl₂ in refluxing benzene yielded **1a** and **1b** in 87% and 78% yield, respectively.

The generality of the process was examined with a variety of substituted anthranilic acids (**6**) and various lactams (**9**) to provide the corresponding 2,3-polymethylene-4(3*H*)-quinazolinone derivatives (**1**) in 78–89% yields (Table 1).

The electronic properties of the substituent on anthranilic acid did not seem to greatly affect the reaction (compounds **1ab**, **1ac**, **1ad**, **1ag** vs. **1ae**, **1af**). It is worthy to note that the




Fig. 1. Simple Alkaloids Derived from 4(3*H*)-Quinazolinone






Chart 1

* To whom correspondence should be addressed. e-mail: ydjahng@ynu.ac.kr

Table 1. Synthesis of 2,3-polymethylene-4(3*H*)-quinazolinones

Compd.	R	<i>n</i>	Yield (%)	mp (°C)	Compd.	R	<i>n</i>	Yield (%)	mp (°C)
1aa	H	1	87	109 [104 ¹⁾]	1bb	2-Cl	2	87	107 [106 ⁴⁷⁾]
1ab	2-Cl	1	85	177 [178—179 ⁴⁵⁾]	1bc	2-Br	2	87	163—164
1ac	2-Br	1	86	158 [157—158 ⁴⁵⁾]	1bd	2-I	2	84	286—287
1ad	2-I	1	87	150 [148—150 ⁴⁵⁾]	1be	2-OH	2	85	257—259
1ae	2-OH	1	86	289—290	1bf	2-CH ₃	2	88	90 [204—206 ^{a)46)}]
1af	2-CH ₃	1	89	110 [174—175 ^{a)46)}]	1bg	2-NO ₂	2	88	172 [171—172 ⁴⁶⁾]
1ag	2-NO ₂	1	85	187 [187—188 ⁴⁵⁾]	1bh	3-Cl	2	82	133 [132—133 ¹⁸⁾]
1ah	3-Cl	1	86	188 [186—188 ¹⁸⁾]	1ca	H	3	78	97 [95—97 ⁶⁾]
1ba	H	2	78	99 [98.5—99.5 ²⁾]	1da	H	4	82	112 [109—110 ¹⁶⁾]

a) Melting point of corresponding HCl salt.



present method is applicable to compounds with an OH group. Although the compound with an OH group, 2-hydroxy-7,8-dihydro-6*H*,11*H*-pyrrolo[2,1-*b*]quinazolin-11-one (**1ae**), was previously prepared in two steps: condensation of corresponding 5-methoxyanthranilic acid with *O*-ethylbutyrolactim followed by acid catalyzed demethylation,⁴⁷⁾ the compounds with an OH as a substituent were prepared in 85—86% yields (compounds **1ae**, **1be**).

Following the reactions shown above, known lactams, 3-oxo-1*H*-pyrrolo[3,4-*b*]quinoline (**11**),⁴⁸⁾ isatin (**12**), and 1-oxo-1,2,3,4-tetrahydropyrido[3,4-*b*]indole (**13**),⁴⁹⁾ were converted to the corresponding alkaloids luotonin A (**3**), tryptanthrin (**4**), and rutaecarpine (**5**), respectively, in 78—85% yields (Chart 2).

It should be noted that the reactions of anthranilic acid or its esters with lactams in refluxing POCl₃ have already been attempted, but the yields were generally low.^{6,45,50)} This has limited the utility for the alkaloids derived from 4(3*H*)-quinazolinone.

In conclusion, a one-pot procedure for the preparation of various 2,3-polymethylene-4(3*H*)-quinazolinones has been established by a reaction of anthranilic acid, lactam, and SOCl₂. The described procedure was applied to the preparation of three simple 4(3*H*)-quinazolinone alkaloids, luotonin A, tryptanthrin, and rutaecarpine.

Experimental

Melting points were determined using a Fischer–Jones melting points apparatus and are not corrected. IR spectra were obtained using a Perkin-Elmer 1330 spectrophotometer. NMR spectra were obtained using a Bruker-250 spectrometer 250 MHz for ¹H-NMR and 62.5 MHz for ¹³C-NMR and are re-

ported as parts per million (ppm) from the internal standard tetramethylsilane. Chemicals and solvents were commercial reagent grade and used without further purification. The compounds **11**⁴⁹⁾ and **13**⁵⁰⁾ were prepared by employing previously reported methods. Electrospray ionization (ESI) mass spectrometry (MS) experiments were performed on a LCQ advantage-trap mass spectrometer (Thermo Finnigan, San Jose, CA, U.S.A.). Elemental analyses were performed with a Hewlett-Packard Model 185B elemental analyzer.

General Procedure of 2,3-Polymethylene-4(3*H*)-quinazolinones A solution of anthranilic acid (1.37 g, 0.01 mol), lactam (0.12 mol) and SOCl₂ (2 ml, *ca.* 2 eq) in dry benzene or pyridine (20 ml) was refluxed for 4 h. The reaction mixture was poured to ice-water and made basic with NH₄OH (100 ml), and the precipitate was collected as a crude product.

2-Bromo-7,8-dihydro-6*H*,11*H*-pyrrolo[2,1-*b*]quinazolin-11-one (1ac**)** Yellow needles: mp 158 °C. Unreported spectral data are as follows: ¹H-NMR (DMSO-*d*₆, 250 MHz) δ: 8.20 (1H, d, *J*=1.8 Hz, H1), 7.99 (1H, dd, *J*=8.5, 1.8 Hz, H3), 7.65 (1H, d, *J*=8.5 Hz, H4), 4.08 (2H, t, *J*=7.3 Hz), 3.17 (2H, d, *J*=7.3 Hz), 2.20 (2H, quintet, *J*=7.3 Hz). ESI-MS *m/z*: 266 [Calcd for C₁₁H₉BrN₂O: 266 (M+1)]. *Anal.* Calcd for C₁₁H₉BrN₂O: C, 49.84; H, 3.42; N, 10.57. Found: C, 49.81; H, 3.43; N, 10.58.

2-Iodo-7,8-dihydro-6*H*,11*H*-pyrrolo[2,1-*b*]quinazolin-11-one (1ad**)** Yellow needles: mp 150 °C. Unreported spectral data are as follows: ¹H-NMR (DMSO-*d*₆, 250 MHz) δ: 8.37 (1H, s, H1), 8.11 (1H, dd, *J*=8.5, 2.0 Hz, H3), 7.46 (1H, d, *J*=8.5 Hz, H4), 4.07 (2H, t, *J*=7.3 Hz), 3.14 (2H, d, *J*=7.3 Hz), 2.18 (2H, quintet, *J*=7.3 Hz). ESI-MS *m/z*: 313 [Calcd for C₁₁H₉I₂N₂O: 313 (M+H)]. *Anal.* Calcd for C₁₁H₉I₂N₂O: C, 42.33; H, 2.91; N, 8.98. Found: C, 42.27; H, 3.03; N, 8.98.

2-Hydroxy-7,8-dihydro-6*H*,11*H*-pyrrolo[2,1-*b*]quinazolin-11-one (1ae**)** White needles: mp 289—290 °C. ¹H-NMR (DMSO-*d*₆, 250 MHz) δ: 10.50 (br s, OH), 7.65 (1H, d, *J*=8.8 Hz, H4), 7.45 (1H, d, *J*=2.5 Hz, H1), 7.38 (1H, dd, *J*=8.5, 2.5 Hz, H3), 4.09 (2H, t, *J*=7.0 Hz), 3.26 (2H, d, *J*=7.6 Hz), 2.23 (2H, quintet, *J*=7.6 Hz). ESI-MS *m/z*: 203 [Calcd for C₁₁H₁₀N₂O₂: 203 (M+H)]. *Anal.* Calcd for C₁₁H₁₀N₂O₂: C, 65.34; H, 4.98; N, 13.85. Found: C, 65.35; H, 5.01; N, 13.84.

2-Nitro-7,8-dihydro-6*H*,11*H*-pyrrolo[2,1-*b*]quinazolin-11-one (1ag**)** Yellow needles: mp 187 °C. Unreported spectral data are as follows: ¹H-NMR (DMSO-*d*₆, 250 MHz) δ: 8.78 (1H, d, *J*=2.0 Hz, H1), 8.52 (1H, dd, *J*=8.5, 2.0 Hz, H3), 7.79 (1H, d, *J*=8.5 Hz, H4), 4.09 (2H, t, *J*=7.3 Hz), 3.16 (2H, d, *J*=7.3 Hz), 2.20 (2H, quintet, *J*=7.3 Hz). ESI-MS *m/z*: 232 [Calcd for C₁₁H₉N₃O₃: 232 (M+H)]. *Anal.* Calcd for C₁₁H₉N₃O₃: C, 57.14; H, 3.92; N, 18.17. Found: C, 57.19; H, 3.93; N, 18.08.

2-Bromo-6,7,8,9-tetrahydro-11*H*-pyrido[2,1-*b*]quinazolin-11-one (1bc**)** White needles: mp 163—164 °C. ¹H-NMR (DMSO-*d*₆, 250 MHz) δ: 8.22 (1H, d, *J*=2.3 Hz, H1), 8.04 (1H, dd, *J*=8.8, 2.3 Hz, H3), 7.71 (1H, d, *J*=8.8 Hz, H4), 3.93 (2H, t, *J*=6.0 Hz), 3.06 (2H, d, *J*=6.0 Hz), 1.98—1.82 (4H, m). ESI-MS *m/z*: 279 [Calcd for C₁₂H₁₁BrN₂O: 279 (M+H)]. *Anal.* Calcd for C₁₂H₁₁BrN₂O: C, 51.63; H, 3.97; N, 10.04. Found: C, 51.65; H, 3.95; N, 10.04.

2-Iodo-6,7,8,9-tetrahydro-11*H*-pyrido[2,1-*b*]quinazolin-11-one (1bd**)** Yellow needles: mp 286—287 °C. ¹H-NMR (DMSO-*d*₆, 250 MHz) δ: 8.41 (1H, d, *J*=2.0 Hz, H1), 8.20 (1H, dd, *J*=8.8, 2.3 Hz, H3), 7.58 (1H, d, *J*=8.8 Hz, H4), 3.93 (2H, t, *J*=6.0 Hz), 3.08 (2H, d, *J*=6.0 Hz), 1.97—1.82

(4H, m). ESI-MS m/z : 327 [Calcd for $C_{12}H_{11}N_2O$: 327 (M+H)]. *Anal.* Calcd for $C_{12}H_{11}N_2O$: C, 44.19; H, 3.40; N, 8.59. Found: C, 44.21; H, 3.42; N, 8.61.

2-Hydroxy-6,7,8,9-tetrahydro-11H-pyrido[2,1-b]quinazolin-11-one (1be) White needles: mp 257—259 °C. 1H -NMR (DMSO- d_6 , 250 MHz) δ : 10.67 (br s, OH), 7.69 (d, 1H, $J=8.3$ Hz, H4), 7.45 (d, 1H, $J=0.8$ Hz, H1), 7.43 (dd, 1H, $J=8.3, 0.8$ Hz, H3), 3.95 (t, 2H, $J=5.6$ Hz), 3.11 (d, 2H, $J=5.6$ Hz), 1.92 (quintet, 2H, $J=5.6$ Hz), 1.87 (quintet, 2H, $J=5.6$ Hz). ESI-MS m/z : 217 [Calcd for $C_{12}H_{12}N_2O_2$: 217 (M+H)]. *Anal.* Calcd for $C_{12}H_{12}N_2O_2$: C, 66.65; H, 5.59; N, 12.96. Found: C, 66.62; H, 5.61; N, 12.97.

Luotonin A (3) The crude product was recrystallized from $CHCl_3$ to yield pale yellow needles (78%): mp 281—283 °C (dec.) (lit.²³) mp 252 °C (dec.), lit.⁵¹ mp 281—283 °C. Spectral data were identical to those reported previously.²³

Tryptanthrin (4) The crude product was recrystallized from EtOH to yield 4 (85%) as yellow needles: mp 266—267 °C (lit.²⁶) mp 267—268 °C. Spectral data were identical to those reported previously.¹¹

Rutaecarpine (5) The crude product was recrystallized from CH_3OH to yield 5 (82%) as white needles: mp 259—260 °C (lit.²⁷) mp 258 °C. Spectral data were identical to those reported previously.⁵²

10b: A solution of anthranilic acid (13.72 g, 0.10 mol) and $SOCl_2$ (50 g) in dry benzene (500 ml) was refluxed for 2 h. The solvent was then evaporated under reduced pressure at room temperature to afford iminoketene as an oily liquid, to which a solution of 2-pyrrolidinone (7.20 g, 0.085 mol) in dry benzene (100 ml) was added. After setting overnight at room temperature, the precipitate formed was collected. Recrystallization from EtOH gave 20.0 g (91%) of white needles: mp 180 °C. IR (KBr) cm^{-1} : 3200—2500, 1702, 1644. 1H -NMR (250 MHz, $CDCl_3$) δ : 10.92 (br s, D_2O exchangeable, OH), 8.22 (1H, dd, $J=7.5, 1.3$ Hz, H6), 8.19 (1H, d, $J=7.8$ Hz, H3), 7.78 (1H, td, $J=8.5, 1.3$ Hz, H5), 7.54 (1H, td, $J=8.5, 1.3$ Hz, H4), 4.15 (2H, t, $J=6.2$ Hz), 3.63 (2H, t, $J=6.2$ Hz), 2.18 (2H, quintet, $J=6.2$ Hz), 2.05 (2H, quintet, $J=6.2$ Hz). ^{13}C -NMR (62.5 MHz, $CDCl_3$) δ : 160.55, 158.51, 136.87, 136.55, 128.98, 127.49, 119.99, 117.85, 43.62, 27.82, 21.08, 17.92, 17.16. ESI-MS m/z : 219 [Calcd for $C_{12}H_{14}N_2O_2$ (M+H): 219]. MS m/z : 218 (M^+ , 12) 200 ($M^+ - H_2O$, 100), 173 ($M^+ - COOH$, 4), 160 (11). *Anal.* Calcd for $C_{12}H_{14}N_2O_2$: C, 66.04; H, 6.47; N, 12.84. Found: C, 66.08; H, 6.43; N, 12.89.

2,3-Tetramethylene-4(3H)-quinazolinone (6,7,8,9-Tetrahydro-11H-pyrido[2,1-b]quinazolin-11-one, 1ba) A mixture of **10b** (4.36 g, 0.02 mol) in polyphosphoric acid (40 ml) was heated at 100—120 °C for 1 h. The reaction mixture was poured to ice (400 g) and the resulting mixture was made basic with 1 N NaOH. The precipitate formed was collected and recrystallized from EtOH to afford white needles (3.48 g, 87%): mp 98—99 °C (lit.²) mp 98.5—99.5 °C. 1H -NMR (250 MHz, $CDCl_3$) δ : 8.23 (1H, dd, $J=8.0$ Hz, H1), 7.68 (1H, td, $J=8.3, 1.3$ Hz, H2), 7.55 (1H, d, $J=8.0$ Hz, H4), 7.39 (1H, td, $J=8.0, 1.0$ Hz, H3), 4.05 (2H, t, $J=6.2$ Hz), 2.97 (2H, t, $J=6.5$ Hz), 2.01—1.87 (4H, m).

Acknowledgement Financial support from the Korean Research Foundation Grant (KRF-2005-041-E00496) is gratefully acknowledged. DHK and CSS are recipients of the BK-21 scholarship. Currently, KCJ is a student at the Department of Biochemistry, University of Iowa, Iowa City, IA 52242, U.S.A.

References and Notes

- Chatterjee A., Ganguly M. G., *Phytochemistry*, **7**, 307—311 (1968).
- Fitzgerald J. S., Johns S. R., Lamberton J. A., Redcliffe A. H., *Aust. J. Chem.*, **19**, 151—159 (1966).
- Johns S. R., Lamberton J. A., *Chem. Commun.*, **1965**, 267 (1965).
- Späth E., Platzer N., *Ber. Dtsch. Chem. Ges. B*, **68**, 2221—2226 (1935).
- Stephen T., Stephen H., *J. Chem. Soc.*, **1956**, 4694—4695 (1956).
- Petersen S., Tietze E., *Liebigs Ann. Chem.*, **623**, 166—176 (1959).
- Kamal A., Shankaraiah N., Devaiah V., Reddy K. L., *Tetrahedron Lett.*, **47**, 9025—9028 (2006).
- Hamid A., Elomri A., Daich A., *Tetrahedron Lett.*, **47**, 1777—1781 (2006).
- Gil C., Braese S., *Chemistry-A European Journal*, **11**, 2680—2688 (2005).
- Liu J.-F., Ye P., Sprague K., Sargent K., Yohannes D., Baldino C. M., Wilson C. J., Ng S.-C., *Org. Lett.* **7**, 3363—3366 (2005).
- Lee E.-S., Park J. G., Jahng Y., *Tetrahedron Lett.*, **44**, 1883—1886 (2003).
- Yadav J. S., Reddy B. V. S., *Tetrahedron Lett.*, **43**, 1905—1907 (2002).
- Nishiyama Y., Hirose M., Kitagaito W., Sonoda N., *Tetrahedron Lett.*, **43**, 1855—1858 (2002).
- Akazone M., Kondo T., Watanabe Y., *J. Org. Chem.*, **58**, 310—312 (1993).
- Mhsaske S. B., Argade N. P., *J. Org. Chem.*, **66**, 9038—9040 (2001).
- Takeuchi H., Matsushita Y., Eguchi S., *J. Org. Chem.*, **56**, 1535—1537 (1991).
- Dunn A. D., Kinnear K. I., *J. Heterocycl. Chem.*, **23**, 53—57 (1986).
- Kamal A., Ramana K. V., Rao M. V., *J. Org. Chem.*, **66**, 997—1001 (2001).
- Kametani T., Loc C. V., Higa T., Koizumi M., Ihara M., Fukumoto K., *J. Am. Chem. Soc.*, **99**, 2306—2309 (1977).
- Onaka T., *Tetrahedron Lett.*, **1971**, 4387—4390 (1971).
- Amin A. H., Mehta D. R., *Nature (London)*, **183**, 1317 (1959).
- Wu X., Qin G., Cheung K. K., Cheng K. F., *Tetrahedron*, **53**, 13323—13328 (1997).
- Ma Z. Z., Hano Y., Nomura T., Chen Y.-J., *Heterocycles*, **46**, 541—546 (1997).
- Brufani M., Fedeli W., Mazza F., Gerhard A., Keller-Schierlein W., *Experientia*, **27**, 1249—1250 (1971).
- Schindler F., Zähler H., *Arch. Microbiol.*, **79**, 187—203 (1971).
- Bergman J., Egestad B., Lindström J. O., *Tetrahedron Lett.*, **1977**, 2625—2626 (1977).
- Asahina Y., Kashiwaki K., *J. Pharm. Soc. Jpn.*, **405**, 1293 (1915).
- Bergman J., "The Alkaloids, Chemistry and Pharmacology," Vol. XXI, ed. by Brossi A., Academic Press, New York, 1983, pp. 29—54.
- von Nientowski S., *J. Prakt. Chem.*, **51**, 564—572 (1895).
- Brown D. J., "Quinazolines, Supplement I," Interscience Publishers, New York, 1996, pp. 1—150.
- Armarego W. L. F., "Fused Pyrimidines," ed. by Brown D. J., Interscience Publishers, New York, 1967, pp. 89—218.
- Morris R. S., Handford W. F., Adams R., *J. Am. Chem. Soc.*, **57**, 951—954 (1935).
- Möhrle H., Seidel C. M., *Chem. Ber.*, **106**, 1595—1600 (1973).
- Nesterova I. N., Radkevich T. P., Granik V. G., *Khim.-Farm. Zhur.*, **25**, 28—30 (1991) [*Chem. Abstr.*, **116**, 41412 (1992)].
- Japan Kokai Tokkyo Koho, JP 83-190614 (1984).
- Manhas M. S., Amin S. G., *J. Heterocycl. Chem.*, **13**, 903—906 (1976).
- Levy P. R., Stephen H., *J. Chem. Soc.*, **1956**, 985—988 (1956).
- Mason J. J., Bergman J., *Org. Biomol. Chem.*, **5**, 2486—2490 (2007).
- Zhou H.-B., Liu G.-S., Yao Z.-J., *J. Org. Chem.*, **72**, 6270—6272 (2007).
- Nacro K., Zha C., Guzzo P. R., Herr R. J., Peace D., Friedrich T. D., *Bioorg. Med. Chem.*, **15**, 4237—4246 (2007).
- Servais A., Azzouz M., Lopes D., Courillon C., Malacria M., *Angew. Chem. Int. Ed. Engl.*, **46**, 576—579 (2007).
- Bowman W. R., Elsegood M. R. J., Stein T., Weaver G. W., *Org. Biomol. Chem.*, **5**, 103—113 (2007).
- Kamal A., Shankaraiah N., Devaiah V., Reddy K. L., *Tetrahedron Lett.*, **47**, 9025—9028 (2006).
- Low yield of this type of reaction has been reported: see Bergman J., Bergman S., *J. Org. Chem.*, **50**, 1246—1255 (1985).
- Shakhidoyatov Kh. M., Irisbaev A., Yun L. M., Oripov E., Kadyrov Ch. Sh., *Khim. Get. Soed.*, **11**, 1286—1291 (1976).
- Koizumi M., Matsuura I., Murakami Y., Japan Patent 52077093 (1977).
- Devi G., Kapil R. S., Poli S. P., *Ind. J. Chem. B*, **14**, 354—356 (1976).
- Sugasawa T., Toyota T., Sasakura K., Hidaka T., *Chem. Pharm. Bull.*, **19**, 1971—1974 (1971).
- Abramovitch R. A., Shapiro D., *J. Chem. Soc.*, **1956**, 4589—4592 (1956).
- Asahina Y., Manske R. H. F., Robinson R., *J. Chem. Soc.*, **1927**, 1708—1710 (1927).
- Ma Z. Z., Hano Y., Nomura T., Chen Y.-J., *Heterocycles*, **51**, 1593—1596 (1999).
- Lee S. H., Kim S. I., Park J. G., Lee E.-S., Jahng Y., *Heterocycles*, **55**, 1555—1560 (2001).