

# UC San Diego

## UC San Diego Previously Published Works

### Title

Open-Source Sequence Clustering Methods Improve the State Of the Art.

### Permalink

<https://escholarship.org/uc/item/25b0j91w>

### Journal

mSystems, 1(1)

### ISSN

2379-5077

### Authors

Kopylova, Evguenia  
Navas-Molina, Jose A  
Mercier, Céline  
et al.

### Publication Date

2016

### DOI

10.1128/msystems.00003-15

Peer reviewed



# Open-Source Sequence Clustering Methods Improve the State Of the Art

Evguenia Kopylova,<sup>a</sup> Jose A. Navas-Molina,<sup>a,b</sup> Céline Mercier,<sup>c</sup> Zhenjiang Zech Xu,<sup>a</sup> Frédéric Mahé,<sup>d</sup> Yan He,<sup>e</sup> Hong-Wei Zhou,<sup>e</sup> Torbjørn Rognes,<sup>f,g</sup> J. Gregory Caporaso,<sup>h</sup> Rob Knight<sup>a,b</sup>

Department of Pediatrics, UCSD School of Medicine, La Jolla, California, USA<sup>a</sup>; Department of Computer Science and Engineering, University of California, San Diego, La Jolla, California, USA<sup>b</sup>; Laboratoire d'Ecologie Alpine (LECA), CNRS UMR 5553, Université Grenoble Alpes, Grenoble, France<sup>c</sup>; Department of Ecology, University of Kaiserslautern, Kaiserslautern, Germany<sup>d</sup>; Department of Environmental Health, State Key Laboratory of Organ Failure Research, Guangdong Provincial Key Laboratory of Tropical Disease Research, School of Public Health and Tropical Medicine, Southern Medical University, Guangzhou, Guangdong, China<sup>e</sup>; Department of Informatics, University of Oslo, Oslo, Norway<sup>f</sup>; Department of Microbiology, Oslo University Hospital, Rikshospitalet, Oslo, Norway<sup>g</sup>; Department of Biological Sciences, Northern Arizona University, Flagstaff, Arizona, USA<sup>h</sup>

**ABSTRACT** Sequence clustering is a common early step in amplicon-based microbial community analysis, when raw sequencing reads are clustered into operational taxonomic units (OTUs) to reduce the run time of subsequent analysis steps. Here, we evaluated the performance of recently released state-of-the-art open-source clustering software products, namely, OTUCLUST, Swarm, SUMACLUSt, and SortMeRNA, against current principal options (UCLUST and USEARCH) in QIIME, hierarchical clustering methods in mothur, and USEARCH's most recent clustering algorithm, UPARSE. All the latest open-source tools showed promising results, reporting up to 60% fewer spurious OTUs than UCLUST, indicating that the underlying clustering algorithm can vastly reduce the number of these derived OTUs. Furthermore, we observed that stringent quality filtering, such as is done in UPARSE, can cause a significant underestimation of species abundance and diversity, leading to incorrect biological results. Swarm, SUMACLUSt, and SortMeRNA have been included in the QIIME 1.9.0 release.

**IMPORTANCE** Massive collections of next-generation sequencing data call for fast, accurate, and easily accessible bioinformatics algorithms to perform sequence clustering. A comprehensive benchmark is presented, including open-source tools and the popular USEARCH suite. Simulated, mock, and environmental communities were used to analyze sensitivity, selectivity, species diversity (alpha and beta), and taxonomic composition. The results demonstrate that recent clustering algorithms can significantly improve accuracy and preserve estimated diversity without the application of aggressive filtering. Moreover, these tools are all open source, apply multiple levels of multithreading, and scale to the demands of modern next-generation sequencing data, which is essential for the analysis of massive multidisciplinary studies such as the Earth Microbiome Project (EMP) (J. A. Gilbert, J. K. Jansson, and R. Knight, *BMC Biol* 12:69, 2014, <http://dx.doi.org/10.1186/s12915-014-0069-1>).

**KEYWORDS:** sequence clustering, operational taxonomic units, microbial community analysis, amplicon sequencing

Current DNA sequencing technologies generate hundreds of gigabytes of data in a single run and have enabled new detailed investigations into the human microbiome (1–3) and initiatives to characterize the Earth ecosystem's microbiome, such as the EMP. Analysis of microbiome datasets typically begins by clustering raw biological

Received 4 October 2015 Accepted 10 January 2016 Published 9 February 2016


**Citation** Kopylova E, Navas-Molina JA, Mercier C, Xu ZZ, Mahé F, He Y, Zhou H-W, Rognes T, Caporaso JG, Knight R. 2016. Open-source sequence clustering methods improve the state of the art. *mSystems* 1(1):e00003-15. doi: [10.1128/mSystems.00003-15](https://doi.org/10.1128/mSystems.00003-15).

**Editor** Nicola Segata, University of Trento

**Copyright** © 2016 Kopylova et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Evguenia Kopylova, [jenya.kopylov@gmail.com](mailto:jenya.kopylov@gmail.com).

For a commentary on this article, see <http://doi.org/10.1128/mSystems.00027-16>

 Open-source sequence clustering methods improve the state of the art

sequence reads into operational taxonomic units (OTUs) based on sequence similarity, a process frequently referred to as OTU clustering or delineating. Sequencing costs are dropping faster than Moore's law (4), increasing the need for efficient and accurate OTU clustering software. QIIME (5) has been using UCLUST (6) as the default clustering method since UCLUST's publication (corresponding to QIIME version 1.0.0), due to its increase in performance over other popular tools, such as BLAST (7), DOTUR (8), or CD-HIT (9–11). However, UCLUST and USEARCH are closed-source software (the 64-bit versions, which are needed to handle large datasets, require an expensive license, even for academic use) and have limited documentation (<http://www.drive5.com/usearch/>). Moreover, based on UCLUST documentation ([http://www.drive5.com/uclust/uclust\\_userguide\\_1\\_1\\_579.pdf](http://www.drive5.com/uclust/uclust_userguide_1_1_579.pdf)), the allegedly serial implementation is impractical for massive high-throughput sequencing data. More accurate, faster, community-accessible tools are needed to overcome these challenges.

Within the previous 2 years, four new sequence-clustering tools have emerged: OTUCLUST from the Micca package (12), Swarm (13, 14), SUMACLUSt (C. Mercier, F. Boyer, E. Kopylova, P. Taberlet, A. Bonin, and E. Coissac, submitted for publication), and SortMeRNA (15). These tools include open-source implementation, and the latter three implement multilevel parallelization, providing excellent potential alternatives to UCLUST. In this study, we evaluated these new open-source tools and compared them against UCLUST and USEARCH, two commonly used options available in QIIME, UPARSE (16), the latest USEARCH amplicon analysis pipeline, and the three hierarchical clustering algorithms available in mothur (17).

## RESULTS

**Software description.** OTU clustering can be performed in three different ways (18): closed reference, *de novo*, and open reference. In the closed-reference approach, the input sequences are clustered against a reference sequence database. In *de novo* clustering, the input sequences are grouped based on pairwise similarity among all sequences in the data set. The open-reference approach (19) begins by running a closed-reference step, which is followed by a *de novo* step that clusters the sequences that fail closed-reference assignment.

Swarm (13, 14) is a *de novo* clustering algorithm based on an unsupervised single-linkage-clustering method that reduces the impact of clustering parameters on the resulting OTUs by avoiding arbitrary global clustering thresholds and input sequence ordering dependence. Swarm builds OTUs in two steps: (i) an initial set of OTUs is constructed by iteratively agglomerating similar amplicons, and (ii) amplicon abundance values are used to reveal OTUs' internal structures and to break them into sub-OTUs, if necessary.

OTUCLUST (12) and SUMACLUSt are also *de novo* clustering algorithms; both are based on a greedy strategy in which the clusters are constructed incrementally by comparing an abundance-ordered list of input sequences against the representative set of already-chosen sequences (initially empty) (20). A similar approach is also used by UCLUST and CD-HIT, but OTUCLUST and SUMACLUSt have been designed to perform exact sequence alignment, rather than relying on fast heuristics. In addition, OTUCLUST performs its own sequence dereplication and chimera removal (via UCHIME [21]).

mothur (17) implements three *de novo* clustering algorithms (nearest neighbor, furthest neighbor, and average neighbor) which cluster sequences based on genomic distance. In nearest neighbor (single linkage), a sequence is linked to an OTU if it is similar to any other sequence in that OTU, in furthest neighbor (complete linkage), a sequence is linked to an OTU if it is similar to all other sequences in that OTU, and in average neighbor, a sequence is linked to an OTU if it is similar to the averaged differences between all other sequences in that OTU. More details on these algorithms are available in references 8 and 22.

SortMeRNA (15) is suited for closed-reference OTU clustering. It is a local sequence alignment tool, in that it searches for optimal regions of similarity between two sequences. Query sequences (e.g., rRNA amplicons) are searched against a reference

**TABLE 1** Description of studies used in analysis

Data set	QIIME identity	Reference	Gene	Region	No. of reads	No. of samples	Read length	Platform
Simulated								
sim_even		24	16S	V4	107,600	1	150	ART
sim_staggered		24	16S	V4	107,025	1	150	ART
Mock								
Bokulich_2	1685	25	16S	V4	6,938,836	4	189–251	MiSeq
Bokulich_3	1686	25	16S	V4	3,594,237	4	114–151	MiSeq
Bokulich_6	1688	25	16S	V4	250,903	1	114–150	MiSeq
mock_nematodes		26	18S	V4	9,061	1	54–305	GS FLX
Genuine								
canadian_soil	632	27	16S	V4	2,966,053	13	76–100	HiSeq
body_sites	449	28	16S	V2	886,630	602	117–351	GS FLX
global_soil	2107	29	18S	V9	9,252,764	57	119–151	HiSeq

database, and an  $E$  value threshold is applied to evaluate the quality of resulting alignments. In SortMeRNA 2.0, the reference sequence achieving the lowest  $E$  value when aligned with a query sequence is chosen as the OTU centroid for that query. In addition to passing the  $E$  value threshold, the query must also have sufficient percent identity and coverage (both set to 97% by default). Contrary to UCLUST, the run time of SortMeRNA is not affected by reducing these thresholds (e.g., clustering at 60% identity).

UCLUST and USEARCH (versions 5.2 and 6.1) are supported in QIIME (v1.8.0). Both tools can perform *de novo*, closed-reference, and open-reference (except for USEARCH 5.2) clustering. In QIIME's implementation, USEARCH 5.2 is executed via a pipeline closely shadowing otupipe (6, 21) to cluster OTUs, and USEARCH 6.1 performs chimera checking in an external script. UPARSE (16) is the latest *de novo* amplicon analysis pipeline from USEARCH; it applies stringent quality filtering and length trimming to remove erroneous reads and implements a novel greedy algorithm that performs OTU clustering and chimera removal concurrently.

**Experimental design.** Swarm 1.2.19, SUMACLUSt 1.0.00, and SortMeRNA 2.0 have been integrated into QIIME 1.9.0 and can be used through QIIME's three different OTU clustering commands (18): `pick_closed_reference_otus.py`, `pick_de_novo_otus.py`, and `pick_open_reference_otus.py`.

A variety of datasets were chosen to evaluate the performance of these open-source OTU clustering approaches relative to QIIME's UCLUST/USEARCH-based OTU clustering approaches as well as UPARSE (see Table 1 for details). Two 16S rRNA gene simulated datasets were generated as FASTQ files. The first one (`sim_even`) represents an even distribution of 1,076 species, randomly subsampled from the Greengenes 97% database and computationally amplified at the same depth (100 reads/amplicon) and length (150 bp) using PrimerProspector (23) for extracting the V4 region and the ART (24) simulator for amplification and sequencing simulation. The second data set (`sim_staggered`) represents the same 1,076 species as the `sim_even` data set but amplified at different (random) species abundance levels. We used four different previously published mock community data sets: three 16S rRNA gene mock community data sets (`Bokulich_2`, `Bokulich_3`, and `Bokulich_6`) from Bokulich et al. (25) and an 18S gene (`mock_nematodes`) data set from Porazinska et al. (26). Finally, we also used three previously published natural data sets: a 16S rRNA gene soil data set (`canadian_soil`) from Neufeld et al. (27), a 16S rRNA gene human data set (`body_sites`) from Costello et al. (28), and an 18S rRNA gene soil data set (`global_soil`) from Ramirez et al. (29).

**Performance.** All tools were run with default parameters. Input FASTA files for Swarm, SUMACLUSt, and SortMeRNA were generated using QIIME's demultiplexing and quality filtering workflow. Input FASTA files for OTUCLUST, mothur, and UPARSE were demultiplexed using QIIME and quality filtered using each tool's recommended

**TABLE 2** Benchmark summary<sup>a</sup>

Software	Data set																										
	Simulated						Mock						Genuine														
	sim_even (V4)		sim_staggered (V4)		Bokulich_2 (V4)		Bokulich_3 (V4)		Bokulich_6 (V4)		body_sites (V2)		canadian_soil (V4)		global_soil (V9, 18S)												
OTUs	PD	F <sub>1</sub>	OTUs	PD	F <sub>1</sub>	OTUs	PD	F <sub>1</sub>	OTUs	PD	F <sub>1</sub>	OTUs	M <sup>2</sup>	ρ	OTUs	M <sup>2</sup>	ρ										
<i>de_novo</i>	swarm	1,042	101.50	0.84	1,035	104.00	0.83	7,084	[4-50]	0.48	6,349	[4-35]	0.50	1,223	39.41	0.54	14,184	0.19	0.96	59,688	0.16	0.94	80,321	0.87	0.98		
	sumacrust	1,031	104.06	0.83	1,022	109.92	0.83	9,575	[4-157]	0.38	13,982	[4-190]	0.41	3,317	90.80	0.52	7,103	0.18	0.99	74,284	0.14	0.87	60,781	0.50	0.96		
	uparse_q3	1,013	104.02	0.84	997	110.57	0.84						199	9.22	0.59	156	0.38	0.29	11,259	0.03	0.85						
	uparse_q16	972	100.74	0.84	806	93.28	0.78						31	3.53	0.45	108	0.36	0.26	6,275	0.06	0.75						
	uclust	1,045	105.37	0.83	1,035	110.42	0.83	20,084	[5-234]	0.40	21,929	[5-236]	0.40	4,397	105.37	0.52	11,204	0.00	1.00	91,143	0.00	1.00	82,642	0.00	1.00		
	usearch52	1,035	106.09	0.83	1,015	110.76	0.81	1,522	[3-22]	0.50	2,602	[4-28]	0.55	798	22.86	0.55	3,903	0.17	0.94	47,679	0.05	0.94	41,668	0.93	0.98		
	usearch61	1,049	104.85	0.84	1,034	110.68	0.83	22,987	[7-313]	0.39	24,704	[7-292]	0.41	4,635	123.04	0.51	14,483	0.18	0.99	102,435	0.06	0.99	102,211	0.48	0.98		
	otuclust_q3	996	111.03	0.84	953	106.88	0.81						438	[2-8]	0.61	228	10.36	0.61	2,753	0.18	0.85	18,373	0.08	0.82			
	otuclust_q20	996	111.03	0.84	953	106.88	0.81						314	[2-6]	0.65	113	7.20	0.58	2,654	0.16	0.85	18,373	0.07	0.81			
	mothur_near	957	110.09	0.82	949	110.45	0.81						1,600	[2-51]	0.44	447	23.63	0.54	806	0.45	0.12	31,546	0.06	0.76	11,440	0.53	0.76
	mothur_fur	978	109.22	0.82	970	109.86	0.81						28,808	[5-263]	0.40	5,159	75.05	0.51	3,358	0.22	0.23	92,887	0.03	0.86	32,378	0.56	0.78
	mothur_avg	963	109.99	0.82	959	110.98	0.82						13,255	[4-176]	0.41	2,314	55.90	0.51	2,491	0.26	0.11	83,664	0.05	0.86	20,809	0.49	0.72
	<i>closed_ref</i>	usearch61	F <sub>1</sub> tax	1,275	129.19	0.83	1,267	127.50	0.82	1,027	[5-26]	0.53	614	[4-18]	0.59	631	26.02	0.61	5,982	0.06	0.98	13,808	0.06	0.96	3,784	0.50	0.55
		F <sub>1</sub> OTUs			0.68			0.69																			
uclust		F <sub>1</sub> tax	1,238	127.59	0.83	1,225	126.02	0.84	1,053	[5-27]	0.53	557	[5-18]	0.57	547	25.03	0.60	5,446	0.00	1.00	13,659	0.00	1.00	305	0.00	1.00	
		F <sub>1</sub> OTUs			0.69			0.70																			
sortmerma		F <sub>1</sub> tax	1,072	122.75	0.82	1,067	121.89	0.81	396	[4-15]	0.53	290	[4-13]	0.61	382	19.47	0.57	6,174	0.06	0.99	13,281	0.06	0.98	255	0.34	0.75	
<i>open_ref</i>	usearch52	F <sub>1</sub> OTUs	1,001	115.38	0.80	980	113.39	0.78	571	[5-30]	0.54	331	[5-22]	0.64	315	18.24	0.59	3,355	0.08	0.97	4,121	0.04	0.79	5,763	0.48	0.19	
		F <sub>1</sub> tax			0.70			0.68																			
	uclust		1,262	106.12	0.83	1,245	111.29	0.83	10,169	[3-97]	0.40	4,170	[3-104]	0.42	4,109	93.67	0.48	12,442	0.00	1.00	87,936	0.00	1.00	37,380	0.00	1.00	
sortmerma		1,072	104.77	0.82	1,085	111.80	0.81	9,272	[3-132]	0.39	2,649	[3-140]	0.41	2,727	88.56	0.51	10,242	0.06	0.98	79,363	0.03	0.82	35,345	0.12	0.92		
sumacrust		1,304	106.04	0.83	1,293	112.36	0.83	9,414	[3-108]	0.40	3,966	[3-126]	0.41	3,421	80.89	0.53	12,807	0.06	0.97	87,300	0.06	0.80	43,175	0.10	0.94		

<sup>a</sup>OTU counts do not include singletons. F measure (F<sub>1</sub>) is for assigned taxonomies at the genus level. The phylogenetic diversity (PD) whole-tree column for Bokulich\_2 and Bokulich\_3 represent PD intervals across various sampling depths. Procrustes M<sup>2</sup> (the sum of the squared deviations or the dissimilarity of two datasets for UniFrac PCoA) and rho (Pearson's correlation coefficient for taxonomies at genus level) values are with respect to UCLUST (default for QIIME versions 1.0.0 to 1.9.1). Monte Carlo P values were not included, since all values were <0.05 except for *de novo* usearch52 versus uclust (P = 0.09). The darkest blue shades represent the highest F<sub>1</sub> scores, while the darkest red shades represent results closest to those obtained with UCLUST.

standard operation procedure (SOP) (see Materials and Methods). Sequence filtering for OTUCLUST was performed with default quality score cutoffs of 20 (labeled as OTUCLUST\_q20) and 3 (default in QIIME based on the results reported in reference 25, labeled as OTUCLUST\_q3). UPARSE was run using the recommended settings with truncation lengths of 150 bp and 250 bp; similarly to OTUCLUST, runs were performed with a default quality score cutoff of 16 (labeled as UPARSE\_q16) and additionally with a quality score cutoff of 3 (labeled as UPARSE\_q3). Biological observation matrix (BIOM) format (30) tables were used as input to post clustering analyses. Taxonomy for reported OTUs was assigned using the RDP Classifier (31) against the 97% representative databases for Greengenes (32, 33) (version 13.8) and Silva (34) (version 111) for all methods. Performance was evaluated using a variety of metrics, including the accuracy of OTU and taxonomic assignments, alpha diversity (within-sample diversity), beta diversity (between-sample diversity), and taxonomic correlation. All tools showed increased precision after the removal of singleton OTUs (OTUs consisting of only one sequence), so all results presented here have had singleton OTUs removed. Table 2 summarizes basic performance results for all software.

**Expected community composition: sensitivity and specificity. (i) Simulated data.** For *de novo* clustering, most tools report F measures (or F<sub>1</sub> score, a metric that assesses the accuracy of taxonomic composition and observed OTUs, with a range from 0 to 1, where 1 is the best score) of 0.82 to 0.84 (sim\_even) and 0.81 to 0.83 (sim\_staggered) at the genus level (Table 2). Variation in results was emphasized in sim\_staggered, where UPARSE\_q16 reported the lowest F measure (0.78) as a result of stringent read filtering that removed nearly 95% of the reads prior to clustering. UPARSE\_q3 removed roughly 4% of the reads and reported improved results on a par with those of Swarm, SUMACRUST, UCLUST, and USEARCH61. The highest F measure (0.83 to 0.84) and number of OTUs closest to the expected one (1,076) were reported for software using input files from QIIME's method of sequence filtering (Swarm, SUMACRUST, UCLUST, and USEARCH61). All tools except UPARSE\_q16 reported comparable alpha diversity phylogenetic diversity (PD) whole-tree (35) (a measure of diversity which considers the phylogenetic differences between species) values (mean of 109.21 with a standard deviation of 2.16 [Table 2]).

Among the closed-reference methods, SortMeRNA yields the fewest OTUs while achieving comparable or higher F measures for assigned taxonomy (F<sub>1</sub> tax in Table 2) and OTUs (F<sub>1</sub> OTUs in Table 2) and reported a phylogenetic diversity (121.89) closest to

the ground truth (123.75) in comparison to UCLUST (126.02), USEARCH 5.2 (113.39), and USEARCH 6.1 (127.50). This is a result of SortMeRNA's more exhaustive search for better alignments, which can increase run time but becomes imperative when short reads are aligned against a highly conservative set of sequences, such as the rRNA gene. The complete Greengenes database contains over a million rRNAs, and almost 73% of all full-length V4 regions (~250 nucleotides) are not unique. This emphasizes the highly conservative nature of rRNA, even in this hypervariable region, and suggests the need for thorough searches to ensure higher-quality alignments (especially for read lengths that do not cover the entire region). At the genus level of taxonomy, all tools report F measures of 0.80 to 0.83 (sim\_even) and 0.78 to 0.84 (sim\_staggered), which can be attributed to the many-to-one relationship between OTUs and taxonomy strings for the Greengenes 97% database.

For open-reference clustering, QIIME's subsampling pipeline combining SortMeRNA and SUMACLUSt reports the fewest OTUs in comparison to UCLUST and USEARCH 6.1. The F measure is 0.82 to 0.83 (sim\_even) and 0.81 to 0.83 (sim\_staggered) for all tools, which is in agreement with *de novo* and closed-reference results. These results are expected given the nature of open-reference clustering, which combines the closed-reference approach with the *de novo* approach.

**(ii) Mock communities.** Results for Bokulich\_2 (and Bokulich\_3 for UPARSE) are unavailable for UPARSE, OTUCLUST, and mothur due to significant memory, run time, and disk space requirements, respectively. All other methods were compared against the expected taxonomic composition for each data set. In addition, Pearson's correlation coefficient was computed to measure the relatedness of taxonomic assignment between all pairs of tools (see column rho in Table 2 for all tools versus UCLUST). Values can range between -1 and 1, with -1 indicating a negative correlation, 0 indicating no correlation, and 1 indicating a positive correlation (strong relationship).

For *de novo* clustering, USEARCH 5.2, UPARSE\_(q3, q16), OTUCLUST\_(q3, q20), and mothur\_nearest frequently reported the lowest number of OTUs, the lowest number of observed taxa, and the highest F measure (Table 2). Since the F measure is computed using true-positive taxonomies based on the expected composition, possible contamination species (false positives) are unaccounted for. However, false-positive taxonomies can also arise from OTUs formed by chimeric sequences (sequences from two organisms that bind together during PCR and are subsequently sequenced as a single read) or incorrect assignment by taxonomy assignment tools. To investigate the origins of false-positive taxonomies reported by the tools, we checked all OTUs for chimeras using UCHIME (20) and mapped the nonchimeric OTUs against BLAST's NT database using MEGABLAST. Most of the false-positive taxa were not wholly comprised of chimeric OTUs (meaning that the collection of OTUs mapping to the same taxa was composed of chimeric and genuine sequences), and the majority of such nonchimeric taxa consisted of OTUs mapping with an *E* value of  $<1e-50$  to BLAST's NT database (e.g., in Table 3 there are 57 false-positive taxa reported by SUMACLUSt, but only 4 of those taxa are fully comprised of chimeric OTUs [FP-chimeric], and 99% of OTUs representing the remaining 53 nonchimeric taxa mapped with high similarity to BLAST's NT database). Not surprisingly, all false-positive taxa whose OTUs mapped with  $<97\%$  similarity (FP-other) are less abundant than the taxa whose OTUs map with  $\geq 97\%$  similarity (FP-known) and significantly less abundant than true-positive taxa (see Fig. S1 to S3 in the supplemental material). In fact, false-positive taxonomies (especially FP-other and FP-chimeric) comprise few and low-abundance OTUs, which can be analyzed and filtered out if necessary after clustering. Since UPARSE filters out a large fraction of presumably erroneous reads (even prior to chimera checking), it can detect the most abundant species (as can other tools) but also potentially overlook low-abundance species. For the Bokulich\_2 and Bokulich\_3 data sets, the top 20 most abundant genera follow a similar relative abundance distribution for all *de novo* tools, which is a direct reflection of hundreds of thousands of reads representing each expected genus in these data sets. However, for the much smaller data set Bokulich\_6,

**TABLE 3** Sensitivity and selectivity statistics for assigned taxonomies at genus level, Bokulich\_2<sup>a</sup>

Software	No. of OTUs (no singletons)	P	R	F1	No. of taxonomies					
					FP			TP	FN	Total
<i>De novo</i>										
usearch52	1,522	0.34	1	0.5	18	0	35	5	13	17
Swarm	7,084	0.32	1	0.48	18	0	38	7	22	9
uclust	20,084	0.25	1	0.4	18	0	53	4	15	34
usearch61	22,987	0.24	1	0.39	18	0	56	4	18	34
sumaclust	9,575	0.24	1	0.38	18	0	57	4	15	38
Closed reference										
usearch52	571	0.37	1	0.54	18	0	30	3	13	14
sortmerna	396	0.36	1	0.53	18	0	31	4	26	1
uclust	1,053	0.36	1	0.53	18	0	32	6	26	0
usearch61	1,027	0.36	1	0.53	18	0	32	4	28	0
Open reference										
uclust	10,169	0.25	1	0.4	18	0	52	4	19	29
usearch61	9,414	0.25	1	0.4	18	0	53	4	18	31
sortmerna_sumaclust	9,272	0.24	1	0.39	18	0	55	5	16	34

<sup>a</sup>P, precision; R, recall; F1, F measure, TP, true positive; FN, false negative; FP, false positive. The last three columns represent a refined breakdown of FP data, including false-positive taxonomies for which all comprising OTUs were classified as chimeric (using UCHIME) (chimeric), mapped to BLAST's NT database with  $\geq 97\%$  similarity (known), or mapped to BLAST's NT database with  $< 97\%$  similarity (other).


UPARSE\_q16 reported only half of the expected genera relative to all other tools, and the relative abundance of some of the genera significantly decreased (Fig. 1; PD values in Table 2). OTUCLUST, mothur\_nearest, mothur\_average, Swarm, and SUMACLUSt reported significantly fewer OTUs than UCLUST and USEARCH 6.1, as well as a lower alpha diversity (Tables 2 and 3). Thus, tools with lower false-positive rates accomplish this by more stringent quality control, but they are less suitable for finding lower-abundance genera.

For closed-reference clustering, SortMeRNA reported up to 60% fewer OTUs and a PD of about half that of UCLUST and USEARCH 6.1 (Table 2). On the genus taxonomy level, USEARCH 5.2 reported a high F measure (due to a lower number of false-positive genera), but unlike all other tools, the majority of false-positive genera are composed of reads mapping with  $\leq 97\%$  identity and coverage to BLAST's NT database (FP-other). In fact, all other tools filtered out a large portion of these false-positive reads due to insufficient identity matches to the reference database. The difference appears to be caused by USEARCH 5.2's identity definition (which does not consider insertions or deletions), which scores alignments higher than other tools. SortMeRNA generates taxonomic profiles similar to those obtained with other tools, with a Pearson's correlation coefficient of  $> 0.93$  (Fig. 1).

As expected for open-reference clustering, SortMeRNA combined with SUMACLUSt reported fewer OTUs than UCLUST and USEARCH 6.1 while preserving high accuracy and lower alpha diversity for both the number of observed OTUs and the phylogenetic diversity. Specific details can be found in the supplemental material.

The Pearson coefficient for comparisons between all tools and methods remained relatively stable, from  $\sim 0.99$  (Bokulich\_2) to 0.97 to 1 (Bokulich\_3) to 0.92 to 0.99 (Bokulich\_6), showing a strong relationship between all algorithms. The coefficient was lower in the cases where the taxonomy could not be assigned (e.g., 0.0273 for SortMeRNA versus UCLUST for data set Bokulich\_3) or significant filtering of sequences (e.g., 0.3719 for UCLUST versus UPARSE\_q16 for data set Bokulich\_6).


**Natural community composition.** Results for UPARSE\_q4 and UPARSE\_q16 are unavailable for the global\_soil data set due to memory limitations in the 32-bit version of UPARSE and for OTUCLUST due to significant run time (limited to one thread).



**FIG 1** Layered bar chart showing top 20 abundant genera, Bokulich\_6. The bars do not reach 1, since only a fraction (top 20) of taxonomies was illustrated.

In contrast to mock communities, the Pearson correlation for natural communities was much more variable (0.70 to 0.94 for the canadian\_soil data set, 0.28 to 0.99 for the body\_sites data set, and 0.19 to 0.98 for the global\_soil data set) (Table 2), highlighting differences between all clustering algorithms in a complex environment that are not immediately visible in either simulated or mock communities. These ranges do not take





**FIG 2** Taxonomic composition graph illustrating top 50 (per software) abundant genera, *body\_sites*. The bars do not reach 1, since only a fraction (top 50) of taxonomies was illustrated. *mothur* was run using recommended filtering (*trim.seqs* function) for 454 SOP and with QIIME's *split\_libraries\_fastq.py* to highlight the effect of different filtering methods.

into account outliers that were caused by an inconsistency with RDP assignments for the most abundant taxa (*Bokulich\_3*) and stringent filtering of reads by *UPARSE\_q16* (*Bokulich\_6*) (Fig. 1). QIIME, *UPARSE*, *OTUCLUST*, and *mothur* include different sequence filtering methods, which could be the major reason behind inconsistent taxonomic compositions (Pearson's correlation in Table 2; Fig. 2). As illustrated in Fig. 2, running *mothur* with sequences that were quality-filtered by *mothur* and QIIME produced significantly different taxonomic compositions. As expected, the highest correlation exists for studies with the longest reads and the largest number of reads per sample, showing that clustering results converge to the same conclusions with longer, higher-quality reads and deep sequencing (Fig. 3).


As with the mock-community results, all tools frequently reported fewer OTUs and lower alpha diversities than *UCLUST* and *USEARCH 6.1* (Table 2 and Fig. 4; also, see Fig. S4 and S5 in the supplemental material). Procrustes analysis (36) was used to compare unweighted UniFrac (37, 38) principal coordinates analysis (PCoA) (22) generated by all methods versus *UCLUST* (the current default OTU picker in QIIME). The Procrustes  $M^2$  metric for *body\_sites* and *canadian\_soil* was  $<0.3$  for most software (Table 2), indicating that beta diversity patterns are similar irrespective of the OTU clustering method used. Neither recommended nor relaxed quality filtering parameters for *UPARSE* worked well for the *body\_sites* data set, where 98.5% and 99.2% of reads were filtered out for *UPARSE\_q3* and *UPARSE\_q16* (with a trim length of 250 bp), respectively, resulting in very few remaining samples and high  $M^2$  values (Table 2; also, see Fig. S4 in the supplemental material). Although read quality filtering is an important preprocessing step, more work is required to regulate these parameters (perhaps by an automated estimation of optimal truncation length and quality), as they can be very sensitive to different types of data.

## DISCUSSION

We evaluated the performance of four recently published open-source sequence clustering tools against the widely used *mothur*, *UCLUST*, and *USEARCH* tools using

k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Propionibacteriaceae_g_Propionibacterium
k_Bacteria_p_Firmicutes_c_Bacillo_Bacillales_Other_Other
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Ruminococcaceae_g_Faecalibacterium
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Williamsiaceae_g_Williamsia
k_Bacteria_p_Proteobacteria_c_Alphaproteobacteria_o_Sphingomonadales_f_Sphingomonadaceae_Other
k_Bacteria_p_Proteobacteria_c_Alphaproteobacteria_o_Sphingomonadales_f_Sphingomonadaceae_g_Sphingobium
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Tissotellaceae_g_WAL_18550
k_Bacteria_k_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_Other_Other
k_Bacteria_p_Actinobacteria_c_Corinobacteriia_Corinobacteriales_f_Corinobacteriaceae_g_
k_Bacteria_p_Proteobacteria_c_Alphaproteobacteria_o_Rhodospirales_f_Rhodospiraceae_Other_Other
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Micromonosporaceae_g_Actinoplasma
k_Bacteria_p_Actinobacteria_c_MB_A2-108_o_0319-7114_f_g_
k_Bacteria_p_Proteobacteria_c_Alphaproteobacteria_o_Caulobacteriales_f_Caulobacteriaceae_Other
k_Bacteria_k_Bacteroidetes_c_Cytophagia_o_Cytophagales_f_Cytophagaceae_g_Dyadobacter
k_Bacteria_p_Actinobacteria_c_Corinobacteriia_Corinobacteriales_f_Corinobacteriaceae_g_Atopium
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Lachnospiraceae_Other
k_Bacteria_p_Terricoccales_f_Mollisutes_f_995_f_g_
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Verruococcaceae_Other
k_Bacteria_p_Proteobacteria_c_Alphaproteobacteria_o_Rhodospirales_f_Methylobacteriaceae_g_Methylobacterium
k_Bacteria_k_Actinobacteria_c_Actinobacteriia_Corinobacteriales_f_Corinobacteriaceae_g_Citriella
k_Bacteria_k_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Nocardiodaceae_Other
k_Bacteria_k_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Bifidobacteriaceae_g_
k_Bacteria_p_Acidobacteria_c_Solibacteriia_o_Solibacteriales_Other_Other
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Tissotellaceae_g_148
k_Bacteria_p_Proteobacteria_c_Gammaproteobacteria_o_Pseudomonadales_f_Moraxellaceae_g_Moraxella
k_Bacteria_k_Bacteroidetes_Other_Other_Other
k_Bacteria_p_Proteobacteria_c_Gammaproteobacteria_o_Pasturellales_f_Pasturellaceae_g_Haemophilus
k_Bacteria_p_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Paraprevotellaceae_g_Prevotella
k_Bacteria_p_Acidobacteria_c_Corinobacteriia_o_RBK11_f_BH6070_g_
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Lachnospiraceae_g_Mayella
k_Bacteria_k_Bacteroidetes_c_Sipropirales_o_Sipropirales_f_Chitinophagaceae_g_Flavobacter
k_Bacteria_p_Proteobacteria_c_Betaproteobacteria_o_Burkholderiales_f_Oxalobacteraceae_Other
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Actinomycetaceae_g_
k_Bacteria_k_Bacteroidetes_c_Cytophagia_o_Cytophagales_f_Cytophagaceae_g_Hymenobacter
k_Bacteria_p_Actinobacteria_c_Corinobacteriia_Corinobacteriales_f_Corinobacteriaceae_g_Adercreutzia
k_Bacteria_p_Proteobacteria_c_Gammaproteobacteria_o_Xanthomonadales_f_Xanthomonadaceae_Other
k_Bacteria_k_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Yuakusamiaceae_g_Yuakusamia
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Intropirangaceae_g_Introbacter
k_Bacteria_k_Chloroflexi_c_Anardiniaceae_f_CFB_264_f_g_
k_Bacteria_k_Bacteroidetes_c_Sphingobacteriia_o_Sphingobacteriales_f_Sphingobacteriaceae_g_Pedobacter
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Ruminococcaceae_Other
k_Bacteria_p_Firmicutes_c_Bacilli_Other_Other
k_Bacteria_k_Proteobacteria_c_Alphaproteobacteria_o_Rhodospirales_f_Rhodospiraceae_g_Picrococcus
k_Bacteria_p_Proteobacteria_c_Epsilonproteobacteria_o_Campylobacteriales_f_Campylobacteriaceae_g_Campylobacter
k_Bacteria_k_Firmicutes_c_Clostridia_Clostridiales_f_Verruococcaceae_g_Verruococcus
k_Bacteria_p_Firmicutes_c_Bacillo_Lactobacillales_Other_Other
k_Bacteria_k_Bacteroidetes_c_Sipropirales_o_Sipropirales_f_Chitinophagaceae_Other
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Lachnospiraceae_g_Roodbira
k_Bacteria_p_SRL_c_f_g_
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Micromonosporaceae_Other
k_Bacteria_p_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Bifidobacteriaceae_Other
k_Bacteria_k_Fusobacteria_c_Fusobacteriia_o_Fusobacteriales_f_Lentitrichaceae_g_Lentitrichia
k_Bacteria_p_Firmicutes_c_Erysipelotrichi_o_Erysipelotrichales_f_Erysipelotrichaceae_g_Buflidia
k_Bacteria_k_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Micrococcaceae_g_
k_Bacteria_p_Proteobacteria_c_Alphaproteobacteria_o_Rhodospirales_f_Rhodospiraceae_g_Agrobacterium
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Ruminococcaceae_g_Oxiflexia
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Brevibacteriaceae_g_Brevibacterium
k_Bacteria_k_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Paraprevotellaceae_g_Paraprevotella
k_Bacteria_k_Bacteroidetes_c_Gammaproteobacteria_o_Enterobacteriales_f_Enterobacteriaceae_g_Citrobacter
k_Bacteria_p_Firmicutes_c_Erysipelotrichi_o_Erysipelotrichales_f_Erysipelotrichaceae_g_Eubacter
k_Bacteria_p_Proteobacteria_c_Alphaproteobacteria_o_Rhodospirales_f_Rhodospiraceae_g_Rhodobacter
k_Bacteria_p_Proteobacteria_c_Alphaproteobacteria_o_932_f_g_
k_Bacteria_p_Firmicutes_c_Bacillo_Gemmatiales_f_g_
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Micrococcaceae_g_Antrobacter
k_Bacteria_k_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_524_f_g_
k_Bacteria_k_Bacteroidetes_c_Flavobacteriia_o_Flavobacteriales_f>Weissbacteriaceae_g_Chryso bacterium
k_Bacteria_p_Proteobacteria_c_Gammaproteobacteria_o_Pseudomonadales_f_Moraxellaceae_g_Erythrobacter
k_Bacteria_k_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Dorvakiaceae_g_Dorvaki
k_Bacteria_k_Terricoccales_f_Mollisutes_o_Mycoplasmatales_f_Mycoplasmataceae_g_Mycoplasma
k_Bacteria_p_Firmicutes_c_Bacillo_Lactobacillales_f_Streptococcaceae_g_Lactococcus
k_Bacteria_p_Acidobacteria_c_Acidobacteriia_o_Acidobacteriales_f_Korarchaeaceae_g_
k_Bacteria_p_Proteobacteria_c_Gammaproteobacteria_o_Pseudomonadales_f_Nocarditellaceae_g_Acnetobacter
k_Bacteria_k_Thermi_c_Deinococcu_o_Deinococcales_f_Deinococcaceae_g_Deinococcus
k_Bacteria_k_Bacteroidetes_c_Sipropirales_o_Sipropirales_f_Chitinophagaceae_g_Sedimentibacterium
k_Bacteria_p_Proteobacteria_c_Sphingomonadales_f_Sphingomonadales_f_Sphingomonadaceae_g_Sphingomonas
k_Bacteria_k_Bacteroidetes_c_Flavobacteriia_o_Flavobacteriales_f_Flavobacteriaceae_g_Capocytophaga
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Nocarditellaceae_g_
k_Bacteria_k_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Neotomabacteriaceae_g_Neotomabacter
k_Bacteria_k_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Bacteroidaceae_Other
k_Bacteria_k_Proteobacteria_c_Betaproteobacteria_o_Burkholderiales_f_Oxalobacteriaceae_g_
k_Bacteria_p_Proteobacteria_c_Alphaproteobacteria_o_Rhodospirales_f_Rhodospiraceae_Other
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Dermabacteriaceae_g_Dermabacter
k_Bacteria_k_Bacteroidetes_c_Actinobacteriia_Actinomycetales_f_Micrococcaceae_g_Rothia
k_Bacteria_k_Bacteroidetes_c_Sphingobacteriia_o_Sphingobacteriales_f_Sphingobacteriaceae_Other
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Actinomycetaceae_g_Varillaculum
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Tissotellaceae_g_Pingotia
k_Bacteria_k_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Kanamurellaceae_g_
k_Bacteria_p_Proteobacteria_c_Alphaproteobacteria_o_Rhodospirales_f_g_
k_Bacteria_p_Verrucomicrobiia_c_Verrucomicrobiales_f_Verrucomicrobiaceae_g_Alkermansia
k_Bacteria_p_Firmicutes_c_Bacillo_Lactobacillales_f_Streptococcaceae_Other
k_Bacteria_k_Bacteriia_Other_Other_Other
k_Bacteria_p_Bacteroidetes_c_Flavobacteriia_o_Flavobacteriales_f>Weissbacteriaceae_Other
k_Bacteria_k_Proteobacteria_c_Betaproteobacteria_o_Rhodocyclales_f_Rhodocyclaceae_g_Hydrogenophilus
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Verruococcaceae_g_Dialster
k_Bacteria_p_Proteobacteria_c_Epsilonproteobacteria_o_Campylobacteriales_f_Campylobacteriaceae_Other
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Propionibacteriaceae_g_Propionimicrobium
k_Bacteria_k_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Fusobacteriaceae_g_
k_Bacteria_k_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Porphyrinonadaceae_g_Porphyrinomas
k_Bacteria_p_Cyanobacteria_c_Cytophagia_o_Sirotophyta_f_g_
k_Bacteria_p_Proteobacteria_c_Gammaproteobacteria_o_Pseudomonadales_f_Pseudomonadaceae_g_Pseudomonas
k_Bacteria_k_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Porphyrinonadaceae_g_Panabacteriis
k_Bacteria_p_Acidobacteria_c_Acidobacteriia_o_ii-15_Other
k_Bacteria_p_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Bacteroidiaceae_g_Bacteroides
k_Bacteria_p_Firmicutes_c_Bacillo_Lactobacillales_f_Aerococcaceae_g_Aerococcus
k_Bacteria_p_Proteobacteria_c_Betaproteobacteria_o_Burkholderiales_f_Aliqualitaceae_g_Suttrella
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Frankiaceae_Other
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Lachnospiraceae_g_Lachnospira
k_Bacteria_p_Terricoccales_f_Mollisutes_o_Mycoplasmatales_f_Mycoplasmataceae_g_Uraplasma
k_Bacteria_p_Firmicutes_c_Bacillo_Lactobacillales_f_Campylobacteriales_f_Granulicatella
k_Bacteria_k_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Paraprevotellaceae_Other
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Micrococcaceae_g_Kocuria
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Corynebacteriaceae_g_Corynebacterium
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_Other_Other
k_Bacteria_p_Proteobacteria_c_Deltaproteobacteria_o_Neisseriales_f_Neisseriaceae_g_Neisseria
k_Bacteria_p_Proteobacteria_c_Alphaproteobacteria_o_Rhodospirales_f_Rhodospiraceae_Other
k_Bacteria_p_Actinobacteria_c_Rubrobacteriia_o_Rubrobacteriales_f_Rubrobacteriaceae_g_Rubrobacter
k_Bacteria_k_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Intropirangaceae_g_
k_Bacteria_p_Gemmatimonadetes_c_Gemm_3_o_f_g_
k_Bacteria_k_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Prevotellaceae_g_Prevotella
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Tissotellaceae_g_Polyphibius
k_Bacteria_k_Bacteroidetes_c_Cytophagia_o_Cytophagales_f_Cytophagaceae_g_
k_Bacteria_p_Cyanobacteria_Other_Other_Other
k_Bacteria_k_Actinobacteria_c_Actinobacteriia_o_Fusobacteriales_f_Fusobacteriaceae_g_Fusobacterium
k_Bacteria_p_Acidobacteria_c_Acidobacteriia_o_Acidobacteriales_f_Acibacteriaceae_g_
k_Bacteria_p_Fusobacteriia_c_Fusobacteriia_o_Fusobacteriales_f_Lentitrichiaceae_Other
k_Bacteria_k_Proteobacteria_c_Betaproteobacteria_o_Burkholderiales_f_Commonwealthaceae_Other
k_Bacteria_p_Proteobacteria_c_Gammaproteobacteria_Other_Other
k_Bacteria_p_Proteobacteria_c_Betaproteobacteria_Other_Other
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Micrococcaceae_Other
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Lachnospiraceae_g_Dyadobacterium
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_Other
k_Bacteria_p_Cyanobacteria_c_C03_2_o_152_f_g_
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Lachnospiraceae_g_Ctenococcus
k_Bacteria_p_Firmicutes_c_Bacillo_Lactobacillales_f_Streptococcaceae_g_Streptococcus
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Lachnospiraceae_g_Dorea
k_Bacteria_p_Firmicutes_Other_Other_Other
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Actinomycetaceae_g_Actinomycetes
k_Bacteria_p_Proteobacteria_c_Gammaproteobacteria_o_Enterobacteriales_f_Enterobacteriaceae_Other
k_Bacteria_k_Bacteroidetes_c_Cytophagia_o_Cytophagales_f_Cytophagaceae_g_Sirosonia
k_Bacteria_Other_Other_Other
k_Bacteria_p_Fusobacteriia_c_Fusobacteriia_o_Fusobacteriales_f_Fusobacteriaceae_Other
k_Bacteria_p_Firmicutes_c_Bacillo_Bifidales_f_Staphylococcaceae_g_Staphylococcus
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Micromonosporaceae_g_
k_Bacteria_p_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Dorvakiaceae_g_Buflitrimonas
k_Bacteria_k_Proteobacteria_c_Alphaproteobacteria_Other_Other
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_g_
k_Bacteria_k_Chloroflexi_c_Thermomicrobiia_o_JG30-47_CM451_f_g_
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Lachnospiraceae_g_Catiniella
k_Bacteria_k_Bacteroidetes_c_Bacteroidia_o_Bacteroidales_f_Chironomobacteriaceae_g_
k_Bacteria_p_Proteobacteria_c_Betaproteobacteria_o_Burkholderiales_f_Oxalobacteriaceae_g_Janthrobacterium
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Micrococcaceae_g_Micrococcus
k_Bacteria_p_Firmicutes_c_Bacillo_Lactobacillales_f_Lactobacillaceae_g_Lactobacillus
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Other_Other
k_Bacteria_p_Actinobacteria_c_Actinobacteriia_Actinomycetales_f_Sporichthyaceae_g_
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Ruminococcaceae_g_Ruminococcus
k_Bacteria_p_Firmicutes_c_Clostridia_Clostridiales_f_Tissotellaceae_g_Anerococcus

FIG 2 continued




**FIG 3** Taxonomic composition graph illustrating top 50 (per software) abundant genera, canadian\_soil. The bars do not reach 1, since only a fraction (top 50) of taxonomies was illustrated.

simulated data, mock communities, and natural microbial communities. We found that Swarm, SUMACLUSt, UCLUST, and UPARSE (with relaxed parameters) performed equally well on simulated datasets where the ground truth was well established, with mothur\_average and OTUCLUST closely behind. Despite this controlled chimera-free environment, UPARSE with recommended parameters reported the lowest accuracy for the sim\_staggered data set, implying that stringent quality filtering can cause a significant underestimation of species abundance and diversity and lead to incorrect biological results. For the mock communities, most tools were able to correctly detect the expected number and identity of genera, but only UPARSE reported significantly fewer false-positive taxa (followed by OTUCLUST and USEARCH). For UPARSE, this was expected, as a large proportion of reads was filtered out prior to clustering, leaving evidence of only the most abundant taxa (OTUs comprised of hundreds of thousands of reads). The majority of false-positive taxa reported by other tools were low-abundance OTUs that could be mapped to BLAST's NT database with very high similarity ( $E$  value,  $<1e-50$ ). If the user's primary goal is to focus on the most abundant microbial profiles, low-abundance OTUs may be filtered out postclustering, but care should be taken, as such low-abundance OTUs can be important members of communities (39).


In terms of accurately predicted taxonomic composition for *de novo* tools, Swarm performed well across all simulated and mock datasets, followed closely by SUMACLUSt and UCLUST. However, both Swarm and SUMACLUSt reported significantly fewer OTUs and lower alpha diversities than UCLUST. The performance of other *de novo* methods, such as mothur and OTUCLUST, showed more variation across datasets; however, these results were largely influenced by the preliminary sequence filtering step, where both tools removed more data than QIIME's method. We found that QIIME's filtering approach worked well across all datasets, rendering the most data for clustering tools to work with. For closed-reference tools, SortMeRNA significantly outperformed UCLUST and USEARCH for predicting OTUs and performed as well or better in terms of predicted taxonomic composition. Several studies could not be processed with mothur, OTUCLUST, or the free academic distribution of UPARSE due to their large sizes, either because of an unreasonable disk space requirement in the case of mothur, unreasonable run time in the case of OTUCLUST (no multithreading support), or a relatively small

	k_Bacteria;p__Proteobacteria;c__Deltaproteobacteria;o__Desulfuromonadales;f__Desulfuromonadaceae;g__
	k_Bacteria;p__Actinobacteria;c__Thermoleophilia;o__Gaiellales;f__Gaiellaceae;g__
	k_Bacteria;p__Proteobacteria;c__Alphaproteobacteria;o__Sphingomonadales;f__Sphingomonadaceae;Other
	k_Bacteria;p__Acidobacteria;Other;Other;Other;Other
	k_Bacteria;p__Verrucomicrobia;c__[Spartobacteria];o__[Chthoniobacterales];f__[Chthoniobacteraceae];g__DA101
	k_Bacteria;p__Proteobacteria;c__Alphaproteobacteria;o__Rhizobiales;Other;Other
	k_Bacteria;p__Proteobacteria;c__Alphaproteobacteria;o__Rhizobiales;f__Bradyrhizobiaceae;g__Bradyrhizobium
	k_Bacteria;p__Acidobacteria;c__Solibacteres;o__Solibacterales;Other;Other
	k__Archaea;p__Crenarchaeota;c__Thaumarchaeota;o__Nitrososphaerales;f__Nitrososphaeraceae;g__Candidatus Nitrososphaera
	k_Bacteria;p__Chloroflexi;c__Ellin6529;o__f__g__
	k_Bacteria;p__Bacteroidetes;c__Cytophagia;o__Cytophagales;f__Cytophagaceae;g__
	k_Bacteria;p__Bacteroidetes;Other;Other;Other;Other
	k_Bacteria;p__Proteobacteria;c__Deltaproteobacteria;o__Desulfuromonadales;Other;Other
	k_Bacteria;p__Gemmatimonadetes;c__Gemmatimonadetes;o__N1423WL;f__g__
	k_Bacteria;p__Gemmatimonadetes;c__Gemm-1;o__f__g__
	k_Bacteria;p__Acidobacteria;c__[Chloracidobacteria];o__RB41;f__Ellin6075;g__
	k_Bacteria;p__Bacteroidetes;c__[Saprosirae];o__[Saprosirales];f__Chitinophagaceae;g__Flavisolibacter
	k_Bacteria;p__Proteobacteria;c__Betaproteobacteria;o__Burkholderiales;Other;Other
	k_Bacteria;p__Acidobacteria;c__Acidobacteria-6;o__iii1-15;f__RB40;g__
	k_Bacteria;p__Verrucomicrobia;c__[Pedosphaerae];o__[Pedosphaerales];Other;Other
	k_Bacteria;p__Nitrospirae;c__Nitrospira;o__Nitrospirales;f__Nitrospiraceae;g__Nitrospira
	k_Bacteria;p__Proteobacteria;c__Betaproteobacteria;o__A21b;f__EB1003;g__
	k_Bacteria;p__Acidobacteria;c__Acidobacteria;o__Acidobacteriales;f__Acidobacteriaceae;g__
	k_Bacteria;p__Acidobacteria;c__Acidobacteria-5;o__f__g__
	k_Bacteria;p__Acidobacteria;c__Holophagae;o__Holophagales;f__Holophagaceae;g__Geothrix
	k_Bacteria;p__Bacteroidetes;c__[Saprosirae];o__[Saprosirales];f__Chitinophagaceae;Other
	k_Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__Xanthomonadales;f__Xanthomonadaceae;Other
	k_Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__[Marinicellales];f__[Marinicellaceae];g__Marinicella
	k_Bacteria;p__Acidobacteria;c__Solibacteres;o__Solibacterales;f__Solibacteraceae;g__Candidatus Solibacter
	k_Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__Alteromonadales;f__Colwelliaceae;g__
	k_Bacteria;p__Verrucomicrobia;c__[Pedosphaerae];o__[Pedosphaerales];f__Ellin517;g__
	k_Bacteria;p__Proteobacteria;c__Betaproteobacteria;o__MND1;f__g__
	k_Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__Legionellales;f__Coxiellaceae;g__Aqicella
	k_Bacteria;p__Bacteroidetes;c__Cytophagia;o__Cytophagales;Other;Other
	k_Bacteria;p__Proteobacteria;c__Alphaproteobacteria;o__Ellin329;f__g__
	k_Bacteria;p__Bacteroidetes;c__Flavobacteriia;o__Flavobacteriales;f__Flavobacteriaceae;g__Flavobacterium
	k_Bacteria;p__Proteobacteria;c__Deltaproteobacteria;Other;Other;Other
	k_Bacteria;p__Acidobacteria;c__Acidobacteria;o__Acidobacteriales;f__Koribacteraceae;g__
	k_Bacteria;p__Proteobacteria;c__Alphaproteobacteria;o__Rhodobacterales;f__Rhodobacteraceae;g__Phaeobacter
	k_Bacteria;p__Proteobacteria;c__Betaproteobacteria;o__Burkholderiales;f__Comamonadaceae;g__Rhodoferrax
	k_Bacteria;p__Acidobacteria;c__DA052;o__Ellin6513;f__g__
	k_Bacteria;p__Proteobacteria;c__Alphaproteobacteria;o__Sphingomonadales;f__Sphingomonadaceae;g__Kaistobacter
	k_Bacteria;p__Proteobacteria;c__Alphaproteobacteria;o__Rhodobacterales;f__Rhodobacteraceae;g__Octadecabacter
	k_Bacteria;p__OD1;c__ZB2;o__f__g__
	k_Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__Alteromonadales;Other;Other
	k_Bacteria;p__Proteobacteria;c__Alphaproteobacteria;o__Rhodospirillales;f__Acetobacteraceae;Other
	k_Bacteria;p__Bacteroidetes;c__Sphingobacteriia;o__Sphingobacteriales;f__Sphingobacteriaceae;Other
	k_Bacteria;p__Proteobacteria;c__Alphaproteobacteria;o__Rhodobacterales;f__Rhodobacteraceae;Other
	k_Bacteria;p__Proteobacteria;Other;Other;Other;Other
	k_Bacteria;p__Proteobacteria;c__Alphaproteobacteria;o__Rhizobiales;f__Hyphomicrobiaceae;g__Rhodoplanes
	k_Bacteria;p__Bacteroidetes;c__Cytophagia;o__Cytophagales;f__Cytophagaceae;Other
	k_Bacteria;p__Proteobacteria;c__Betaproteobacteria;o__Burkholderiales;f__Oxalobacteraceae;g__Janthinobacterium
	k_Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__Pseudomonadales;f__Pseudomonadaceae;g__Pseudomonas
	k_Bacteria;p__Verrucomicrobia;c__[Spartobacteria];o__[Chthoniobacterales];f__[Chthoniobacteraceae];Other
	k_Bacteria;p__Acidobacteria;c__Acidobacteria-6;o__iii1-15;Other;Other
	k_Bacteria;p__Acidobacteria;c__Solibacteres;o__Solibacterales;f__g__
	k_Bacteria;p__Proteobacteria;c__Deltaproteobacteria;o__Myxococcales;Other;Other
	k_Bacteria;p__Verrucomicrobia;c__[Pedosphaerae];o__[Pedosphaerales];f__auto67_4W;g__
	k_Bacteria;p__Proteobacteria;c__Betaproteobacteria;o__IS-44;f__g__
	k_Bacteria;p__Acidobacteria;c__Acidobacteria-6;o__iii1-15;f__g__
	Unclassified;Other;Other;Other;Other;Other
	k_Bacteria;p__Proteobacteria;c__Betaproteobacteria;o__Burkholderiales;f__Comamonadaceae;Other
	k_Bacteria;p__Proteobacteria;c__Gammaproteobacteria;Other;Other;Other
	k_Bacteria;p__Proteobacteria;c__Betaproteobacteria;Other;Other;Other
	k_Bacteria;p__Actinobacteria;c__Actinobacteria;o__Actinomycetales;Other;Other
	k_Bacteria;p__Acidobacteria;c__Acidobacteria;o__Acidobacteriales;f__Acidobacteriaceae;Other
	k_Bacteria;p__Proteobacteria;c__Deltaproteobacteria;o__Desulfuromonadales;f__Desulfuromonadaceae;Other
	k_Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__Alteromonadales;f__Colwelliaceae;Other
	k_Bacteria;p__Verrucomicrobia;Other;Other;Other;Other
	k_Bacteria;Other;Other;Other;Other;Other
	k_Bacteria;p__Proteobacteria;c__Deltaproteobacteria;o__Desulfuromonadales;f__Geobacteraceae;Other
	k_Bacteria;p__Acidobacteria;c__[Chloracidobacteria];o__RB41;f__g__
	k_Bacteria;p__Proteobacteria;c__Alphaproteobacteria;Other;Other;Other
	k_Bacteria;p__Acidobacteria;c__Acidobacteria;o__Acidobacteriales;f__Koribacteraceae;g__Candidatus Koribacter
	k_Bacteria;p__Acidobacteria;c__;o__f__g__
	k_Bacteria;p__Proteobacteria;c__Gammaproteobacteria;o__PYR10d3;f__g__
	k_Bacteria;p__Proteobacteria;c__Deltaproteobacteria;o__NB1-j;f__NB1-i;g__

FIG 3 continued



**FIG 4** Alpha diversity for tools at different sampling depths (order: *de novo*, closed reference, and open reference), canadian\_soil.



**FIG 5** Run time performance for all benchmarked software. All tests were performed using 1 to 32 cores on Intel Xeon CPU E5-2640 v3 at 2.60 GHz. Input files contained reads subsampled from the Global Gut. For serial performance, some tools do not show results for 10<sup>8</sup> reads due to exceeding wall time limit (230 h) or failed memory allocation. For parallel performance, a single file containing 1 million Illumina sequences was used over multiple threads.

memory limit in the case of UPARSE. Regarding UPARSE, the small memory limit makes it necessary to purchase the 64-bit license in order to process large projects (e.g., see Yatsunenکو et al.'s work [40], which contains 500 GB of raw sequence data generated on 17 HiSeq lanes) or use open-source alternatives. QIIME's current open-source, open-reference pipeline (based on SortMeRNA and SUMACLUSt) was able to process this quantity of data within 24 h using 64 threads on Intel Xeon CPU E5-4620 v2 at 2.60GHz or within 3 days using 64 threads on AMD Opteron Processor 6276.

Although most open-source tools report an increased run time in comparison to UCLUST and USEARCH (Fig. 5), they provide the benefit of finding significantly fewer OTUs. In the case of SortMeRNA, longer reads (~150 bp) are quicker to align than the same number of shorter reads (~100 bp) due to many fewer high-scoring candidate reference sequences to analyze. Moreover, all of these tools support multilevel multi-

threading and can easily scale to modern big-data processing demands. An alternative to reducing run time is to filter out a substantial number of reads, as done by UPARSE; unfortunately, the filtering parameters are sensitive to different data, and choosing them manually by trial and error can be a time-consuming task with unpredictable outcomes in diversity.

The three open-source software products, Swarm, SUMACLUSt, and SortMeRNA, are now accessible through the widely used QIIME software package (release 1.9.0). Swarm 2 was released in reference 14 and reported to be faster and more memory efficient than Swarm 1; however, as of this writing, only Swarm 1 has been integrated into QIIME. Ongoing work to improve the QIIME OTU clustering workflows that use these tools includes adding a targeted gene prefilter for *de novo* clustering to remove (prior to clustering) any sequences not matching a specific gene model (e.g., 16S rRNA) and a refined reference database for targeted hypervariable regions (e.g., V4 at 97% identity) to improve alignment quality (41). Furthermore, research is in progress to introduce an open-source implementation of chimera detection directly within QIIME. Both of these improvements will further reduce the number of unrelated or erroneous reads recruited into OTUs, a known problem with both the UCLUST- and USEARCH-based OTU clustering illustrated here, without underestimating diversity.

## MATERIALS AND METHODS

All steps taken to generate the analyses presented in this article are documented and implemented as shell or python scripts, available at <https://github.com/ekopylova/OTU-clustering>.

**Performance benchmarks.** Open-source with multilevel parallelization tools tested in this paper—Swarm, SUMACLUSt, and SortMeRNA—have been integrated into QIIME 1.9.0. For these tools, all benchmarks were launched through QIIME. For UPARSE, the recommended workflow ([http://www.drive5.com/usearch/manual/uparse\\_cmds.html](http://www.drive5.com/usearch/manual/uparse_cmds.html)) was run. For OTUCLUSt, the script micca-preproc was used for sequence filtering and the command otuclust for clustering. For mothur, the MiSeq SOP (42) (website accessed 27 October 2015) and 454 SOP (43) (website accessed 27 October 2015) were run. The shell scripts commands\_16S.sh and commands\_18S.sh were used to launch all tools, and the open-source project (<https://github.com/josenavas/QIIME-Scaling>) was used for measuring their run time performance. All run time performance tests were performed using 1 to 32 threads on Intel Xeon CPU E5-2640 v3 at 2.60 GHz.

**Precision and recall.** For simulated and mock datasets, false-positive (FP; taxonomy/OTU string exists in observed but not expected), false-negative (FN; taxonomy/OTU string exists in expected but not observed), and true-positive (TP; taxonomy/OTU string exists in both observed and expected) measures were computed between the pickers' results (observed) and the ground truth or expected taxonomic composition (expected). The following definitions were used: precision = TP/(TP + FP); recall = TP/(TP + FN); F measure =  $2 \times \text{precision} \times \text{recall} / (\text{precision} + \text{recall})$ .

The python script run\_compute\_precision\_recall.py was used to compute TP, FP, FN, precision, recall, F measure, the number of false-positive taxa whose complete set of OTUs are identified as chimeric (FP-chimeric) by UCHIME, the number of false-positive taxa whose complete set of OTUs map with  $\geq 97\%$  identity and coverage to BLAST's NT database (FP-known), and the number of false-positive taxa whose complete set of OTUs map with  $< 97\%$  identity and coverage to BLAST's NT database (FP-other). The script plot\_tp\_fp\_distribution.py was used to generate Fig. S1, S2, and S3 in the supplemental material.

**Simulating reads (even and staggered).** All of the following steps can be executed using the shell script simulate\_reads.sh.

Reads were simulated using PrimerProspector (23) and the ART simulator (24). For the even data set, the following steps were taken. (i) Use PrimerProspector to extract V4 regions from the Greengenes 97% representative database (version 13.8); (ii) subsample 0.011% of the sequences from the resulting V4 region database; and (iii) simulate even abundance reads with ART simulator using the subsampled V4 sequences.

Amplicon sequencing simulation in ART (version VanillaIceCream-03-11-2014) could generate only evenly distributed communities. To simulate the staggered data set, a staggered distribution of template sequences was passed (for example, 3 duplicates of OTU1, 10 duplicates of OTU2, etc.). To simulate the staggered data set, the following steps were taken. (i) Generate a random staggered distribution FASTA file of template V4 sequences using the list of OTU identifications from the even data set and the V4 subsampled sequences and (ii) simulate staggered abundance reads with ART using the staggered subsampled V4 sequences.

For both even and staggered reads, QIIME's split\_libraries\_fastq.py script was run to filter simulated reads based on quality scores and format FASTA labels to be compatible with QIIME (reads for UPARSE, mothur, and OTUCLUSt were not filtered; only FASTA labels were reformatted).

**Building ground-truth BIOM tables.** Ground-truth OTU maps and BIOM tables were constructed using the simulate\_reads.sh script that was used for simulating reads. OTU maps were generated using the reads' origin information stored in the FASTA labels of ART-simulated reads. BIOM tables were generated using QIIME's make\_otu\_table.py script together with Greengenes 97% taxonomy strings.

**Construction of a Silva 97% representative OTU tree.** A eukaryotic/18S rRNA sequence set tree was built using QIIME's `filter_alignment.py` and `make_phylogeny.py` scripts:

```
filter_alignment.py -i Silva_111_post/rep_set_aligned/97_Silva_111_rep_set.fasta -e 0.0005 -g 0.80 -o Silva_111_post/trees; make_phylogeny.py -i Silva_111_post/trees/97_Silva_111_rep_set_pfiltered.fasta -o Silva_111_post/trees/97_Silva_111_rep_set_pfiltered.tre.
```

**Calculating alpha diversity, beta diversity, and taxonomic correlation.** Custom scripts iterating over all benchmarking results were used to launch QIIME's alpha and beta diversity analyses. The script `run_single_rarefaction_and_plot.py` was used to compute and plot alpha diversity as shown in Fig. 4 and in Fig. S4 and S5 in the supplemental material. The script `run_beta_diversity_and_procrustes.py` was used to compute beta diversity and run Procrustes analysis.

## SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <http://dx.doi.org/10.1128/mSystems.00003-15>.

Figure S1, PDF file, 0.1 MB.  
 Figure S2, PDF file, 0.1 MB.  
 Figure S3, PDF file, 0.1 MB.  
 Figure S4, PDF file, 0.2 MB.  
 Figure S5, PDF file, 0.2 MB.  
 Table S1, PDF file, 0.04 MB.  
 Table S2, PDF file, 0.02 MB.  
 Table S3, PDF file, 0.02 MB.  
 Table S4, PDF file, 0.02 MB.  
 Table S5, PDF file, 0.04 MB.

## ACKNOWLEDGMENTS

We thank William Walters, Amnon Amir, Amanda Birmingham, Embriette Hyde, and Daniel McDonald for their time and valuable suggestions to improve the quality of the manuscript.

## FUNDING INFORMATION

HHS | National Institutes of Health (NIH) provided funding to Evguenia Kopylova, José Antonio Navas-Molina, and Rob Knight under grant number 1S10OD012300. Deutsche Forschungsgemeinschaft (DFG) provided funding to Frédéric Mahé under grant number DU1319/1-1.

This work was partially supported by the Howard Hughes Medical Institute and the Alfred P. Sloan Foundation.

## REFERENCES

1. **The Human Microbiome Project Consortium.** 2012. A framework for human microbiome research. *Nature* **486**:215–221. <http://dx.doi.org/10.1038/nature11209>.
2. **Human Microbiome Project Consortium.** 2012. Structure, function and diversity of the healthy human microbiome. *Nature* **486**:207–214. <http://dx.doi.org/10.1038/nature11234>.
3. **Turnbaugh PJ, Ley RE, Hamady M, Fraser-Liggett CM, Knight R, Gordon JI.** 2007. The human microbiome project. *Nature* **449**:804–810. <http://dx.doi.org/10.1038/nature06244>.
4. **Wetterstrand KA.** 2013. DNA sequencing costs: data from the NHGRI genome sequencing program (GSP). <http://www.genome.gov/sequencingcosts>. Accessed 15 November 2014.
5. **Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Peña AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh PJ, Walters WA, Widmann J, Yatsunencko T, Zaneveld J, Knight R.** 2010. QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* **7**:335–336. <http://dx.doi.org/10.1038/nmeth.f.303.7>.
6. **Edgar RC.** 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**:2460–2461. <http://dx.doi.org/10.1093/bioinformatics/btq461>.
7. **Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ.** 1990. Basic local alignment search tool. *J Mol Biol* **215**:403–410. <http://dx.doi.org/10.1006/jmbi.1990.9999>.
8. **Schloss PD, Handelsman J.** 2005. Introducing DOTUR, a computer program for defining operational taxonomic units and estimating species richness. *Appl Environ Microbiol* **71**:1501–1506. <http://dx.doi.org/10.1128/AEM.71.3.1501>.
9. **Li W, Jaroszewski L, Godzik A.** 2001. Clustering of highly homologous sequences to reduce the size of large protein databases. *Bioinformatics* **17**:282–283. <http://dx.doi.org/10.1093/bioinformatics/17.3.282>.
10. **Li W, Jaroszewski L, Godzik A.** 2002. Tolerating some redundancy significantly speeds up clustering of large protein databases. *Bioinformatics* **18**:77–82. <http://dx.doi.org/10.1093/bioinformatics/18.1.77>.
11. **Li W, Godzik A.** 2006. Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* **22**:1658–1659. <http://dx.doi.org/10.1093/bioinformatics/btl158>.
12. **Albanese D, Fontana P, De Filippo C, Cavalieri D, Donati C.** 2015. Micca: a complete and accurate software for taxonomic profiling of metagenomic data. *Sci Rep* **5**:9743. <http://dx.doi.org/10.1038/srep09743>.
13. **Mahé F, Rognes T, Quince C, de Vargas C, Dunthorn M.** 2014. Swarm: robust and fast clustering method for amplicon-based studies. *PeerJ* **2**:e593. <http://dx.doi.org/10.7717/peerj.593>.
14. **Mahé F, Rognes T, Quince C, de Vargas C, Dunthorn M.** 2015. Swarm v2: highly-scalable and high-resolution amplicon clustering. *PeerJ* **3**:e1420. <http://dx.doi.org/10.7717/peerj.1420>.
15. **Kopylova E, Noé L, Touzet H.** 2012. SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics* **28**:3211–3217. <http://dx.doi.org/10.1093/bioinformatics/bts611>.



16. **Edgar RC.** 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat Methods* **10**:996–998. <http://dx.doi.org/10.1038/nmeth.2604>.
17. **Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA, Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ, Weber CF.** 2009. Introducing mothur: open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl Environ Microbiol* **75**:7537–7541. <http://dx.doi.org/10.1128/AEM.01541-09>.
18. **Navas-Molina JA, Peralta-Sánchez JM, González A, McMurdie PJ, Vázquez-Baeza Y, Xu Z, Ursell LK, Lauber C, Zhou H, Song SJ, Huntley J, Ackermann GL, Berg-Lyons D, Holmes S, Caporaso JG, Knight R.** 2013. Advancing our understanding of the human microbiome using QIIME. *Methods Enzymol* **531**:371–444. <http://dx.doi.org/10.1016/B978-0-12-407863-5.00019-8>.
19. **Rideout JR, He Y, Navas-Molina JA, Walters WA, Ursell LK, Gibbons SM, Chase J, McDonald D, Gonzalez A, Robbins-Pianka A, Clemente JC, Gilbert JA, Huse SM, Zhou H-W, Knight R, Caporaso JG.** 2014. Subsampled open-reference clustering creates consistent, comprehensive OTU definitions and scales to billions of sequences. *PeerJ* **2**:e545. <http://dx.doi.org/10.7717/peerj.545>.
20. **Hobohm U, Scharf M, Schneider R, Sander C.** 1992. Selection of representative protein data sets. *Protein Sci* **1**:409–417. <http://dx.doi.org/10.1002/pro.5560010313>.
21. **Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R.** 2011. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* **27**:2194–2200. <http://dx.doi.org/10.1093/bioinformatics/btr381>.
22. **Legendre P, Legendre L.** 1998. Numerical ecology, 2nd ed. Developments in environmental modelling, vol 20, p. Elsevier Science, Amsterdam, The Netherlands.
23. **Walters WA, Caporaso JG, Lauber CL, Berg-Lyons D, Fierer N, Knight R.** 2011. PrimerProspector: de novo design and taxonomic analysis of barcoded polymerase chain reaction primers. *Bioinformatics* **27**:1159–1161. <http://dx.doi.org/10.1093/bioinformatics/btr087>.
24. **Huang W, Li L, Myers JR, Marth GT.** 2012. ART: a next-generation sequencing read simulator. *Bioinformatics* **28**:593–594. <http://dx.doi.org/10.1093/bioinformatics/btr708>.
25. **Bokulich NA, Subramanian S, Faith JJ, Gevers D, Gordon JI, Knight R, Mills DA, Caporaso JG.** 2013. Quality-filtering vastly improves diversity estimates from Illumina amplicon sequencing. *Nat Methods* **10**:57–59. <http://dx.doi.org/10.1038/nmeth.2276>.
26. **Porazinska DL, Giblin-Davis RM, Faller L, Farmerie W, Kanzaki N, Morris K, Powers TO, Tucker AE, Sung W, Thomas WK.** 2009. Evaluating high-throughput sequencing as a method for metagenomic analysis of nematode diversity. *Mol Ecol Resour* **9**:1439–1450. <http://dx.doi.org/10.1111/j.1755-0998.2009.02611.x>.
27. **Neufeld JD, Engel K, Cheng J, Moreno-Hagelsieb G, Rose DR, Charles TC.** 2011. Open resource metagenomics: a model for sharing metagenomic libraries. *Stand Genomic Sci* **5**:203–210. <http://dx.doi.org/10.4056/sigs.1974654>.
28. **Costello EK, Lauber CL, Hamady M, Fierer N, Gordon JI, Knight R.** 2009. Bacterial community variation in human body habitats across space and time. *Science* **326**:1694–1697. <http://dx.doi.org/10.1126/science.1177486>.
29. **Ramirez KS, Leff JW, Barberán A, Bates ST, Betley J, Crowther TW, Kelly EF, Oldfield EE, Shaw EA, Steenbock C, Bradford MA, Wall DH, Fierer N.** 2014. Biogeographic patterns in below-ground diversity in New York City's Central Park are similar to those observed globally. *Proc R Soc B Biol Sci* **281**:1–9. <http://dx.doi.org/10.1098/rspb.2014.1988>.
30. **McDonald D, Clemente JC, Kuczynski J, Rideout JR, Stombaugh J, Wendel D, Wilke A, Huse S, Hufnagle J, Meyer F, Knight R, Caporaso JG.** 2012. The biological observation matrix (BIOM) format or: how I learned to stop worrying and love the ome-ome. *Gigascience* **1**:7. <http://dx.doi.org/10.1186/2047-217X-1-7>.
31. **Wang Q, Garrity GM, Tiedje JM, Cole JR.** 2007. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* **73**:5261–5267. <http://dx.doi.org/10.1128/AEM.00062-07>.
32. **DeSantis TZ, Hugenholtz P, Larsen N, Rojas M, Brodie EL, Keller K, Huber T, Dalevi D, Hu P, Andersen GL.** 2006. Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol* **72**:5069–5072. <http://dx.doi.org/10.1128/AEM.03006-05>.
33. **McDonald D, Price MN, Goodrich J, Nawrocki EP, DeSantis TZ, Probst A, Andersen GL, Knight R, Hugenholtz P.** 2012. An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. *ISME J* **6**:610–618. <http://dx.doi.org/10.1038/ismej.2011.139>.
34. **Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glöckner FO.** 2007. SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res* **35**:7188–7196. <http://dx.doi.org/10.1093/nar/gkm864>.
35. **Faith DP.** 1992. Conservation evaluation and phylogenetic diversity. *Biol Conserv* **61**:1–10. [http://dx.doi.org/10.1016/0006-3207\(92\)91201-3](http://dx.doi.org/10.1016/0006-3207(92)91201-3).
36. **Gower JC.** 1975. Generalized Procrustes analysis. *Psychometrika* **40**:33–51. <http://dx.doi.org/10.1007/BF02291478>.
37. **Lozupone C, Lladser ME, Knights D, Stombaugh J, Knight R.** 2011. UniFrac: an effective distance metric for microbial community comparison. *ISME J* **5**:169–172. <http://dx.doi.org/10.1038/ismej.2010.133>.
38. **Hamady M, Lozupone C, Knight R.** 2010. Fast UniFrac: facilitating high-throughput phylogenetic analyses of microbial communities including analysis of pyrosequencing and PhyloChip data. *ISME J* **4**:17–27. <http://dx.doi.org/10.1038/ismej.2009.97>.
39. **Shade A, Jones SE, Caporaso JG, Handelsman J, Knight R, Fierer N, Gilbert JA.** 2014. Conditionally rare taxa disproportionately contribute to temporal changes in microbial diversity. *mBio* **5**:e01371-14. <http://dx.doi.org/10.1128/mBio.01371-14>.
40. **Yatsunenko T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, Magris M, Hidalgo G, Baldassano RN, Anokhin AP, Heath AC, Warner B, Reeder J, Kuczynski J, Caporaso JG, Lozupone CA, Lauber C, Clemente JC, Knights D, Knight R, Gordon JI.** 2012. Human gut microbiome viewed across age and geography. *Nature* **486**:222–227. <http://dx.doi.org/10.1038/nature11053>.
41. **Westcott SL, Schloss PD.** 2015. De novo clustering methods outperform reference-based methods for assigning 16S rRNA gene sequences to operational taxonomic units. *PeerJ* **3**:e14872. <http://dx.doi.org/10.7717/peerj.1487>.
42. **Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD.** 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the miseq Illumina sequencing platform. *Appl Environ Microbiol* **79**:5112–5120. <http://dx.doi.org/10.1128/AEM.01043-13>.
43. **Schloss PD, Gevers D, Westcott SL.** 2011. Reducing the effects of PCR amplification and sequencing artifacts on 16S rRNA-based studies. *PLoS One* **6**:e27310. <http://dx.doi.org/10.1371/journal.pone.0027310>.