

REVIEW

Open Access



Opportunities to improve the seasonal dynamics of water use in lentil (*Lens culinaris* Medik.) to enhance yield increase in water-limited environments

Michel Edmond Ghanem^{1*}, Fatima ez-zahra Kibbou¹, Julie Guiguitant^{1,2} and Thomas R. Sinclair³

Abstract

Lentil (*Lens culinaris* Medikus) is one of the most important annual food legumes that plays an important role in the food and nutritional security of millions in the world. Lentil is mainly grown under rainfed environments, where drought is one of the most challenging abiotic stresses that negatively impacts lentil production in the arid and semi-arid areas. Therefore, development of drought-adapted cultivars is one of the major objectives of national and international lentil breeding programs. The goal of this review is to provide a report on the current status of traits of lentil that might result in yield increases in water-limited environments and identify opportunities for research on other traits. Lately, traits that are either related to developmental plasticity and/or altered rooting and shoot characteristics have received considerable attention in the efforts to increase lentil yield in water-limited environments. However, two traits that have recently been proven to be especially useful in other legumes are still missing in lentil drought research: early partial stomatal closure under soil drying, and limited-transpiration under high atmospheric vapor pressure deficit. This review provides suggestions for further exploitation of these two soil–water-conservation traits in lentil.

Keywords: Lentil, Drought, Limited transpiration, Vapor pressure deficit, Soil drying, Root

Introduction

Cultivated lentil (*Lens culinaris* Medikus) is an important cool-season legume, particularly within countries in North America, Australia, South Asia, West Asia, and North Africa [9]. The seeds are a cheap source of protein with a high vitamin and mineral content, thus contributing to the fight against hidden hunger. However, lentil yields are generally low. The most recent available data show that globally in 2014, 4.5 million tons of lentils was produced on 4.8 million hectares of land [10] for an average yield of 0.94 tons ha⁻¹. This average yield is comparable to chickpea (*Cicer arietinum* L.), which has a global

average yield of 0.96 tons ha⁻¹. On the other hand, legumes like soybean [*Glycine max* (L.) Merr.] have average yields that can reach 3 tons ha⁻¹ or more, highlighting an apparent potential for lentil yield improvement.

In addition to the nutritional benefits of lentil, lentil can be useful in improved cropping systems due to its symbiotic nitrogen fixation capability. It has been estimated that lentils reduce dependency on inorganic nitrogen fertilizer and improve soil health because they fix nitrogen at an average rate of 80 kg N ha⁻¹ year⁻¹ [8, 35]. Moreover, with its ability for nitrogen fixation and carbon sequestration in soils, lentil helps in enhancing the sustainability of the cereal-based farming when it is included as a rotation crop. Lastly, the lentil straw is valued as animal feed in developing countries [6].

Lentil, however, is commonly grown in developing countries on marginal lands with poor soil in rain-fed environments [14]. Like many other crops that grow in

*Correspondence: m.ghanem@cgiar.org

¹ Crop Physiology Laboratory, International Center for Agricultural Research in the Dry Areas (ICARDA), North-Africa Platform, Rabat, Morocco

Full list of author information is available at the end of the article

the Mediterranean and semi-arid zones, lentil often faces terminal drought stress during the reproductive phase as a consequence of diminishing rainfall or plant available water and rising temperatures. In such cases, what is usually called “drought tolerance” could be, in part, the consequence of plant constitutive traits that affect how soil water is used earlier in the growing season when water is non-limiting to plant transpiration [66]. The present yield of lentil in India is reported to be low [25] due to terminal drought and high temperatures, particularly during flowering and seed growth [32]. Climate change is expected to increase temperatures and an increase in extremes of rainfall as well as an increased risk of drought in many areas where lentil is grown [5]. Therefore, development of drought-adapted cultivars is one of the major objectives of national and international lentil breeding programs [8].

This paper reviews the current status of research on traits that can alter the pattern of crop water use resulting in the possibility of yield increase of lentil under water-deficit conditions. For decades, two main topics have received much attention in that area that are developmental plasticity, and altered rooting and shoot characteristics. This review will briefly highlight investigations on individual traits that are either related to developmental plasticity (phenology) or below ground traits. Two additional water-conservation traits considered in this review that have been missing in lentil drought studies are initiation of partial stomatal closure early in the soil drying cycle, and partial stomatal closure at elevated atmospheric vapor pressure deficit. This review considers the opportunities for lentil yield increase in water-limited environments by addressing these two traits linked to temporal dynamics of water use.

Developmental plasticity

Lentil is an annual herbaceous plant with indeterminate growth exhibiting high variation in its growth habit: single stem, erect, semi-erect, compact growth or much-branched low bushy forms [13, 14, 31, 42]. Developmental plasticity can involve one or all of these plant characteristics. In particular, the ability of plants to adjust the duration of different growth phases in response to the availability of soil water during the growing season enables a plant to produce higher yields when the growing period is longer [52]. Such plasticity is an important mechanism in unpredictable climates and under unfavorable soil water regimes. Studies have shown that drought escape (early flowering and pod setting) is a common drought strategy for lentil in environments where water deficits and high temperatures at the reproductive stage induce senescence and early maturity [7, 51, 54, 64, 69].

Siddique et al. [54] have shown that while early flowering and accelerated phenological development gave higher seed yields and higher yield stability than late flowering under terminal drought. Thus, drought-induced early maturity could be advantageous in some dry seasons, but achieving a higher yield under well-watered conditions is often associated with longer growth duration, late flowering and greater water use [51]. Therefore, drought escape is not always a viable breeding strategy for lentil if increasing yield under a range of water conditions is the objective.

Root and shoot characteristics

The value of root and shoot characteristics for drought adaption in lentil has often been judged based on the sole criterion of yield increase. Although yield is definitely the ultimate criterion for any breeding effort, yield itself cannot be considered as a trait since it is an integrated parameter, which involves many different traits at totally different organization levels of a plant: subcellular-, cellular-, tissue-, organ-, whole plant-, and stand-level [17]. Therefore, data on “yield phenotypes” alone are inappropriate in efforts to advance specific traits that can ultimately contribute to yield under defined environments [17].

The root system of lentil is characterized by a slender taproot with a mass of fibrous lateral roots that may be shallow, intermediate or deep [43]. High genetic variation has been reported for root traits such as taproot length, lateral root number, total root length, and total root weight for lentil germplasm from different origins [13, 14, 25, 26, 31, 42]. The high heritability estimates that were reported by these authors indicate the feasibility of making use of this genetic variability for the development of drought-adapted cultivars. Deep, well-developed roots and vigorous shoots at early-seedling stage were associated with drought escape and tolerance in lentil [21, 42] as a way to ensure uptake of water and nutrients. Sarker et al. [42] reported high correlations between stem length, taproot length, and lateral root number with lentil grain yield under drought.

Given the complexity of drought, it is not surprising that root and shoot traits have been found to interact in the expression of tolerance. Kumar et al. [25] assessed genetic variability for 12 attributes including root and shoot traits among lentil genotypes originating from rainfed areas adapted to short-season environments. They found correlations between the individual traits and achieving a higher yield under drought conditions. More recently, Mishra et al. [32] found correlations between chlorophyll content and stability, increased accumulation of osmotically active solutes, soluble sugars and proline, lower H₂O₂ and malondialdehyde (MDA) contents, and

lower carbon isotope discrimination (ΔC_{13}) values and drought resistance in lentil. This study was conducted on a very limited number of genotypes (two) and drought was imposed at three different phenological stages (vegetative, flowering initiation and pod development). It is likely that the results cannot be extrapolated since the reported correlations are most probably related to the particular conditions of the study.

The above-mentioned studies remain descriptive and correlative and allow limited mechanistic explanation of the underlying functional processes, especially in terms of water uptake and water-use dynamics. For instance, although the selection for deeper rooting to access water at depth could be potentially interesting, it might lead to faster soil water depletion, which would be a problem for crops depending on stored soil moisture. Likewise, rapid development of leaf area is likely to result in more rapid depletion of soil water. This functional aspect of the changes in root and shoot and root characteristics on the temporal dynamics of water use have been overlooked. Thus, it is not possible to obtain a clear and univocal indication of root and shoot functionality when just relying on structural features.

A better integration of different, yet interactive, traits in their role in the plant's adaptation to water limitation is highly needed by having a better understanding of the dynamics of plant water use under both under well-watered conditions and upon exposure to water deficits [66].

Early transpiration decrease with soil drying

Studies indicate that sufficient amounts of water at key times during the plant cycle may be more important than total water availability across the whole cycle [30, 37,

66]. One plant trait to achieve conservative water use to increase water availability during reproductive develop is partial stomatal closure at a higher soil water content during the soil drying cycle than normally occurs in most plants [28, 57]. If there is a late-season water deficit, genotypes with a conservative water behavior have the possibility of using the conserved soil water to sustain physiological activity during seed fill and generate a greater yield than genotypes that do not have the trait [56].

A study conducted on chickpea (*Cicer arietinum* L.), a crop that is grown in similar regions as lentil, showed that plants that exhibit early stomatal closure to reduce transpiration rates (TR) when soil water deficit occurs, save water for more critical plant stages like reproduction and grain-filling under rain-fed environments [68]. Transpiration does not decrease immediately after water is withheld, but rather there is a threshold soil water content at which transpiration rate declines [34]. The initiation of a decline in transpiration rate with soil drying occurs at certain fraction of transpirable soil water (FTSW) thresholds [2], which indicates a significant and exploitable water-saving trait. Figure 1a illustrates the changes in transpiration as soil water content decreases and shows a plot of transpiration rate of plants subjected to soil drying relative to well-watered plants versus the FTSW remaining in the soil for three different cultivars displaying early, normal, and delayed transpiration decrease with soil drying [41].

The soil water is represented by a reservoir characterized by its total transpirable soil water (TTSW), representing the difference between maximum and minimum (extractable) water content, the actual transpirable soil water (ATSW) and the fraction of transpirable soil water




Fig. 1 **a** Plot of transpiration rate of plant subjected to soil drying relative to well-watered plants versus the fraction of transpirable soil water (FTSW) remaining in the soil (adapted from Sadok and Sinclair 2011). **b** Plot of transpiration rate versus vapor pressure deficit (VPD) for two genotypes with contrasting response to VPD: on where transpiration rate increases linearly with increasing VPD (*continuous line*) and the other showing the limited transpiration rate (*dotted line*) (adapted from [61])

FTSW is calculated as the ratio of ATSW to the TTSW remaining at any time during the season [58]. For example, in chickpea (*Cicer arietinum* L.), a conservative transpiration decline, occurring at a high FTSW when the soil is still relatively wet distinguished drought-adapted and drought-sensitive genotypes [68].

In lentil, experiments conducted in Western Australia, with a Mediterranean-type climate, showed that seed yield was not correlated with total water use or with water use before flowering [27], but was positively correlated with post-flowering water use [53]. Evidence in groundnut (*Arachis hypogaea* L.) [37], pearl millet (*Pennisetum glaucum* L.) [22, 23], and wheat (*Triticum aestivum* L.) [11] indicates that lower vegetative rates of water use leave more water available for grain filling. Under rainfed environments in Nepal and Jordan, early-sown lentils produced greater yield as a result of adequate vegetative mass, longer grain-filling periods, and higher pod number [32, 35, 49]. It is, however, surprising that these findings have not been extended to study seasonal dynamics of water use in regard to early-season soil water conservation and increased water availability to complete seed filling under drought conditions [28, 59].

Partial stomatal closure under high atmospheric vapor pressure deficit (VPD)

A specific trait that is especially promising for allowing conservative soil water use is one in which transpiration rate is limited under high, midday vapor pressure deficit (VPD). Figure 1b shows a plot of transpiration rate versus vapor pressure deficit for two genotypes with contrasting responses to high VPD: a linear increase in transpiration rate with increasing VPD and a limited-transpiration rate above a certain VPD breakpoint. The partial restriction of transpiration rate under high VPD limits the rate of soil water use, and raises the transpiration efficiency, allowing the crop to conserve water to support plant growth later in the season when drought develops [4, 62].

Sinclair et al. [59] examined the possible benefits for sorghum (*Sorghum bicolor* L.) from limiting transpiration rate to a constant, maximum transpiration rate value under high levels of air VPD even when soil moisture contents were high. Using a crop model that simulated sorghum growth in Australia, they reported the possibility of a yield increase in about 75% of the seasons over 100 years at four different locations.

Considerable evidence has confirmed that the limited-transpiration trait, assessed under well-watered conditions, is expressed in selected genotypes of several crop species, including soybean [12, 20, 40, 45, 60], peanut (*Arachis hypogaea* L.) [3, 47], sorghum [19, 24, 39, 48], chickpea (*Cicer arietinum* L.) [68], pearl millet [22], cowpea (*Vigna unguiculata* L.) [1], maize (*Zea mays* L.) [18,

67], and wheat [38, 44]. With regard to lentil, no information is available to date on diversity among lentil genotypes in the transpiration response to vapor pressure deficit (VPD). Since lentil is grown in environments with high VPD conditions (hot and dry areas) in the post-rainy season (South-Asia) where scanty rainfall is frequently observed, the limited-transpiration trait might be especially important in this particular region.

Model assessment of water conservation traits

The impact of water-conservation traits on lentil productivity is likely to vary across growing seasons with geography, environment type, and level of expression of the trait. The positive consequence of partial stomatal closure with soil drying or under high VPD is that transpiration rate is decreased early in the growing season so that there is conservation of soil water for use later in the season to complete reproductive growth. The initiation of soil water conservation will have the double benefit of increasing transpiration water use efficiency, and conserving soil water for use later in the growing season as drought develops in comparison with those plants that delay their stomatal closure [62]. If the period of water-deficit is sufficiently long, the plant genotype with partial stomatal closure with soil drying will be better positioned to sustain CO₂ assimilation and to produce greater yield. However, if there is late-season rainfall or if the decrease of photosynthetic rate due to partial stomatal closure is too high, the benefit of conserved water would not be obtained, thus, lines with water-conservation traits lead to equal or lower yield than line that does not express these traits [62]. Considering the breadth of geographical area and environments in which lentil is grown, the assessment of the potential benefits of the water-saving traits on lentil yield can only be done effectively by using crop simulation modeling.

Not only is there no information among lentil genotypes about the expression of limited-transpiration traits, there is no indication of the possible yield benefit of developing genotypes that express these traits. The complexity associated with genotype × environment × management interactions can be explored in a quantitative assessment using a mechanistic simulation model [30, 63]. Simulation studies have proved useful to evaluate the impact of a limited-transpiration rate at high VPD in sorghum [59], soybean [61, Sinclair et al. 2014] and maize [30]. Simulations of chickpea—another cool-season legume—have shown that the limited-transpiration trait can result in 3–7% increased grain yield in Iran depending on location and soil depth [63].

Therefore, simulation modeling efforts are also needed to do a geospatial assessment of the likely effect of genotypic variation in the limited-transpiration traits on yield performance [61, 63] of lentil at a regional scale. A lentil

version of the SSM-Legumes model was developed and proven robust in evaluating variation in phenological development and yield of lentil in a range of environments, with different rainfall patterns, in the Middle East [15], and investigating the roles of changing phenology and sowing dates on the possible expansion of lentil culture in East Africa [16]. This model needs to be applied to assessing the potential values of these traits on water-limited lentil.

Conclusion

Two water-conservation traits seem to be especially promising in increasing lentil yield in water—(1) limited environments early partial stomatal closure under soil drying, resulting in soil water conservation and (2) limited-transpiration under high atmospheric vapor pressure deficit. A multi-level, multi-faceted approach needs to be applied to study these traits in lentil involving simulation studies using a mechanistic crop simulation modeling and physiological screenings.

Authors' contributions

MEG, TRS conceived the paper. MEG, TRS, FK, and JG wrote the paper. All authors read and approved the final manuscript.

Author details

¹ Crop Physiology Laboratory, International Center for Agricultural Research in the Dry Areas (ICARDA), North-Africa Platform, Rabat, Morocco. ² Montpellier SupAgro, Montpellier, France. ³ Department of Crop Science, North Carolina State University (NCSU), Raleigh, NC, USA.

Acknowledgements

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

All authors agreed on publication.

Funding

This work was supported by the CGIAR Research Program on Grain Legumes and the USAID/CGIAR-US Universities Linkage Program.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 15 December 2016 Accepted: 12 April 2017

Published online: 23 August 2017

References

- Belko N, Zaman-Allah M, Diop NN, Cisse N, Zombre G, Ehlers JD, Vadez V. Restriction of transpiration rate under high vapor pressure deficit and non-limiting water conditions is important for terminal drought tolerance in cowpea. *Plant Biol.* 2013;15:304–16.
- Devi MJ, Sinclair TR, Vadez V, Krishnamurthy L. Peanut genotypic variation in transpiration efficiency and decreased transpiration during progressive soil drying. *F Crop Res.* 2009;114(2):280–5. doi:10.1016/j.fcr.2009.08.012.
- Devi JM, Sinclair TR, Vadez V. Genotypic variation in peanut for transpiration response to vapor pressure deficit. *Crop Sci.* 2010;50:191–6.
- Devi JM, Taliercio EW, Sinclair TR. Leaf expansion of soybean subjected to high and low atmospheric vapour pressure deficits. *J Exp Bot.* 2015;66:1845–50.
- Donat MG, Lowry AL, Alexander LV, O'Gorman PA, Maher N. More extreme precipitation in the world's dry and wet regions. *Nat Clim Ch.* 2016;6:508–13.
- Erskine W, Rihawi S, Capper BS. Variation in lentil straw quality. *Anim Feed Sci Technol.* 1990;28:61–9.
- Erskine W, Tufail M, Russell A, Tyagi MC, Rahman MM, Saxena MC. Current and future strategies in breeding lentil for resistance to biotic and abiotic stresses. *Euphytica.* 1994;73:127–35.
- Erskine W, Muehlbauer FJ, Sarker A, Sharma B. The lentil: botany production and uses. Wallingford: CAB International; 2009. p. 447.
- Erskine W, Sarker A, Kumar S. Crops that feed the world: investing in lentil improvement toward a food secure world. *Food Secur.* 2011;3:127–39. doi:10.1007/s12571-011-0124-5.
- FAO (2015) FAOSTAT. <http://faostat3.fao.org/browse/Q/QC/E>. Accessed 17 Feb 2016.
- Fischer RA. Growth and water limitation to dryland wheat in Australia: a physiological framework. *J Aust Inst Agric Sci.* 1979;45:83–94.
- Fletcher AL, Sinclair TR, Allen LH Jr. Transpiration responses to vapor pressure deficit in well-watered slow wilting and commercial soybean. *Environ Exp Bot.* 2007;61:145–51.
- Gahoonia TS, Omar A, Sarker A, Rahman MM, Erskine W, et al. Root traits, nutrient uptake, multi location grain yield and benefit-cost ratio of two lentil (*Lens culinaris*, Medikus) varieties. *Plant Soil.* 2005;272:153–61.
- Gahoonia TS, Ali O, Sarker A. Genetic variation in root traits and nutrient acquisition of lentil genotypes. *J Plant Nutr.* 2006;29:643–55.
- Ghanem ME, Marrou H, Soltani A, Kumar S, Sinclair TR. Lentil variation in phenology and yield evaluated with a model. *Agron J.* 2015;107:1967–77.
- Ghanem ME, Marrou H, Biradar C, Sinclair TR. Production potential of lentil (*Lens culinaris* Medik.) in East Africa. *Agric Syst.* 2015;137:24–38.
- Ghanem ME, Marrou H, Sinclair TR. Physiological phenotyping of plants for crop improvement. *Trends Plant Sci.* 2015;20:139–44.
- Gholipour M, Choudhary S, Sinclair TR, Messina CD, Cooper M. Transpiration response of maize hybrids to atmospheric vapour pressure deficit. *J Agron Crop Sci.* 2013;199:155–60.
- Gholipour M, Prasad PV, Mutava RN, Sinclair TR. Genetic variability of transpiration response to vapor pressure deficit among sorghum genotypes. *F Crop Res.* 2010;119:85–90.
- Gilbert ME, Zwieniecki MA, Holbrook NM. Independent variation in photosynthetic capacity and stomatal conductance leads to differences in intrinsic water use efficiency in 11 soybean genotypes before and during mild drought. *J Exp Bot.* 2011;62:2875–87.
- Idrissi O, Udupa SM, Houasli C, De Keyser E, Van Damme P, De Riek J. Genetic diversity analysis of moroccan lentil (*Lens culinaris* Medik.) landraces using simple sequence repeat and amplified fragment length polymorphisms reveals functional adaptation towards agro-environmental origins. *Plant Breed.* 2015;134:322–32. doi:10.1111/pbr.12261.
- Kholová J, Hash CT, Kakkera A, Kocova M, Vadez V. Constitutive water-conserving mechanisms are correlated with the terminal drought tolerance of pearl millet [*Pennisetum glaucum* (L.) R. Br.]. *J Exp Bot.* 2010;61(2):369–77. doi:10.1093/jxb/erp314.
- Kholová J, Hash CT, Kumar PL, Yadav RS, Kocova M, Vadez V. Terminal drought-tolerant pearl millet [*Pennisetum glaucum* (L.) R. Br.] have high leaf ABA and limit transpiration at high vapour pressure deficit. *J Exp Bot.* 2010;61(5):1431–40. doi:10.1093/jxb/erq013.
- Kholová J, Tharanya M, Kaliamoorthy S, Malayee S, Baddam R, Hammer GL, McLean G, Deshpande S, Hash CT, Craufurd PQ, Vadez V. Modelling the effect of plant water use traits on yield and stay-green expression in sorghum. *Funct Plant Biol.* 2014;41(10–11):1019–34.
- Kumar J, Basu PS, Srivastava E, Chaturvedi SK, Nadarajan N, Kumar S. Phenotyping of traits imparting drought tolerance in lentil. *Crop Pasture Sci.* 2012;63:547–54.
- Kumar SK, Barpete S, Kumar J, Gupta P, Sarker A. Global lentil production: constraints and strategies. *SATSA Mukhapatra—Annu. Tech Issue.* 2013;17:1–13.
- Leport L, Turner NC, French RJ, Tennant D, Thomson BD, Siddique KHM. Water relations, gas exchange and growth of cool-season grain legumes in a Mediterranean-type environment. *Eur J Agron.* 1998;9(4):295–303. doi:10.1016/s1161-0301(98)00042-2.

28. Ludlow MM, Muchow RC. A critical evolution of traits for improving crop yields in water-limited environments. *Adv Agron.* 1990;43:107–53.
29. Meisner CA, Karnok KJ. Peanut root response to drought stress. *Agron J.* 1992;84:159–65.
30. Messina CD, Sinclair TR, Hammer GL, Curan D, Thompson J, Oler Z, Gho C, Cooper M. Limited-transpiration trait may increase maize drought tolerance in the US Corn Belt. *Agron J.* 2015;107:1978–86.
31. Mia WM, Yamaguchi A, Kono Y. Root system structure of six food legume species: interand intraspecific variation. *Jpn J Crop Sci.* 1996;65(1):131–40.
32. Mishra BK, Singh AK, Lal JP, Srivastava JP. Drought stress resistance in lentil. National symposium on vegetable legumes for soil and human health, Varanasi; 2016.
33. Neupane S, Neupane RK, Shrestha R. Report on varietal and agronomical studies on lentil. Paper presented at the Winter Crop Workshop in Nepal Agricultural Research Council (NARC), Kathmandu, Nepal, 16–20 September; 1991.
34. Pang J, Neil C, Turner Khan T, Yan-Lei D, Xiong JL, Colmer TD, Devilla R, Stefanova K, Kadambot HM, Siddique. Response of chickpea (*Cicer arietinum* L.) to terminal drought: leaf stomatal conductance, pod abscisic acid concentration, and seed set. *J Exp Bot.* 2016. doi:10.1093/jxb/erw153.
35. Quinn MA. Biological nitrogen fixation and soil health improvement. In: Erskine W, Muehlbauer FJ, Sarker A, Sharma B, editors. The lentil—botany, production and uses. Wallingford: Comm Agric Bureau Int; 2009. p. 229–47.
36. Rahman A, Tawaha M, Turk MA, et al. Effect of dates and rates of sowing on yield and yield components of lentil (*Lens culinaris* Medik.) under semi-arid conditions. *Pak J Biol Sci.* 2002;5:531–2.
37. Ratnakumar P, Vadez V, Nigam SN, Krishnamurthy L. Assessment of transpiration efficiency in peanut (*Arachis hypogaea* L.) under drought by lysimetric system. *Plant Biol.* 2009;11:124–30.
38. Rebetzke GJ, Condon AG, Richards RA, Farquhar GD. Genetic control of leaf conductance in three wheat crosses. *Aust J Agric Res.* 2003;54:381–7.
39. Riar MK, Sinclair TR, Vara Prasad PV. Persistence of limited-transpiration-rate trait in sorghum at high temperature. *Environ Exp Bot.* 2015;115:58–62.
40. Sadok W, Sinclair TR. Genetic variability of transpiration response to vapor pressure deficit among soybean (*Glycine max* [L] Merr.) genotypes selected from a recombinant inbred line population. *F Crop Res.* 2009;113:156–60.
41. Sadok W, Sinclair TR. Crops yield increase under water limited conditions: review of recent physiological advances in soybean genetic improvement. *Adv Agron.* 2011;113:313–37.
42. Sarker A, Erskine W, Singh M. Variation in shoot and root characteristics and their association with drought tolerance in lentil landraces. *Genet Resour Crop Evol.* 2005;52(1):89–97. doi:10.1007/s10722-005-0289-x.
43. Saxena MC. Plant morphology, anatomy and growth habit. In: Erskine W, Muehlbauer F, Sarker A, Sharma B, editors. The lentil: botany, production and uses. 1st ed. London: CABI Publishing; 2009. p. 34–46.
44. Schoppach R, Sadok W. Differential sensitivities of transpiration to evaporative demand and soil water deficit among wheat elite cultivars indicate different strategies for drought tolerance. *Environ Exp Bot.* 2012;84:1–10.
45. Seversike TM, Sermons SM, Sinclair TR, Carter TE Jr, Rufty TW. Temperature interactions with transpiration response to vapor pressure deficit among cultivated and wild soybean genotypes. *Physiol Plant.* 2013;148:62–73.
46. Sharma SN, Prasad RP. Effect of soil moisture regimes on the yield and water use of lentil (*Lens culinaris* Medic). *Irrig Sci.* 1984;5:285–93.
47. Shekoofa A, Devi JM, Sinclair TR, Holbrook CC, Isleib TG. Divergence in drought-resistance traits among parents of recombinant 1 peanut inbred lines. *Crop Sci.* 2013;53:2569–76.
48. Shekoofa A, Balota M, Sinclair TR. Limited-transpiration trait evaluated in growth chamber and field for sorghum genotypes. *Environ Exp Bot.* 2014;99:175–9.
49. Shrestha R. Effect of planting dates and seed rates on lentil (var. Simal) at Khumaltar (1993/94–1995/96). National Winter Crop Workshop. Regional Agricultural Research Station, Bhairahawa. Nepal Agricultural Research Council; 1996.
50. Shrestha R. Adaptation of lentil (*Lens culinaris* Medikus ssp. *culinaris*) to rainfed environments—response to water deficits. Crawley: The University of Western Australia; 2005.
51. Shrestha R, Turner NC, Siddique KHM, Turner DW. Physiological and seed yield responses to water deficits among lentil genotypes from diverse origins. *Aust J Agric Res.* 2006;57(8):903–15. doi:10.1071/ar05204.
52. Shrestha R, Siddique KHM, Turner DW, Turner NC. Breeding and management to minimize the effects of drought and improve water use efficiency. In: Erskine W, Muehlbauer F, Sarker A, Sharma B, editors. The lentil: botany, production and uses. 1st ed. London, UK: CABI Publishing; 2009.
53. Siddique KHM, Regan KL, Tennant D, Thomson BD. Water use and water use efficiency of cool season grain legumes in low rainfall Mediterranean-type environments. *Eur J Agron.* 2001;15:267–80.
54. Siddique KHM, Loss SP, Thomson BD, et al. Cool season grain legumes in dryland Mediterranean environments of Western Australia: significance of early flowering. In: Saxena NP, editor. Management of agricultural drought: agronomic and genetic options. Enfield: Science Publishers Inc.; 2003. p. 151–61.
55. Silim SN, Saxena MC, Erskine W. Adaptation of lentil to the Mediterranean environment. I. Factors affecting yield under drought conditions. *Exp Agric.* 1993;29(01):9. doi:10.1017/s0014479700020366.
56. Sinclair TR. Theoretical analysis of soil and plant traits influencing daily plant water flux on drying soil. *Agron J.* 2005;97:1148–52.
57. Sinclair TR. Is transpiration efficiency a viable plant trait in breeding for crop improvement? *Funct Plant Biol.* 2012;39:359–65.
58. Sinclair TR, Ludlow MM. Influence of soil water supply on the plant water balance of four tropical grain legumes. *Aust J Plant Physiol.* 1986;13:329–41.
59. Sinclair TR, Hammer GL, van Oosterom EJ. Potential yield and water-use efficiency benefits in sorghum from limited maximum transpiration rate. *Funct Plant Biol.* 2005;32:945–52.
60. Sinclair TR, Zwieniecki MA, Holbrook NM. Low leaf hydraulic conductance associated with drought tolerance in soybean. *Physiol Plant.* 2008;132:446–51.
61. Sinclair TR, Messina CD, Beatty A, Samples M. Assessment across the United States of the benefits of altered soybean drought traits. *Agron J.* 2010;102:475–82.
62. Sinclair TR, Devi JM, Carter TE Jr. Limited-transpiration trait for increased yield for water-limited soybean: from model to phenotype to genotype to cultivars. In: Yin X, Struik PC, editors. Crop systems biology. Berlin: Springer; 2016. p. 129–46.
63. Soltani A, Sinclair TR. Modeling physiology of crop development, growth and yield. CABI ed. Wallingford; 2012.
64. Thomson BD, Siddique KHM, Barr MD, Wilson JM. Grain legume species in low rainfall mediterranean-type environments I. Phenology and seed yield. *Field Crops Res.* 1997;54:173–87.
65. Turner NC, Wright GC, Siddique KHM. Adaptation of grain legumes (pulses) to water-limited environments. *Adv Agron.* 2001. doi:10.1016/s0065-2113(01)71015-2.
66. Vadez V, Berger JD, Warkentin T, Asseng S, Ratnakumar P, Poorna Chandra Rao K, Gaur PM, Munier-Jolain N, Larmure A, Voisin AS, Sharma HC, Pande S, Sharma M, Krishnamurthy L, ZamanAllah MA. Adaptation of grain legumes to climate change: a review. *Agron Sustain Dev.* 2012;32(1):31–44.
67. Yang Z, Sinclair TR, Messina CD, Cooper M, Hammer GL. Temperature effect on transpiration response of maize plants to vapour pressure deficit. *Environ Exp Bot.* 2012;78:157–62.
68. Zaman-Allah M, Jenkinson DM, Vadez V. Chickpea genotypes contrasting for seed yield under terminal drought stress in the field differ for traits related to control of water use. *Funct Plant Biol.* 2011;38:270–328.
69. Zhang H, Pala M, Oweis T, Harris H. Water use and water-use efficiency of chickpea and lentil in a Mediterranean environment. *Aust J Agric Res.* 2000;51:295–304.