

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

OPTICAL ABSORPTION INTENSITIES OF RARE-EARTH IONS

### Permalink

<https://escholarship.org/uc/item/0hd516pd>

### Author

Judd, B.R.

### Publication Date

1962

**University of California**

**Ernest O. Lawrence  
Radiation Laboratory**

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

**Berkeley, California**

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory

Berkeley, California

Contract No. W-7405-eng-48

OPTICAL ABSORPTION INTENSITIES OF RARE-EARTH IONS

B. R. Judd

January, 1962

## OPTICAL ABSORPTION INTENSITIES OF RARE-EARTH IONS

Contents

Abstract . . . . .	iv
I. Introduction . . . . .	1
II. Matrix Elements . . . . .	2
III. Approximations . . . . .	6
IV. Solutions of Rare Earth Ions. . . . .	11
V. Other Contributions to P . . . . .	13
VI. Comparison with Experiment. . . . .	16
VII. Variation of $T_2$ . . . . .	19
VIII. Environment of a Rare-Earth Ion in Solution. . . . .	20
IX. Calculation of the Parameters $T_\lambda$ . . . . .	23
X. Discussion . . . . .	28
XI. Conclusion . . . . .	31
Acknowledgments . . . . .	32
References . . . . .	41

## OPTICAL ABSORPTION INTENSITIES OF RARE EARTH IONS

B. R. Judd

Lawrence Radiation Laboratory  
 University of California  
 Berkeley, California

January, 1962

## ABSTRACT

Electric dipole transitions within the  $4f$  shell of a rare-earth ion are permitted if the surroundings of the ion are such that its nucleus is not situated at a center of inversion. An expression is found for the oscillator strength of a transition between two states of the ground configuration  $4f^N$ , on the assumption that the levels of each excited configuration of the type  $4f^N n'd$  or  $4f^N n'g$  extend over an energy range small as compared to the energy of the configuration above the ground configuration. On summing over all transitions between the components of the ground level  $\psi_J$  and those of an excited level  $\psi'_{J'}$ , both of  $4f^N$ , the oscillator strength  $P$  corresponding to the transition  $\psi_J \rightarrow \psi'_{J'}$  is found to be given by

$$P = \sum T_\lambda \nu (\psi_J \| U^{(\lambda)} \| \psi'_{J'})^2,$$

where  $U^{(\lambda)}$  is a tensor operator of rank  $\lambda$ , and the sum runs over the three values 2, 4, and 6 of  $\lambda$ . Transitions that also involve changes in the vibrational modes of the complex comprising a rare-earth ion and its surroundings, provide a contribution to  $P$  of precisely similar form. It is shown that sets of parameters  $T_\lambda$  can be chosen to give a good fit with the experimental data on aqueous solutions of  $NdCl_3$  and  $ErCl_3$ . A calculation

on the basis of a model, in which the first hydration layer of the rare-earth ion does not possess a center of symmetry, leads to parameters  $T_\lambda$  that are smaller than those observed for  $\text{Nd}^{3+}$  and  $\text{Er}^{3+}$  by factors of 2 and 8 respectively. Reasons for the discrepancies are discussed.

## OPTICAL ABSORPTION INTENSITIES OF RARE EARTH IONS

B. R. Judd

Lawrence Radiation Laboratory  
University of California  
Berkeley, California

## I. INTRODUCTION

The last decade has witnessed a remarkable growth in our knowledge of the spectroscopic properties of triply ionized rare-earth atoms. The interplay of experiment and theory has led to the elucidation of appreciable parts of the term schemes of many ions, and the splittings in the levels that arise when a rare earth ion is situated in a crystal lattice are now understood rather well. From the present vantage point, Van Vleck's classic paper The Puzzle of the Rare Earths<sup>1</sup> makes interesting reading, published as it was at a time when even the configurations involved in the spectral transitions had not been definitely established. The arguments remain essentially valid. The sharp absorption lines of rare-earth crystals in the visible and infra-red regions of the spectrum do correspond to transitions within the configurations of the type  $4f^N$ , and the so-called extra levels have their origin in the interplay of electronic and vibrational effects.

Van Vleck's paper discusses the nature of the electronic transitions, that is, whether they can be classified as electric dipole, magnetic dipole, or electric quadrupole. His conclusion, that all three types play a role, was later criticized by Broer, Gorter, and Hoogschagen, who showed that the observed intensities of the transitions are in almost all cases too intense for magnetic dipole or electric quadrupole radiation to be important.<sup>2</sup> They also demonstrated that electric dipole transitions could be sufficiently



strong to match the experimental intensities, but their calculations can at best be described as semi-quantitative. The difficulty in estimating intensities of electric dipole transitions is that they arise from the admixture into  $4f^N$  of configurations of opposite parity. To calculate such admixtures, not only must the energies and eigenfunctions of configurations such as  $4f^{N-1} 5d$  be known, but also that part of the crystal field potential responsible for the admixing. The problem of obtaining these data has proved complicated enough to restrain the performance of further theoretical work on the intensities of the absorption lines of the rare earths, though considerable advances have been made in the last few years on the similar problem of estimating the intensities of lines of transition-metal ions (see, for example, Griffith<sup>3</sup>). An added reason for the absence of a detailed theory may be the comparative lack of experimental data; for, apart from a few isolated cases,<sup>4</sup> the only oscillator strengths measured at present appear to be for solutions of rare-earth ions.<sup>5,6</sup> However, the situation will undoubtedly be remedied shortly. This expectation, taken with the information gained in the last decade on the properties, both experimental and theoretical, of the rare-earth ions, makes a fresh examination of the intensities of the absorption lines an attractive venture.

## II. MATRIX ELEMENTS

The oscillator strength  $P'$  of a spectral line, corresponding to the electric dipole transition from the component  $i$  of the ground level of an ion to the component  $f$  of an excited level, is given by

$$P' = \chi [8\pi^2 m\nu/h] | (i | D_q^{(1)} | f) |^2. \quad (1)$$

In this equation,  $m$  is the mass of an electron,  $h$  is Planck's constant, and

$\nu$  is the frequency of the line. The factor  $\chi$  makes allowance for the refractive index of the medium in which the ion is embedded: according to Broer et al., for water  $\chi = 1.19$ .<sup>2</sup> In terms of the polar coordinates  $(r_j, \theta_j, \phi_j)$  of electron  $j$ ,

$$D_q^{(k)} = \sum_j r_j^k C_q^{(k)}(\theta_j, \phi_j),$$

where

$$C_q^{(k)}(\theta_j, \phi_j) = [4\pi/(2k+1)]^{\frac{1}{2}} Y_{kq}(\theta_j, \phi_j),$$

$Y_{kq}$  being a spherical harmonic. The choice of  $q$  in Eq. (1) depends on the polarization of the incident light. Eq. (1) can be regarded as a slight elaboration of Eq. (6-58) of Slater.<sup>7</sup>

In order to evaluate the matrix element of Eq. (1), we need detailed descriptions of the states  $i$  and  $f$ . Owing to the comparatively small splittings of the levels produced by the crystal field, it is usually a good approximation to assume at first that the quantum number  $J$ , corresponding to the total angular momentum of the electron system of the rare-earth ion, remains a good quantum number. Corresponding to the component  $i$  of the ground level of the configuration  $\ell^N$ , there exists, to the first approximation, a linear combination

$$(A| \equiv \sum_M (\ell^N \psi_{J M} | a_M, \quad (2)$$

where  $M$  denotes the quantum number of the projection  $J_z$  of  $\underline{J}$ . The symbol  $\psi$  stands for the additional quantum numbers that may be necessary to define the level uniquely; if RS (Russell-Saunders) coupling were strictly followed, it would incorporate a definite  $S$  and  $L$ , the quantum numbers corresponding to the total spin and total orbit, respectively, of the electron system. However, it is unnecessary at this point to assume RS coupling. By analogy with Eq. (2),

we may write

$$|A'\rangle \equiv \sum_{M'} a'_{M'} |l^N \psi' J' M'\rangle \quad (3)$$

for the first approximation to the upper state.

It might be thought that rare-earth ions in solution would be subject to rapidly fluctuating electric fields, and that the coefficients  $a_M$  and  $a'_{M'}$  would therefore vary with time. While this may be true to a slight extent, it is now virtually certain that for aqueous solutions the immediate surroundings of the ions are rigidly locked in position. Evidence for this will be presented later; we mention it here to eliminate the possible misapprehension that the linear combinations (2) and (3) might have a well-defined significance only for ions imbedded in crystal lattices.

The states  $|A\rangle$  and  $|A'\rangle$ , being constructed from the same configuration  $l^N$ , possess the same parity. However, under the replacement  $r_j \rightarrow -r_j$ , we find  $\underline{D}^{(1)} \rightarrow -\underline{D}^{(1)}$ . The equation

$$\langle A | \underline{D}_q^{(1)} | A' \rangle = 0$$

follows, and hence to the first approximation, no electric dipole transition occur. This is merely a statement of the Laporte rule, of course. To obtain nonvanishing matrix elements of the components of  $\underline{D}^{(1)}$ , it is necessary to admix into  $|A\rangle$  and  $|A'\rangle$  states built from configurations of opposite parity to  $l^N$ . For the moment, we consider only those configurations of the type  $l^{N-1} l'$ ; these are certainly the most important. To distinguish such configurations, we augment  $l'$  with the principal quantum number  $n'$ ; the symbol  $n$  is reserved for the analogous quantum number for the electrons of the ground configuration  $l^N$ , but we shall give it explicitly only when an ambiguity threatens.

The admixing of configurations of opposite parity can come about if the contribution  $V$  to the Hamiltonian arising from the interaction of the electrons of the ion with the electric field of the lattice, here assumed to be static, contains terms of odd parity. On making the expansion

$$V = \sum_{t,p} A_{tp} D_p(t),$$

this condition becomes equivalent to the demand that not all  $A_{tp}$ , for which  $t$  is odd, vanish. The states (2) and (3) are now replaced by

$$\begin{aligned} |B\rangle \equiv \sum_M (\ell^N \psi J M | a_M \\ + \sum_K (\ell^{N-1} (n'l') \psi'' J'' M'' | b(n'l', \psi'' J'' M'')) \end{aligned}$$

and

$$\begin{aligned} |B'\rangle \equiv \sum_{M'} a'_{M'} | \ell^N \psi' J' M' \rangle \\ + \sum_K b'(n'l', \psi'' J'' M'') | \ell^{N-1} (n'l') \psi'' J'' M'' \rangle, \end{aligned}$$

where

$$\begin{aligned} b(n'l', \psi'' J'' M'') = \sum_M a_M (\ell^N \psi J M | V | \ell^{N-1} (n'l') \psi'' J'' M'') \\ \times [E(\psi J) - E(n'l', \psi'' J'')]^{-1}, \end{aligned} \quad (4)$$

and

$$\begin{aligned} b'(n'l', \psi'' J'' M'') = \sum_{M'} a'_{M'} (\ell^{N-1} (n'l') \psi'' J'' M'' | V | \ell^N \psi' J' M') \\ \times [E(\psi' J') - E(n'l', \psi'' J'')]^{-1}. \end{aligned} \quad (5)$$

The symbol  $\sum_K$  stands for the sum over  $\psi''$ ,  $J''$ ,  $M''$   $l'$  and over those values of  $n'$  for which  $\ell^{N-1} (n'l')$  is an excited configuration. In Eqs. (4) and (5),  $E(\psi J)$  and  $E(\psi' J')$  denote the energies of the levels  $\psi J$  and  $\psi' J'$  of  $\ell^N$ ;

similarly,  $E(n'l', \psi''J'')$  stands for the energy of the level  $\psi''J''$  of  $l^{N-1}(n'l')$ .

It is now a simple matter to obtain the equation

$$\begin{aligned}
 (B|D_q^{(1)}|B') &= \sum a_M a'_{M'} A_{tp} \{ (l^N \psi J M | D_q^{(1)} | l^{N-1}(n'l') \psi'' J'' M'') \\
 &\quad \times (l^{N-1}(n'l') \psi'' J'' M'' | D_p^{(t)} | l^N \psi' J' M') \\
 &\quad \times [E(\psi' J') - E(n'l', \psi'' J'')]^{-1} \\
 &+ (l^N \psi J M | D_p^{(t)} | l^{N-1}(n'l') \psi'' J'' M'') \\
 &\quad \times (l^{N-1}(n'l') \psi'' J'' M'' | D_q^{(1)} | l^N \psi' J' M') \\
 &\quad \times [E(\psi J) - E(n'l', \psi'' J'')]^{-1} \}, \tag{6}
 \end{aligned}$$

the sum running over  $M, M', t, p$ , and those quantum numbers implied by the symbol  $\kappa$ .

### III. APPROXIMATIONS

For all but the most trivial configurations, the nine-fold sum of Eq. (6) is quite unwieldy. We must therefore search for approximate methods, taking care to make them as realistic as possible. The occurrence of the structure

$$|l^{N-1}(n'l') \psi'' J'' M''\rangle \langle l^{N-1}(n'l') \psi'' J'' M''|$$

in Eq. (6) suggests that it might be possible to adapt the closure procedure in some way, thereby uniting  $D_q^{(1)}$  and  $D_p^{(t)}$  into a single operator that acts

between states of  $\ell^N$ . For a description and analysis of this method, see Griffith.<sup>8</sup> The number of summations we wish to absorb into the closure depends on how far we are prepared to assume  $E(n'l', \psi'' J'')$  is invariant with respect to  $n'$ ,  $\ell'$ ,  $\psi''$ , or  $J''$ . For example, the mildest approximation we can make is to suppose that the splittings within multiplets of the excited configurations are negligible compared with the energies that the configurations as a whole lie above  $\ell^N$ . This amounts to supposing  $E(n'l', \psi'' J'')$  is independent of  $J''$ . If the states of  $\ell^N$  are expanded as linear combinations of perfect RS-coupled states of the type

$$(\ell^N \gamma S L J M |,$$

we can perform the sums over  $J''$  and  $M''$  in Eq. (6) by making use of equations such as

$$\begin{aligned} & \sum_{J'', M''} (\ell^N \gamma S L J M | D_q^{(1)} | \ell^{N-1} \ell' \gamma'' S L'' J'' M'') \\ & \times (\ell^{N-1} \ell' \gamma'' S L'' J'' M'' | D_p^{(t)} | \ell^N \gamma' S L' J' M') \\ & = \sum_{\lambda} (-1)^{p+q+L+L'} (2\lambda+1) \begin{pmatrix} 1 & \lambda & t \\ q & -q-p & p \end{pmatrix} \left\{ \begin{matrix} 1 & \lambda & t \\ L' & L'' & L \end{matrix} \right\} \\ & \times (\ell^N \gamma S L J M | T_{p+q}^{(\lambda)} | \ell^N \gamma' S L' J' M'), \end{aligned} \quad (7)$$

where  $T_{p+q}^{(\lambda)}$  is a tensor whose amplitude is determined by

$$(L || T^{(\lambda)} || L') = (L || D^{(1)} || L'')(L'' || D^{(t)} || L'). \quad (8)$$

The easiest way to verify Eq. (7) is to express all the matrix elements in terms of reduced matrix elements of the type involved in Eq. (8); it is then found that Eq. (7) is equivalent to the Biedenharn-Elliott sum rule (see Edmonds<sup>9</sup>).

It is at once evident that the simplifications afforded by using Eqs. (7) and (8) are very slight. The degree of closure must therefore be extended. The least severe extension is to suppose that  $E(n'l', \psi'' J'')$  is invariant with respect to  $\psi''$  as well as to  $J''$ . This is equivalent to regarding the excited configuration  $l^{N-1} (n'l')$  as completely degenerate. A glance at diagrams giving the approximate positions and extensions of low-lying configurations of the rare-earth ions, such as Figs. 4 and 5 of Dieke, Crosswhite and Dunn,<sup>10</sup> indicates at once that this assumption is only moderately fulfilled. It therefore constitutes a weak link in the theory. However, we may hope that the very complexity of configurations of the type  $l^{N-1} (n'l')$  might reduce the possible error; for if there are a great many terms  $\psi''$  in  $l^{N-1} (n'l')$ , it would not be unreasonable to expect that the entire sum over  $\psi''$ , if broken up into smaller sums over groups of closely lying terms, would decompose into a number of parts that, for various  $\psi'$  and  $J'$ , were roughly proportional to one another.

Be this as it may, the approximation leads to a great simplification in the mathematics. The analogue of Eq. (7) is

$$\begin{aligned}
 & \sum_{J'', M'' \psi''} (l^N \psi_{J M} | D_q^{(1)} | l^{N-1} (n'l') \psi'' J'' M'') \\
 & \times (l^{N-1} (n'l') \psi'' J'' M'' | D_p^{(t)} | l^N \psi' J' M') \\
 & = \sum_{\lambda} (-1)^{p+q} (2\lambda+1) \begin{pmatrix} 1 & \lambda & t \\ q & -p-q & p \end{pmatrix} \left\{ \begin{matrix} 1 & \lambda & t \\ l & l' & l \end{matrix} \right\} \\
 & \times (n l | r | n'l') (n l | r^t | n'l') (l || C^{(1)} || l') \\
 & \times (l' || C^{(t)} || l) (l^N \psi_{J M} | U_{p+q}^{(\lambda)} | l^N \psi' J' M'), \quad (9)
 \end{aligned}$$

where  $\underline{u}^{(\lambda)}$  is the sum over all the electrons of the single-electron tensors  $u^{(\lambda)}$ , for which

$$(\ell \parallel \underline{u}^{(\lambda)} \parallel \ell) = 1.$$

In Eq. (9), the abbreviation

$$(n \ell | r^k | n' \ell') = \int_0^\infty \mathcal{R}(n \ell) r^k \mathcal{R}(n' \ell') dr \quad (10)$$

is introduced, where  $\mathcal{R}/r$  is the radial part of the appropriate single-electron eigenfunction. A straightforward way of deriving Eq. (9) is to expand the matrix elements of Eq. (7) by means of Eq. (27) of Racah,<sup>11</sup> to perform the sum over  $\psi''$ , and then to pass from RS to intermediate coupling. Alternatively, both Eqs. (7) and (9) can be obtained from Eq. (7.1.1) of Edmonds,<sup>9</sup> provided the symbol  $\gamma''$  of that equation is interpreted judiciously.

Equation (9) is excellent for the purposes we have in mind; however, the closure procedure can be extended even further. If we assume  $E(n' \ell'; \psi'' J'')$  to be invariant with respect to  $n'$  as well as to  $\psi''$  and  $J''$ , and if the full description of the ground configuration contains no electrons with azimuthal quantum number  $\ell'$ , then the fact that the radial functions  $\mathcal{R}(n' \ell')$ , for all  $n'$  form a complete set allows us to write

$$\sum_{n'} (n \ell | r | n' \ell') (n' \ell' | r^t | n \ell) = (n \ell | r^{t+1} | n \ell), \quad (11)$$

and the problem of calculating interconfiguration radial integrals disappears. Since for rare-earth ions both the 3d and 4d shells are filled, this technique could not be used for  $\ell' = 2$ ; on the other hand, there is no objection to applying it to electrons for which  $\ell' = 4$ , since no g orbital is occupied in the ground configuration. The possible occupation of  $\ell'$  orbitals precludes our extending the closure to all four quantum numbers  $n'$ ,  $\ell'$ ,  $\psi''$  and  $J''$ .



Equation (9) can be used immediately to simplify the first product on the right-hand side of Eq. (6). A precisely similar substitution may be made for the second product; but owing to the relation

$$\begin{pmatrix} 1 & \lambda & t \\ q & -p-q & p \end{pmatrix} = (-1)^{1+\lambda+t} \begin{pmatrix} t & \lambda & 1 \\ p & -p-q & q \end{pmatrix}$$

the two parts cancel to a large extent if  $1+\lambda+t$  is odd. For the right-hand side of Eq. (9) not to vanish,  $t$  must be odd; hence the condition is fulfilled if  $\lambda$  is odd. The cancellation would be perfect if, for a given  $n'$  and  $l'$ , the energy denominators

$$\begin{aligned} E(\psi J) - E(n' l', \psi' J''), \\ E(\psi' J') - E(n' l', \psi'' J''), \end{aligned} \tag{12}$$

which are supposed to be independent of  $\psi''$  and  $J''$ , could be assumed equal. This is equivalent to the supposition that the configurations  $l^{N-1} (n' l')$  lie far above the states involved in the optical transitions. Although the theory could no doubt be developed without making this assumption, a considerable simplification in the mathematics results if it is made. We therefore replace both differences (12) with the single expression  $\Delta(n' l')$ . Equation (6) can now be written as

$$\begin{aligned} & (B | D_q^{(1)} | B') \\ = & \sum_{p,t,\text{even } \lambda} (2\lambda+1)(-1)^{p+q} A_{tp} \begin{pmatrix} 1 & \lambda & t \\ q & -p-q & p \end{pmatrix} \\ & \times (A | U_{p+q}^{(\lambda)} | A') E(t, \lambda), \end{aligned} \tag{13}$$

where

$$\begin{aligned} E(t, \lambda) = & 2 \sum (2\ell+1)(2\ell'+1)(-1)^{\ell+\ell'} \\ & \times \begin{Bmatrix} 1 & \lambda & t \\ \ell & \ell' & \ell \end{Bmatrix} \begin{pmatrix} \ell & 1 & \ell' \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} \ell' & t & \ell \\ 0 & 0 & 0 \end{pmatrix} \\ & \times (n \ell | r | n' \ell') (n \ell | r^t | n' \ell') / \Delta(n' \ell'). \end{aligned} \quad (14)$$

The summation of Eq. (14) runs over all values of  $n'$  and  $\ell'$  consistent with  $\ell^{N-1} (n' \ell')$  being an excited configuration. In Eq. (13), the operator  $U_{p+q}^{(\lambda)}$  connects states of  $\ell^N$ ; its matrix elements can therefore be calculated by standard tensor-operator techniques.

#### IV. SOLUTIONS OF RARE-EARTH IONS

If, for a rare earth crystal, one wished to limit one's investigation of the intensities of the lines in some way, for example, to study the relative intensities of a group of lines corresponding to the transitions between the components of just a pair of levels, then no doubt Eq. (13) could be manipulated to throw the relevant quantum numbers into sharper relief. However, for general purposes, it seems unlikely that Eq. (13) could be simplified much further. In order, then, to calculate the oscillator strength of the transition from the component corresponding to  $|A\rangle$  to that corresponding to  $|A'\rangle$ , the radial integrals and crystal field parameters  $A_{pt}$  must be estimated, the sums of Eqs. (13) and (14) carried out, and the resulting matrix element

$$(B | D_q^{(1)} | B')$$

substituted for

$$(i | D_q^{(1)} | f)$$

in Eq. (1).

Apart from a few important exceptions, the fine structures of the absorption lines for rare-earth ions in solution, in contrast to those for ions in crystals, have not been resolved. Each broad absorption line corresponds to a transition from the ground level to an excited level. The measured oscillator strength of such a line is therefore the sum of the oscillator strengths of the various component lines, suitably weighted to allow for the differential probability of occupation of the components of the ground level. In the absence of detailed knowledge of the surroundings of a rare-earth ion in solution, the energies of the components of the ground level, and hence their probabilities of occupation, cannot be calculated. However, the splittings of the ground levels of rare earth ions in crystals, such as have been observed<sup>12</sup> or calculated,<sup>13</sup> seldom exceed  $250 \text{ cm}^{-1}$ . For a level where this splitting obtains, the ratio of the probabilities of occupation of the highest to the lowest component is as high as 0.3 at room temperature; therefore not too great an error should be introduced if we assume all the components of the ground level are equally likely to be occupied. Allowing for the arbitrary orientation of the rare-earth ions, Eq. (1) is replaced by

$$P = \chi [8\pi^2 m\nu/3h (2J+1)] \sum |(i | D_q^{(1)} | f)|^2, \quad (15)$$

where the sum runs over  $q$  and all components  $i$  and  $f$  of the ground and excited level. An equivalent formula has been given by Broer et al.<sup>2</sup> Using Eq. (13), we see the sum over  $i$  and  $f$  reduces to a sum over certain states of the type  $(A |$  and  $|A')$ . It is, of course, unnecessary to introduce the eigenfunctions of Eqs. (2) and (3); we can simply take the states  $(\ell^N \psi J M |$  and  $|\ell^N \psi' J' M')$

for the components of the ground and excited levels respectively, and sum over M and M'. As is to be expected, all quantum numbers and suffixes that depend on a fixed direction in space disappear, and we obtain

$$P = \sum_{\text{even } \lambda} T_{\lambda} \nu(\ell^N \psi_J \parallel U^{(\lambda)} \parallel \ell^N \psi'_{J'})^2, \quad (16)$$

where

$$T_{\lambda} = \lambda[8\pi^2 m/3h] (2\lambda+1) \sum_t (2t+1) B_t \Xi^2(t, \lambda)/(2J+1), \quad (17)$$

and

$$B_t = \sum_p |A_{tp}|^2 / (2t+1)^2. \quad (18)$$

#### V. OTHER CONTRIBUTIONS TO P

Before using Eq. (16) to make a direct comparison between experiment and theory, it is convenient to discuss briefly some effects that have so far been ignored. In the first place, no closed shells have been disturbed in the construction of the perturbing configurations; but it is clear that for Nd IV  $4f^2$ , for example, the tensors  $\underline{D}^{(k)}$  can couple the ground configuration  $4f^3$  to the configurations  $3d^9 4f^4$  and  $4d^9 4f^4$ , as well as to configurations such as  $4f^2 5d$  or  $4f^2 5g$ . However, owing to the symmetry about the double closed shell

$$(n'' \ell'')^{4\ell''+2} (n \ell)^{4\ell+2},$$

all matrix elements of the type

$$((n'' \ell'')^{4\ell''+2} (n \ell)^N \psi_{J M} \parallel D_q^{(k)} \parallel (n'' \ell'')^{4\ell''+1} (n \ell)^{N+1} \psi''_{J'' M''})$$

can differ from the corresponding quantities

$$((n\ell)^{4\ell+2-N} \psi_{J M} | D_q^{(k)} | (n\ell)^{4\ell+1-N} (n''\ell'') \psi''_{J'' M''})$$

by a phase factor at most. In view of the relation

$$\begin{aligned} |(\ell^{4\ell+2-N} \psi_{J} || U^{(\lambda)} || \ell^{4\ell+2-N} \psi'_{J'})|^2 \\ = |(\ell^N \psi_{J} || U^{(\lambda)} || \ell^N \psi'_{J'})|^2, \end{aligned}$$

Eq. (16) remains valid; but the sum over  $n'$  and  $\ell'$  of Eq. (14), which determines  $T_\lambda$  in virtue of Eq. (17), has to be augmented by those quantum numbers  $n''$  and  $\ell''$  corresponding to electrons in closed shells in the ground state of the ion. The large energy  $\Delta(n''\ell'')$  required to remove an electron from a closed shell, together with the expectation that the radial integrals

$$(n \ell | r^k | n''\ell'')$$

for  $k > 0$  are small, leads us to anticipate that the required modifications to the coefficients  $T_\lambda$  are insignificant. It is interesting to observe that if we assume that all configurations of both types

$$(n\ell)^{N-1} (n'\ell')$$

and

$$(n''\ell'')^{4\ell''+1} (n\ell)^{N+1}$$

coalesce into a single highly degenerate level, the objection in Sec. III to extending the closure procedure to all quantum numbers disappears.

There exists a second and intrinsically more interesting mechanism that can contribute to the intensities of rare-earth ions in solution. So far, the electric field acting on an ion has been considered to be completely static. As mentioned in Sec. I, however, lines exist in the spectra of rare-earth crystals that correspond to the excitation of vibrational quanta. If the immediate surroundings of a rare-earth ion in solution form a stable complex, as seems likely, vibrational modes may exist, the excitation of which could contribute to the intensities of the broad absorption lines. To examine this idea in more detail, we follow Griffith and denote the normal co-ordinates of the vibrating complex by  $Q_i$ .<sup>3</sup> Further, let  $\eta$  stand for the totality of the vibrational quantum numbers. For our purposes, the basic eigenfunctions of the system are taken to be simple products of harmonic oscillator eigenfunctions with the electronic eigenfunctions of the rare-earth ion. If we suppose the parameters  $A_{tp}$  of Sec. II correspond to some equilibrium arrangement of the complex, then allowance for small vibrations can be made by replacing  $V$  by

$$V' = \sum_{t,p} \left[ A_{tp} + \sum_i \frac{\partial A_{tp}}{\partial Q_i} Q_i \right] D_p^{(t)}.$$

For  $\eta \neq \eta'$ , it is a simple matter to obtain the equation

$$(B, \eta | D_q^{(1)} | B', \eta') = \sum_i (\eta | Q_i | \eta') \frac{\partial}{\partial Q_i} (B | D_q^{(1)} | B').$$

At a given temperature there is a certain probability that the vibrating complex is in the state defined by the set of quantum numbers  $\eta$ . If we denote this probability by  $\rho(\eta)$ , then the assumption made in Sec. IV regarding the population of the purely electronic components of the ground level leads in this case to a contribution  $P''$  to  $P$  given by

$$P'' = \sum_{\text{even } \lambda} T_{\lambda}' \nu(\ell^N \psi_J \parallel U^{(\lambda)} \parallel \ell^N \psi_{J'})^2, \quad (19)$$

where

$$T_{\lambda}' = \chi [8\pi^2 m/3h](2\lambda+1) \\ \times \sum_t (2t+1) B_t' E^2(t, \lambda)/(2J+1),$$

and

$$B_t' = \sum_{p, i, \eta, \eta'} \left| \frac{\partial A_{tp}}{\partial Q_i} \right|^2 |(\eta | Q_i | \eta')|^2 \rho(\eta)/(2t+1)^2.$$

The importance for us of these results lies in the fact that Eq. (19) is of precisely the same form as Eq. (16). If, then, the  $T_{\lambda}$  are treated as parameters to be adjusted to fit the experimental data, a good fit is no guarantee that the lines are purely electronic in origin. Indeed, if the surroundings of a rare-earth ion in solution are such that all  $A_{tp}$  for odd  $t$  vanish, then  $P''$  gives the sole contribution to  $P$ .

## VI. COMPARISON WITH EXPERIMENT

The absorption data for aqueous solutions of rare-earth ions are too extensive to be analyzed completely within a reasonable length of time. It was therefore decided to limit the investigations to  $\text{Nd}^{3+}$  and  $\text{Er}^{3+}$ , corresponding to three 4f electrons and three holes in a complete 4f shell, respectively. Both ions exhibit a sufficiently complex absorption spectrum to provide a good test of the theory. Moreover, Wybourne has recently fitted the energies of the levels of these ions to a detailed and reasonably complete theory,<sup>14</sup> thereby providing extremely accurate eigenfunctions for us to work with. The availability of tables of reduced matrix elements of the type

$$(f^3 \gamma S L \parallel U^{(\lambda)} \parallel f^3 \gamma' S L')$$

is an added incentive for choosing these particular ions.<sup>15</sup>

The procedure for calculating the reduced matrix elements of Eq. (16) runs as follows:

(i) Carry out expansions of the type

$$(\ell^N \psi J | = \sum_{\gamma, S, L} h(\gamma S L) (\ell^N \gamma S L J |.$$

The coefficients  $h$  of the RS-coupled states are given by Wybourne.<sup>14</sup>

(ii) Express every matrix element in Eq. (16) as a sum over reduced matrix elements involving RS coupled states.

(iii) Evaluate the new matrix elements by means of the formula

$$\begin{aligned} & (\ell^N \gamma S L J \parallel U^{(\lambda)} \parallel \ell^N \gamma' S' L' J') \\ &= \delta(S, S') (-1)^{S+L'+J+\lambda} [(2J+1)(2J'+1)]^{\frac{1}{2}} \\ & \times \left\{ \begin{array}{ccc} L & \lambda & L' \\ J' & S & J \end{array} \right\} (\ell^N \gamma S L \parallel U^{(\lambda)} \parallel \ell^N \gamma' S L'), \end{aligned} \quad (20)$$

which can easily be obtained from Eq. (7.1.8) of Edmonds.<sup>9</sup>

(iv) Use the tables of 6- $j$  symbols<sup>16</sup> and the tables of reduced matrix elements<sup>15</sup> to calculate the right-hand side of Eq. (20).

The results of the calculations are given in Table I for  $Nd^{3+}$  and in Table II for  $Er^{3+}$ . The levels are labelled by their principal components; the spectroscopic symbols are enclosed in square brackets to emphasize that the SL designations are not exact. All the levels listed in Tables I and II, with the exceptions of  ${}^4D_{7/2}$ ,  ${}^2L_{17/2}$ ,  ${}^2I_{13/2}$ ,  ${}^2L_{15/2}$ ,  ${}^4D_{1/2}$ ,  ${}^4D_{5/2}$ , and  ${}^2I_{11/2}$  of  $Nd^{3+}$ , have been identified with levels observed experimentally; the seven exceptions are included because their energies correspond closely to two broad bands measured by Hoogschagen.<sup>5</sup>



Since all reduced matrix elements of  $U_{mm}^{(\lambda)}$  for  $\lambda > 6$  vanish between f-electron states, the oscillator strengths  $P$  depend only on the three parameters  $T_2$ ,  $T_4$ , and  $T_6$ . It is a simple matter to take the experimental data for a given solution and choose the three parameters that give the best fit with experiment. This has been done for Hoogschagen's data for aqueous solutions of  $\text{NdCl}_3$  and  $\text{ErCl}_3$ , and also for the analogous data of Stewart on aqueous solutions of  $\text{Nd}(\text{ClO}_4)_3$ . A least-squares procedure is used in all three cases, although the variation of  $P$  over almost three orders of magnitude suggests that some other scheme might be more appropriate. The results are set out in Tables III, IV, and V. Hoogschagen found that for quite concentrated solutions (about 0.1 M), the oscillator strengths for the chloride and nitrate solutions of a given rare-earth ion vary only very slightly; the nitrate data are included in Tables III and IV, but additional fitting procedures have not been carried out for them. The excellence of the agreement can be taken in at a glance by referring to Figs. 1 and 2, where the experimental and theoretical data for the solutions of  $\text{NdCl}_3$  and  $\text{ErCl}_3$ , given in the second and third columns of Tables III and IV, are drawn out. For  $\text{Nd}^{3+}$ , even the feeble transitions  $^4I_{9/2} \rightarrow ^2P_{3/2}$ ,  $^4I_{9/2} \rightarrow ^2D_{5/2}$ , and  $^4I_{9/2} \rightarrow ^2P_{1/2}$  are well accounted for. Only the band in the region 29 600-31 250  $\text{cm}^{-1}$  is in significant disagreement with the theory, perhaps indicating that the assumed level assignments are incorrect.

In addition to the data given in Table III, Stewart has recorded the oscillator strengths of a number of weak lines in the ultraviolet range for solutions of neodymium perchlorate. These have not been included in the analysis, partly because of the difficulty of identifying the upper levels, and partly because these levels are quite close to the lower levels of  $4f^25d$ , thus vitiating the assumptions made in the derivation of Eq. (13).

VII. VARIATION OF  $T_2$ 

Owing to the selection rules

$$\Delta L, \Delta J \leq 2$$

on the matrix elements of  $U^{(2)}$ , the parameter  $T_2$  often plays only a minor roll in determining the oscillator strengths  $P$ . Bearing in mind that  $P$  depends on the squares of the reduced matrix elements, we see from Table II that only the transition  ${}^4I_{15/2} \rightarrow {}^2H_{11/2}$  in erbium salts is at all sensitive to  $T_2$ . Curiously enough, it is only this transition that exhibits an intensity difference between the chloride and the nitrate solutions (see Table IV).

If we now turn to the data for solutions of  $NdCl_3$  and  $Nd(NO_3)_3$ , we find very similar effects. The largest difference in intensity between the corresponding lines in the two solutions occurs for the transition to the two virtually coincident levels  ${}^4G_{5/2}$  and  ${}^2G_{7/2}$ ; and a glance at Table I reveals that of all the matrix elements

$$(f^3 [{}^4I_{9/2}] \parallel U^{(2)} \parallel f^3 [S'L'J']),$$

the ones for

$$[S'L'J'] \equiv [{}^4G_{5/2}]$$

and for

$$[S'L'J'] \equiv [{}^2G_{7/2}]$$

are the two largest in magnitude. Again, the second-largest difference occurs for transitions to the group of levels  ${}^2K_{13/2}$ ,  ${}^4G_{7/2}$ , and  ${}^4G_{9/2}$ ; and the matrix elements for which

$$[S'L'J'] \equiv [{}^4G_{7/2}]$$

is the third largest in magnitude. Lesser differences do not appear to be simply related to matrix elements of  $U_{\lambda}^{(2)}$ ; but in spite of this, the evidence is sufficiently strong to leave little doubt that of the three parameters,  $T_2$  is peculiarly sensitive to changes in the anion. Of course, Stewart's work with the perchlorate provides a third set of data to compare with the chloride and the nitrate; but it is felt that the differences between the fourth and seventh columns of Table III, and hence also between the first two rows of Table V, are to be ascribed mainly to differences in experimental technique rather than to any real change in the parameters  $T_{\lambda}$ .

#### VIII. ENVIRONMENT OF A RARE-EARTH ION IN SOLUTION

So far, the quantities  $T_{\lambda}$  have been treated purely as variable parameters, to be adjusted to fit experiment. To account for their values, we must construct a model for at least the immediate surroundings of a rare-earth ion. Unfortunately, little is known about the form such a model should take. That the nearest neighbors of a rare-earth ion occupy well defined positions is clear from the mere existence of fine structure in the spectra of solutions of europium salts.<sup>17</sup> The occurrence of an identical fine structure in aqueous solutions of europium chloride and very dilute europium nitrate indicates that in these two cases the nearest neighbors, and probably the next-nearest neighbors too, are water molecules. Examination of the lines corresponding to the transitions  ${}^7F_0 \rightarrow {}^5D_1$ ,  ${}^7F_0 \rightarrow {}^5D_2$ ,  ${}^5D_0 \rightarrow {}^7F_1$ , and  ${}^5D_0 \rightarrow {}^7F_2$  reveals that all the degeneracies of the levels involved are lifted; the point symmetry at a rare-earth ion must therefore be quite low. Several lines show a marked increase of intensity when alcohol is used in place of water as a solvent, and Sayre, Miller, and Freed regarded this as demonstrating that the complex comprising a rare-earth ion and its immediate surroundings possesses a center of inversion, in contrast to

the situation for alcoholic solvents.<sup>18</sup> Taken with the splittings of the levels, this interpretation limits the immediate point symmetry at a rare-earth ion in aqueous solution to  $D_{2h}$ . Miller subsequently proposed a model possessing this symmetry, using the criterion that the water molecules around the rare-earth ion should be arranged as in a fragment of a high-pressure ice.<sup>19</sup> He chose a configuration in which the eight water molecules nearest the ion lie at the vertices of two rectangles, whose planes are perpendicular and whose centers coincide with the nucleus of the rare-earth ion. This arrangement was consistent with the structure of ice III derived by McFarlan.<sup>20</sup>

A different configuration was proposed later by Brady, who carried out x-ray diffraction experiments on aqueous solutions of  $\text{ErCl}_3$ .<sup>21</sup> He found that six or possibly seven water molecules cluster around the rare earth ion, the distance between the erbium nucleus and the nuclei of the oxygen atoms being about  $2.3 \text{ \AA}$ . He interpreted the finer points of the diffraction pattern in terms of a model in which the rare-earth ion is at the center of an octahedron of water molecules. Two chlorine atoms are supposed to be on opposite sides of the octahedron such that their nuclei are coplanar with four oxygen nuclei and the erbium nucleus.

Objections can be raised to both Miller's and Brady's models. As Miller himself pointed out to the writer, a recent re-examination of the structure of ice III has shown that there are no fragments with  $D_{2h}$  symmetry;<sup>22</sup> also, there appears to be no position where a rare-earth ion can be placed interstitially and have six or seven oxygen atoms as close as the diffraction data demand. Until the structures of denser forms of ice become known, no further progress along the lines suggested by Miller seems possible.

Turning now to Brady's model, we note first that the superposition of an axial and an octahedral field splits a level for which  $J=1$  into only two components, in disagreement with experiment. Secondly, since the splittings

of levels are largely determined by the nearest neighbors of the rare-earth ion, and since no octahedral field leaves all levels for which  $J=1$  degenerate, we should expect the splitting of such a level to be extremely small. However, the splittings of  ${}^7F_1$  and  ${}^5D_1$  in aqueous solutions of  $\text{EuCl}_3$  are as large as those of the corresponding levels of  $\text{EuCl}_3 \cdot 6\text{H}_2\text{O}$ , a crystal where the immediate point symmetry at a europium ion is as low as  $C_2$ .

In the absence of a satisfactory model for the surroundings of a rare-earth ion, we must modify the project of calculating accurate values of the parameters  $T_\lambda$ . A feature shared by both Brady's and Miller's models is the presence of a center of inversion: the occurrence of any transitions at all must be ascribed either to vibrational effects, represented by Eq. (19), or else to the absence of a center of inversion in the outer hydration layers of the rare-earth ion. No vibrational structure of the kind typical of crystals has been seen in aqueous solutions of  $\text{EuCl}_3$ ,<sup>18</sup> though the excitation of very low-frequency oscillations, corresponding to the rare-earth ion and its immediate surroundings moving as a unit, would not be distinguishable from the main electronic lines. Even accepting this possibility, we should expect a calculation of the parameters  $T_\lambda$  on the basis of a model that did not possess a center of inversion to give values far in excess of experiment if the first hydration layer actually possessed such a center. We can, then, at least test to see whether the observed values of  $T_\lambda$  given in Table V are consistent with the hypothesis of Sayre et al. that a center of inversion exists. From the many models of the rare-earth ion and its surroundings that we might construct, it seems proper to choose one that reproduces, approximately at any rate, the observed splittings of the levels  ${}^5D_1$ ,  ${}^5D_2$ ,  ${}^7F_1$  and  ${}^7F_2$ . The striking similarity of these splittings to those of the corresponding levels of  $\text{Eu}^{3+}$  in  $\text{EuCl}_3 \cdot 6\text{H}_2\text{O}$  has already been remarked. The detailed analysis of the isomorphic crystal  $\text{GdCl}_3 \cdot 6\text{H}_2\text{O}$  indicates that the europium ion is surrounded by

six water molecules and two chlorine ions.<sup>23</sup> If we remove the latter without disturbing the former, the crystal splittings of the levels should not be too greatly affected, and the resulting complex  $\text{Eu}(\text{OH}_2)_6^{3+}$  does not possess a center of inversion. Moreover, the number of water molecules is consistent with the x-ray diffraction data. We should stress at this point that it is not suggested that the actual configuration of water molecules in aqueous solutions is the same as that in crystals of  $\text{EuCl}_3 \cdot 6\text{H}_2\text{O}$ ; we are merely constructing a model that should reproduce, within, say an order of magnitude, values of the parameters  $T_\lambda$  characteristic of a configuration of water molecules that does not possess a center of inversion.

#### IX. CALCULATION OF THE PARAMETERS $T_\lambda$

The configuration of water molecules surrounding a rare-earth ion influences the parameters  $T_\lambda$  through the quantities  $B_t$ , defined in Eq. (18). For a configuration of charges  $q_i$  at coordinates  $(R_i, \theta_i, \phi_i)$ , the crystal field parameters  $A_{tp}$  are given by

$$A_{tp} = (-1)^{p+1} \sum_i q_i R_i^{-t-1} C_{-p}^{(t)}(\theta_i, \phi_i), \quad (21)$$

provided it is assumed that the electrons on the rare-earth ion spend a negligible time at radial distances greater than the smallest  $R_i$ . If each charge  $q_i$  is replaced by a dipole of strength  $\mu$ , directed in the same sense towards the origin, and lying a distance  $R$  from it, the substitution

$$q_i R_i^{-t-1} \rightarrow \mu e (t+1) R^{-t-2}$$

should be made in Eq. (21). If we use the spherical harmonic addition theorem

(for example, see Edmonds<sup>9</sup>), we find

$$B_t = [\mu e(t+1)/(2t+1)R^{t+2}]^2 \sum_{i,j} P_t(\cos \omega_{ij}), \quad (22)$$

where  $\omega_{ij}$  denotes the angle between the radial vectors leading to dipoles  $i$  and  $j$ . Terms for which  $i=j$  in the sum must be included.

For the proposed model of the rare-earth complex, the angles  $\omega_{ij}$  can be easily calculated from the known positions of the oxygen atoms in  $\text{GdCl}_3 \cdot 6\text{H}_2\text{O}$ . In doing this, it is to be noted that Marezio et al. use an oblique coordinate scheme.<sup>23</sup> The only values of  $t$  of interest to us are 1, 3, 5, and 7; for all other values, the  $3-j$  symbols in Eq. (14) vanish on putting  $l=3$ . We find that

$$\sum_{i,j} P_t(\cos \omega_{ij})$$

assumes the values 1.671, 14.44, and 1.726 for  $t=3, 5,$  and  $7$  respectively. The analogous sum for  $t=1$  is also non-zero, implying that there is a finite electric field at the origin, and hence that the rare-earth ion is not in a position of equilibrium. This blemish in our model is not serious, however. Our aim is solely to obtain approximate values for the quantities  $B_t$ , and not to construct a perfectly self-consistent model. In any case, we could easily take advantage of the much lower power of  $1/R$  associated with  $P_1$  in Eq. (22) to pass the responsibility of ensuring that  $B_1$  is zero to the second hydration layer of a more elaborate model, while only slightly affecting  $B_3, B_5,$  and  $B_7$ .

The next step is to estimate the dipole moment  $\mu$ . For  $\text{Gd}(\text{OH}_2)_6^{3+}$ , the average distance  $R'$  of the six oxygen nuclei from the nucleus of the ion  $\text{Gd}^{3+}$  is  $2.412 \text{ \AA}$ . Accepting the atomic radii given by Templeton and Dauben,<sup>24</sup> we find this distance should be increased to  $2.469 \text{ \AA}$  for  $\text{Nd}^{3+}$ , and decreased to  $2.355 \text{ \AA}$  for  $\text{Er}^{3+}$ . These distances are not the distances  $R$  to the centers of the water dipoles, of course. If we make the simplifying assumption that the

negative charge of a water molecule in the complex coincides with the oxygen nucleus, then with a knowledge of the polarizability  $\alpha$  and the ordinary dipole moment  $\mu'$  of the water molecule, we can calculate  $\mu$  by solving the equations

$$\mu = \mu' + 3|e|\alpha/R^2,$$

$$\mu = 4|e|x,$$

and

$$R = R' + x.$$

Taking  $\mu' = 1.85 \times 10^{-10}$  e.s.u., and  $\alpha = 1.48 \times 10^{-24}$ ,<sup>25</sup> we obtain

$$\mu = 4.72 \times 10^{-18} \text{ e.s.u.},$$

and

$$R = 2.716 \text{ \AA}$$

for  $\text{Nd}^{3+}$ , and

$$\mu = 4.95 \times 10^{-18} \text{ e.s.u.},$$

and

$$R = 2.615 \text{ \AA}$$

for  $\text{Er}^{3+}$ . The calculation of  $B_3$ ,  $B_5$ , and  $B_7$  may now be readily completed.

The remaining factor in Eq. (17) involves  $E(t,\lambda)$ . As can be seen from a glance at Eq. (14), the calculation of quantities of this type entails the estimation of some radial integrals and energy denominators. It is to be expected that the term in the sum of Eq. (14) for which  $l' = 2$  and  $n' = 5$  predominates; partly because  $\Delta(5d)$  is the smallest of all energy denominators, as can be seen from Fig. 5 of Dieke et al.,<sup>10</sup> and partly because the smaller degree of overlap between a 4f eigenfunction and other orbitals of the type  $n'd$  for which  $n' \geq 6$  should result in greatly reduced radial integrals. Let us therefore concentrate on excited configurations of the type  $4f^{N-1} 5d$  and, for the moment, neglect all others.



Dieke et al. have found that for  $Ce^{3+}$  the configuration  $5d$  lies about  $50000\text{ cm}^{-1}$  above  $4f$ , whereas for  $Yb^{3+}$ ,  $4f^{12}5d$  lies about  $100000\text{ cm}^{-1}$  above  $4f^{13}$ . A linear interpolation gives the values  $58000\text{ cm}^{-1}$  and  $92000\text{ cm}^{-1}$  for  $Nd^{3+}$  and  $Er^{3+}$  respectively. It seems reasonable to take one of these values, appropriate to the ion under investigation, for the denominator  $\Delta(5d)$ .

The radial integrals would be difficult to estimate were it not for the work of Rajnak, who has recently calculated  $5d$  eigenfunctions for  $Pr^{3+}$  and  $Tm^{3+}$ .<sup>26</sup> She made the assumption that the central field  $5d$  electron moves in is the same as that for a  $4f$  electron; the latter can be obtained from a self-consistent calculation carried out by Ridley for the ground states of these two ions.<sup>27</sup> The radial integrals and their interpolated values are set out in the first three rows of Table VI. Strictly, the use of Eq. (10) to calculate all the radial integrals is not consistent with the assumption  $r < R$ ; but the errors introduced in doing so are sufficiently small to be neglected.

It is now a straightforward matter to collect the various parts of the calculation together. The values of  $T_\lambda$  thus obtained are given in the third column of Table VII. Some of the entries of Table V are included so that a direct comparison between experiment and theory can be made. It is immediately seen that  $T_4$  and  $T_6$  for  $Nd^{3+}$  agree to within a factor of 3, a result that must be regarded as satisfactory. However,  $T_2$  for  $Nd^{3+}$  and all three parameters for  $Er^{3+}$  are too small by an order of magnitude. It might be thought that the neglect of configurations of the type  $4f^{N-1}n'd$ , where  $n' \geq 6$ , is largely responsible for the discrepancies. However, this is unlikely.

In the first place, it can be seen from Table VI that the products

$$(4f|r^t|5d)(4f|r|5d)$$

are almost equal to the corresponding quantities

$$(4f|r^{t+1}|4f).$$

Hence, even if we used Eq. (11) to perform a closure over all configurations of the type  $4f^{N-1}n'd$ ,  $3d^9 4f^{N+1}$ , and  $4d^9 4f^{N+1}$  (for a fixed  $N$ ), placing them as low as  $4f^{N-1}5d$  in energy, our results would differ insignificantly from those already obtained.

The irrelevance of higher configurations of the type  $4f^{N-1}n'd$  can be seen in another way. If  $l' = 2$ , then  $T_2$ ,  $T_4$ , and  $T_6$  depend on  $B_3$  and  $B_5$  only. The linear relationship between the parameters  $T_\lambda$  that this implies is

$$T_2/8 = T_4/55 - T_6/3575.$$

Since we have  $T_\lambda > 0$ , the inequality  $T_2/T_4 < 8/55$  follows. The observed ratios are much larger than  $8/55$ , and hence they cannot be accounted for by including the effects of  $4f^{N-1}6d$ ,  $4f^{N-1}7d$ ,  $4d^{N-1}7d$ ,  $4d^9 4f^{N+1}$ , etc.

Configurations of the type  $4f^{N-1}n'g$  remain to be considered. Their comparative proximity to the ionizing limit suggests the validity of a closure procedure over all  $n'$ ; but the large radial extension of the  $g$  eigenfunctions makes it difficult to decide where the ionizing limit is for an ion in solution. For the free ion  $\text{Pr}^{3+}$ , even the nodeless eigenfunctions  $\mathcal{R}(5g)$  attains its maximum value as far as  $3.3 \text{ \AA}$  from the nucleus; it follows that the eigenfunctions for ions in solution are determined by conditions beyond the first hydration layer. If the six dipoles of this layer are replaced by an equivalent uniform dipole shell, the classical electrostatic potential difference between points inside and just outside is  $6\mu/R^2$ , which is equivalent to approximately  $100\,000 \text{ cm}^{-1}$ . Presumably, the ionizing limit of a free rare-earth ion should be reduced by at least this amount for an ion in solution. Interpolating

between Ridley's energies  $\epsilon$  for  $\text{Pr}^{3+}$  and  $\text{Tm}^{3+}$ ,<sup>27</sup> and assuming the energies of the configurations  $4f^{N-1} n'g$  coincide at the corrected ionizing limit, we find  $\Delta(n'g)$  to be  $167\,000\text{ cm}^{-1}$  for  $\text{Nd}^{3+}$  and  $207\,000\text{ cm}^{-1}$  for  $\text{Er}^{3+}$ . Terms in Eq. (17) that involve  $B_7$  are negligible, and the new values of  $E(t,\lambda)$ , for which  $(n'l')$  runs over (5d) and all pairs of the type  $(n'g)$ , can be obtained from the old by multiplication by

$$1 + \frac{(t+7)!(6-t)!}{(\lambda+7)!(6-\lambda)!} \frac{(4f|r^{t+1}|4f)}{(4f|r|5d)(4f|r^t|5d)} \frac{\Delta(5d)}{\Delta(n'g)}.$$

The results of the calculation are given in the fourth column of Table VII.

The parameters  $T_\lambda$  are in all cases increased, but they are still too small by factors of 2 and 8 for  $\text{Nd}^{3+}$  and  $\text{Er}^{3+}$  respectively.

## X. DISCUSSION

In searching for the causes of the discrepancies between experiment and theory, we should not lose sight of the fact that, for  $\text{Nd}^{3+}$ , the agreement is perhaps better than we might reasonably have anticipated. Owing to the dependence of  $T_\lambda$  on  $R^{-10}$  or  $R^{-14}$ , the discrepancies for  $\text{Nd}^{3+}$  could be easily accounted for by decreasing  $R$  by as little as  $0.1\text{ \AA}$ . The tendency of the negative charge on the oxygen atoms to be drawn towards the  $3+$  rare-earth ion might easily be large enough to require a correction to  $R$  of this order of magnitude, though other reasons can be easily thought up. For example, the protons of the water molecules presumably take part in bonding to the second hydration layer, and their positions are not solely determined by the charge on the rare-earth ion, as we have assumed.

Increasing  $T_\lambda$  by a factor of 2 would still leave a discrepancy of a factor of 4 for  $\text{Er}^{3+}$ . Now, throughout the entire analysis, it has been assumed that the configuration of water molecules surrounding  $\text{Er}^{3+}$  is essentially the

same as that surrounding  $\text{Nd}^{3+}$ , save for a radial scaling factor. However, the change in ionic radii is sufficiently large for many crystals containing  $\text{Nd}^{3+}$  or  $\text{Er}^{3+}$ , for example,  $\text{NdCl}_3$  and  $\text{ErCl}_3$ , to exist in dissimilar forms. If this were the case here, then we should expect the calculated reduction of the parameters  $T_\lambda$  to reproduce the observed reduction to within an order of magnitude only. Other causes may be the source of the discrepancy, of course: the shapes of  $\mathcal{R}(4f)$  and  $\mathcal{R}(5d)$  are changing with atomic number much more rapidly in the region of  $\text{Pr}^{3+}$  than near  $\text{Tm}^{3+}$ , and the linear interpolation method used to derive the third and fourth columns of Table VI may be unreliable. Again, the use of free-ion eigenfunctions as modestly extended as  $\mathcal{R}(5d)$  might constitute too gross an approximation for ions in solution; corrections as small as 10% to the entries of Table VI, if applied in the most telling directions, reduce the calculated drop in the parameters  $T_\lambda$  in going from  $\text{Nd}^{3+}$  to  $\text{Er}^{3+}$  by a factor of more than 2. It seems unlikely that lines involving the excitation of vibrational modes are responsible for the discrepancy in the relative intensities of lines of  $\text{Nd}^{3+}$  and  $\text{Er}^{3+}$ ; for, even if they are quite strong, their intensities depend on the radial integrals of Table VI, and should therefore decrease in step with the electronic lines.

The fact that the calculated parameters  $T_\lambda$  of Table VII are smaller than the experimental ones is, of itself, an important result. For, as indicated in Sec. VII, we would have expected the calculations to grossly overestimate the  $T_\lambda$  if the system comprising the rare-earth ion and the first hydration layer possessed a center of inversion. We can therefore conclude that this system does not possess a center of inversion. The deduction of Sayre et al. that the opposite is true was based on the feebleness of certain lines in the spectrum of aqueous solutions of  $\text{EuCl}_3$  compared to alcoholic solutions.<sup>18</sup> One of these lines, corresponding to the transition  $7F_0 \rightarrow 5D_2$ ,

has been measured by Hoogschagen, and has an oscillator strength of 0.009.<sup>5</sup> Small as this number is, it is consistent with the absence of a center of inversion. To demonstrate this, we use perturbation theory and calculate the appropriate matrix elements of  $\underline{U}^{(\lambda)}$  by means of the formula

$$\begin{aligned} & ([^7F_0] \parallel U^{(\lambda)} \parallel [^5D_2]) \\ &= ({}^7F_0 | \Lambda | ^5D_0) ({}^5D_0 \parallel U^{(\lambda)} \parallel ^5D_2) [E({}^7F_0) - E({}^5D_0)]^{-1} \\ &+ ({}^7F_0 \parallel U^{(\lambda)} \parallel {}^7F_2) ({}^7F_2 | \Lambda | ^5D_2) [E({}^7F_2) - E({}^5D_2)]^{-1}, \end{aligned}$$

where  $\Lambda$  is the spin-orbit interaction. The matrix elements can be evaluated by the methods of Elliott et al.<sup>28</sup> The oscillator strength  $P$  depends solely on  $T_2$ , which can be obtained by linearly interpolating the observed values for aqueous solutions of  $NdCl_3$  and  $ErCl_3$ . The result,  $P = 0.006$ , is in satisfactory agreement with experiment.

We are now left with the problem of explaining why certain lines in the spectrum of alcoholic solutions of  $EuCl_3$  are anomalously strong. We shall not explore this problem in detail here: it is worth noting, however, that the transitions that are much more intense in alcoholic solvents, such as  ${}^7F_0 \rightarrow {}^5D_2$  of  $Eu^{3+}$ , and what appear to be transitions of the type  ${}^8S_{7/2} \rightarrow {}^6D_J$  of  $Gd^{3+}$ ,<sup>29</sup> depend solely on  $T_2$ . On the other hand, the transitions that do not undergo striking changes of intensity, such as  ${}^3H_4 \rightarrow {}^3P_J$  of  $Pr^{3+}$ ,<sup>29</sup> depend on  $T_4$  and  $T_6$  only, at least in the lowest order of perturbation theory. Since both  $T_2$  and  $T_4$  depend strongly on  $B_3$ , an apparent increase of  $T_2$  in alcoholic solutions without a corresponding increase of  $T_4$  could occur only through the excitation of vibrational modes, which might be undetected as such if sufficiently low in

frequency. Only  $T_2'$  of Eq. (19) depends on  $B_1'$ ; hence, in fitting experiment to the parameters  $T_2$ ,  $T_4$  and  $T_6$ , a large value of  $B_1'$  would make itself felt through  $T_2$  alone. It is possible that the peculiar variation of  $T_2$  described in Sec. VI has its origin in a mechanism of this sort; but, from what is known of the relative intensities of electronic lines and their accompanying vibrational structures, it seems hard to account for a change of  $T_2$  by an order of magnitude in this way. Quantitative measurements on alcoholic solutions of  $\text{ErCl}_3$  or  $\text{NdCl}_3$  would decide whether it is correct to interpret the anomalously strong lines in alcoholic solutions of  $\text{EuCl}_3$  in terms of an increase of the parameter  $T_2'$ , or whether some other mechanism should be sought.

## XI. CONCLUSION

Although the theory of Secs. II and III is applicable to a rare-earth ion in a crystalline environment, the absence of experimental data on the oscillator strengths of lines in rare-earth crystals has obliged us to discuss the theory in terms of solutions of rare-earth ions. The difficulty of distinguishing between the pure electronic parts of the line intensities from contributions coming from transitions in which vibrational modes are simultaneously excited is not present for the spectra of rare earth crystals, or at least for those spectra that have been analyzed. Data for crystals, when available, will therefore permit more rigorous tests of the theory to be carried out.

ACKNOWLEDGMENTS

I wish to thank Mrs. K. Rajnak for allowing me to make use of her eigenfunctions for  $\text{Pr}^{3+}$  and  $\text{Tm}^{3+}$  prior to their publication. Several conversations with Dr. D. G. Miller and Professor B. B. Cunningham proved helpful.

This work was performed under the auspices of the U. S. Atomic Energy Commission.

Table I. Reduced matrix elements of  $U^{(\lambda)}$  for  $\text{Nd}^{3+}$ 

S'L'J'	Calculated Energy (in $\text{cm}^{-1}$ above ${}^4\text{I}_{9/2}$ ) <sup>a</sup>	$(f^3[{}^4\text{I}_{9/2}] \parallel U^{(\lambda)} \parallel f^3 [S' L' J'])$		
		$\lambda = 2$	$\lambda = 4$	$\lambda = 6$
${}^4\text{D}_{7/2}$	31004	0.0004	0.0607	-0.0893
${}^2\text{L}_{17/2}$	30932	0	-0.0318	-0.0358
${}^2\text{I}_{13/2}$	30073	0.0123	0.0370	-0.0424
${}^2\text{L}_{15/2}$	29413	0	0.1612	0.1006
${}^4\text{D}_{1/2}$	29276	0	-0.5093	0
${}^4\text{D}_{5/2}$	28836	0.0111	-0.2390	0.1659
${}^2\text{I}_{11/2}$	28694	-0.0717	-0.1238	0.0594
${}^4\text{D}_{3/2}$	28641	0	0.4417	-0.1299
${}^2\text{P}_{3/2}$	26348	0	0.0345	-0.0271
${}^2\text{D}_{5/2}$	23880	-0.0002	0.0116	0.0479
${}^2\text{P}_{1/2}$	23147	0	-0.1884	0
${}^4\text{G}_{11/2}$	21826	-0.0023	-0.0741	0.0906
${}^2\text{G}_{9/2}$	21255	-0.0358	0.1381	-0.1318
${}^2\text{D}_{3/2}$	21247	0	-0.1367	0.0121
${}^2\text{K}_{15/2}$	21027	0	-0.0747	-0.1228
${}^4\text{G}_{9/2}$	19720	-0.0662	0.2388	-0.1926
${}^4\text{G}_{7/2}$	19320	0.2529	-0.4257	0.2488
${}^2\text{K}_{13/2}$	18978	-0.0846	0.0157	0.1810
${}^4\text{G}_{5/2}$	17356	-0.9471	0.6399	-0.1885
${}^2\text{G}_{7/2}$	17354	-0.2580	0.4009	-0.1539
${}^2\text{H}_{11/2}$	15985	-0.0073	0.0515	-0.1020
${}^4\text{F}_{9/2}$	14903	0.0275	-0.0936	-0.2114



Table I. (Cont)

S'L'J'	Calculated Energy (in $\text{cm}^{-1}$ above ${}^4\text{I}_{9/2}$ ) <sup>a</sup>	$(f^3[{}^4\text{I}_{9/2}] \parallel U^{(\lambda)} \parallel f^3[S' L' J'])$		
		$\lambda = 2$	$\lambda = 4$	$\lambda = 6$
${}^4\text{F}_{7/2}$	13611	0.0337	0.2034	0.6525
${}^4\text{S}_{3/2}$	13454	0	0.0549	0.4862
${}^2\text{H}_{9/2}$	12612	0.0986	-0.0914	0.3392
${}^4\text{F}_{5/2}$	12607	0.0303	-0.4862	-0.6299
${}^4\text{F}_{3/2}$	11524	0	0.4778	0.2317

<sup>a</sup>See reference 14.

Table II. Reduced matrix elements of  $U^{(\lambda)}$  for  $\text{Er}^{3+}$ 

S'L'J'	Calculated Energy (in $\text{cm}^{-1}$ above ${}^4\text{I}_{15/2}$ ) <sup>a</sup>	$(f^{11} [{}^4\text{I}_{15/2}] \  U^{(\lambda)} \  f^{11} [S' L' J'])$		
		$\lambda = 2$	$\lambda = 4$	$\lambda = 6$
${}^2\text{G}_{9/2}$	24893	0	0.1314	0.4809
${}^4\text{F}_{3/2}$	22701	0	0	-0.3658
${}^4\text{F}_{5/2}$	22321	0	0	-0.4712
${}^4\text{F}_{7/2}$	20717	0	-0.3816	-0.7909
${}^2\text{H}_{11/2}$	19407	0.8456	-0.6420	0.3127
${}^4\text{S}_{3/2}$	18525	0	0	0.4624
${}^4\text{F}_{9/2}$	15449	0	-0.7107	-0.6717
${}^4\text{I}_{9/2}$	12496	0	-0.4567	-0.1302
${}^4\text{I}_{11/2}$	10415	0.1856	-0.0290	-0.6259

<sup>a</sup> See reference 14.

Table III. Oscillator Strengths for  $\text{Nd}^{3+}$ 

Spectral Region ( $\text{cm}^{-1}$ )	Upper Levels involved in transition	$P \times 10^6$				
		$\text{NaCl}_3$		$\text{Na}(\text{NO}_3)_3$	$\text{Na}(\text{ClO}_4)_3$	
		Theory	Expt <sup>a</sup>	Expt <sup>a</sup>	Theory	Expt <sup>b</sup>
29600 - 31250	${}^4\text{D}_{7/2}, {}^2\text{I}_{13/2}, {}^2\text{L}_{17/2}$	0.46	2.36		0.40	1.7
26750 - 29600	$\left[ \begin{array}{l} {}^4\text{D}_{3/2}, {}^2\text{I}_{11/2}, {}^4\text{D}_{5/2} \\ {}^4\text{D}_{1/2}, {}^2\text{L}_{15/2} \end{array} \right]$	10.13	9.52		10.16	9.8
25750 - 26750	${}^2\text{P}_{3/2}$	0.04	0.05	0.05	0.03	0.02
24250 - 25750	c		0.03	0.03		
23500 - 24250	${}^2\text{D}_{5/2}$	0.06	0.08	0.08	0.05	0.06
22750 - 23500	${}^2\text{P}_{1/2}$	0.43	0.38	0.38	0.45	0.30
20250 - 22750	$\left[ \begin{array}{l} {}^2\text{K}_{15/2}, {}^2\text{D}_{3/2}, \\ {}^2\text{G}_{9/2}, {}^4\text{G}_{11/2} \end{array} \right]$	1.47	2.31	2.35	1.32	1.9
18250 - 20250	${}^2\text{K}_{13/2}, {}^4\text{G}_{7/2}, {}^4\text{G}_{9/2}$	5.48	6.58	6.78	4.92	5.8
16250 - 18250	${}^2\text{G}_{7/2}, {}^4\text{G}_{5/2}$	10.6	10.5	11.7	8.38	8.3
15250 - 16250	${}^2\text{H}_{11/2}$	0.20	0.39	0.39	0.17	0.14
14250 - 15250	${}^4\text{F}_{9/2}$	0.78	0.83	0.83	0.65	0.51
13000 - 14250	${}^4\text{F}_{7/2}, {}^4\text{S}_{3/2}$	9.84	8.88	8.78	8.17	7.6

Table III (Cont)

Spectral Region ( $\text{cm}^{-1}$ )	Upper Levels involved in transition	$P \times 10^6$				
		$\text{NdCl}_3$		$\text{Nd}(\text{NO}_3)_3$	$\text{Nd}(\text{ClO}_4)_3$	
		Theory	Expt <sup>a</sup>	Expt <sup>a</sup>	Theory	Expt <sup>b</sup>
11900 - 13000	$^4\text{F}_{5/2}, ^2\text{H}_{9/2}$	8.48	9.22	9.17	7.32	7.7
11000 - 11900	$^4\text{F}_{3/2}$	2.02	3.02	2.93	1.96	2.3

<sup>a</sup> From Ref. 5

<sup>b</sup> From Ref. 6

<sup>c</sup> The line in this region reported by Hoogschagen has not been observed by Stewart, and no corresponding level occurs in the theoretical scheme. It is certainly spurious.

Table IV. Oscillator strengths for  $\text{Er}^{3+}$ 

Spectral Region ( $\text{cm}^{-1}$ )	Upper level involved in transition	$P \times 10^6$		
		$\text{ErCl}_3$		$\text{Er}(\text{NO}_3)_3$
		Theory	Expt <sup>a</sup>	Expt <sup>a</sup>
23900 - 25100	${}^2\text{G}_{9/2}$	0.89	0.74	0.74
21500 - 23100	$\left[ \begin{array}{l} {}^4\text{F}_{5/2} \\ {}^4\text{F}_{3/2} \end{array} \right]$	1.15	1.31	1.31
20000 - 21500	${}^4\text{F}_{7/2}$	2.34	2.22	2.22
18700 - 20000	${}^2\text{H}_{11/2}$	2.91	2.91	3.14
17500 - 18700	${}^4\text{S}_{3/2}$	0.57	0.83	0.83
14600 - 16400	${}^4\text{F}_{9/2}$	2.27	2.37	2.37
12000 - 12900	${}^4\text{I}_{9/2}$	0.47	0.34	0.34
9900 - 10400	${}^4\text{I}_{11/2}$	0.63	0.50	0.50

<sup>a</sup> From Ref. 5.

Table V. Observed values of the parameters  $T_\lambda$  (in units of  $10^{-21}$  sec.)

Solute	$T_2$	$T_4$	$T_6$
$\text{NdCl}_3$	8.7	17.3	35.3
$\text{Nd}(\text{ClO}_4)_3$	4.2	18.0	29.1
$\text{ErCl}_3$	3.2	5.6	4.8

Table VI. Radial integrals ( in atomic units)

Integral	$\text{Pr}^{3+}$	$\text{Nd}^{3+}$	$\text{Er}^{3+}$	$\text{Tm}^{3+}$
$(4f   r   5d)$	0.900	0.869	0.615	0.583
$(4f   r^3   5d)$	5.47	5.17	2.75	2.45
$(4f   r^5   5d)$	50.5	47.1	19.9	16.5
$(4f   r^2   4f)$	1.464	1.394	0.831	0.761
$(4f   r^4   4f)$	5.34	4.96	1.95	1.57
$(4f   r^6   4f)$	39.6	36.4	10.5	7.31
$(4f   r^8   4f)$	500	450	100	62

Table VII. Theoretical and selected experimental values of the parameters  $T_{\lambda}$  (in units of  $10^{-21}$  sec.).

Ion	Parameter	Calculated		Observed in Chloride Solutions (From Table V)
		$(n'l) \equiv (5d)$ only	$(n'l') \equiv (5d)$ , and all $(n'g)$	
$\text{Nd}^{3+}$	$T_2$	0.94	3.62	8.7
	$T_4$	6.73	9.96	17.3
	$T_6$	16.4	17.2	35.3
$\text{Er}^{3+}$	$T_2$	0.05	0.28	3.2
	$T_4$	0.38	0.59	5.6
	$T_6$	0.68	0.72	4.8

## REFERENCES


1. J. H. Van Vleck, *J. Phys. Chem.* 41, 67 (1937).
2. L. J. F. Broer, C. J. Gorter, and J. Hoogschagen, *Physica* 11, 231 (1945).
3. J. S. Griffith, *The Theory of Transition-Metal Ions* (Cambridge University Press, Cambridge, England, 1961).
4. A. Merz, *Ann Phys.* 28, 569 (1937); H. Ewald, *Z. Physik*, 110, 428 (1938).
5. J. Hoogschagen, *Thesis*, Leyden (1947). Most of the material in this work is summarized in: J. Hoogschagen and C. J. Gorter, *Physica* 14, 197 (1948).
6. D. C. Stewart, *Absorption Spectra of Lanthanide and Actinide Rare Earths, II Transition Probabilities for +3 Ions in the Two Series*, Argonne National Laboratory Report ANL-4812 1952 (unpublished).
7. J. C. Slater, *Quantum Theory of Atomic Structure* (McGraw-Hill Book Company, New York, 1960), Vol. I.
8. J. S. Griffith, *Mol. Phys.* 3, 477 (1960).
9. A. R. Edmonds, *Angular Momentum in Quantum Mechanics* (Princeton University Press, Princeton, New Jersey, 1957).
10. G. H. Dieke, H. M. Crosswhite, and B. Dunn, *J. Opt. Soc. Am.* 51, 820 (1961).
11. G. Racah, *Phys. Rev.* 63, 367 (1943).
12. R. A. Satten, *J. Chem. Phys.* 21, 637 (1953); H. Lammermann, *Z. Physik* 150, 551 (1958); E. H. Carlson and G. H. Dieke, *J. Chem. Phys.* 34, 1602 (1961).
13. B. R. Judd, *Proc. Roy. Soc. (London) A* 241, 414 (1957); M. J. D. Powell and R. Orbach, *Proc. Phys. Soc. (London)* 78, 753 (1961).
14. B. G. Wybourne, *J. Chem. Phys.* 32, 639 (1960); 34, 279 (1961).

Orthogonality checks brought to light a number of errors in the eigenfunctions tabulated in the second paper. In the  $J=1/2$  part of his Table II, 0.2426 should be -0.2426; 0.9760 should be 0.9701; and 0.2178




should be 0.2426. The entry +0.0076 in the first line of the  $J=7/2$  part of Table II should be -0.0076. No checks have been made in his Table III for levels other than those actually observed, but it is clear from inspection that the entries of the  $J=1/2$  part are inconsistent.

15. B. R. Judd, Proc. Roy. Soc. A 250, 562 (1959).
16. R. Bivins, N. Metropolis, M. Rotenberg, and J. K. Wooton, The 3-j and 6-j symbols (Technology Press, Massachusetts Institute of Technology, Cambridge, Massachusetts, 1959).
17. S. Freed, Rev. Mod. Phys. 14, 105 (1942).
18. E. V. Sayre, D. G. Miller, and S. Freed, J. Chem. Phys. 28, 109 (1957).
19. D. G. Miller, J. Am. Chem. Soc. 80, 3576 (1958).
20. R. L. McFarlan, J. Chem. Phys. 4, 253 (1936).
21. G. W. Brady, J. Chem. Phys. 33, 1079 (1960).
22. W. B. Kamb and S. K. Datta, Nature 187, 140 (1960).
23. M. Marezio, H. A. Plettinger, and W. H. Zachariasen, Acta. Cryst. 14, 234 (1961).
24. D. H. Templeton and C. H. Dauben, J. Am. Chem. Soc. 76, 5237 (1954).
25. D. Polder, Physica 9, 709 (1942).
26. K. Rajnak, private communication (December, 1961).
27. E. C. Ridley, Proc. Cambridge Phil. Soc. 56, 41 (1960).
28. J. P. Elliott, B. R. Judd, and W. A. Runciman, Proc. Roy. Soc. (London) A 240, 509 (1957).
29. D. G. Miller, E. V. Sayre, and S. Freed, J. Chem. Phys. 29, 454 (1958).



MU-25824

Fig. 1. A comparison between experimental and theoretical oscillator strengths of transitions in aqueous solutions of  $\text{NdCl}_3$ . The lengths of the horizontal lines running from the central vertical line give a measure of  $-\log P$ ; theoretical values are given on the left, experimental values on the right. The ordinate of a horizontal line gives the approximate energy of the corresponding transition, in  $\text{cm}^{-1}$ .



MU-25825

Fig. 2. A comparison between experimental and theoretical oscillator strengths of transitions in aqueous solutions of  $\text{ErCl}_3$ . The design of the figure is the same as that of Fig. 1.

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.