

Accepted Manuscript

Optical tweezers and their applications

Paolo Polimeno, Alessandro Magazzù, Maria Antonia Iati, Francesco Patti, Rosalba Saija, Cristian Degli Esposti Boschi, Maria Grazia Donato, Pietro G. Gucciardi, Philip H. Jones, Giovanni Volpe, Onofrio M. Maragò

PII: S0022-4073(18)30445-X
DOI: [10.1016/j.jqsrt.2018.07.013](https://doi.org/10.1016/j.jqsrt.2018.07.013)
Reference: JQSRT 6162



To appear in: *Journal of Quantitative Spectroscopy & Radiative Transfer*

Received date: 18 June 2018
Revised date: 13 July 2018
Accepted date: 16 July 2018

Please cite this article as: Paolo Polimeno, Alessandro Magazzù, Maria Antonia Iati, Francesco Patti, Rosalba Saija, Cristian Degli Esposti Boschi, Maria Grazia Donato, Pietro G. Gucciardi, Philip H. Jones, Giovanni Volpe, Onofrio M. Maragò, Optical tweezers and their applications, *Journal of Quantitative Spectroscopy & Radiative Transfer* (2018), doi: [10.1016/j.jqsrt.2018.07.013](https://doi.org/10.1016/j.jqsrt.2018.07.013)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Highlights

- We review optical tweezers and their applications
- T-matrix methods for calculations of optical forces
- Overview of typical optical tweezers calibration methods
- Applications in biology, nanotechnology, spectroscopy, stochastic thermodynamics, critical Casimir forces, and active matter

ACCEPTED MANUSCRIPT

Optical tweezers and their applications

Paolo Polimeno^{a,1}, Alessandro Magazzù^{b,1}, Maria Antonia Iati^{a,*}, Francesco Patti^{a,c}, Rosalba Saija^{a,c}, Cristian Degli Esposti Boschi^d, Maria Grazia Donato^a, Pietro G. Gucciardi^a, Philip H. Jones^e, Giovanni Volpe^b, Onofrio M. Maragò^{a,*}

^a*CNR-IPCF, Istituto per i Processi Chimico-Fisici, Viale F. Stagno d'Alcontres 37, I-98158 Messina, Italy*

^b*Göteborgs Universitet, Institutionen för Fysik, Origovägen 6 B, I-41296 Göteborg, Sweden*

^c*Università di Messina, Dipartimento di Scienze Matematiche e Informatiche, Scienze Fisiche e Scienze della Terra, Viale F. Stagno d'Alcontres, 31, I-98166 Messina, Italy*

^d*CNR-IMM sezione di Bologna, Istituto per la Microelettronica e Microsistemi, Via P. Gobetti 101, I-40129 Bologna, Italy*

^e*University College London, Department of Physics and Astronomy, Gower Street, London WC1E 6BT, UK*

Abstract

Optical tweezers, tools based on strongly focused light, enable optical trapping, manipulation, and characterisation of a wide range of microscopic and nanoscopic materials. In the limiting cases of spherical particles either much smaller or much larger than the trapping wavelength, the force in optical tweezers separates into a conservative gradient force, which is proportional to the light intensity gradient and responsible for trapping, and a non-conservative scattering force, which is proportional to the light intensity and is generally detrimental for trapping, but fundamental for optical manipulation and laser cooling. For non-spherical particles or at intermediate (meso)scales, the situation is more complex and this traditional identification of gradient and scattering force is more elusive. Moreover, shape and composition can have dramatic consequences for optically trapped particle dynamics. Here, after an introduction to the theory and practice of optical forces with a focus on the role of shape and composition, we give an overview of some recent applications to biology, nan-

*Corresponding authors

Email addresses: mariaantonia.iati@cnr.it (Maria Antonia Iati), onofrio.marago@cnr.it (Onofrio M. Maragò)

¹These authors contributed equally to this work.

otechnology, spectroscopy, stochastic thermodynamics, critical Casimir forces, and active matter.

Keywords: Optical trapping, optical manipulation, T-matrix, spectroscopic tweezers, critical Casimir forces, active matter

1. Introduction

Mechanical effects of light are known since Kepler's explanation of comets tail [1]. The advent of the laser has brought to novel concepts and applications of optical forces yielding a real scientific revolution [2, 3, 4]. *Optical tweezers* [5] (OT), in their simplest configuration, are instruments based on a tightly focused laser beam that is capable to trap and manipulate a wide range of particles in its focal spot [6]. Since their first demonstration in 1986 by Sir Arthur Ashkin [5], they have become a key technique for the trapping, manipulation, and characterisation of atoms [7], microscopic [8, 9] and nanoscopic particles [10], as well as biomolecules, viruses, bacteria and cells [11, 12, 13]. When used as a force transducer, optical tweezers are able to measure forces in the femtonewton range [6, 10]. The concept of photonic force microscopy has been also developed by scanning a trapped particle over surfaces in a liquid environment and sensing the force interaction between trapped probe and surface [14, 15].

Despite the tremendous progress in optical trapping techniques, the development of an accurate theoretical modeling for optical tweezers has been slower and often based on approximations [6]. In the limiting cases of spherical particles either much smaller (*dipole approximation*) [16] or much larger (*ray optics approximation*) [17] than the trapping wavelength, the force in optical tweezers separates into two different contributions: a conservative *gradient force*, proportional to the light intensity gradient, responsible for trapping, that can be related to an effective potential, and a non-conservative *scattering force*, that has a dissipative nature, cannot be related to an effective potential, is proportional to the light intensity and is generally detrimental for trapping, but fundamental for optical manipulation and laser cooling [6, 10]. However, for non-spherical

particles or at intermediate (meso)scales, the situation is more complex and this traditional identification of gradient and scattering force is more elusive [18]. Moreover, shape and composition can have dramatic consequences for the dynamics of optically trapped particles [10, 19, 20, 21].

30 Here, we first give a review on the theory of optical forces and optical trapping with a focus on modeling optical tweezers with *T-matrix* methods. Then we discuss some of the most common experimental implementation and methods for optical force calibration. Finally, we discuss a selection of recent applications in biology, nanotechnology, spectroscopy, critical Casimir forces, stochastic
35 thermodynamics, and active matter.

2. Optical forces and optical tweezers

Optical trapping of particles is a consequence of the radiation force that stems from the conservation of electromagnetic momentum upon scattering [6, 22]. Historically, the theoretical understanding of this physical process has
40 been investigated through the use of suitable approximations that depend on the size of the particle with respect to the light wavelength [4]. While calculations based on a full electromagnetic theory and the integration of the Maxwell stress tensor [6] can be rather complex and computationally intensive, exploiting approximations can be an advantage, *e.g.*, when combining optical force modeling
45 with stochastic simulations [23, 24].

For calculating optical forces acting on spherical or quasi-spherical particles, it is customary to identify several regimes which depend on the particle size [6]. For each regime, simplifications and approximations have been made for a better and more quantitative understanding and calculation of optical forces.
50 The *size parameter*, $x = k_m a$, is crucially used to determine the range of validity of these approximations, where $k_m = 2\pi n_m / \lambda_0$ is the light wavenumber in the medium surrounding the particle, a is the particle radius, λ_0 is the laser wavelength in vacuum used for trapping and n_m is the refractive index of the surrounding dielectric, non-magnetic, non-absorbing medium. When the parti-

55 cle size is much bigger than the wavelength of the laser beam, that is $x \gg 1$,
 optical trapping forces can be calculated with the *ray optics approximation* [25].
 The accuracy of this approximation increases with the size parameter, whereas
 electromagnetic scattering theories might become unpractical due to increasing
 computational complexity. This makes ray optics very useful when dealing with
 60 large particles. However, we note that ray optics is approximate by definition
 as it ignores the information on the phase of the incident electromagnetic wave
 and all related interference effects [26]. Thus, the lower size limit of validity is
 often dependent on the computed quantity [27, 28, 29]. If the size of the particle
 is much smaller than λ_0 ($x \ll 1$), we can adopt the *dipole approximation* and
 65 consider the particle as a dipole [30, 16, 31]. This means we are considering
 the electromagnetic fields homogeneous inside the particle under the condition
 $|n_p/n_m|x \ll 1$, where n_p is the refractive index of the particle. This condition
 has to be considered with care when dealing with high refractive index dielectric
 particles (*e.g.*, silicon) or noble metal (*e.g.*, gold, silver) nanoparticles, where
 70 the presence of plasmonic resonances dominate the optical response [32]. In the
 intermediate regime, that is when the particle size is comparable with the light
 wavelength ($x \simeq 1$) or for highly non-spherical or non-homogeneous particles,
 we need a complete wave-optical modeling of the particle-light interaction to
 calculate the trapping forces and different methods can be considered [6].

75 2.1. Ray optics

In the ray optics regime [17] the optical field is described by considering it
 as a collection of N light rays and employing the tools of geometrical optics [6].
 Each ray carries with it a portion of the incident power, P_i , so that the total
 power is $P_i = N \sum_m P_i^{(m)}$, and a linear momentum per second $n_m P_i^{(m)}/c$. When
 80 a ray impinges on a surface with an incident angle θ_i , it will be partly reflected
 and partly transmitted with a transmitted angle θ_t , according to the Snell's
 law [33]. As a consequence of energy conservation, the power is split between
 the reflected and transmitted part of the ray following Fresnel coefficients [23].
 Moreover, at each scattering event, the ray changes its direction and hence its

85 momentum change causes a reaction force on the center of mass of the particle. Most of the power carried by the incident ray is delivered to the transmitted ray that travels inside the particle until it impinges on the opposite surface. Here it will be reflected and transmitted again and a large portion of the power will be transmitted outside the sphere. The process will continue until all light escapes
90 from the sphere.

In general, if more than one ray interacts with a particle, the total force is given by the sum of the forces generated by the reflection and refraction of each ray. For simplicity we consider in the following a highly focused laser beam impinging on a spherical particle. To calculate the optical force of the
95 focused beam, we model it as a set of many rays that converge at a very large angle in the focal spot. Thus, we need to sum up all force contributions, $\mathbf{F}_{\text{ray}}^{(m)}$, associated to the reflection and transmission of each m -th ray with power $P_i^{(m)}$ in the direction $\hat{\mathbf{r}}_i^{(m)}$ forming the beam. Thus, the total force acting on the centre of mass of the sphere is $\mathbf{F}_{\text{GO}} = \sum_m \mathbf{F}_{\text{ray}}^{(m)}$, and more explicitly [6, 23]:

$$\mathbf{F}_{\text{GO}} = \sum_m \left[\frac{n_m P_i^{(m)}}{c} \hat{\mathbf{r}}_i^{(m)} - \frac{n_m P_r^{(m)}}{c} \hat{\mathbf{r}}_{r,0}^{(m)} - \sum_{j=1}^{+\infty} \frac{n_p P_{t,j}^{(m)}}{c} \hat{\mathbf{r}}_{t,j}^{(m)} \right]. \quad (1)$$

100 where $\hat{\mathbf{r}}_{r,0}^{(m)}$, $P_r^{(m)}$ and $\hat{\mathbf{r}}_{t,j}^{(m)}$, $P_{t,j}^{(m)}$ are the unit vector and power in the direction of the reflected 0-ray and the transmitted j -th ones, respectively. Each optical force, associated with the m -th ray, has components only in the incidence plane and can be split in two perpendicular components (note that this is valid only for spheres). The component in the direction of the incoming ray represents the
105 scattering force that pushes the particle away from the center of the trap. The component perpendicular to the incoming ray is the gradient force, that pulls the particle towards the optical axis when $n_m < n_p$. Instead, if $n_m > n_p$ the particle is pushed away from the high intensity focal region and different optical trapping strategies must be used [34]. It is often useful to define the dimension-
110 less quantities (trapping efficiencies) obtained dividing these force components by $n_m P_i^{(m)}/c$, that quantify how efficiently the momentum is transferred from

the ray to the particle.

For a single-beam optical tweezers, the focused rays will generate a restoring force proportional to the particle's displacement from an equilibrium point, that is for small displacements optical trapping can be modeled as an harmonic response. Due to the scattering force, the particle is displaced away from the focal point to an equilibrium position $\mathbf{C}_{\text{eq}} = [x_{\text{eq}}, y_{\text{eq}}, z_{\text{eq}}]$. Thus, for small displacements optical trapping forces are approximated as Hookean restoring forces, *i.e.*, $F_x \approx -\kappa_x(x - x_{\text{eq}})$, that can be related to an effective harmonic potential with trap stiffnesses κ_x , κ_y and κ_z .

A ray optics approach can be employed to study more complex particle shapes as long as the characteristic particle dimensions are much larger than the wavelength. For non-spherical particles two main differences occur. First, an optical torque can be generated from the difference between the angular momentum carried by the incoming rays and the outgoing rays with respect to the particle center-of-mass [6]. This shape-dependent optical torque can yield alignment [35] or rotation [36] of the particle. The second difference is the occurrence of a transverse component of the radiation pressure, *i.e.*, a set of parallel rays on a non-spherical particle can produce a force transverse to the incident light propagation direction, yielding an *optical lift effect* [37].

2.2. Dipole approximation

When the particle size parameter is small, $x \ll 1$, optical trapping forces can be calculated exploiting a dipole approximation, *i.e.*, the particle can be approximated as a small induced dipole immersed in an electromagnetic field $\mathbf{E}(\mathbf{r}, t)$, which can be considered homogeneous inside the particle ($x|n_p/n_m| \ll 1$) [30]. An induced dipole is thus generated and experiences electrostatic forces arising from its interaction with the inducing electric field. Consequently, an oscillating electromagnetic field, such as that of the laser beam used for an optical tweezers, induces an oscillating dipole, which also experiences forces arising from its interaction with the inducing electromagnetic field. Furthermore, an oscillating dipole radiates an electromagnetic field that can produce a mechanical effect on

other induced dipoles leading to, in some cases, *optical binding* [38].

This picture can be extended to a small particle, so that if the external field is not too large, the induced dipole moment, $\mathbf{p}(\mathbf{r}, t)$, can be expressed in terms of a linear complex polarizability, α_p , with respect to the surrounding medium, given by [39]:

$$\alpha_p = \alpha_0 \left(1 - i \frac{k_m^3 \alpha_0}{6\pi \varepsilon_m} \right)^{-1} \quad (2)$$

with ε_m dielectric permittivity of the medium and α_0 being the static Clausius-Mossotti polarizability, $\alpha_0 = 3V\varepsilon_m(\varepsilon_p - \varepsilon_m)/(\varepsilon_p + 2\varepsilon_m)$, where V is the particle volume and ε_p dielectric permittivity of the particle. 145

The polarizability in the dipole regime is linked to the cross-sections. In fact, the light-particle interaction, including optical forces, can be described in terms of extinction, σ_{ext} , scattering, σ_{scat} , and absorption, $\sigma_{\text{abs}} = \sigma_{\text{ext}} - \sigma_{\text{scat}}$. For a small particle of polarizability α_p , we can write the extinction and scattering cross-sections as [6]: 150

$$\sigma_{\text{ext,d}} = \frac{k_m}{\varepsilon_m} \Im\{\alpha_p\}, \quad \sigma_{\text{scat,d}} = \frac{k_m^4}{6\pi\varepsilon_m^2} |\alpha_p|^2. \quad (3)$$

Thus, we can consider the time-averaged optical force experienced by a small particle when illuminated by a time-varying electromagnetic field [16, 31, 40]:

$$\langle \mathbf{F} \rangle_{\text{DA}} = \frac{1}{2} \Re \left\{ \sum_i \alpha_p E_i \nabla E_i^* \right\}. \quad (4)$$

where E_i are the electric field components. Starting from this expression, one can explicitly write the optical force in terms of extinction cross-section and particle's polarizability [16, 31, 40, 41, 42]: 155

$$\langle \mathbf{F} \rangle_{\text{DA}} = \frac{1}{2} \frac{n_m}{c \varepsilon_m} \Re\{\alpha_p\} \nabla I(\mathbf{r}) + \frac{n_m}{c} \sigma_{\text{ext}} \langle \mathbf{S} \rangle - \frac{1}{2} c n_m \sigma_{\text{ext}} \nabla \times \langle \mathbf{s} \rangle \quad (5)$$

where $I(\mathbf{r}) = \frac{1}{2} n_m c |\mathbf{E}(\mathbf{r})|^2$ is the intensity of the electric field, $\langle \mathbf{S} \rangle = \frac{1}{2} \Re\{\mathbf{E} \times \mathbf{H}^*\}$ is the time-averaged Poynting vector of the incoming wave and $\langle \mathbf{s} \rangle = i \frac{\varepsilon_m}{2\omega} \mathbf{E} \times \mathbf{E}^*$ is the time-averaged spin angular momentum density [40, 41, 42].

The first term in Eq. (5) represents the gradient force and is responsible for particle confinement in optical tweezers. Arising from the potential energy of a dipole immersed in the electric field, it is conservative. Particles with a positive $\Re\{\alpha_p\}$ will be attracted toward the high intensity region of the optical field. Conversely, when the real part of the polarizability is negative the particles are repelled by the high intensity region. As an example, for an incident paraxial laser beam with a typical Gaussian intensity profile which propagates along the z axis, the trap stiffnesses $\kappa_{G,\rho}$ in the polarisation plane and $\kappa_{G,z}$ along the direction of propagation related to the gradient force are [43]:

$$\kappa_{G,\rho} = 2 \frac{\Re\{\alpha_p\}}{cn_m} \frac{I_0}{w_0^2}, \quad \kappa_{G,z} = \frac{\Re\{\alpha_p\}}{cn_m} \frac{I_0}{z_0^2}. \quad (6)$$

where I_0 is the maximum intensity at the center of the beam, w_0 is the Gaussian beam waist and z_0 is the beam Rayleigh range[6].

The second term in Eq. (5) is the scattering force. It is responsible for the radiation pressure and is non-conservative. Furthermore, it is directed along the propagation direction of the laser beam [3]. The last term in Eq. (5) is a spin-dependent force [40, 41]. This term is also non-conservative and dependent on the extinction cross-section. It can be generated by polarisation gradients in the electromagnetic field, but usually does not play a major role in optical trapping because it is either zero or very small compared to the other contributions. However, it may play a more significant role when considering optical trapping with optical beams of higher order with inhomogeneous polarisation patterns such as cylindrical vector beams [44, 45, 46] or superpositions of circularly polarised Hermite-Gauss beams [47]. Note that the time-averaged Poynting vector can be decomposed into the sum of an orbital and spin momentum density [48] and hence the non-conservative optical forces are related to the orbital component of the field momentum directed as the local wavevector [40, 49]. Thus, spin-dependent optical forces occur when the Poynting vector is not directed as the local wavevector and a transverse spin-dependent force component occurs [50, 51].

2.3. Electromagnetic theory and modeling

Particles illuminated by a radiation field experience a radiation force \mathbf{F}_{rad} and torque $\mathbf{\Gamma}_{\text{rad}}$, which contribute to determining their dynamical behaviour. Since the interaction between radiation and matter is regulated by conservation laws, it is possible to derive \mathbf{F}_{rad} and $\mathbf{\Gamma}_{\text{rad}}$ using the conservation of linear and angular momentum. So, the time-averaging optical force and torque exerted by a monochromatic light on a particle is given by [22, 52, 53, 54]:

$$\mathbf{F}_{\text{rad}} = \oint_S \hat{\mathbf{n}} \cdot \langle \mathbf{T}_M \rangle dS, \quad \mathbf{\Gamma}_{\text{rad}} = - \oint_S (\langle \mathbf{T}_M \rangle \times \mathbf{r}) \cdot \hat{\mathbf{n}} dS \quad (7)$$

where the integration is carried out over the surface S surrounding the scattering particle, $\hat{\mathbf{n}}$ is the outward normal unit vector, \mathbf{r} is the vector position, and $\langle \mathbf{T}_M \rangle$ is the averaged Maxwell stress tensor which describes the mechanical interaction of light with matter [55]. The general expression of the Maxwell stress tensor in a medium in the Minkowski form [6] can be simplified since we consider always harmonic fields, at angular frequency ω in a homogeneous, linear, and non-dispersive medium. In fact, writing the real physical fields, *e.g.* $\mathcal{E}(\mathbf{r}, t) = \Re \{ \mathbf{E}(\mathbf{r}) e^{-i\omega t} \}$, in terms of the complex amplitudes, *e.g.* $\mathbf{E} = \mathbf{E}(\mathbf{r})$, the averaged Maxwell stress tensor simplifies as [6, 18]:

$$\langle \mathbf{T}_M \rangle = \frac{\epsilon_m}{2} \Re \left\{ \mathbf{E} \otimes \mathbf{E}^* + \frac{c^2}{n_m^2} \mathbf{B} \otimes \mathbf{B}^* - \frac{1}{2} \left(|\mathbf{E}|^2 + \frac{c^2}{n_m^2} |\mathbf{B}|^2 \right) \mathbf{I} \right\} \quad (8)$$

where the fields, $\mathbf{E} = \mathbf{E}_i + \mathbf{E}_s$ and $\mathbf{B} = \mathbf{B}_i + \mathbf{B}_s$, are the total electric and magnetic fields, superposition of the incident ($\mathbf{E}_i, \mathbf{B}_i$) and scattered ($\mathbf{E}_s, \mathbf{B}_s$) fields and \mathbf{I} is the dyadic unit.

Several techniques have been proposed for computing electromagnetic scattering by non spherical particles and actually there is no single universal method that provides the best results in all cases. Depending on the specific particle parameters, one particular technique may prove to be the most appropriate in terms of efficiency, accuracy, and applicability [26]. While a comprehensive overview of numerical methods for optical tweezers it is beyond the scope of this work, some cover of the topic can be found elsewhere [56, 57, 58, 59]. Here we

give a very brief description of some common computational methods that have
205 been applied to the modeling of optical trapping of irregular shaped particles,
such as discrete dipole approximation [60, 61] (DDA), finite-difference time-
domain [62, 63] (FDTD), and we focus our attention to transition (T-)matrix
[64] methods with some notes on the connection with generalised Lorenz-Mie
theories [65] (GLMTs).

210 The DDA is a finite element method originally devised by Purcell and Penny-
packer [66] and later improved by Draine and Flatau [67]. In the DDA method,
a particle is split into a series of dipoles, each of which interacts with the inci-
dent electromagnetic wave and with the electromagnetic waves re-radiated by
all the other dipoles [60, 61]. DDA methods have been applied successfully to
215 optical tweezers modeling of both single-beam optical trapping of non-spherical
particles [68] and holographic multiple trapping of complex shaped structures
[69, 70]. The FDTD, instead, is based on the numerical integration of the
Maxwell equations in the time domain [62, 63]. The electric and magnetic fields
are sampled at discrete times and positions and, therefore, do not assume a har-
220 monic time-dependence. Applications of FDTD to optical trapping problems
have been explored for single-beam [71] and holographic [72] optical tweezers.
More recently, finite elements methods have been also used to model optical
trapping by near-field and plasmonic structures [73, 74]. The DDA and FDTD
methods, although more computationally intensive than T-matrix or GLMT,
225 can be readily applied to particles of any shape and composition, and to any
light field configuration.

T-matrix [64] methods have been widely used in optical tweezers modeling.
They are particularly suitable to describe optical trapping of composite and
arbitrary-shaped particles. Such approach provides a compact formalism based
230 on the multipole expansion of the fields suitable for dealing with scatterers of
(almost) arbitrary morphology [22, 26], *e.g.*, aggregates of spheres or clusters
[75, 76, 18, 19], plasmonic particles [77, 78, 79], stratified spheres [80, 81], and
spheres with internal aggregation (spheres within spheres) [22, 82]. The T-
matrix approach has been widely used for rigorously computing electromagnetic

235 scattering by single and composite particles. It yields a computational procedure
 computationally less intensive than those required by finite elements methods.
 Moreover, the transformation properties [75, 83, 84, 85, 26] under rotation and
 translation of the T-matrix make it possible several simplifications in optical
 force calculations as it requires the T-matrix to be computed only once, with
 240 respect to the particle reference frame thus saving much computational time.

For spherical particles, accurate modeling of optical tweezers can be obtained
 by generalizing Mie theory [86] through the so-called generalised Lorenz-Mie
 theories (GLMTs). In this approach a generic laser beam is expanded on a
 base of functions, *e.g.*, vector spherical harmonics, and the scattering problem
 245 is solved for spheres, so that separation of variables can be used to get the
 expansion coefficients of the scattered fields [65, 87]. The precise connection
 between the T-matrix formulation and GLMTs can be found in [88], while a
 description of the use of GLMTs for calculations in optical tweezers can be
 found, *e.g.*, in [89, 90, 91, 92, 93].

250 2.4. T-matrix methods

In order to calculate radiation forces and torques acting on an optically
 trapped particle through Eqs. (7), we give as an example a description of a
 T-matrix approach [18]. For simplicity we consider a generic non-magnetic
 ($\mu = 1$) particle with refractive index n_p in a medium of refractive index n_m
 255 illuminated by an incident field $\mathbf{E}_i(\mathbf{r})$ (the case $\mu \neq 1$ can be treated con-
 sidering the corresponding multipole expansion of the magnetic fields). The
 scattered electric field is $\mathbf{E}_s(\mathbf{r})$, so that the total electric field outside the parti-
 cle is $\mathbf{E}(\mathbf{r}) = \mathbf{E}_i(\mathbf{r}) + \mathbf{E}_s(\mathbf{r})$, while inside the particle is $\mathbf{E}_p(\mathbf{r})$. The key point of
 T-matrix methods is the expansion of the fields into a basis of vector spherical
 260 harmonics and the consequent application of the boundary conditions across the
 particle surface [22, 26]. The incident, $\mathbf{E}_i(\mathbf{r})$, and internal, $\mathbf{E}_p(\mathbf{r})$, fields have to
 be regular at the origin (fixed inside the scattering particle). Thus, they can
 be expanded in \mathbf{J} -multipoles that are a combination of vector spherical har-
 monics whose radial function is a spherical Bessel function, $j_l(kr)$, ensuring the

265 finiteness at the origin [6, 22], *e.g.*:

$$\mathbf{E}_i(r, \hat{\mathbf{r}}) = E_0 \sum_{p=1,2} \sum_{lm} W_{i,lm}^{(p)} \mathbf{J}_{lm}^{(p)}(kr, \hat{\mathbf{r}}). \quad (9)$$

where E_0 is the field amplitude, $W_{i,lm}^{(p)}$ are expansion coefficients, the label p is referred to the multipolar components of magnetic ($p = 1$) and electric ($p = 2$) nature and it is linked with parity [83]. In analogy to the incident and internal fields, the scattered wave is expanded in \mathbf{H} -multipoles, whose radial function is a spherical Hankel function, $h_l(kr)$, of the first kind because the scattered field has to satisfy the radiation condition at infinity [94]:

$$\mathbf{E}_s(r, \hat{\mathbf{r}}) = E_0 \sum_{p=1,2} \sum_{lm} A_{s,lm}^{(p)} \mathbf{H}_{lm}^{(p)}(kr, \hat{\mathbf{r}}), \quad (10)$$

where $A_{s,lm}^{(p)}$ are the amplitudes of the scattered fields which are determined by imposing the boundary conditions across the surface of the particle. In general, they depend on the orientation of the scattering particle with respect to the incident field. The multipole expansion of the normalized scattering amplitude is obtained by taking the limit of the \mathbf{H} -multipole fields for $kr \rightarrow \infty$ from which the asymptotic form of the scattered field is obtained [22].

The T-matrix of the scattering particle is the operator that, acting on the known multipole amplitudes of the incident field $W_{i,lm}^{(p)}$, gives the amplitudes of the scattered field, $A_{s,lm}^{(p)}$. Because of the linearity of Maxwell's equations and of the boundary conditions, the T-matrix is a linear operator, \mathcal{T} , so that $\mathbf{E}_s = \mathcal{T}\mathbf{E}_i$. Consequently, if both \mathbf{E}_i and \mathbf{E}_s are expanded on suitable bases, \mathcal{T} relates the coefficients of such expansions, encompassing all the information on the morphology and orientation of the particle with respect to the incident field [64]. The transition matrix $\mathcal{T} = \{T_{l'm'lm}^{(p'p)}\}$ of the scattering particle acts on the known multipole amplitudes of the incident field $W_{i,lm}^{(p)}$ to give the unknown amplitudes of the scattered field $A_{s,l'm'}^{(p')}$, *i.e.*,

$$A_{s,\eta l' m'}^{(p')} = \sum_{plm} T_{l'm'lm}^{(p'p)} W_{i,\eta lm}^{(p)}. \quad (11)$$

The quantities $T_{l'm'lm}^{(p'p)}$ take into account the morphology of the particle, but are independent of the state of polarisation of the incident field. Therefore, Eq. (11) holds true whatever the polarisation and it relates the basis-polarised amplitudes of the incident and of the scattered field.

The special case of a homogeneous sphere (Mie theory) of radius a yield a diagonal T-matrix, independent of m , whose coefficients are linked to the Mie coefficients, $a_l = -A_{s,lm}^{(2)}/W_{i,lm}^{(2)}$ and $b_l = -A_{s,lm}^{(1)}/W_{i,lm}^{(1)}$ [22].

By substituting the expansions of the incident [Eq. (9)] and scattered waves [Eq. (10)] in terms of multipoles taken in the asymptotic limit ($r \rightarrow \infty$), the general expression for the radiation force along the direction of a unit vector $\hat{\mathbf{u}}$, *i.e.*, $F_{\text{rad}}(\hat{\mathbf{u}}) = \mathbf{F}_{\text{rad}} \cdot \hat{\mathbf{u}}$ can be obtained [18]:

$$F_{\text{rad}}(\hat{\mathbf{u}}) = -\frac{\varepsilon_m E_0^2}{2k_m^2} \Re \left\{ \sum_{plm} \sum_{p'l'm'} i^{l-l'} I_{lm'l'm'}^{(pp')}(\hat{\mathbf{u}}) \left[A_{s,lm}^{(p)*} A_{s,l'm'}^{(p')} + W_{i,lm}^{(p)*} A_{s,l'm'}^{(p')} \right] \right\}, \quad (12)$$

where the integrals $I_{lm'l'm'}^{(pp')}(\hat{\mathbf{u}})$ can be expressed in closed form in terms of spherical harmonics [18]. Note how the force expressed by Eq. (12) can be separated into two parts related to extinction and scattering terms, $F_{\text{rad}}(\hat{\mathbf{u}}) = F_{\text{ext}}(\hat{\mathbf{u}}) - F_{\text{scat}}(\hat{\mathbf{u}})$, where:

$$F_{\text{ext}}(\hat{\mathbf{u}}) = -\frac{\varepsilon_m E_0^2}{2k_m^2} \Re \left\{ \sum_{plm} \sum_{p'l'm'} W_{i,lm}^{(p)*} A_{s,l'm'}^{(p')} i^{l-l'} I_{lm'l'm'}^{(pp')}(\hat{\mathbf{u}}) \right\} \quad (13)$$

and

$$F_{\text{scat}}(\hat{\mathbf{u}}) = \frac{\varepsilon_m E_0^2}{2k_m^2} \Re \left\{ \sum_{plm} \sum_{p'l'm'} A_{s,lm}^{(p)*} A_{s,l'm'}^{(p')} i^{l-l'} I_{lm'l'm'}^{(pp')}(\hat{\mathbf{u}}) \right\}. \quad (14)$$

$F_{\text{ext}}(\hat{\mathbf{u}})$ depends both on $A_{s,lm}^{(p)}$ and on the amplitudes $W_{i,lm}^{(p)}$ of the incident field, while $F_{\text{scat}}(\hat{\mathbf{u}})$ depends on the amplitudes $A_{s,lm}^{(p)}$ of the scattered field only. The subscript are used since this dependence is analogous to that on the extinction and scattering cross sections for the force exerted by a plane wave on a sphere [52]. In fact, the generic expressions for the radiation force are valid

305 for any scatterer and for any direction of the incident field. In the particular case of a linearly polarised plane wave that impinges on a spherical particle, the radiation force is simplified as [52]:

$$\mathbf{F}_{\text{rad}} = \frac{n_m}{c} I_0 \left[\sigma_{\text{ext}} \hat{\mathbf{k}}_i - \oint_{\Omega} \frac{d\sigma_{\text{scat}}}{d\Omega} \hat{\mathbf{r}} d\Omega \right], \quad (15)$$

in which σ_{ext} and $d\sigma_{\text{scat}}/d\Omega$ are the extinction and differential cross-sections, respectively, that can be expressed in terms of Mie coefficients [6], I_0 is the 310 intensity and $\hat{\mathbf{k}}_i$ is the propagation direction of the incident plane wave.

A similar procedure can be followed for the expression of the torque that can be split into extinction and scattering terms [95, 96]. Thus, for the axial z -component, $\Gamma_{\text{rad},z} = \mathbf{\Gamma}_{\text{rad}} \cdot \hat{\mathbf{z}} = \Gamma_{\text{ext},z} - \Gamma_{\text{scat},z}$, we have that:

$$\Gamma_{\text{rad},z} = \underbrace{-\frac{\varepsilon_m E_0^2}{2k_m^3} \sum_{plm} m \text{Re} \left\{ W_{i,lm}^{(p)} A_{s,lm}^{(p)*} \right\}}_{\text{extinction}} - \underbrace{\frac{\varepsilon_m E_0^2}{2k_m^3} \sum_{plm} m |A_{s,lm}^{(p)}|^2}_{\text{scattering}}, \quad (16)$$

where we have explicitly distinguished the extinction and scattering contribu- 315 tions. This relation obtained by Borghese *et al.* [95, 96] is a generalisation of the result obtained by Marston and Crichton [97] for a spherical particle illuminated by a circularly polarised plane wave, for which the longitudinal component of the torque is simply proportional to the absorption cross section, σ_{abs} :

$$\Gamma_{\text{rad},z} = \pm \frac{I_0}{\omega} (\sigma_{\text{ext}} - \sigma_{\text{scat}}) = \pm \frac{I_0}{\omega} \sigma_{\text{abs}} \quad (17)$$

where the \pm sign is related to left-handed or right-handed helicity of the inci- 320 dent circularly polarised light, respectively. The transverse components of the radiation torque, *i.e.*, $\Gamma_{\text{rad},x} = \mathbf{\Gamma}_{\text{rad}} \cdot \hat{\mathbf{x}}$ and $\Gamma_{\text{rad},y} = \mathbf{\Gamma}_{\text{rad}} \cdot \hat{\mathbf{y}}$, can be calculated

in a similar way [6, 95], obtaining:

$$\begin{aligned} \Gamma_{\text{rad},x} = & -\frac{\varepsilon_m E_0^2}{4k_m^3} \sum_{plm} \Re \left\{ \underbrace{s_{lm}^{(-)} W_{i,l,m+1}^{(p)} A_{s,lm}^{(p)*} + s_{lm}^{(+)} W_{i,l,m-1}^{(p)} A_{s,lm}^{(p)*}}_{\text{extinction}} \right. \\ & \left. - \underbrace{\frac{\varepsilon_m E_0^2}{4k_m^3} \sum_{plm} \Re \left\{ s_{lm}^{(-)} A_{s,l,m+1}^{(p)} A_{s,lm}^{(p)*} + s_{lm}^{(+)} A_{s,l,m-1}^{(p)} A_{s,lm}^{(p)*} \right\}}_{\text{scattering}} \right\} \end{aligned} \quad (18)$$

and

$$\begin{aligned} \Gamma_{\text{rad},y} = & -\frac{\varepsilon_m E_0^2}{4k_m^3} \sum_{plm} \Im \left\{ \underbrace{-s_{lm}^{(-)} W_{i,l,m+1}^{(p)} A_{s,lm}^{(p)*} + s_{lm}^{(+)} W_{i,l,m-1}^{(p)} A_{s,lm}^{(p)*}}_{\text{extinction}} \right. \\ & \left. - \underbrace{\frac{\varepsilon_m E_0^2}{4k_m^3} \sum_{plm} \Im \left\{ -s_{lm}^{(-)} A_{s,l,m+1}^{(p)} A_{s,lm}^{(p)*} + s_{lm}^{(+)} A_{s,l,m-1}^{(p)} A_{s,lm}^{(p)*} \right\}}_{\text{scattering}} \right\} \end{aligned} \quad (19)$$

where $s_{lm}^{(-)} = \sqrt{(l-m)(l+1+m)}$ and $s_{lm}^{(+)} = \sqrt{(l+m)(l+1-m)}$.

325 2.5. Focused fields

Finally, to calculate the radiation forces in an optical tweezers we need to consider the multipole amplitudes $\tilde{W}_{i,lm}^{(p)}$ of a highly focused beam. The expansion of a focused beam around the focal point is obtained by exploiting the angular spectrum representation [6, 18, 90]:

$$\mathbf{E}_f(x, y, z) = \frac{ik_t f e^{-ik_t f}}{2\pi} \int_0^{\theta_{\max}} \sin \theta \int_0^{2\pi} \mathbf{E}_{f,t}(\theta, \varphi) e^{i[k_t x \cos \theta + k_t y \sin \theta \cos \varphi]} e^{ik_t z} d\varphi d\theta, \quad (20)$$

in which f is the focal length, k_t is the wavenumber transmitted through the objective lens, and each transmitted plane wave, $\mathbf{E}_{f,t}(\theta, \varphi)$, can be expanded into multipoles according to Eq. (9). Therefore, the amplitudes of the focused field are:

$$\tilde{W}_{i,lm}^{(p)}(\mathbf{P}) = \frac{ik_t f e^{-ik_t f}}{2\pi} \int_0^{\theta_{\max}} \sin \theta \int_0^{2\pi} E_i(\theta, \varphi) W_{i,lm}^{(p)}(\hat{\mathbf{k}}_i, \hat{\mathbf{e}}_i) e^{i\mathbf{k}_t \cdot \mathbf{P}} d\varphi d\theta, \quad (21)$$

where the centre around which the expansion is performed is considered displaced by \mathbf{P} with respect to the focal point \mathbf{O} and the amplitudes $\tilde{W}_{i,lm}^{(p)}(\mathbf{P})$ define the focal fields and can be numerically calculated once the characteristics of the optical system are known. The expression for the radiation force along the direction of a unit vector $\hat{\mathbf{u}}$, *i.e.*, $F_{\text{rad}}(\hat{\mathbf{u}}) = \mathbf{F}_{\text{rad}} \cdot \hat{\mathbf{u}}$ can be obtained through the knowledge of the scattered amplitudes $\tilde{A}_{s,lm}^{(p)}$ related to the incident focal fields through the particle T-matrix [Eqs. (11)]. In practice, the expression of the optical force in the focal region is obtained from the correspondent one for the plane wave, Eq. (12), by changing $E_0 W_{i,lm}^{(p)} \rightarrow \tilde{W}_{i,lm}^{(p)}(\mathbf{P})$ and $E_0 A_{s,lm}^{(p)} \rightarrow \tilde{A}_{s,lm}^{(p)}$ [18].

2.6. Size scaling in optical trapping for spherical particles

As application of the T-Matrix approach we investigate the size scaling of the optical trapping for non-absorbing polystyrene spheres. Size scaling is a paradigm of nanoscience [98] as it characterises condensed matter systems for many applications in the most different research fields [99]. Crucial properties of materials and interactions change dramatically with size [100] influencing a wide variety of technological applications [101]. The size scaling properties of optical forces help us understand the important features of optical trapping in a wide size range and their comparison with experiments [102].

Polystyrene spheres are a typical sample used in optical tweezers experiments since they are easily purchased in accurate spherical shape and are used as size standard in microscopy techniques. They are easily functionalized and are well suited for optical tweezers applications in a biological environment [13]. Thus, they often represent the standard sample used for optical force calibration in optical tweezers [6, 102]. The particle refractive index is $n_p(\lambda_0 = 830 \text{ nm}) = 1.57$ [103]. We fix the wavelength of the linearly polarised trapping laser beam at $\lambda_0 = 830 \text{ nm}$ and we consider the particles immersed in water ($n_m = 1.33$). We consider a Gaussian laser beam focused by an ideal aplanatic lens which does not absorb and does not produce any aberration. The numerical aperture of the objective lens is taken as $\text{NA} = 1.3$.

We are interested in the exact calculation of optical forces, related trap stiffnesses, their scaling with particle size, and comparison with dipole approximation and ray optics calculations. We cover a wide size range with particle radius ranging from 50 nm up to 1.4 μm . Therefore, the size parameter range is [0.5 – 14.1] and only at the extreme values of the considered interval ray optics or dipole approximation calculations are safe to be used. This is indeed the typical size range of optical tweezers experiments [6]. The direction of propagation of the beam is taken along z and the polarisation axis along x .

A graph of the calculated strongly focused incident field intensity is provided in Figs. (1a - 1b - 1c). In these field intensity maps it is evident how the most elongated region is along the propagation (axial) direction, z , as expected for a focused Gaussian beam. Moreover, in the xy plane the linear polarisation of the incident beam breaks the cylindrical symmetry in the focal plane producing a tighter spot size along y with respect to the polarisation direction, x . This has observable consequences also on the symmetry of the optical trap and the transverse stiffnesses, κ_x and κ_y , particularly when dealing with nanostructures [10, 104, 20].

Within the T-matrix approach we have computed the (x, y, z) components of the optical force on the points of a computational grid employed with appropriate resolution. These components are calculated in a micron-sized range, $[-1.4 \mu\text{m}, 1.4 \mu\text{m}]$, around the paraxial nominal focus of the beam. So, we can plot the force as a function of particle displacement in each spatial direction, x, y, z . The trapping position of the particle in the longitudinal (z) direction is typically offset from the centre of the coordinate system due to the offset of the optical scattering force. To calculate the transverse force on the particle at the equilibrium position, the z (longitudinal) coordinate at which the axial force vanishes must first be found. The force plots in the transverse directions (x, y) can then be calculated. It is often convenient to calculate the dimensionless force efficiencies along the three cartesian directions, $Q_i = cF_i/n_m P$ with $i = x, y, z$. In Fig. (1d) we show them as calculated for a polystyrene sphere with radius $a = 300$ nm. The graphs present a relative maximum and minimum




Figure 1: The figures (a), (b) and (c) are maps of the focused field intensity normalized to the field entering the objective lens in the xy (a), xz (b), and yz (c) planes. We have considered a strongly focused field ($\text{NA} = 1.3$) that is linearly polarised along x and propagates along the z direction. Therefore, in the xy plane the cylindrical symmetry of the intensity map is broken by the polarisation, while in xz and yz planes the shape of the field appears elongated because of the beam propagation. With this incoming field, in the figure (d), the trapping efficiencies (Q_x , Q_y , Q_z) for a polystyrene ($n_p = 1.57$) sphere (radius of 300 nm) immersed in water ($n_m = 1.33$) are graphed in the transverse x, y (black and blue lines respectively) and longitudinal z (red line and dots) directions, as function of the particle displacement from the nominal paraxial focus. In (e), we compare the trapping efficiency along x direction for spheres of different radii: 200 nm (magenta line), 300 nm (blue line), 400 nm (green line) and 600 nm (red line).

approximately at the particle radius. As expected, it is here at these extremal points that the greatest restoring force is exerted because the sphere occupies approximately the entire focal region. To confirm this, we can see the trend of trap efficiencies as the radius changes, *i.e.* for Q_x , Fig. (1e). Instead, for larger dimensions of the particle, the slope of the curves gets smaller around the equilibrium point. The reason is that for larger particles the volume of the particle exceeds the interaction volume related with the laser spot, and the corresponding stiffness tends to zero as the particle radius increases.

In proximity of the equilibrium point the optical force can be linearized as an elastic restoring force with negative slope, *e.g.*, $F_x \approx -\kappa_x x$ for the x -direction. Thus, optical tweezers can be approximated with an effective harmonic potential with spring constants or trap stiffnesses κ_x , κ_y , κ_z . These quantities are of crucial importance in experiments because represent quantities measured when performing optical tweezers calibration [105]. In order to calculate the optical trap stiffnesses, we simply get the slope of the force-displacement graphs at the equilibrium position, where the force vanishes. In Figs. (2a - 2b - 2c) we show the calculated stiffnesses as a function of particle radius, a .

In the three graphs, an absolute maximum is present corresponding to a radius of $a \approx 300$ nm. This is the radius for which the volume of the scatterer overlaps the laser spot, optimizing the interaction region [27, 28]. However, the peak in the axial direction, Fig. (2c), is more depressed than in the transverse ones, see Figs. (2a - 2b). This is due to the shape of the focal spot shown in Figs. (1a - 1b - 1c), representing the maps associated with the incident field, which show a tighter profile in the transverse direction than in the axial one, yielding a tighter effective potential and a larger stiffness in x, y with respect to z . For size larger than the one corresponding to this maximum, the scaling of the force constants follows a hyperbolic scaling behaviour, $\kappa \propto a^{-1}$, this is also consistent with calculations in the ray optics approximation [6], blue lines in Figs. (3a - 3b). Using the T-matrix approach, we can reproduce the full electromagnetic theory of optical forces with great precision. In fact, when particles become larger than the interaction region, we also find the onset of




Figure 2: Optical trap stiffnesses, κ_x (a), κ_y (b), κ_z (c), for a polystyrene ($n_p = 1.57$) sphere immersed in water ($n_m = 1.33$) in the transverse, x (a) - y (b), and longitudinal z (c) directions, as a function of the particle radius a . The figures (d) and (e) show respectively the polarisation anisotropy, κ_{pol} , and trap aspect ratio, κ_{ar} , as a function of a . The radius of the considered spheres spans in the interval [50 nm - 1400 nm]. The symbols, excluding zero, are the calculated points by a T-matrix approach.




Figure 3: Optical trap stiffnesses along the x -direction for a polystyrene ($n_p = 1.57$) sphere immersed in water ($n_m = 1.33$) as a function of the particle radius a . The radius of the considered spheres spans in the interval [50 nm - 1400 nm]. The black data are obtained using the T-matrix approach. The red line is, instead, referred to the dipole approximation calculations considering a diffraction limited Gaussian beam. While the blue line represent optical trapping calculations in the ray optics approximation. In (a) we show a linear scale plot, while in (b) we show the same data in logarithmic scale to explicitly follow the size scaling.

modulation in the stiffnesses caused by the interference between the different multipoles [27, 28]. An exhaustive explanation of the interplay between the multipole interference for mesoscale particles is described by Nieminen et al. [29]. In fact, the gradient force from which we obtain the trap stiffnesses appears to be only weakly affected by this interference and good agreement between T-matrix and ray optics calculations is found for values of particle size well below the typical ray optics validity range ($x \gtrsim 50$) [27, 28, 29]. Note that this agreement is not occurring for other relevant quantities such as the scattering force where ray optics (that by definition neglects interference effects) fails to predict its strong variation with size and it only yields an average value [29].

On the other hand, for smaller values, the behaviour of the trap constants increases with a cubic power law, $\kappa \propto a^3$, in agreement with the volumetric scaling of the particle polarizability in the dipole approximation. As an example, in Fig. (3a), we compare the scaling of the transverse stiffness, κ_x , calculated with the T-matrix approach, Fig. (2a), with what is analytically obtained by a dipole approximation and considering a diffraction limited Gaussian beam spot. In our analytical calculation, the paraxial Gaussian beam waist w_0 is provided by the Abbe criterion [33, 106] so that $w_0 = 0.5\lambda_0/\text{NA}$. As shown by a logarithmic scale plot in Fig. (3b), the comparison between values calculated in dipole approximation and T-matrix is very good up to 200 nm. The same type of comparison and considerations hold for the size scaling of κ_y and κ_z . Such behaviour has been experimentally verified by measuring trap stiffnesses in a wide range of particle size from nano to the microscale [102].

Finally, in Figs. (2d - 2e) we represent respectively the polarisation anisotropy, $\kappa_{\text{pol}} \equiv 1 - \kappa_x/\kappa_y$, and the aspect ratio $\kappa_{\text{ar}} \equiv (\kappa_x + \kappa_y)/\kappa_z$. These quantities are related to both laser beam (polarisation, propagation) and particle properties (size, shape, asymmetry) and can be easily measured in experiments and linked to theoretical predictions [102, 104, 20].




Figure 4: Scheme of a typical OT setup. Trapping is achieved by a tightly focussed laser beam using a high-numerical-aperture objective lens (OBJ), which is also used to image the sample on a camera. The beam produced by the laser source (LS) is enlarged through a telescope lens system (lenses L1 and L2) to overfill the back aperture of the objective. A dichroic mirror (DM1) is used to reflect the laser light to the objective and to transmit the visible light to the camera. The sample is held and moved by a sample holder (SH). Inset (a) represents the real image of a SiNW optically trapped and aligned along the propagating axis of the laser beam [20, 107]. When the laser is switched off the SiNW performs Brownian motion and its alignment is randomized. Scattered and transmitted lights are collected and overlapped by a condenser lens (C) and projected on a QPD by a second dichroic mirror (DM2) and a lens (L4). Signals from the QPD are analyzed by a PC in order to calculate the calibration factors and to reconstruct the 3D Brownian motion of the trapped sample.

3. Experimental practice

An optical tweezers setup is an ideal tool to trap and manipulate micro and nanoparticles in liquid solutions with nanometer accuracy and to measure and apply piconewtonian forces [14]. Experimentally, optical trapping is achieved by focussing a laser beam to the diffraction limit by an objective lens with a high numerical aperture. For this reason optical tweezers setups are usually realized coupling a laser light source with an optical microscope, where the objective is also used to image the sample on a camera. An optical tweezers setup can be based on different kinds of optical microscopes: normal or inverted, commercial or customized, according to user requirements and budget. Usually, customized microscopes have lower costs and a bigger flexibility compared to commercial ones. Furthermore, more stable traps are obtained using optical tweezers based on inverted microscopes, because the laser beam is directed upwards and gravity works in an opposite direction to the radiation pressure. Since the realization of the first single-beam optical tweezers in 1986 [5], several technologies have been developed and merged with optical tweezers to customize setups for several purposes[105]. Currently, thanks to the new generation of digital sensors it is possible to investigate phenomena having spatial and time scales not reachable until few years ago. Liquid crystal displays (LCDs) with high resolution have been employed as reflective elements in holographic optical tweezers (HOT) to shape laser beams. In doing this, it is possible to create and move multiple traps in real time and to generate vector beams able to carry optical spin and angular momentum [108]. Furthermore, real time tracking and feedback position systems are now available [109]. Although there are several kinds of optical tweezers setups, their principle of operation can be understood by considering a standard optical tweezers setup (fig. 4). This is a forward-scattering setup mounted around an inverted microscope. A laser source generates a collimated beam, which is expanded by a telescope lens system and reflected by a dichroic mirror to overfill the back aperture of an oil immersion objective having an high numerical aperture. The overfilling creates the maximal optical field gradient

in the focal spot for a more efficient trapping. The dichroic mirror acts like a short pass filter, it reflects the laser light and transmits the visible light to a camera, preventing the saturation of the CMOS detector and allowing the view of the sample on a monitor. The sample is held and moved by micrometric translational stage mounted on an inverted microscope. Forward scattered light from trapped particle is collected by a condenser and reflected by a dichroic mirror to a quadrant photo detector (QPD) [6]. Thus, particle tracking and optical tweezers calibration can be achieved either by video microscopy either by photonic force microscopy, *i.e.*, by the temporal or frequency analysis of the light scattered onto the QPD. Digital video microscopy is based on the analysis of the images from a digital camera to track the trajectories and displacements of the imaged objects with a subpixel resolution and a sampling rate depending on the camera performances [110]. Photonic force microscopy allows to track the displacements of a single trapped particle, using a photodetector, with a typical spatial resolution of 0.1 nm and sampling rate above 100 kHz [111].

3.1. Optical tweezers calibration

The main goal of calibrating an optical tweezers is to determine their trap stiffness κ and to measure the absolute displacements of the trapped particles. For small displacements from the equilibrium position (e.g. along x axis), the trapping potential is considered to be harmonic with a Hookean restoring force $F_x = -\kappa_x x$ and potential energy $E_x = \frac{1}{2}\kappa_x x^2$ (we will refer only to the x component in order to shorten the notation). In doing this, once the trap stiffness κ_x is known, the x component of an external force acting on the particle can be quantified by measuring the particle displacement x . For spherical particles measuring these three stiffnesses, $\kappa_x, \kappa_y, \kappa_z$, fully characterise the optical trap. However, this is strictly valid for spherical particles where only translational degrees of freedom can be considered. For non-spherical particles optical torque, polarisation, and angular fluctuations play a role and must be included in the calibration analysis and the reconstruction of the effective trapping potential [20, 21, 45, 104, 107, 112] (see for example fig. 5). In fact, the dynamics of




Figure 5: Data analysis for a SiNW with a diameter of 10 nm and a length of 3.6 μm . a) x component of Brownian motion. b) Probability distribution function $\rho(x)$. c) Potential well profile of the trap $U(x)$. Red lines represents a quadratic fit, from which the trap stiffness κ_x can be calculated as fitting parameter.

non-spherical particles in optical traps can be very complicated due to the occurrence of transverse components of the scattering force and optical torques [54, 56]. For example elongated rod-shaped (*e.g.* nanowires) or ellipsoidal particles align with the optical (z -)axis of the trap (see fig. 4) due to the optical torque and show small angular thermal fluctuations about their stability axis that can be measured together with optical aligning torques [20, 104]. Moreover, the translation-rotation coupling in non-spherical particles can yield regular biased orbital motion that has been the subject of recent experimental and theoretical investigations [107, 113, 114, 115, 116].

Typical optical tweezers setups, use cameras or photo-detectors to detect the motion of trapped particles. In both cases, their measurements units (pixels and volts) are proportional, by a calibration constant and around the equilibrium position, to the absolute particle displacement [6, 117]. There are several active or passive calibration protocols, but here we briefly discuss only the most common passive ones as applied to spherical particles: potential analysis, autocorrelation function analysis, power spectrum and mean square displacement analysis [6, 117]. The simultaneous application of different calibration methods is a good check to confirm their consistency and the quality of the acquired experimental data. Spatial calibration with digital video microscopy is easier than QPD since it is possible to image directly reference targets having well known dimensions to find the conversion factor pixel/ μm . Once this factor is known it is possible to proceed to the force calibration.

Optical potential method.— Considering the Brownian motion of a trapped particle (fig. 5a) it is possible to calculate its positional distribution functions $\rho(x)$ (fig. 5b). This can be used to determine the profile of the trapping potential $U(x) = k_B T \log \rho(x)$ (fig. 5c). In doing this, the trap stiffness κ_x can be obtained as fit parameter of the potential $U(x)$ (fig. 5c).

Autocorrelation method.— In this case the starting point of the calibration procedure is the Langevin equation describing displacement $x(t)$ of an optically

535 trapped particle in liquid solution [118, 119, 120]:

$$m \frac{d^2 x(t)}{dt^2} + \gamma \frac{dx(t)}{dt} + \kappa_x x(t) = F_x(t) \quad (22)$$

where $\gamma \frac{dx}{dt}$ is the viscous damping term, $\kappa_x x(t)$ is the restoring force term, $F_x(t)$ is a random fluctuating force and γ is the hydrodynamic viscous coefficient related to the medium viscosity η and to the particle size and shape, which for spherical particles with radius r becomes $\gamma = 6\pi\eta r$, according to the Stokes law. In the low Reynolds number limit, applicable to microparticles suspended in water, the system can be considered overdamped and the first term of eq. (22) can be neglected when compared to the others. In this special case, the particle motion $x(t)$ can be well described by the overdamped Langevin equation:

$$\frac{dx(t)}{dt} + \omega x(t) = \xi_x(t) \quad (23)$$

where $\xi(t) = \frac{F(t)}{\gamma}$.

545 This equation can be properly recast to determine the analytic expression of $C_{xx}(\tau)$, which is the autocorrelation of the displacement $x(t)$ [6, 117]. In particular, for a spherical particle $C_{xx}(\tau)$ decays exponentially with a characteristic time τ dependings on the trap spring constant κ_x and on the hydrodynamic drag coefficient γ [121]:

$$C_{xx}(\tau) = C_{xx}(0) \exp\left(-\frac{\kappa_x}{\gamma} \tau\right) \quad (24)$$

550 Since the QPD produces voltage signals $V_x(t) = \beta_x x(t)$ proportional by the position calibration factor β_x to $x(t)$, the autocorrelation function of the experimental displacements can be expressed as $C_{xx}^V(\tau) = \langle V_x(t)V_x(t+\tau) \rangle = \beta_x^2 C_{xx}(\tau)$, where according to the equipartition theorem $C_{xx}^V(0) = \beta_x^2 C_{xx}(0) = \beta_x^2 k_B T / \kappa_x$ and the calibration factor β_x can be expressed as:

$$\beta_x = \sqrt{\frac{C_{xx}^V(0) \kappa_x}{k_B T}} \quad (25)$$

555 The experimental autocorrelation function $C_{xx}^V(\tau)$ can be fitted by eq. (24) to obtain the stiffness κ_x which, together with the experimental value of $C_{xx}^V(0)$, are employed to calculate the calibration factor β_x according to the eq. (25).

Power spectrum method.– The power spectral density (PSD) of a signal, describing how its energy is distributed with frequency, can be also employed to 560 calibrate optical tweezers. Considering the Fourier transform of the overdamped Langeven equation (eq. 23):

$$-i\omega\tilde{x}(\omega) + \omega_x\tilde{x}(\omega) = \tilde{\xi}_x(\omega) \quad (26)$$

where we defined $\omega_x = \kappa_x/\gamma$ and solved for $\tilde{x}(\omega) = \tilde{\xi}_x(\omega)/(-i\omega + \omega_x)$.

The corresponding power spectrum shows a Lorentzian profile:

$$S_x(\omega) = |\tilde{x}(\omega)|^2 = \frac{2D}{\omega^2 + \omega_x^2} \quad (27)$$

where the half-width of the distribution is the relaxation frequency of the trap ω_x . By fitting the power spectrum of the signal, it is possible to obtain the corner frequency ω_x as fit parameter and hence deduce the trap spring constant κ_x . Also in this case the power spectrum of the signal from a QPD is proportional to the power spectrum of the particle fluctuation:

$$S_x^V(\omega) = \beta_x^2 S_x(\omega) = \beta_x^2 \frac{2D}{\omega^2 + \omega_x^2} \quad (28)$$

Setting $\omega = 0$ the calibration factor β_x can be expressed as:

$$\beta_x = \sqrt{\frac{S_x^V(0)\omega_x^2}{2D}} \quad (29)$$

Mean square displacement.– This calibration method is based on the analysis of the mean square displacement (MSD) of the trapped particle: a measure of 565 the deviation of the particle position from its initial position in a time interval τ [122, 117]:

$$\text{MSD}_x(\tau) = \langle [x(t+\tau) - x(t)]^2 \rangle = 2 \frac{k_B T}{\kappa_x} [1 - e^{-\omega_x |\tau|}]. \quad (30)$$

Considering this equation it is possible to notice that $\text{MSD}(0) = 0$, $\text{MSD}(\tau)$ presents a transition from a linear growth corresponding to a free diffusion

behaviour at low short time scale ($\tau \ll \omega_x^{-1}$) to a plateau value of $2\frac{k_B T}{\kappa_x}$ for
 570 $\tau \gg \omega_x^{-1}$, due to the particle confinement at long timescales. Also in this case
 the MSD of the QPD signals is proportional to that of the particle displacements,
 $\text{MSD}_x^V(\tau) = \beta_x^2 \text{MSD}_x(\tau)$. Combining the relations above and setting $\tau \rightarrow \infty$
 we get $\text{MSD}_x^V(\infty) = 2\beta_x^2 / \frac{k_B T}{\kappa_x}$
 from which the calibration factor can be calculated as:

$$\beta_x = \sqrt{\frac{\text{MSD}_x^V(\infty)\kappa_x}{2k_B T}} \quad (31)$$

575 It is noteworthy that all the calibration methods used to calibrate optical
 tweezers based on photodetectors, can be also applied to camera ones, in this
 case instead of volt there will be pixel units.

4. Applications

In this section we give an overview of some recent applications of optical
 580 tweezers and optical forces in liquid environment. The contactless manipula-
 tion of particles promotes the use of OT in a wide variety of research fields.
 Here, we describe a selection of systems where OT has enabled advances in
 biology, microbubbles manipulation, chiral optomechanics, nanotechnology, op-
 tical binding, spectroscopy, critical Casimir forces, stochastic thermodynamics,
 585 and active matter.

4.1. Mechanical properties of red blood cells

Optical tweezers have numerous applications in biological sciences, ranging
 from single molecule studies [123, 124] to cell biophysics [125, 126]. In this
 section we will consider one particularly successful application, namely the in-
 590 vestigation and determination of the mechanical and elastic properties of red
 blood cells (RBCs). Very early in the history of optical tweezers Ashkin *et al.*
 showed that RBCs (and many other biological species) could be trapped with-
 out optical damage using an infra-red laser beam [11]. Subsequent experiments
 aimed at determining mechanical properties of RBCs have used both the direct

595 trapping method, or applied forces to the cell indirectly using optically trapped
microbeads bound to the cell as ‘handles’. The first method has the advantage
of simplicity of set-up and sample preparation, in the second calibration of the
applied forces is more straightforward.

Hénon *et al.* attempted to determine the shear modulus of the RBC mem-
600 brane by stretching cells using microbead handles in calibrated optical traps [127].
Experiments were performed on cells in the native biconcave discoid shape, and
cells that were osmotically swollen to a spherical shape, which permitted an ex-
act algebraical solution to the deformation geometry. In both cases for applied
forces below 15 pN a linear (with increasing force) deformation of the cell was
605 observed, producing a shear modulus of $\mu = 2.5 \pm 0.4 \mu\text{N} \cdot \text{m}^{-1}$. At higher
applied forces the cell entered a nonlinear deformation regime. Later, Dao *et al.*
[128] used the same experimental geometry in a system capable of applying
much higher forces of up to 340 pN. These experiments were complemented by
computed simulations, and suggested that at high applied stress where defor-
610 mations as large as 50% were observed, the membrane shear modulus could be
as high as $\mu = 11.1 \mu\text{N} \cdot \text{m}^{-1}$.

Bronkhorst *et al.* showed that optical tweezers could be used to deform a red
blood cell under direct trapping using a line of three optical traps to ‘fold’ the
cell and observe the relaxation time [129]. Later Liao *et al.* showed that RBCs
615 could also be stretched under direct trapping, using a single beam that jumped
rapidly between two locations to grip the cell [130]. As the separation between
the two trap locations was increased the cell was stretched along the line joining
them. In this work the elongation of the cell was explained qualitatively using a
two-dimensional model. Later theoretical work accurately calculated the optical
620 stress distribution over the surface of an RBC osmotically swollen to adopt a
spherical shape and its resulting deformation in the dual tweezers stretching
experiment [131]. By solving for the dynamic deformation of the RBC, that
is, accounting for the change in shape and redistribution of optical stress as
the cell deforms towards the stretched state the authors could determine the
625 membrane mechanical properties (Young’s modulus and shear modulus) before

and after treatment with N-ethylmaleimide, which is known to cause a decrease in cellular deformability. This work was further refined for the RBC in its native biconcave discoid shape [132]. When predicting the deformation of the RBCs under optical stress it is necessary to account for both the non-linear [133, 134] and viscoelastic [135] properties of the cell. By combining optical tweezers stretching with Raman spectroscopy, Raj *et al.* found that at large deformation the RBC must undergo a structural transformation in order to bear the high load [136].

An alternative technique for deforming cells using optical forces is the optical stretcher [137]. Rather than using strongly focused laser beams as is the case in optical tweezers, the optical stretcher employs two weakly diverging counter-propagating beams to trap particles. The distribution of optical stress over the surface of the trapped object leads to elongation along the beam propagation axis [137]. When complemented by deformation modelling [138, 139], the optical stretcher can be used to quantify cell mechanical properties in a high throughput manner [140] as the cells flow in a microchannel through the stretching beams rather than being captured individually in optical tweezers. In some situations, extreme stretching can give rise to irrecoverable shape changes of the cell into a so-called “columnar” form [141].

Altered mechanical properties of RBCs has been linked with a number of pathological conditions [142, 143], and optical tweezers have been used to test several of these including malaria (*plasmodium falciparum*) [144, 145], diabetic retinopathy [146] and birdshot chorioretinopathy [147]. Similarly optical tweezers have been used to show that the drug Atorvastatin softens the RBC membrane [148].

4.2. Microbubbles

Optical trapping of gas microbubbles presents additional challenges due to their buoyancy, and to their refractive index which is lower than the suspending medium. Due to their low relative refractive index the optical gradient force acts to repel them from the high intensity part of a laser beam and conse-

quently alternative strategies are needed for optical trapping. One possibility is to use a Laguerre-Gaussian (LG) beam [149] whose annular intensity profile provides transverse confinement of the bubble to the beam's dark core. In the vertical direction the position of the bubble is determined by the balance
660 between optical forces and buoyancy, and so the LG beam does not form a three-dimensional optical trap. When implemented in an upright microscope with the trapping LG beam propagating downwards it is a form of 'counter-levitation'. Three-dimensional optical trapping requires the addition of a second counter-propagating hollow beam [34]. In principle, structured light such as optical
665 bottle beams [150] can be also used for three-dimensional optical trapping of microbubbles.

A second strategy is to use a 'time-averaged' optical trap, created by scanning a focussed Gaussian around the circumference of the microbubble. If the scanning frequency is high enough, that is, if the beam rotates around the bubble
670 faster than it can diffuse away then the bubble remains trapped on the axis of rotation [151]. Such a trapping scheme was implemented in an upright microscope, and the dependence and scaling of the trapping force on the bubble radius and radius of the trap beam scanning circle has been investigated [152]. A theoretical analysis of the optical forces in this configuration [153] has shown
675 that that there is a sufficient optical gradient force in all directions to provide a restoring force on the bubble, and hence it is a true three-dimensional optical trap. Indeed, the axial gradient force can be enough to confine a microbubble in an inverted microscope optical tweezers [154].

Microbubbles are also amenable to trapping using acoustic fields. The combination
680 of acoustic and optical trapping presents a great advantage in the differing length scales over which they operate: the acoustic field may be used for long-range manipulation before an optical trap is used for high-resolution control of position [155]. The optical and acoustic traps also complement each other as, once calibrated [156], the optical trap may be used to measure the
685 acoustic forces on the bubble [157]. Such a system may be used to characterise an individual microbubble, and in particular its acoustic response, with a view to

reducing the uncertainties from population averaging [158]. Further applications of optically trapped microbubbles include Raman spectroscopy of the gas enclosed by the bubble [159], measuring the microbubble shell thickness [160],
690 and positioning the bubble adjacent to cells prior to it being irradiated by ultrasound. The cells may then be damaged or destroyed by the oscillations and subsequent collapse of the microbubble [161].

4.3. Chiral optical forces.

Chirality derives from the lack of mirror symmetry of an object. A chiral object exists in left- and right-handed version (enantiomers) that cannot
695 be superimposed by translations and rotations within the space where they are embedded. Also circularly polarised light is chiral, with handedness depending on the electric field sense of rotation with respect to the propagation axis. Besides linear momentum, circularly polarised light may transfer also
700 spin angular momentum [162]. Recently, several works considered chirality-dependent optical forces on small particles aiming at all-optical separation of enantiomers [163, 164, 165]. The optomechanical interaction of chiral light with mesoscopic objects has been studied in optical tweezers, giving the opportunity to investigate the transfer of spin angular momentum to birefringent particles
705 and observe spin-dependent light-induced rotations [166, 167, 168, 169]. Chiral optomechanics on supramolecular chiral particles has been recently investigated on cholesteric liquid crystals (CLC) [170, 171, 172, 173, 174]. In particular, left-handed CLC solid microparticles have been synthesized [175], optically trapped, and chiral rotations observed for the corresponding left-handed circularly polarised light [176, 177]. In fact, the left-handed CLC particles behave as chiral
710 mirrors, which reflect only left-handed light, while maintaining its handedness [176]. On the contrary, right-handed light is transmitted unaffected. Due to the conservation of angular momentum, a reaction torque is transferred from the circularly polarised trapping beam to the chiral microparticle [176]. In case
715 of microparticles in the low chirality regime [178], an additional contribution to the optical torque due to residual birefringence must be taken into account to




Figure 6: Micro-engine powered by critical demixing. An optically trapped colloidal particle, with radius $a = 1.24 \mu\text{m}$, immersed in a water-2.6-lutidine critical solution performs rotations around the trapping beam. The performances of the engine can be tuned by adjusting the criticality of the mixture via the ambient temperature. At low temperature (a), the particle is stably trapped. When the temperature increases (b), the particle starts to rotate around the optical beam (white solid lines trajectories), reaching its maximum value for a temperature of 26° (c) with a very reproducible trajectory. A further increase of the temperature results in a decrease of the engine rotations and performances (d, e). The white bars in (e) correspond to $1 \mu\text{m}$. Figure adapted from [179]

fully explain the rotation dynamics of trapped CLC particles at variable degree of ellipticity of the trapping beam [177]. The combination of chiral and optical retardance properties allows to better control the optomechanics of these microparticles that open perspectives for the study of chiral optical sorting, laser trapping and cooling of chiral mesoscopic particles based on the coupling of their translational and rotational degrees of freedom, and microcavity optomechanics, by embedding at the centre of the chiral microparticle a resonant nanoparticle.

4.4. Nanotechnology applications

During the last few decades much effort has gone into the miniaturization of machines down to microscopic scales, often inspired by biological systems [180]. This miniaturization process is crucial for the development of nanotechnology [181] and, in this context, optical tweezers are a powerful tool to assemble micro- and nanodevices thanks to their ability of contactless manipulation [182, 183, 10]. Moreover, they are capable of applying and detecting extremely small (femtonewton) forces and torques yielding potential for driving nanomachines [184, 185, 186, 187]. At the nanoscale, semiconducting [188, 104, 185, 20, 45, 21], metal [125, 189, 190, 191, 192, 193, 194], and hy-

brid [81, 78] colloidal particles have been trapped and manipulated opening
735 novel exciting possibilities for assembly, characterisation and optical control of
nanodevices and biomolecules [10].

Nanodevices or micro-engines need power to operate and to be controlled. A
solution to this demand can be provided by structured optical beams, carrying
orbital and spin angular momentum (SAM and OAM), generated by holographic
740 optical tweezers (HOT) or similar techniques [195]. In particular, microrotators
and micropumps have been realized by transferring SAM and OAM to micropar-
ticles [196, 166, 197, 176, 177].

Another approach to power nanodevices is to emulate the working principles
of heat engines. The nucleation of vapour bubbles inside silicon micro-cavities
745 has been used to realize several microscopic heat engines with a working vol-
ume of only 0.6 mm^3 [198, 199]. Also, a micro-particle has been employed as
a piston in an optical tweezer, realising a microscopic steam engine powered by
the periodic generation of cavitation bubbles [200]. Microscopic versions of the
Stirling and Carnot cycles have been realized using optically trapped particles
750 to study their stochastic thermodynamic properties [201, 202]. Recently, a mi-
croscopic engine powered by the density fluctuations of a critical solution has
also been proposed [179], where a micron-sized particle performs revolutions
around the optical beam when optically trapped in a water-2,6-lutidine critical
mixture. The work performed by this engine is adjustable by the power of opti-
755 cal trap, the temperature of the environment and the criticality of the mixture
(Fig. 6)[179].

4.5. *Optical binding*

Optical binding describes the spontaneous organisation of large numbers of
colloidal particles in an optical field that arises as a result of multiple optical
760 scattering[38, 203, 204]. It was first observed by Burns et al.[205], who demon-
strated an optical binding interaction between a pair of particles in a line-shaped
focus by showing that there existed discrete values of the particle's separation
in the transverse direction where they formed a stable bound pair. This work

was followed by the observation of a large number of particles that ‘crystallised’
765 in the interference pattern formed in the transverse plane of the intersection of
a number of laser beams[206].

The optical binding interaction can be simply understood by considering a
pair of dipoles in an optical field[6]. Consider the case of two dipolar particles
arranged parallel to the propagation direction of a laser beam. The first dipole
770 scatters radiation, and the scattered field propagates in the forward direction to
impinge on the second dipole but with a fixed phase difference with the incident
field. The net effect is to modify the magnitude of the scattering force in the
forward direction, but as the phase difference between the two fields is constant
the force does not change sign, both particles are pushed in the same direction
775 and the first tends to catch up with the second. Now, if we consider the fields
impinging on the first particle, these are the incident field and the fields that is
backscattered by the second particle. The phase difference between these fields
depends on the relative separation of the particles, and hence the force between
the particles oscillated with a change of sign with a period of half the optical
780 wavelength. This gives rise to the observed optical binding interaction and the
existence of stable inter-particle separations.

Optical binding may be realised experimentally in a number of configura-
tions. The original experiments of Burns *et al.*[205, 206] used shaped optical
fields to produce optical binding in the transverse plane. Longitudinal optical
785 binding can be realised with counter-propagating free-space beams, for example,
emerging from a pair of optical fibres [207, 208], or with counter-propagating
evanescent fields [209, 210]. The evanescent field may be generated by a weakly
focusing a laser beam onto an interface at an angle just greater than the crit-
ical angle for total internal reflection. When the radiation pressure from this
790 beam is balanced by a counter-propagating beam a variety of optically bound
structures form that depend on the optical fields configuration and the particle
size and optical properties [209, 211]. In such experiments video tracking of
the Brownian motion of the particles in the optically bound structure provides
a powerful technique for evaluating the optical forces and structure form and

795 dynamics [212, 213].

An alternative method for creating an evanescent field is to use a waveguide with suitably small dimensions. This scheme was first demonstrated using a channel waveguide in a planar surface, the evanescent field of which can trap and propel microparticles [214, 215]. More recently a tapered optical fibre scheme
800 has been demonstrated for unidirectional transport using a single beam [216, 217], and stable trapping using counter-propagating beams [218, 219]. Tapered optical fibres are fabricated from standard fibre using a ‘heat-and-pull’ technique [220] to draw the fibre to micron- or sub-micron dimensions, at which scale the majority of optical power is carried in the evanescent field that
805 extend into the region surrounding the fibre [221]. The high degree of control that can be exerted over the tapering process permits selective excitation of higher modes of the fibre and an extra degree of control over the trapping and binding [222, 223].

The optical binding phenomenon is not limited to dielectric particles: optical
810 binding of silver [224] and gold [225] has been observed, with a plasmonic enhancement of binding forces. Nor is it limited to spherical particles: binding of rod-shaped particles (carbon nanotube bundles) has also been demonstrated [10]. In this case, similarly to spherical particles, a variety of bound structures is predicted to occur [226, 227].

815 4.6. *Optical force positioning, aggregation, and spectroscopy.*

As already pointed out, the behaviour of a nanoparticle in the focal spot of a laser beam is controlled by the balance between the gradient force and the radiation pressure. In ordinary trapping measurements, the radiation pressure must be minimized, aiming at confining the nanoparticle in the optical trap. However,
820 several other interesting applications can be developed by exploiting the pushing effect of the radiation pressure. Optical force printing [228, 229, 230, 231] is one of these applications. The all-optical patterning of surfaces is allowed by single-beam trapping and subsequent positioning of Au colloids [228]. However, the process can be improved by directly using the strong axial force acting on




Figure 7: (a) Sketch of the optical force position and aggregation operation.

Nanorods-protein complexes intercepted by a focussed laser beam are optically pushed towards the bottom surface of a glass microfluidic chamber. Here they aggregate, yielding efficient SERS-active clusters. Proteins embedded at the clusters hot spots undergo a strong local field giving rise to a SERS emission that allows to reach picomolar sensitivity in biomolecular detection. (b) SERS emission of BSA-NRs complexes increasing as a function of time during the optical force aggregation process.

825 a plasmonic nanosphere at resonance wavelengths, which pushes it toward the substrate [229, 230, 231]. The use of radiation pressure instead of trapping forces to guide the particle toward the substrate avoids the constraint on the size of the plasmonic particle, which to be trapped must be not larger than approximately 200 nm in diameter [77]. Moreover, by using a Spatial Light Modulator
830 a laser beam can be split into several beams creating an optical pattern, for simultaneous and controlled positioning of nanoparticles on substrates [229].

In this context, it has been shown that surface-enhanced Raman scattering [79] (SERS)-active aggregates can be created on glass substrates by optical manipulation of gold nanorods (AuNRs) in a liquid environment [232, 233] containing biomolecules. In fact, due to their very large extinction cross-sections, these
835 AuNRs cannot be optically trapped with visible wavelengths [232]. However, they can be pushed towards the sample chamber glass walls where they aggregate and form hot-spots on which the Raman spectrum of diluted biomolecules can be strongly enhanced. This method combines the great advantage of op-

840 tical force printing with the capability of spectroscopic detection of molecules
directly in liquid environment, which is the ideal condition for bio-inspired re-
search. A number of biomolecules have been studied. First measurements fo-
cused on Bovine Serum Albumin (BSA) molecule, for which the Raman limit
of detection (LOD) in water has been lowered from 10^{-3} M to 5×10^{-8} M
845 [233]. Another protein, hemoglobin, has been detected at concentrations as low
as 1 pM [232], well below the SERS LOD of approximately 100 nM. Results
on amino-acids such as phenylalanine [232] and enzymes such as lysozyme [232]
and catalase [233] have been also reported [232]. This molecular detection tech-
nique can be also made protein-selective by functionalizing the AuNRs with
850 aptamers, which capture a target biomolecule. Proof-of-concept results have
been obtained with ochratoxin A [233], a nephrotic mycotoxin which may con-
taminate food commodities and wine representing potential public health risk.
With this methodology, the toxin has been detected in liquid environment at
concentrations as low as $1 \mu\text{M}$ [233].

855 Optical positioning of AuNRs is particularly interesting because it is a low
cost alternative to much more complex lithographic techniques. As a further
advantage, it allows an easy and fast way to decorate with metallic parti-
cles also not flat substrates, such as three dimensional micro- and nanostruc-
tures obtained with direct laser writing by two-photon photopolymerization
860 (TPP)[234, 235, 236, 237, 238, 239, 240, 241, 242], opening the way to the re-
alization of hybrid metal-polymer devices such as, for example, spectroscopic
probes and sensors.

Optical positioning is not limited to plasmonic particles. Optical forces have
been used to manipulate nanosheets [21] of hexagonal boron nitride (h-BN),
865 molybdenum disulfide (MoS_2) and tungsten disulfide (WS_2) obtained by liquid
phase exfoliation (LPE). Whereas the weakly absorbing h-BN can be optically
trapped, allowing the all-optical measurement of the mean flake size directly in
liquid environment, the strongly absorbing MoS_2 and WS_2 nanosheets cannot
be confined in the optical trap, due to overwhelming contribution of radiation
870 pressure on gradient force. Thanks to the consequent optical pushing effect,

these nanosheets can be used, in association with BSA, to pattern a common microscope slide, without any special surface pre-treatment, in relatively short times (minutes) and directly in liquid environment [21]. Possible applications concern the realization of nanosheet-based devices of interest in photonics [243],
875 optoelectronics [244], and energy storage [245], or of van der Waals heterostructures [246] in which nanosheets of different composition can be stacked on top of each other.

4.7. Critical Casimir Forces

The miniaturisation of devices and machines down to the microscopic and
880 nanoscopic scale involves also forces in a more challenging way. Forces of the order of nano- and piconewton need to be generated and applied with nanometric precision. This is challenging because of the presence of thermal fluctuations. Thermal fluctuations are often seen as a nuisance making the behaviour of micro-engines less deterministic and less predictable [201]. However, under
885 appropriate conditions, fluctuations can produce tunable and localized *critical Casimir forces* to drive nanomachines.

Critical Casimir forces are due to the confinement between multiple objects of the density fluctuations in a critical solution close to its critical temperature. One of the most important and useful features of these forces is their dependence
890 on temperature and on the surface properties of the involved objects. Critical Casimir forces were originally predicted in 1978 da Fisher and De Gennes [248], and they have been measured directly only recently in 2008 [249], which spurred a widespread interest in studying their properties with experiments [250, 251, 252].

895 Optical tweezers represent, again, an important tool for the measurements and characterisation of critical Casimir forces in nanodevices. Critical fluctuations can be confined by trapped objects, using HOT, giving rise to critical Casimir forces. Increasing the temperature of the solution toward the critical point, it is possible to characterise the temperature dependence of these forces
900 by the displacement of the confining objects. This displacement can be quan-




Figure 8: Effects of critical Casimir forces on the interdistance $r(t)$ of a pair of micro particles, diffusing freely in a water-2.6-lutidine solution at different temperatures. (a, b) At the temperature $T=33^\circ\text{C}$ lower than $T_c \approx 34^\circ\text{C}$, Critical Casimir forces are not affecting (a) the time distribution $r(t)$ and (b) its spatial distribution $P(r)$ which is Gaussian. (c, d) Increasing the temperature of the solution to $T=33,6^\circ\text{C}$, critical Casimir forces start to arise affecting (c) $r(t)$ and (d) $P(r)$, which now presents an incipient peak at lower value of r . (e, f) This effect became dominant at the temperature $T=33,9^\circ\text{C}$, very close to T_c . Here critical Casimir forces prevent the free diffusion of the particles, which are stuck together as showed in panel (e) where all the values of the $r(t)$ are confined within a small region resulting from the equilibrium between attractive critical Casimir forces and repulsive electrostatic forces. (f) At this temperature also the spatial distribution $P(r)$ is strongly affected by critical Casimir forces, presenting a narrow and intense peak corresponding to the equilibrium distance between the particle described before. The solid lines in panels (a,c,e) indicate representative experimental interparticle trajectories while the background colors represents the probability distribution of 400 different $r(t)$, while the dashed horizontal line indicates the interparticle distance corresponding to the diameter of the colloids. Solid lines in panels (b,d,f) are the theoretical distribution of r , obtained via Monte Carlo integration (10^6 samples) of two optically trapped particles subjected to the theoretical total potential[247].

tified by digital video microscopy [110] or by photonic force microscopy [111]. Both of these techniques match the spatial and time domains of critical Casimir forces but HOT combined with digital video microscopy provides a more flexible investigation tool.

905 Recently, blinking HOT and digital video microscopy have been employed to investigate the effects of critical Casimir forces on the dynamics of a pair of colloidal particles in the absence of optical potentials and dispersed in the bulk of a water-2,6-lutidine critical mixture [252, 247]. From the measurement of the time dependence of their center-to-center distance, it was determined
910 the relative diffusion and drift velocity of the particles, which are in agreement with the results of numerical simulations of Brownian dynamics based on the critical Casimir forces (Fig. 8) [247]. The comparison with these simulations allowed to infer the correlation length of the critical fluctuations, and therefore the intensity of the critical Casimir forces and the distance from the critical
915 temperature, forming the basis of a protocol for the fine tuning of the critical Casimir force field. This possibility provided new opportunities for the design and the realisation of self-assembled nano-structures and nanodevices.

4.8. Stochastic thermodynamics

Optical tweezers can be also employed as a very powerful tool to unveil and
920 characterise the statistical properties of micro- and nanoscopic systems. Here Brownian noise and large thermal fluctuations play a crucial role by introducing stochasticity [254, 255]. In particular, the dynamics of optically trapped particles results from the interplay between deterministic optical force fields and Brownian motion, which introduces a well-defined noisy background. Therefore,
925 optically trapped particles can be employed as probes to investigate statistical physics phenomena, whose dynamics are driven by both random and deterministic forces, ranging from biomolecules and nanodevices to financial markets and human organisations.

For the last two decades, optical tweezers have contributed to develop nan-
930 othermodynamics, investigating the statical properties of nanosystems to pre-




Figure 9: Emergence of crossover from Boltzmann to non-Boltzmann spatial distributions for a trapped particle in an active bath. Dots represent the distribution of an optically trapped particle, while dashed lines correspond to Gaussian distributions. in the case of not active bath, for increasing values of the stiffness (a) $k = 0.42 \text{ fN nm}^{-1}$, (b) $k = 3.6 \text{ fN nm}^{-1}$ and (c) $k = 22 \text{ fN nm}^{-1}$, the particle becomes more confined and its spatial distribution remains Gaussian. Vertical lines represent the error on six acquired trajectories. When the particle is trapped in an active bath, as the stiffness increases the particle is confined within a length scale comparable to the persistence length in the active bath (L_a red bar). In this case the particle distribution deviate from Gaussian and can be fitted with a heavy-tailed q -Gaussian distribution (solid black line) with (d) $q = 1.013$, (e) $q = 1.023$, and (f) $q = 1.142$ Panels (d-e). Figure adapted from [253]

dict the properties and performance of nanoobjects. For example, using trapped colloidal particles, it has been demonstrated that the presence of thermal fluctuations allows the violation of the second law of thermodynamics for small systems over short time scales, even though the second law still holds on average [255].

One of the biggest advances that can be achieved by applying stochastic thermodynamics to microscopic systems is the possibility to recover information about an equilibrium state of the system from measurements where the system is off equilibrium instead of just averaging the fluctuations out [256]. Stochastic thermodynamics can be successfully applied to different systems, such as living matters [257, 258]. For example, biomolecules are often coupled, with active baths, due to molecular motors inside the cytoplasm. This coupling is thought to lead to anomalous diffusion within the cytoplasm, a phenomenon that is largely not yet understood [257]. Recently, the behaviour of an optically trapped particle, immersed in an active bath, has been investigated by digital video microscopy to show a transition from Boltzmann to non-Boltzmann statistic [253] (see fig. 9). This transition takes place whenever the characteristic scale of an optical trap becomes comparable to the characteristic correlation length of the active noise. A consequence of this transition is that non-equilibrium relations such as the Jarzynski equality [259] and Crooks fluctuation theorem [260] cannot be applied in active baths according to their classical formulation [253]. Although this behaviour is unexpected in active system, its investigation is crucial to develop better models for living and far-from-equilibrium systems.

4.9. Active matter

Recently, optical tweezers have proved extremely useful for the characterisation of active matter systems and the study of the interplay between optical forces and thermophoresis. Active matter systems are those systems constituted by natural and artificial objects capable of self-propulsion [261]. Among this category there are systems with different size scales, from bacterial colonies to human crowds. Thanks to their ability in converting energy to propulsion, these

systems show behaviours like swarming and the emergence of other collective properties due to far-from equilibrium interactions [262]. This self-propelling behaviour is due to the interplay between random fluctuations (responsible for Brownian motion) and active swimming that drives them into a far-from-
965 equilibrium state. Although self-propulsion is a well-known feature in microorganisms to explore environments for nutrients or to run away from toxic substances [263], a lot of effort is still put in the realization of artificial self-propelling nano and microparticles. Two typical examples of self-propelling particles are artificial Janus particles and biological bacteria such as *Escherichia coli*, both
970 of them well characterised by optical tweezers [264, 265]. Furthermore, optical tweezers can be also employed to synchronize collective behaviour of active particles. In particular, this feature has been used to realize a self-assembled fluid pump by trapped active Brownian particles [266]. Optical tweezers can be also employed to perturb the environment of active matter modifying their collective
975 behaviour [267, 268]. Recently, it has been proved that the presence of spatial disorder can switch the behaviour of colloidal active matter from aggregative to dispersive [269]. Colloidal particles at equilibrium are always gathered by any attractive potential, but in presence of non-equilibrium driving forces, the particles disperse when spatial disorder is added to the potential. In particular,
980 it has been shown that disordered optical potentials can alter the long-term dynamics of colloidal particles inside an active bacterial bath changing their collective behaviour from aggregative to dispersive [269].

5. Conclusions

Optical tweezers are nowadays a key tool for the contactless manipulation
985 of a wide variety of samples at the micro and nanoscale. The modeling of optical tweezers with accurate T-matrix methods has brought advances in the understanding of experiments on non-spherical particles and the role of shape. Here we discussed some fundamental aspects of the theory and experimental practice of optical tweezers with a focus on a selection of applications in biology,

990 spectroscopy, optical positioning, and nanothermodynamics. After almost 50
years since the pioneering experiments by Arthur Ashkin on optical forces on
microparticles, this exciting field is still acquiring momentum expanding its
treads to exciting developments from the life sciences [270], to nano-engines
[271], and quantum technologies [272].

995 6. Acknowledgements

This paper is dedicated to the memory of Ferdinando Borghese, great teacher
of light scattering theory, quantum mechanics, and much more.

References

- [1] J. Kepler, *De cometis libelli tres* (1619).
- 1000 [2] A. Ashkin, Atomic-beam deflection by resonance-radiation pressure, *Physical Review Letters* 25 (19) (1970) 1321.
- [3] A. Ashkin, Acceleration and trapping of particles by radiation pressure, *Physical Review Letters* 24 (4) (1970) 156.
- [4] A. Ashkin, History of optical trapping and manipulation of small-neutral
1005 particle, atoms, and molecules, *IEEE Journal of Selected Topics in Quantum Electronics* 6 (6) (2000) 841–856.
- [5] A. Ashkin, J. M. Dziedzic, J. E. Bjorkholm, S. Chu, Observation of a
single-beam gradient force optical trap for dielectric particles, *Optics Letters* 11 (5) (1986) 288–290.
- 1010 [6] P. H. Jones, O. M. Maragò, G. Volpe, *Optical tweezers: Principles and applications*, Cambridge University Press, Cambridge, UK, 2015.
- [7] C. J. Foot, *Atomic physics*, Oxford University Press, 2005.
- [8] K. Dholakia, T. Čižmár, Shaping the future of manipulation, *Nature Photonics* 5 (6) (2011) 335–342.

- 1015 [9] M. Padgett, R. Bowman, Tweezers with a twist, *Nature Photonics* 5 (6) (2011) 343–348.
- [10] O. M. Maragò, P. H. Jones, P. G. Gucciardi, G. Volpe, A. C. Ferrari, Optical trapping and manipulation of nanostructures, *Nature nanotechnology* 8 (11) (2013) 807.
- 1020 [11] A. Ashkin, J. M. Dziedzic, T. Yamane, Optical trapping and manipulation of single cells using infrared laser beams, *Nature* 330 (6150) (1987) 769–771.
- [12] A. Ashkin, J. M. Dziedzic, Optical trapping and manipulation of viruses and bacteria, *Science* 235 (1987) 1517–1521.
- 1025 [13] F. M. Fazal, S. M. Block, Optical tweezers study life under tension, *Nature Photonics* 5 (6) (2011) 318–321.
- [14] O. Maragò, P. Gucciardi, P. Jones, Photonic force microscopy: from femtonewton force sensing to ultra-sensitive spectroscopy, in: *Scanning Probe Microscopy in Nanoscience and Nanotechnology*, Springer, 2010, pp. 23–
- 1030 56.
- [15] S. N. Olof, J. A. Grieve, D. B. Phillips, H. Rosenkranz, M. L. Yallop, M. J. Miles, A. J. Patil, S. Mann, D. M. Carberry, Measuring nanoscale forces with living probes, *Nano Letters* 12 (11) (2012) 6018–6023.
- [16] P. Chaumet, M. Nieto-Vesperinas, Time-averaged total force on a dipolar sphere in an electromagnetic field, *Optics Letters* 25 (15) (2000) 1065–
- 1035 1067.
- [17] A. Ashkin, Forces of a single-beam gradient laser trap on a dielectric sphere in the ray optics regime, *Biophysical Journal* 61 (2) (1992) 569–582.
- 1040 [18] F. Borghese, P. Denti, R. Saija, M. A. Iatì, Optical trapping of nonspherical particles in the T-matrix formalism, *Optics Express* 15 (19) (2007) 11984–11998.

- [19] F. Borghese, P. Denti, R. Saija, M. A. Iatì, O. M. Maragò, Radiation torque and force on optically trapped linear nanostructures, *Physical Review Letters* 100 (16) (2008) 163903. 1045
- [20] A. Irrera, P. Artoni, R. Saija, P. G. Gucciardi, M. A. Iatì, F. Borghese, P. Denti, F. Iacona, F. Priolo, O. M. Marago, Size-scaling in optical trapping of silicon nanowires, *Nano Letters* 11 (11) (2011) 4879–4884.
- [21] M. G. Donato, E. Messina, A. Foti, T. Smart, P. Jones, M. A. Iatì, R. Saija, P. G. Gucciardi, O. M. Maragò, Optical trapping and optical force positioning of two-dimensional materials, *Nanoscale* 10 (2018) 1245–1255. doi:10.1039/C7NR06465A. 1050
- [22] F. Borghese, P. Denti, R. Saija, *Scattering from model nonspherical particles: Theory and applications to environmental physics*, Springer Science & Business Media, 2007. 1055
- [23] A. Callegari, M. Mijalkov, A. B. Gököz, G. Volpe, Computational toolbox for optical tweezers in geometrical optics, *JOSA B* 32 (5) (2015) B11–B19.
- [24] G. Volpe, G. Volpe, Simulation of a brownian particle in an optical trap, *American Journal of Physics* 81 (3) (2013) 224–230.
- [25] A. Ashkin, Forces of a single-beam gradient laser trap on a dielectric sphere in the ray optics regime, *Methods in Cell Biology* 55 (1997) 1–27. 1060
- [26] M. Mishchenko, L. Travis, A. Lacis, *Scattering, absorption, and emission of light by small particles*, Cambridge University Press, 2002.
- [27] P. M. Neto, H. Nussenzveig, Theory of optical tweezers, *EPL (Europhysics Letters)* 50 (5) (2000) 702. 1065
- [28] A. Mazolli, P. M. Neto, H. Nussenzveig, Theory of trapping forces in optical tweezers, in: *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, Vol. 459, The Royal Society, 2003, pp. 3021–3041.

- 1070 [29] T. A. Nieminen, A. B. Stilgoe, N. R. Heckenberg, H. Rubinsztein-Dunlop, Approximate and exact modeling of optical trapping, in: *Optical Trapping and Optical Micromanipulation VII*, Vol. 7762, International Society for Optics and Photonics, 2010, p. 77622V.
- [30] J. P. Gordon, Radiation forces and momenta in dielectric media, *Physical Review A* 8 (1) (1973) 14.
1075
- [31] J. R. Arias-González, M. Nieto-Vesperinas, Optical forces on small particles: attractive and repulsive nature and plasmon-resonance conditions, *JOSA A* 20 (7) (2003) 1201–1209.
- [32] V. Amendola, R. Pilot, M. Frasconi, O. M. Maragó, M. A. Iatì, Surface plasmon resonance in gold nanoparticles: A review, *Journal of Physics: Condensed Matter* 29 (20) (2017) 203002.
1080
- [33] M. Born, E. Wolf, *Principles of optics: Electromagnetic theory of propagation, interference and diffraction of light*, Cambridge University Press, Cambridge, United Kingdom, 1999.
- 1085 [34] S. E. Skelton, M. Sergides, G. Memoli, O. M. Maragó, P. H. Jones, Trapping and deformation of microbubbles in a dual-beam fibre-optic trap, *Journal of Optics* 14 (7) (2012) 075706.
- [35] P. Galajda, P. Ormos, Orientation of flat particles in optical tweezers by linearly polarized light, *Optics Express* 11 (5) (2003) 446–451.
- 1090 [36] P. Galajda, P. Ormos, Complex micromachines produced and driven by light, *Applied Physics Letters* 78 (2) (2001) 249–251.
- [37] G. A. Swartzlander, T. J. Peterson, A. B. Artusio-Glimpse, A. D. Raisanen, Stable optical lift, *Nature Photonics* 5 (1) (2011) 48–51.
- 1095 [38] K. Dholakia, P. Zemánek, Colloquium: Gripped by light: Optical binding, *Reviews of Modern Physics* 82 (2) (2010) 1767.

- [39] B. T. Draine, J. Goodman, Beyond clausius-mossotti-wave propagation on a polarizable point lattice and the discrete dipole approximation, *The Astrophysical Journal* 405 (1993) 685–697.
- [40] D. Gao, W. Ding, M. Nieto-Vesperinas, X. Ding, M. Rahman, T. Zhang, C. Lim, C.-W. Qiu, Optical manipulation from the microscale to the nanoscale: fundamentals, advances and prospects, *Light: Science & Applications* 6 (9) (2017) e17039.
- [41] S. Albaladejo, M. I. M. M. Laroche, J. J. Sáenz, Scattering forces from the curl of the spin angular momentum of a light field, *Physical Review Letters* 102 (11) (2009) 113602.
- [42] M. I. Marqués, J. J. Sáenz, Reply to comment on scattering forces from the curl of the spin angular momentum of a light field, *Physical Review Letters* 111 (5) (2013) 059302.
- [43] P. Zemánek, A. Jonáš, P. Ják, J. Ježek, M. Šerý, M. Liška, Theoretical comparison of optical traps created by standing wave and single beam, *Optics Communications* 220 (4-6) (2003) 401–412.
- [44] G. Volpe, G. P. Singh, D. Petrov, Optical tweezers with cylindrical vector beams produced by optical fibers, in: *Optical Trapping and Optical Micro-manipulation*, Vol. 5514, International Society for Optics and Photonics, 2004, pp. 283–293.
- [45] M. G. Donato, S. Vasi, R. Sayed, P. H. Jones, F. Bonaccorso, A. C. Ferrari, P. G. Gucciardi, M. O. M., Optical trapping of nanotubes with cylindrical vector beams, *Optics Letters* 37 (16) (2012) 3381–3383.
- [46] S. E. Skelton, M. Sergides, R. Saija, M. A. Iatì, O. M. Maragó, P. H. Jones, Trapping volume control in optical tweezers using cylindrical vector beams, *Optics Letters* 38 (1) (2013) 28–30.

- [47] M. I. Marqués, Beam configuration proposal to verify that scattering forces come from the orbital part of the poynting vector, *Optics Letters* 39 (17) (2014) 5122–5125.
- 1125 [48] M. V. Berry, Optical currents, *Journal of Optics A: Pure and Applied Optics* 11 (9) (2009) 094001.
- [49] K. Y. Bliokh, A. Y. Bekshaev, F. Nori, Extraordinary momentum and spin in evanescent waves, *Nature Communications* 5 (2014) 3300.
- 1130 [50] M. Antognozzi, C. R. Bermingham, R. L. Harniman, S. Simpson, J. Senior, R. Hayward, H. Hoerber, M. R. Dennis, A. Y. Bekshaev, K. Y. Bliokh, et al., Direct measurements of the extraordinary optical momentum and transverse spin-dependent force using a nano-cantilever, *Nature Physics* 12 (8) (2016) 731.
- 1135 [51] C. Triolo, A. Cacciola, S. Patanè, R. Saija, S. Savasta, F. Nori, Spin-momentum locking in the near field of metal nanoparticles, *ACS Photonics* 4 (9) (2017) 2242–2249.
- [52] M. I. Mishchenko, Radiation force caused by scattering, absorption, and emission of light by nonspherical particles, *Journal of Quantitative Spectroscopy and Radiative Transfer* 70 (4-6) (2001) 811–816.
- 1140 [53] T. A. Nieminen, H. Rubinsztein-Dunlop, N. R. Heckenberg, Calculation and optical measurement of laser trapping forces on non-spherical particles, *Journal of Quantitative Spectroscopy and Radiative Transfer* 70 (4-6) (2001) 627–637.
- 1145 [54] R. Saija, M. A. Iatì, A. Giusto, P. Denti, F. Borghese, Transverse components of the radiation force on nonspherical particles in the T-matrix formalism, *Journal of Quantitative Spectroscopy and Radiative Transfer* 94 (2) (2005) 163–179.

- [55] R. N. Pfeifer, T. A. Nieminen, N. R. Heckenberg, H. Rubinsztein-Dunlop,
Colloquium: Momentum of an electromagnetic wave in dielectric media,
1150 *Reviews of Modern Physics* 79 (4) (2007) 1197.
- [56] S. H. Simpson, Inhomogeneous and anisotropic particles in optical
traps: Physical behaviour and applications, *Journal of Quantitative Spec-*
troscopy and Radiative Transfer 146 (2014) 81–99.
- [57] T. A. Nieminen, N. D. Preez-Wilkinson, A. B. Stilgoe, V. L. Y. Loke,
1155 A. A. M. Bui, H. Rubinsztein-Dunlop, Optical tweezers: Theory and mod-
elling, *Journal of Quantitative Spectroscopy and Radiative Transfer* 146
(2014) 59–80.
- [58] X. Qi, T. A. Nieminen, A. B. Stilgoe, V. L. Loke, H. Rubinsztein-Dunlop,
Comparison of t-matrix calculation methods for scattering by cylinders in
1160 optical tweezers, *Optics letters* 39 (16) (2014) 4827–4830.
- [59] A. A. Bui, A. B. Stilgoe, I. C. Lenton, L. J. Gibson, A. V. Kashchuk,
S. Zhang, H. Rubinsztein-Dunlop, T. A. Nieminen, Theory and practice
of simulation of optical tweezers, *Journal of Quantitative Spectroscopy*
and Radiative Transfer 195 (2017) 66–75.
- 1165 [60] C. Girard, Near fields in nanostructures, *Reports on progress in physics*
68 (8) (2005) 1883.
- [61] M. A. Yurkin, A. G. Hoekstra, The discrete dipole approximation: an
overview and recent developments, *Journal of Quantitative Spectroscopy*
and Radiative Transfer 106 (1-3) (2007) 558–589.
- 1170 [62] K. Yee, Numerical solution of initial boundary value problems involving
maxwell's equations in isotropic media, *IEEE Transactions on antennas*
and propagation 14 (3) (1966) 302–307.
- [63] A. Z. Elsherbeni, V. Demir, The finite-difference time-domain method for
electromagnetics with MATLAB simulations, *The Institution of Engineer-*
ing and Technology, 2016.
- 1175

- [64] P. Waterman, Symmetry, unitarity, and geometry in electromagnetic scattering, *Physical Review D* 3 (4) (1971) 825.
- [65] G. Gouesbet, J. Lock, G. Gréhan, Generalized lorenz–mie theories and description of electromagnetic arbitrary shaped beams: localized approx-
1180 imations and localized beam models, a review, *Journal of Quantitative Spectroscopy and Radiative Transfer* 112 (1) (2011) 1–27.
- [66] E. M. Purcell, C. R. Pennypacker, Scattering and absorption of light by nonspherical dielectric grains, *The Astrophysical Journal* 186 (1973) 705–714.
- [67] B. T. Draine, P. J. Flatau, Discrete-dipole approximation for scattering
1185 calculations, *JOSA A* 11 (4) (1994) 1491–1499.
- [68] S. H. Simpson, S. Hanna, Application of the discrete dipole approximation to optical trapping calculations of inhomogeneous and anisotropic particles, *Optics Express* 19 (17) (2011) 16526–16541.
- [69] S. H. Simpson, S. Hanna, Holographic optical trapping of microrods and
1190 nanowires, *JOSA A* 27 (6) (2010) 1255–1264.
- [70] D. Phillips, M. Padgett, S. Hanna, Y.-L. Ho, D. Carberry, M. Miles, S. Simpson, Shape-induced force fields in optical trapping, *Nature Photonics* 8 (5) (2014) 400.
- [71] R. C. Gauthier, Computation of the optical trapping force using an FDTD
1195 based technique, *Optics Express* 13 (10) (2005) 3707–3718.
- [72] D. C. Benito, S. H. Simpson, S. Hanna, FDTD simulations of forces on particles during holographic assembly, *Optics express* 16 (5) (2008) 2942–2957.
- [73] X. Yang, Y. Liu, R. F. Oulton, X. Yin, X. Zhang, Optical forces in hybrid
1200 plasmonic waveguides, *Nano Letters* 11 (2) (2011) 321–328.

- [74] M. A. Zaman, P. Padhy, L. Hesselink, A semi-analytical model of a near-field optical trapping potential well, *Journal of Applied Physics* 122 (16) (2017) 163101.
- 1205 [75] F. Borghese, P. Denti, R. Saija, G. Toscano, O. I. Sindoni, Multiple electromagnetic scattering from a cluster of spheres. I. Theory, *Aerosol Science and Technology* 3 (2) (1984) 227–235.
- [76] R. Saija, M. A. Iatì, P. Denti, F. Borghese, A. Giusto, O. I. Sindoni, Efficient light-scattering calculations for aggregates of large spheres, *Applied Optics* 42 (15) (2003) 2785–2793.
- 1210 [77] R. Saija, P. Denti, F. Borghese, O. M. Maragó, M. A. Iatì, Optical trapping calculations for metal nanoparticles: Comparison with experimental data for Au and Ag spheres., *Optics Express* 17 (2009) 10231–10241.
- [78] D. Spadaro, M. Iatì, J. Pérez-Piñero, C. Vazquez-Vazquez, M. Correa-Duarte, M. Donato, P. Gucciardi, R. Saija, G. Strangi, O. Marago, Optical trapping of plasmonic mesocapsules: Enhanced optical forces and sers, *The Journal of Physical Chemistry C* 121 (1) (2016) 691–700.
- 1215 [79] V. Amendola, R. Pilot, M. Frascioni, O. M. Marago, M. A. Iati, Surface plasmon resonance in gold nanoparticles: a review, *Journal of Physics: Condensed Matter* 29 (20) (2017) 203002.
- 1220 [80] P. J. Wyatt, Scattering of electromagnetic plane waves from inhomogeneous spherically symmetric objects, *Physical Review* 134 (7AB) (1964) AB1.
- [81] D. Spadaro, M. A. Iatì, M. G. Donato, P. G. Gucciardi, R. Saija, A. R. Cherlakola, S. Scaramuzza, V. Amendola, O. M. Maragò, Scaling of optical forces on au–peg core–shell nanoparticles, *RSC Advances* 5 (113) (2015) 93139–93146.
- 1225 [82] F. Borghese, P. Denti, R. Saija, O. Sindoni, Optical properties of spheres containing a spherical eccentric inclusion, *JOSA A* 9 (8) (1992) 1327–1335.

- 1230 [83] E. Fucile, P. Denti, F. Borghese, R. Saija, O. I. Sindoni, Optical properties
of a sphere in the vicinity of a plane surface, *Journal of the Optical Society
of America A* 14 (7) (1997) 1505–1514.
- [84] F. Borghese, P. Denti, R. Saija, M. A. Iati, O. I. Sindoni, Optical proper-
ties of a dispersion of anisotropic particles with non-randomly distributed
1235 orientations. The case of atmospheric ice crystals, *Journal of Quantitative
Spectroscopy and Radiative Transfer* 70 (2) (2001) 237–251.
- [85] D. W. Mackowski, M. I. Mishchenko, Calculation of the T-matrix and
the scattering matrix for ensembles of spheres, *JOSA A* 13 (11) (1996)
2266–2278.
- 1240 [86] G. Mie, Beiträge zur optik trüber medien, speziell kolloidaler met-
allösungen, *Annalen der Physik* 330 (3) (1908) 377–445.
- [87] G. Gouesbet, G. Gréhan, *Generalized lorenz-mie theories*, Springer, 2017.
- [88] G. Gouesbet, T-matrix formulation and generalized lorenz-mie theories
in spherical coordinates, *Optics communications* 283 (4) (2010) 517–521.
- 1245 [89] J. A. Lock, Calculation of the radiation trapping force for laser tweezers
by use of generalized lorenz-mie theory. i. localized model description of
an on-axis tightly focused laser beam with spherical aberration, *Applied
optics* 43 (12) (2004) 2532–2544.
- [90] A. A. R. Neves, A. Fontes, L. de Y. Pozzo, A. A. D. Thomaz, E. Chillce,
1250 E. Rodriguez, L. C. Barbosa, C. L. Cesar, Electromagnetic forces for an
arbitrary optical trapping of a spherical dielectric, *Optics Express* 14 (26)
(2006) 13101–13106.
- [91] A. A. R. Neves, A. Fontes, C. L. Cesar, A. Camposeo, R. Cingolani,
1255 D. Pisignano, Axial optical trapping efficiency through a dielectric inter-
face, *Physical review E* 76 (6) (2007) 061917.

- [92] L. A. Ambrosio, H. E. Hernández-Figueroa, Integral localized approximation description of ordinary bessel beams and application to optical trapping forces, *Biomedical Optics Express* 2 (7) (2011) 1893–1906.
- [93] A. A. R. Neves, Photonic nanojets in optical tweezers, *Journal of Quantitative Spectroscopy and Radiative Transfer* 162 (2015) 122–132. 1260
- [94] A. Ishimaru, Wave propagation and scattering in random media and rough surfaces, *Proceedings of the IEEE* 79 (10) (1991) 1359–1366.
- [95] F. Borghese, P. Denti, R. Saija, M. A. Iatì, Radiation torque on nonspherical particles in the transition matrix formalism, *Optics Express* 14 (20) (2006) 9508–9521. 1265
- [96] F. Borghese, P. Denti, R. Saija, M. A. Iatì, On the rotational stability of nonspherical particles driven by the radiation torque, *Optics Express* 15 (14) (2007) 8960–8971.
- [97] P. L. Marston, J. H. Crichton, Radiation torque on a sphere caused by a circularly-polarized electromagnetic wave, *Physical Review A* 30 (5) (1984) 2508. 1270
- [98] E. L. Wolf, *Nanophysics and nanotechnology: An introduction to modern concepts in nanoscience*, John Wiley & Sons, 2015.
- [99] A. P. Alivisatos, Semiconductor clusters, nanocrystals and quantum dots, *Science* 271 (5251) (1996) 933–937. 1275
- [100] M. S. Gudiksen, L. J. Lauhon, J. Wang, D. C. Smith, C. M. Lieber, Growth of nanowire superlattice structures for nanoscale photonics and electronics, *Nature* 415 (6872) (2002) 617.
- [101] K. A. Dick, A review of nanowire growth promoted by alloys and non-alloying elements with emphasis on Au-assisted III–V nanowires, *Progress in Crystal Growth and Characterization of Materials* 54 (3-4) (2008) 138–173. 1280

- 1285 [102] A. Rohrbach, Stiffness of optical traps: Quantitative agreement between
experiment and electromagnetic theory, *Physical review letters* 95 (16)
(2005) 168102.
- [103] N. Sultanova, S. Kasarova, I. Nikolov, Characteristics of optical polymers
in the design of polymer and hybrid optical systems, *Bulgarian Journal of
Physics* 40 (3) (2013) 258–264.
- 1290 [104] O. M. Marago, P. H. Jones, F. Bonaccorso, V. Scardaci, P. G. Gucciardi,
A. G. Rozhin, A. C. Ferrari, Femtonewton force sensing with optically
trapped nanotubes, *Nano Letters* 8 (10) (2008) 3211–3216.
- [105] G. Pesce, G. Volpe, O. M. Maragò, P. H. Jones, S. Gigan, A. Sasso,
G. Volpe, Step-by-step guide to the realization of advanced optical twee-
ers, *Journal of the Optical Society of America B* 32 (5) (2015) B84–B98.
- 1295 [106] A. Briggs, O. Kolosov, *Acoustic microscopy*, Vol. 67, Oxford University
Press, 2010.
- [107] A. Irrera, A. Magazzù, P. Artoni, S. H. Simpson, S. Hanna, P. H. Jones,
F. Priolo, P. G. Gucciardi, O. M. Maragó, Photonic torque microscopy
of the nonconservative force field for optically trapped silicon nanowires,
1300 *Nano Letters* 16 (7) (2016) 4181–4188.
- [108] J. E. Curtis, B. A. Koss, D. G. Grier, Dynamic holographic optical twee-
ers, *Optics communications* 207 (1-6) (2002) 169–175.
- 1305 [109] O. Otto, F. Czerwinski, J. L. Gornall, G. Stober, L. B. Oddershede, R. Sei-
del, U. F. Keyser, Real-time particle tracking at 10,000 fps using optical
fiber illumination, *Optics express* 18 (22) (2010) 22722–22733.
- [110] J. C. Crocker, D. G. Grier, Methods of digital video microscopy for col-
loidal studies, *Journal of colloid and interface science* 179 (1) (1996) 298–
310.

- [111] E.-L. Florin, A. Pralle, E. Stelzer, J. Hörber, Photonic force microscope calibration by thermal noise analysis, *Applied Physics A: Materials Science & Processing* 66 (1998) S75–S78.
- [112] O. M. Maragó, F. Bonaccorso, R. Saija, G. Privitera, P. G. Gucciardi, M. A. Iati, G. Calogero, P. H. Jones, F. Borghese, P. Denti, et al., Brownian motion of graphene, *ACS Nano* 4 (12) (2010) 7515–7523.
- [113] S. H. Simpson, S. Hanna, First-order nonconservative motion of optically trapped nonspherical particles, *Physical Review E* 82 (2010) 031141.
- [114] W. J. Toe, I. Ortega-Piwonka, C. N. Angstmann, Q. Gao, H. H. Tan, C. Jagadish, B. I. Henry, P. J. Reece, Nonconservative dynamics of optically trapped high-aspect-ratio nanowires, *Physical Review E* 93 (2) (2016) 022137.
- [115] B. Mihiretie, P. Snabre, J.-C. Loudet, B. Pouligny, Optically driven oscillations of ellipsoidal particles. part i: Experimental observations, *The European Physical Journal E* 37 (12) (2014) 124.
- [116] Z. Cheng, T. G. Mason, P. Chaikin, Periodic oscillation of a colloidal disk near a wall in an optical trap, *Physical Review E* 68 (5) (2003) 051404.
- [117] A. Magazzú, D. Spadaro, M. Donato, R. Sayed, E. Messina, C. D’Andrea, A. Foti, B. Fazio, M. Iatí, A. Irrera, R. Saija, P. G. Gucciardi, O. M. Marago, Optical tweezers: A non-destructive tool for soft and biomaterial investigations, *Rendiconti Lincei* 26 (2) (2015) 203–218.
- [118] A. Buosciolo, G. Pesce, A. Sasso, New calibration method for position detector for simultaneous measurements of force constants and local viscosity in optical tweezers, *Optics communications* 230 (4) (2004) 357–368.
- [119] W. Coffey, Y. P. Kalmykov, J. T. Waldron, *The Langevin equation: with applications to stochastic problems in physics, chemistry, and electrical engineering*, Vol. 14, World Scientific, 2004.

- [120] M. Le Bellac, F. Mortessagne, G. G. Batrouni, Equilibrium and non-equilibrium statistical thermodynamics, Cambridge University Press, 2004.
- [121] R. Bar-Ziv, A. Meller, T. Tlusty, E. Moses, J. Stavans, S. A. Safran,
1340 Localized dynamic light scattering: probing single particle dynamics at
the nanoscale, *Physical Review Letters* 78 (1) (1997) 154.
- [122] S. Henderson, S. Mitchell, P. Bartlett, Position correlation microscopy:
probing single particle dynamics in colloidal suspensions, *Colloids and
Surfaces A: Physicochemical and Engineering Aspects* 190 (1) (2001) 81–
1345 88.
- [123] A. D. Mehta, M. Rief, J. A. Spudich, D. A. Smith, R. M. Simmons, Single
molecule biomechanics with optical methods, *Science* 283 (1999) 1689–
1695.
- [124] T. T. Perkins, Optical traps for single molecule biophysics: a primer, *Laser
1350 Photon. Rev.* 3 (2009) 203–220.
- [125] K. Svoboda, S. M. Block, Biological applications of optical forces, *Annu.
Rev. Biophys. Biomol. Struct.* 23 (1994) 247–285.
- [126] H. D. Ou-Yang, M. T. Wei, Complex fluids: probing mechanical properties
of biological systems with optical tweezers, *Ann. Rev. Phys. Chem.* 61
1355 (2010) 421–440.
- [127] S. Hénon, G. Lenormand, A. Richert, F. Gallet, A new determination
of the shear modulus of the human erythrocyte membrane using optical
tweezers, *Biophysical Journal* 76 (2) (1999) 1145 – 1151.
- [128] M. Dao, C. T. Lim, S. Suresh, Mechanics of the human red blood cell
1360 deformed by optical tweezers, *J. Mech. Phys. Solids* 51 (2003) 2259–2280.
- [129] P. J. Bronkhorst, G. J. Streekstra, J. Grimbergen, E. J. Nijhof, J. J. Sixma,
G. J. Brakenhoff, A new method to study shape recovery of red blood cells
using multiple optical trapping, *Biophys. J.* 69 (5) (1995) 1666–1673.

- 1365 [130] G.-B. Liao, P. B. Bareil, Y. Sheng, A. Chiou, One-dimensional jumping
optical tweezers for optical stretching of bi-concave human red blood cells,
Opt. Express 16 (3) (2008) 1996–2004.
- [131] S. Rancourt-Grenier, M.-T. Wei, J.-J. Bai, A. Chiou, P. P. Bareil, P.-
L. Duval, Y. Sheng, Dynamic deformation of red blood cell in dual-trap
optical tweezers, Opt. Express 18 (10) (2010) 10462–10472.
- 1370 [132] L. Yu, Y. Sheng, A. Chiou, Three-dimensional light-scattering and defor-
mation of individual biconcave human blood cells in optical tweezers, Opt.
Express 21 (10) (2013) 12174–12184.
- [133] Y. Yoon, J. Kotar, G. Yoon, P. Cicuta, The nonlinear mechanical response
of the red blood cell, Physical Biology 5 (3) (2008) 036007.
- 1375 [134] Y. Z. Yoon, J. Kotar, A. T. Brown, P. Cicuta, Red blood cell dynamics:
from spontaneous fluctuations to non-linear response, Soft Matter 7 (2011)
2042–2051.
- [135] L. Yu, Y. Sheng, Effect of the object 3d shape on the viscoelastic testing
in optical tweezers, Opt. Express 23 (5) (2015) 6020–6028.
- 1380 [136] S. Raj, M. Marro, M. Wojdyla, D. Petrov, Mechanochemistry of single
red blood cells monitored using raman tweezers, Biomed. Opt. Express
3 (4) (2012) 753–763.
- [137] J. Guck, R. Ananthakrishnan, H. Mahmood, T. J. Moon, C. C. Cunning-
ham, J. Käs, The optical stretcher: A novel laser tool to micromanipulate
1385 cells, Biophysical Journal 81 (2) (2001) 767 – 784.
- [138] P. B. Bareil, Y. Sheng, Y.-Q. Chen, A. Chiou, Calculation of spherical
red blood cell deformation in a dual-beam optical stretcher, Opt. Express
15 (24) (2007) 16029–16034.
- 1390 [139] A. E. Ekpenyong, C. L. Posey, J. L. Chaput, A. K. Burkart, M. M.
Marquardt, T. J. Smith, M. G. Nichols, Determination of cell elastic-

- ity through hybrid ray optics and continuum mechanics modeling of cell deformation in the optical stretcher, *Appl. Opt.* 48 (32) (2009) 6344–6354.
- [140] J. M. A. Mauritz, A. Esposito, T. Tiffert, J. N. Skepper, A. Warley, Y.-Z. Yoon, P. Cicuta, V. L. Lew, J. R. Guck, C. F. Kaminski, Biophotonic techniques for the study of malaria-infected red blood cells, *Medical & Biological Engineering & Computing* 48 (10) (2010) 1055–1063.
- [141] S. Lee, B. Joo, P. J. Jeon, S. Im, K. Oh, Columnar deformation of human red blood cell by highly localized fiber optic bessel beam stretcher, *Biomed. Opt. Express* 6 (11) (2015) 4417–4432.
- [142] S. Suresh, Mechanical response of human red blood cells in health and disease: Some structure-property-function relationships, *Journal of Materials Research* 21 (8) (2006) 1871–1877.
- [143] R. Agrawal, J. Sherwood, J. Chhablani, A. Ricchariya, S. Kim, P. H. Jones, S. Balabani, D. Shima, Red blood cells in retinal vascular disorders, *Blood Cells, Molecules, and Diseases* 56 (1) (2016) 53 – 61.
- [144] S. Suresh, J. Spatz, J. Mills, A. Micoulet, M. Dao, C. Lim, M. Beil, T. Seufferlein, Single-cell biomechanics and human disease states: Gastrointestinal cancer and malaria, *Acta Biomater.* 1 (2005) 16.
- [145] J. M. Mauritz, T. Tiffert, R. Seear, F. Lautenschläger, A. Esposito, V. L. Lew, J. R. Guck, C. F. Kaminski, Detection of plasmodium falciparum-infected red blood cells by optical stretching, *Journal of Biomedical Optics* 15 (3) (2010) 030517.
- [146] R. Agrawal, T. J. Smart, J. Nobre-Cardoso, C. J. Richards, R. Bhatnagar, A. Tufail, D. Shima, P. H. Jones, C. Pavesio, Assessment of red blood cell deformability in type 2 diabetes mellitus and diabetic retinopathy by dual optical tweezers stretching technique, *Scientific Reports* 6 (2016) 15873.
- [147] R. Agrawal, B. Ang, J. Nobre-Cardoso, C. J. Richards, T. J. Smart, R. Bhatnagar, D. Shima, P. H. Jones, C. Pavesio, Non-occlusive retinal

- vascular inflammation and role of red blood cell deformability in birdshot
1420 chorioretinopathy Accepted for publication in Ocular Immunology and In-
flammation.
- [148] V. Sheikh-Hasani, M. Babaei, A. Azadbakht, H. Pazoki-Toroudi,
A. Mashaghi, A. A. Moosavi-Movahedi, S. N. S. Reihani, Atorvastatin
treatment softens human red blood cells: an optical tweezers study,
1425 Biomed. Opt. Express 9 (3) (2018) 1256–1261.
- [149] P. A. Prentice, M. P. MacDonald, T. G. Frank, A. Cuschieri, G. C. Spald-
ing, W. Sibbett, P. A. Campbell, K. Dholakia, Manipulation and filtration
of low index particles with holographic Laguerre-Gaussian optical trap ar-
rays, Opt. Express 12 (4) (2004) 593–600.
- 1430 [150] J. Arlt, M. Padgett, Generation of a beam with a dark focus surrounded
by regions of higher intensity: the optical bottle beam, Optics Letters
25 (4) (2000) 191–193.
- [151] P. H. Jones, E. Stride, N. Saffari, Trapping and manipulation of micro-
scopic bubbles with a scanning optical tweezer, Applied Physics Letters
1435 89 (8) (2006) 081113.
- [152] P. H. Jones, O. M. Maragó, E. P. J. Stride, Parametrization of trapping
forces on microbubbles in scanning optical tweezers, Journal of Optics A:
Pure and Applied Optics 9 (8) (2007) S278.
- [153] S.-Y. Sung, Y.-G. Lee, Trapping of a micro-bubble by non-paraxial Gaus-
sian beam: computation using the FDTD method, Opt. Express 16 (5)
1440 (2008) 3463–3473.
- [154] T. J. Smart, M. B. Unlu, P. H. Jones, Microbubble trapping in inverted
optical tweezers, Vol. 10347, 2017, pp. 1034731–1 – 1034731–11.
- 1445 [155] C. Fury, C. Harfield, P. H. Jones, E. Stride, G. Memoli, Experimental
characterisation of holographic optical traps for microbubbles, Vol. 9126,
2014, p. 91263L.

- [156] C. Fury, P. H. Jones, G. Memoli, Multiscale manipulation of microbubbles employing simultaneous optical and acoustical trapping, Vol. 9164, 2014, pp. 9164 – 9164 – 11.
- 1450 [157] G. Memoli, C. R. Fury, K. O. Baxter, P. N. Glat, P. H. Jones, Acoustic force measurements on polymer-coated microbubbles in a microfluidic device, *The Journal of the Acoustical Society of America* 141 (5) (2017) 3364–3378.
- 1455 [158] C. J. Harfield, C. R. Fury, G. Memoli, P. H. Jones, N. Ovenden, E. P. J. Stride., Analysis of the uncertainty in microbubble characterisation, *Ultrasound in Medicine and Biology* 42 (2016) 1412–1418.
- [159] M. Lankers, J. Popp, G. Rosling, W. Kiefer, Raman investigations on laser-trapped gas bubbles, *Chemical Physics Letters* 277 (4) (1997) 331 – 334.
- 1460 [160] F. Saglimbeni, S. Bianchi, G. Bolognesi, G. Paradossi, R. Di Leonardo, Optical characterization of an individual polymer-shelled microbubble structure via digital holography, *Soft Matter* 8 (2012) 8822–8825.
- [161] P. Prentice, A. Cuschieri, K. Dholakia, M. Prausnitz, P. Campbell, Membrane disruption by optically controlled microbubble cavitation, *Nature Physics* 1 (2005) 107–110.
- 1465 [162] R. A. Beth, Mechanical detection and measurement of the angular momentum of light, *Physical Review* 50 (2) (1936) 115.
- [163] A. Canaguier-Durand, J. A. Hutchison, C. Genet, T. W. Ebbesen, Mechanical separation of chiral dipoles by chiral light, *New Journal of Physics* 15 (12) (2013) 123037.
- 1470 [164] A. Hayat, J. B. Mueller, F. Capasso, Lateral chirality-sorting optical forces, *Proceedings of the National Academy of Sciences* 112 (43) (2015) 13190–13194.

- [165] I. D. Rukhlenko, N. V. Tepliakov, A. S. Baimuratov, S. A. Andronaki,
1475 Y. K. Gunko, A. V. Baranov, A. V. Fedorov, Completely chiral optical
force for enantioseparation, *Scientific Reports* 6 (2016) 36884.
- [166] M. E. J. Friese, T. A. Nieminen, N. R. Heckenberg, H. Rubinsztein-
Dunlop, Optical alignment and spinning of laser-trapped microscopic par-
ticles, *Nature* 394 (1998) 348–350.
- 1480 [167] A. I. Bishop, T. A. Nieminen, N. R. Heckenberg, H. Rubinsztein-Dunlop,
Optical microrheology using rotating laser-trapped particles, *Physical Re-
view Letters* 92 (19) (2004) 198104.
- [168] J. S. Bennett, L. J. Gibson, R. M. Kelly, E. Brousse, B. Baud-
isch, D. Preece, T. A. Nieminen, T. Nicholson, N. R. Heckenberg,
1485 H. Rubinsztein-Dunlop, Spatially-resolved rotational microrheology with
an optically-trapped sphere, *Scientific reports* 3 (2013) 1759.
- [169] Y. Arita, M. Mazilu, K. Dholakia, Laser-induced rotation and cooling of
a trapped microgyroscope in vacuum, *Nature communications* 4 (2013)
2374.
- 1490 [170] G. Tkachenko, E. Brasselet, Spin controlled optical radiation pressure,
Physical Review Letters 111 (2013) 033605.
- [171] G. Tkachenko, E. Brasselet, Optofluidic sorting of material chirality by
chiral light, *Nature Communications* 5.
- [172] G. Tkachenko, E. Brasselet, Helicity-dependent three-dimensional optical
1495 trapping of chiral microparticles, *Nature communications* 5 (2014) 4491.
- [173] R. J. Hernández, A. Mazzulla, A. Pane, K. Volke-Sepúlveda, G. Cipparrone,
Attractive-repulsive dynamics on light-responsive chiral microparti-
cles induced by polarized tweezers, *Lab on a Chip* 13 (2013) 459–467.
- 1500 [174] R. Hernández, A. Mazzulla, C. Provenzano, P. Pagliusi, G. Cipparrone,
Chiral resolution of spin angular momentum in linearly polarized and
unpolarized light, *Scientific reports* 5 (2015) 16926.

- [175] G. Cipparrone, A. Mazzulla, A. Pane, R. J. Hernandez, R. Bartolino, Chiral self-assembled solid microspheres: A novel multifunctional microphonic device, *Advanced Materials* 23 (2011) 5773–5778.
- 1505 [176] M. G. Donato, J. Hernandez, A. Mazzulla, C. Provenzano, R. Saija, R. Sayed, S. Vasi, A. Magazzù, P. Pagliusi, R. Bartolino, P. G. Gucciardi, O. M. Maragó, G. Cipparrone, Polarization-dependent optomechanics mediated by chiral microresonators, *Nature Communications* 5 (2014) 3656.
- [177] M. Donato, A. Mazzulla, P. Pagliusi, A. Magazzù, R. Hernandez, 1510 C. Provenzano, P. Gucciardi, O. Maragó, G. Cipparrone, Light-induced rotations of chiral birefringent microparticles in optical tweezers, *Scientific Reports* 6 (2016) 31977.
- [178] D. Seč, T. Porenta, M. Ravnik, S. Žumer, Geometrical frustration of chiral ordering in cholesteric droplets, *Soft Matter* 8 (48) (2012) 11982–11988.
- 1515 [179] F. Schmidt, A. Magazzù, A. Callegari, L. Biancofiore, F. Cichos, G. Volpe, Microscopic engine powered by critical demixing, *Physical Review Letters* 120 (6) (2018) 068004.
- [180] L. K. Abdelmohsen, F. Peng, Y. Tu, D. A. Wilson, Micro- and nano-motors for biomedical applications, *Journal of Materials Chemistry B* 2 (17) (2014) 2395–2408. 1520
- [181] W. R. Browne, B. L. Feringa, Making molecular machines work, *Nature Nanotechnology* 1 (1) (2006) 25.
- 1525 [182] G. Sinclair, P. Jordan, J. Courtial, M. Padgett, J. Cooper, Z. J. Laczik, Assembly of 3-dimensional structures using programmable holographic optical tweezers, *Optics Express* 12 (22) (2004) 5475–5480.
- [183] D. G. Grier, A revolution in optical manipulation, *Nature* 424 (6950) (2003) 810.

- [184] P. Jones, F. Palmisano, F. Bonaccorso, P. Gucciardi, G. Calogero, A. Ferrari, O. Marago, Rotation detection in light-driven nanorotors, *ACS Nano* 3 (10) (2009) 3077–3084.
- [185] A. A. R. Neves, A. Camposeo, S. Pagliara, R. Saija, F. Borghese, P. Denti, M. A. Iatì, R. Cingolani, O. M. Maragò, D. Pisignano, Rotational dynamics of optically trapped nanofibers, *Optics express* 18 (2) (2010) 822–830.
- [186] A. Lehmuskero, R. Ogier, T. Gschneidner, P. Johansson, M. Kall, Ultrafast spinning of gold nanoparticles in water using circularly polarized light, *Nano Letters* 13 (7) (2013) 3129–3134.
- [187] A. Lehmuskero, P. Johansson, H. Rubinsztein-Dunlop, L. Tong, M. Kall, Laser trapping of colloidal metal nanoparticles, *ACS Nano* 9 (4) (2015) 3453–3469.
- [188] O. Marago, P. Gucciardi, F. Bonaccorso, G. Calogero, V. Scardaci, A. Rozhin, A. Ferrari, P. Jones, R. Saija, F. Borghese, P. Denti, M. Iati, Optical trapping of carbon nanotubes, *Physica E: Low-dimensional Systems and Nanostructures* 40 (7) (2008) 2347–2351.
- [189] P. M. Hansen, V. K. Bhatia, N. Harrit, L. Oddershede, Expanding the optical trapping range of gold nanoparticles, *Nano Letters* 5 (10) (2005) 1937–1942.
- [190] E. Messina, E. Cavallaro, A. Cacciola, M. A. Iatì, P. G. Gucciardi, F. Borghese, P. Denti, R. Saija, G. Compagnini, M. Meneghetti, V. Amendola, O. M. Marago, Plasmon-enhanced optical trapping of gold nanoaggregates with selected optical properties, *ACS nano* 5 (2) (2011) 905–913.
- [191] E. Messina, E. Cavallaro, A. Cacciola, R. Saija, F. Borghese, P. Denti, B. Fazio, C. D’Andrea, P. Gucciardi, M. Iati, M. Meneghetti, G. Compagnini, V. Amendola, O. Marago, Manipulation and raman spectroscopy with optically trapped metal nanoparticles obtained by pulsed laser ab-

- 1555 lation in liquids, *The Journal of Physical Chemistry C* 115 (12) (2011)
5115–5122.
- [192] E. Messina, L. D’Urso, E. Fazio, C. Satriano, M. Donato, C. D’Andrea,
O. Maragò, P. Gucciardi, G. Compagnini, F. Neri, Tuning the structural
and optical properties of gold/silver nano-alloys prepared by laser ablation
1560 in liquids for optical limiting, ultra-sensitive spectroscopy, and optical
trapping, *Journal of Quantitative Spectroscopy and Radiative Transfer*
113 (18) (2012) 2490–2498.
- [193] O. Brzobohatý, M. Šiler, J. Trojek, L. Chvátal, V. Karásek, A. Pataák,
Z. Pokorná, F. Mika, P. Zemánek, Three-dimensional optical trapping of
1565 a plasmonic nanoparticle using low numerical aperture optical tweezers,
Scientific Reports 5 (2015) 8106.
- [194] O. Brzobohaty, M. Šiler, J. Trojek, L. Chvátal, V. Karásek, P. Zemánek,
Non-spherical gold nanoparticles trapped in optical tweezers: shape mat-
ters, *Optics express* 23 (7) (2015) 8179–8189.
- 1570 [195] M. Padgett, R. Di Leonardo, Holographic optical tweezers and their rele-
vance to lab on chip devices, *Lab on a Chip* 11 (7) (2011) 1196–1205.
- [196] N. Simpson, K. Dholakia, L. Allen, M. Padgett, Mechanical equivalence
of spin and orbital angular momentum of light: an optical spanner, *Optics*
Letters 22 (1) (1997) 52–54.
- 1575 [197] K. Ladavac, D. G. Grier, Microoptomechanical pumps assembled and
driven by holographic optical vortex arrays, *Optics Express* 12 (6) (2004)
1144–1149.
- [198] J. Kao, X. Wang, J. Warren, J. Xu, D. Attinger, A bubble-powered micro-
rotor: conception, manufacturing, assembly and characterization, *Journal*
of Micromechanics and Microengineering 17 (12) (2007) 2454.
1580
- [199] C. Lee, M. Liamini, L. G. Frechette, A silicon microturbopump for a
rankine-cycle power-generation microsystem - part ii: Fabrication and

- characterization, *Journal of Microelectromechanical Systems* 20 (1) (2011) 326–338.
- 1585 [200] P. A. Quinto-Su, A microscopic steam engine implemented in an optical tweezer, *Nature Communications* 5 (2014) 5889.
- [201] V. Blickle, C. Bechinger, Realization of a micrometre-sized stochastic heat engine, *Nature Physics* 8 (2) (2012) 143.
- [202] I. A. Martínez, É. Roldán, L. Dinis, D. Petrov, J. M. Parrondo, R. A. 1590 Rica, Brownian carnot engine, *Nature Physics* 12 (1) (2016) 67.
- [203] T. Čižmár, L. C. Dávila Romera, K. Dholakia, D. L. Andrews, Multiple optical trapping and binding: new routes to self-assembly, *J. Phys B: At. Mol. Opt. Phys.* 43 (2010) 102001.
- [204] R. W. Bowman, M. J. Padgett, Optical trapping and binding, *Rep. Prog. 1595 Phys.* 76 (2013) 026401.
- [205] M. M. Burns, J.-M. Fournier, J. A. Golovchenko, Optical binding, *Phys. Rev. Lett.* 63 (1989) 1233–1236.
- [206] M. M. Burns, J.-M. Fournier, J. A. Golovchenko, Optical matter: crystallization and binding in intense optical fields, *Science* 249 (4970) (1990) 1600 749–754.
- [207] W. Singer, M. Frick, S. Bernet, M. Ritsch-Marte, Self-organized array of regularly spaced microbeads in a fiber-optical trap, *J. Opt. Soc. Am. B* 20 (7) (2003) 1568–1574.
- [208] R. Gordon, M. Kawano, J. T. Blakely, D. Sinton, Optohydrodynamic 1605 theory of particles in a dual-beam optical trap, *Phys. Rev. B* 77 (2008) 245125.
- [209] V. Garcés-Chávez, K. Dholakia, G. C. Spalding, Extended-area optically induced organization of microparticles on a surface, *Applied Physics Letters* 86 (3) (2005) 031106.

- 1610 [210] C. D. Mellor, C. D. Bain, Array formation in evanescent waves,
ChemPhysChem 7 (2) (2006) 329–332.
- [211] C. D. Mellor, T. A. Fennerty, C. D. Bain, Polarization effects in optically
bound particles arrays, Opt. Express 14 (21) (2006) 10079–10088.
- [212] X. Han, P. H. Jones, Evanescent wave optical binding forces on spherical
1615 microparticles, Opt. Lett. 40 (17) (2015) 4042–4045.
- [213] X. Han, H. Luo, G. Xiao, P. H. Jones, Optically bound colloidal lattices
in evanescent optical fields, Opt. Lett. 41 (21) (2016) 4935–4938.
- [214] S. Kawata, T. Tani, Optically driven mie particles in an evanescent field
along achanneled waveguide, Opt. Lett. 21 (21) (1996) 1768–1770.
- 1620 [215] T. Tanaka, S. Yamamoto, Optically induced propulsion of small parti-
cles in an evenescent field of higher propagation mode in a multimode,
channeled waveguide, Appl. Phys. Lett. 77 (20) (2000) 3131–3133.
- [216] G. Brambilla, G. S. Murugan, J. S. Wilkinson, D. J. Richardson, Optical
manipulation of microspheres along a subwavelength optical wire, Opt.
1625 Lett. 32 (20) (2007) 3041–3043.
- [217] S. E. Skelton, M. Sergides, R. Patel, E. Karczewska, O. M. Maragó,
P. H. Jones, Evanescent wave optical trapping and transport of micro-
and nanoparticles on tapered optical fibers, Journal of Quantitative Spec-
troscopy and Radiative Transfer 113 (18) (2012) 2512–2520.
- 1630 [218] M. Sergides, S. E. Skelton, E. Karczewska, K. Thorneycroft, O. M.
Maragò, P. H. Jones, Optically bound particle structures in evanescent
wave traps, Vol. 8458, 2012, pp. 8458 – 8458 – 7. doi:10.1117/12.929612.
URL <https://doi.org/10.1117/12.929612>
- [219] M. Daly, V. G. Truong, S. N. Chormaic, Evanescent field trapping of
1635 nanoparticles using nanostructured ultrathin optical fibers, Opt. Express
24 (13) (2016) 14470–14482.

- [220] J. M. Ward, A. Maimaiti, V. H. Le, S. N. Chormaic, Contributed review: Optical micro- and nanofiber pulling rig, *Review of Scientific Instruments* 85 (11) (2014) 111501.
- 1640 [221] M. Daly, M. S. Marios, S. N. Chormaic, Optical trapping and manipulation of micrometer and submicrometer particles, *Laser & Photonics Reviews* 9 (3) (2015) 309–329.
- [222] M. C. Frawley, I. Gusachenko, V. G. Truong, M. Sergides, S. N. Chormaic, Selective particle trapping and optical binding in the evanescent field of an optical nanofiber, *Opt. Express* 22 (13) (2014) 16322–16334.
- 1645 [223] A. Maimaiti, V. G. Truong, M. Sergides, I. Gusachenko, S. N. Chormaic, Higher order microfiber modes for dielectric particle trapping and propulsion, *Sci. Rep.* 5 (2015) 9077. doi:doi:10.1038/srep09077.
- [224] Z. Yan, R. A. Shah, G. Chado, S. K. Gray, M. Pelton, N. F. Scherer, Guiding spatial arrangements of silver nanoparticles by optical binding interactions in shaped light fields, *ACS Nano* 7 (2) (2013) 1790–1802.
- 1650 [225] V. Demergis, E. Florin, Ultrastrong optical binding of metallic nanoparticles, *Nano Letters* 12 (11) (2012) 5756–5760.
- [226] S. H. Simpson, P. H. Jones, O. M. Maragò, S. Hanna, M. J. Miles, Optical binding of nanowires in counterpropagating beams, Vol. 8810, 2013, pp. 8810 – 8810 – 9. doi:10.1117/12.2024466.
- 1655 [227] S. H. Simpson, P. Zemánek, O. M. Maragò, P. H. Jones, S. Hanna, Optical binding of nanowires, *Nano Letters* 17 (6) (2017) 3485–3492.
- [228] M. J. Guffey, N. F. Scherer, All-optical patterning of Au nanoparticles on surfaces using optical traps, *Nano Letters* 10 (11) (2010) 4302–4308.
- 1660 [229] S. Nedev, A. S. Urban, A. A. Lutich, J. Feldmann, Optical force stamping lithography, *Nano Letters* 11 (11) (2011) 5066–5070.

- [230] A. S. Urban, T. Pfeiffer, M. Fedoruk, A. A. Lutich, J. Feldmann, Single-step injection of gold nanoparticles through phospholipid membranes, ACS Nano 5 (5) (2011) 3585–3590. 1665
- [231] J. Gargiulo, I. L. Violi, S. Cerrota, L. Chvatal, E. Cortes, E. M. Perassi, F. Diaz, P. Zemanek, F. D. Stefani, Accuracy and mechanistic details of optical printing of single Au and Ag nanoparticles, ACS Nano 11 (10) (2017) 9678–9688.
- [232] B. Fazio, C. D’Andrea, A. Foti, E. Messina, A. Irrera, M. G. Donato, V. Villari, N. Micali, O. M. Maragò, P. G. Gucciardi, Sers detection of biomolecules at physiological ph via aggregation of gold nanorods mediated by optical forces and plasmonic heating, Scientific Reports 6 (2016) 26952. 1670
- [233] A. Foti, C. D’Andrea, V. Villari, N. Micali, M. G. Donato, B. Fazio, O. M. Maragò, R. Gillibert, M. Lamy de la Chapelle, P. G. Gucciardi, Optical aggregation of gold nanoparticles for sers detection of proteins and toxins in liquid environment: towards ultrasensitive and selective detection, Materials 11 (2018) 440. 1675
- [234] H.-B. Sun, S. Kawata, Two-photon photopolymerization and 3d lithographic microfabrication, Advances in Polymer Science 170 (2004) 169–274. 1680
- [235] C. Liberale, G. Cojoc, P. Candeloro, G. Das, F. Gentile, F. De Angelis, E. Di Fabrizio, Micro-optics fabrication on top of optical fibers using two-photon lithography, IEEE Photonics Technology Letters 22 (7) (2010) 474–476. 1685
- [236] C. Liberale, G. Cojoc, F. Bragheri, P. Minzioni, G. Perozziello, R. La Rocca, L. Ferrara, V. Rajamanickam, E. Di Fabrizio, I. Cristiani, Integrated microfluidic device for single-cell trapping and spectroscopy, Scientific Reports 3 (2013) 1258. 1690

- [237] G. Nelson, R. A. Kirian, U. Weierstall, N. A. Zatsepin, T. Faragó, T. Baumbach, F. Wilde, F. B. Niesler, B. Zimmer, I. Ishigami, et al., Three-dimensional-printed gas dynamic virtual nozzles for x-ray laser sample delivery, *Optics Express* 24 (11) (2016) 11515–11530.
- 1695 [238] S. Hengsbach, A. D. Lantada, Rapid prototyping of multi-scale biomedical microdevices by combining additive manufacturing technologies, *Biomedical microdevices* 16 (4) (2014) 617–627.
- [239] B. Dorin, P. Parkinson, P. Scully, Direct laser write process for 3d conductive carbon circuits in polyimide, *Journal of Materials Chemistry C* 5 (20) (2017) 4923–4930.
- 1700 [240] F. Klein, B. Richter, T. Striebel, C. M. Franz, G. v. Freymann, M. Wegener, M. Bastmeyer, Two-component polymer scaffolds for controlled three-dimensional cell culture, *Advanced Materials* 23 (11) (2011) 1341–1345.
- 1705 [241] B. Richter, V. Hahn, S. Bertels, T. K. Claus, M. Wegener, G. Delaittre, C. Barner-Kowollik, M. Bastmeyer, Guiding cell attachment in 3d micro-scaffolds selectively functionalized with two distinct adhesion proteins, *Advanced Materials* 29 (5).
- [242] F. De Angelis, C. Liberale, M. Coluccio, G. Cojoc, E. Di Fabrizio, Emerging fabrication techniques for 3d nano-structuring in plasmonics and single molecule studies, *Nanoscale* 3 (7) (2011) 2689–2696.
- 1710 [243] L. Britnell, R. Ribeiro, A. Eckmann, R. Jalil, B. Belle, A. Mishchenko, Y.-J. Kim, R. Gorbachev, T. Georgiou, S. Morozov, et al., Strong light-matter interactions in heterostructures of atomically thin films, *Science* 340 (6138) (2013) 1311–1314.
- 1715 [244] Q. H. Wang, K. Kalantar-Zadeh, A. Kis, J. N. Coleman, M. S. Strano, Electronics and optoelectronics of two-dimensional transition metal dichalcogenides, *Nat. Nanotechnol.* 7 (11) (2012) 699–712.

- [245] F. Bonaccorso, L. Colombo, G. Yu, M. Stoller, V. Tozzini, A. C. Ferrari,
1720 R. S. Ruoff, V. Pellegrini, Graphene, related two-dimensional crystals,
and hybrid systems for energy conversion and storage, *Science* 347 (6217)
(2015) 1246501.
- [246] A. K. Geim, I. V. Grigorieva, Van der waals heterostructures, *Nature*
499 (7459) (2013) 419–425.
- 1725 [247] A. Magazzù, A. Callegari, J. P. Staforelli, A. Gambassi, S. Dietrich,
G. Volpe, Controlling the dynamics of colloidal particles by critical casimir
forces, arXiv preprint arXiv:1806.11403.
- [248] M. E. Fisher, P. G. de Gennes, Phénomènes aux parois dans un mélange
binaire critique, *C. R. Acad. Sci. Paris B* 287 (1978) 207–209.
- 1730 [249] C. Hertlein, L. Helden, A. Gambassi, S. Dietrich, C. Bechinger, Direct
measurement of critical casimir forces, *Nature* 451 (7175) (2008) 172.
- [250] A. Gambassi, A. Maciolek, C. Hertlein, U. Nellen, L. Helden, C. Bechinger,
S. Dietrich, Critical casimir effect in classical binary liquid mixtures, *Phys-
ical Review E* 80 (6) (2009) 061143.
- 1735 [251] E. Marino, T. E. Kodger, J. B. ten Hove, A. H. Velders, P. Schall, As-
sembling quantum dots via critical casimir forces, *Solar Energy Materials
and Solar Cells* 158 (2016) 154–159.
- [252] S. Paladugu, A. Callegari, Y. Tuna, L. Barth, S. Dietrich, A. Gambassi,
G. Volpe, Nonadditivity of critical casimir forces, *Nature Communications*
1740 7 (2016) 11403.
- [253] A. Argun, A.-R. Moradi, E. Pinçe, G. B. Bağcı, A. Imparato, G. Volpe,
Non-boltzmann stationary distributions and nonequilibrium relations in
active baths, *Physical Review E* 94 (6) (2016) 062150.
- 1745 [254] D. J. Evans, E. G. D. Cohen, G. P. Morriss, Probability of second law
violations in shearing steady states, *Physical Review Letters* 71 (15) (1993)
2401.

- [255] G. Wang, E. M. Sevick, E. Mittag, D. J. Searles, D. J. Evans, Experimental demonstration of violations of the second law of thermodynamics for small systems and short time scales, *Physical Review Letters* 89 (5) (2002) 050601.
- [256] C. Jarzynski, Equalities and inequalities: Irreversibility and the second law of thermodynamics at the nanoscale, *Annu. Rev. Condens. Matter Phys.* 2 (1) (2011) 329–351.
- [257] E. Barkai, Y. Garini, R. Metzler, Strange kinetics of single molecules in living cells, *Phys. Today* 65 (8) (2012) 29.
- [258] C. Bechinger, R. Di Leonardo, H. Löwen, C. Reichhardt, G. Volpe, G. Volpe, Active particles in complex and crowded environments, *Reviews of Modern Physics* 88 (4) (2016) 045006.
- [259] C. Jarzynski, Nonequilibrium equality for free energy differences, *Physical Review Letters* 78 (14) (1997) 2690.
- [260] G. E. Crooks, Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences, *Physical Review E* 60 (3) (1999) 2721.
- [261] S. Ramaswamy, The mechanics and statistics of active matter, *Ann. Rev. Condens. Matter Phys.* 1 (2010) 323–345.
- [262] F. Schweitzer, *Brownian agents and active particles: collective dynamics in the natural and social sciences*, Springer Science & Business Media, 2007.
- [263] E. Lauga, T. R. Powers, The hydrodynamics of swimming microorganisms, *Reports on Progress in Physics* 72 (9) (2009) 096601.
- [264] H.-R. Jiang, N. Yoshinaga, M. Sano, Active motion of a janus particle by self-thermophoresis in a defocused laser beam, *Physical Review Letters* 105 (26) (2010) 268302.

- [265] H. C. Berg, *E. coli in Motion*, Springer Science & Business Media, 2008.
- 1775 [266] M. Hennes, K. Wolff, H. Stark, Self-induced polar order of active brownian particles in a harmonic trap, *Physical Review Letters* 112 (23) (2014) 238104.
- [267] M. Leoni, J. Kotar, B. Bassetti, P. Cicuta, M. C. Lagomarsino, A basic swimmer at low reynolds number, *Soft Matter* 5 (2) (2009) 472–476.
- 1780 [268] S. M. Block, D. F. Blair, H. C. Berg, Compliance of bacterial flagella measured with optical tweezers, *Nature* 338 (6215) (1989) 514.
- [269] E. Pinçe, S. K. Velu, A. Callegari, P. Elahi, S. Gigan, G. Volpe, G. Volpe, Disorder-mediated crowd control in an active matter system, *Nature Communications* 7 (2016) 10907.
- 1785 [270] T. Ha, Probing nature’s nanomachines one molecule at a time, *Biophysical journal* 110 (5) (2016) 1004.
- [271] I. A. Martínez, É. Roldán, L. Dinis, R. A. Rica, Colloidal heat engines: a review, *Soft Matter* 13 (1) (2017) 22–36.
- 1790 [272] J. Gieseler, J. Millen, Levitated nanoparticles for microscopic thermodynamics - a review, *Entropy* 20 (5) (2018) 326.