

Optimal asymptotic bounds for spherical designs

Andriy Bondarenko, Danylo Radchenko, and Maryna Viazovska

Abstract

In this paper we prove the conjecture of Korevaar and Meyers: for each $N \geq c_d t^d$ there exists a spherical t -design in the sphere S^d consisting of N points, where c_d is a constant depending only on d .

Keywords: Spherical designs, Brouwer degree, Marcinkiewicz–Zygmund inequalities, area-regular partitions

AMS subject classification: 52C35, 41A55, 41A63

1 Introduction

Let S^d be the unit sphere in \mathbb{R}^{d+1} with the Lebesgue measure μ_d normalized by $\mu_d(S^d) = 1$.

A set of points $x_1, \dots, x_N \in S^d$ is called a *spherical t -design* if

$$\int_{S^d} P(x) d\mu_d(x) = \frac{1}{N} \sum_{i=1}^N P(x_i)$$

for all algebraic polynomials in $d + 1$ variables, of total degree at most t . The concept of a spherical design was introduced by Delsarte, Goethals, and Seidel [12]. For each $t, d \in \mathbb{N}$ denote by $N(d, t)$ the minimal number of points in a spherical t -design in S^d . The following lower bound

$$(1) \quad N(d, t) \geq \begin{cases} \binom{d+k}{d} + \binom{d+k-1}{d} & \text{if } t = 2k, \\ 2 \binom{d+k}{d} & \text{if } t = 2k + 1, \end{cases}$$

is proved in [12].

Spherical t -designs attaining this bound are called *tight*. The vertices of a regular $t+1$ -gon form a tight spherical t -design in the circle, so $N(1, t) = t+1$. Exactly eight tight spherical designs are known for $d \geq 2$ and $t \geq 4$. All such configurations of points are highly symmetrical, and optimal from many different points of view (see Cohn, Kumar [8] and Conway, Sloane [11]). Unfortunately, tight designs rarely exist. In particular, Bannai and Damerell [2, 3] have shown that tight spherical designs with $d \geq 2$ and $t \geq 4$ may exist only for $t = 4, 5, 7$ or 11 . Moreover, the only tight 11-design is formed by minimal vectors of the Leech lattice in dimension 24. The bound (1) has been improved by Delsarte's linear programming method for most pairs (d, t) ; see [22].

On the other hand, Seymour and Zaslavsky [20] have proved that spherical t -designs exist for all $d, t \in \mathbb{N}$. However, this proof is nonconstructive and

gives no idea of how big $N(d, t)$ is. So, a natural question is to ask how $N(d, t)$ differs from the tight bound (1). Generally, to find the exact value of $N(d, t)$ even for small d and t is a surprisingly hard problem. For example, everybody believes that 24 minimal vectors of the D_4 root lattice form a 5-design with minimal number of points in S^3 , although it is only proved that $22 \leq N(3, 5) \leq 24$; see [6]. Further, Cohn, Conway, Elkies, and Kumar [7] conjectured that every spherical 5-design consisting of 24 points in S^3 is in a certain 3-parametric family. Recently, Musin [17] has solved a long standing problem related to this conjecture. Namely, he proved that the kissing number in dimension 4 is 24.

In this paper we focus on asymptotic upper bounds on $N(d, t)$ for fixed $d \geq 2$ and $t \rightarrow \infty$. Let us give a brief history of this question. First, Wagner [21] and Bajnok [1] proved that $N(d, t) \leq C_d t^{Cd^4}$ and $N(d, t) \leq C_d t^{Cd^3}$, respectively. Then, Korevaar and Meyers [14] have improved these inequalities by showing that $N(d, t) \leq C_d t^{(d^2+d)/2}$. They have also conjectured that

$$N(d, t) \leq C_d t^d.$$

Note that (1) implies $N(d, t) \geq c_d t^d$. Here and in what follows we denote by C_d and c_d sufficiently large and sufficiently small positive constants depending only on d , respectively.

The conjecture of Korevaar and Meyers attracted the interest of many mathematicians. For instance, Kuijlaars and Saff [19] emphasized the importance of this conjecture for $d = 2$, and revealed its relation to minimal energy problems. Mhaskar, Narcowich, and Ward [16] have constructed positive quadrature formulas in S^d with $C_d t^d$ points having *almost* equal weights. Very recently, Chen, Frommer, Lang, Sloan, and Womersley [9, 10] gave a computer-assisted proof that spherical t -designs with $(t + 1)^2$ points exist in S^2 for $t \leq 100$.

For $d = 2$, there is an even stronger conjecture by Hardin and Sloane [13] saying that $N(2, t) \leq \frac{1}{2}t^2 + o(t^2)$ as $t \rightarrow \infty$. Numerical evidence supporting the conjecture was also given.

In [4], we have suggested a nonconstructive approach for obtaining asymptotic bounds for $N(d, t)$ based on the application of the Brouwer fixed point theorem. This led to the following result:

For each $N \geq C_d t^{\frac{2d(d+1)}{d+2}}$ there exists a spherical t -design in S^d consisting of N points.

Instead of the Brouwer fixed point theorem we use in this paper the following result from the Brouwer degree theory [18, Th. 1.2.6, Th. 1.2.9].

THEOREM A. *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a continuous mapping and Ω an open bounded subset, with boundary $\partial\Omega$, such that $0 \in \Omega \subset \mathbb{R}^n$. If $(x, f(x)) > 0$ for all $x \in \partial\Omega$, then there exists $x \in \Omega$ satisfying $f(x) = 0$.*

We employ this theorem to prove the conjecture of Korevaar and Meyers.

Theorem 1. *For each $N \geq C_d t^d$ there exists a spherical t -design in S^d consisting of N points.*

Note that Theorem 1 is slightly stronger than the original conjecture because it guarantees the existence of spherical t -designs for *each* N greater than $C_d t^d$.

This paper is organized as follows. In Section 2 we explain the main idea of the proof. Then in Section 3 we present some auxiliary results. Finally, we prove Theorem 1 in Section 4.

2 Preliminaries and the main idea

Let \mathcal{P}_t be the Hilbert space of polynomials P on S^d of degree at most t such that

$$\int_{S^d} P(x) d\mu_d(x) = 0,$$

equipped with the usual inner product

$$(P, Q) = \int_{S^d} P(x)Q(x) d\mu_d(x).$$

By the Riesz representation theorem, for each point $x \in S^d$ there exists a unique polynomial $G_x \in \mathcal{P}_t$ such that

$$(G_x, Q) = Q(x) \text{ for all } Q \in \mathcal{P}_t.$$

Then a set of points $x_1, \dots, x_N \in S^d$ forms a spherical t -design if and only if

$$(2) \quad G_{x_1} + \dots + G_{x_N} = 0.$$

For a differentiable function $f : \mathbb{R}^{d+1} \rightarrow \mathbb{R}$ denote by

$$\frac{\partial f}{\partial x}(x_0) := \left(\frac{\partial f}{\partial \xi_1}(x_0), \dots, \frac{\partial f}{\partial \xi_{d+1}}(x_0) \right)$$

the gradient of f at the point $x_0 \in \mathbb{R}^{d+1}$.

For a polynomial $Q \in \mathcal{P}_t$ we define the spherical gradient as follows:

$$(3) \quad \nabla Q(x) := \frac{\partial}{\partial x} Q \left(\frac{x}{|x|} \right),$$

where $|\cdot|$ is the Euclidean norm in \mathbb{R}^{d+1} .

We apply Theorem A to the open subset Ω of a vector space \mathcal{P}_t ,

$$(4) \quad \Omega := \left\{ P \in \mathcal{P}_t \mid \int_{S^d} |\nabla P(x)| d\mu_d(x) < 1 \right\}.$$

Now we observe that the existence of a continuous mapping $F : \mathcal{P}_t \rightarrow (S^d)^N$, such that for all $P \in \partial\Omega$

$$(5) \quad \sum_{i=1}^N P(x_i(P)) > 0, \text{ where } F(P) = (x_1(P), \dots, x_N(P)),$$

readily implies the existence of a spherical t -design in S^d consisting of N points. Consider a mapping $L : (S^d)^N \rightarrow \mathcal{P}_t$ defined by

$$(x_1, \dots, x_N) \xrightarrow{L} G_{x_1} + \dots + G_{x_N},$$

and the following composition mapping $f = L \circ F : \mathcal{P}_t \rightarrow \mathcal{P}_t$. Clearly

$$(P, f(P)) = \sum_{i=1}^N P(x_i(P))$$

for each $P \in \mathcal{P}_t$. Thus, applying Theorem A to the mapping f , the vector space \mathcal{P}_t , and the subset Ω defined by (4), we obtain that $f(Q) = 0$ for some $Q \in \mathcal{P}_t$. Hence, by (2), the components of $F(Q) = (x_1(Q), \dots, x_N(Q))$ form a spherical t -design in S^d consisting of N points.

The most naive approach to construct such F is to start with a certain well-distributed collection of points x_i ($i = 1, \dots, N$), put $F(0) := (x_1, \dots, x_N)$, and then move each point along the spherical gradient vector field of P . Note that this is the most greedy way to increase each $P(x_i(P))$ and make $\sum_{i=1}^N P(x_i(P))$ positive for each $P \in \partial\Omega$. Following this approach we will give an explicit construction of F in Section 4, which will immediately imply the proof of Theorem 1.

3 Auxiliary results

To construct the corresponding mapping F for each $N \geq C_d t^d$ we extensively use the following notion of an area-regular partition.

Let $\mathcal{R} = \{R_1, \dots, R_N\}$ be a finite collection of closed sets $R_i \subset S^d$ such that $\cup_{i=1}^N R_i = S^d$ and $\mu_d(R_i \cap R_j) = 0$ for all $1 \leq i < j \leq N$. The partition \mathcal{R} is called area-regular if $\mu_d(R_i) = 1/N$, $i = 1, \dots, N$. The partition norm for \mathcal{R} is defined by

$$\|\mathcal{R}\| := \max_{R \in \mathcal{R}} \text{diam } R,$$

where $\text{diam } R$ stands for the maximum geodesic distance between two points in R . We need the following fact on area-regular partitions (see Bourgain, Lindenstrauss [5] and Kuijlaars, Saff [15]):

THEOREM B. *For each $N \in \mathbb{N}$ there exists an area-regular partition $\mathcal{R} = \{R_1, \dots, R_N\}$ with $\|\mathcal{R}\| \leq B_d N^{-1/d}$ for some constant B_d large enough.*

We will also use the following spherical Marcinkiewicz–Zygmund type inequality:

THEOREM C. *There exists a constant r_d such that for each area-regular partition $\mathcal{R} = \{R_1, \dots, R_N\}$ with $\|\mathcal{R}\| < \frac{r_d}{m}$, each collection of points $x_i \in R_i$ ($i = 1, \dots, N$), and each algebraic polynomial P of total degree m , the*

inequality

$$(6) \quad \frac{1}{2} \int_{S^d} |P(x)| d\mu_d(x) \leq \frac{1}{N} \sum_{i=1}^N |P(x_i)| \leq \frac{3}{2} \int_{S^d} |P(x)| d\mu_d(x)$$

holds.

Theorem C follows naturally from the proof of Theorem 3.1 in [16].

Corollary 1. *For each area-regular partition $\mathcal{R} = \{R_1, \dots, R_N\}$ with $\|\mathcal{R}\| < \frac{r_d}{m+1}$, each collection of points $x_i \in R_i$ ($i = 1, \dots, N$), and each algebraic polynomial P of total degree m ,*

$$(7) \quad \frac{1}{3\sqrt{d}} \int_{S^d} |\nabla P(x)| d\mu_d(x) \leq \frac{1}{N} \sum_{i=1}^N |\nabla P(x_i)| \leq 3\sqrt{d} \int_{S^d} |\nabla P(x)| d\mu_d(x).$$

Proof. Since $|\nabla P| = \sqrt{P_1^2 + \dots + P_{d+1}^2}$ in S^d , where P_j are polynomials of total degree $m+1$, Corollary 1 is an immediate consequence of (6) applied to P_j , $j = 1, \dots, d+1$. \square

4 Proof of Theorem 1

In this section we construct the map F introduced in Section 2 and thereby finish the proof of Theorem 1.

For $d, t \in \mathbb{N}$, take $C_d > (54dB_d/r_d)^d$, where B_d is as in Theorem B and r_d is as in Theorem C, and fix $N \geq C_d t^d$. Now we are in a position to give an exact construction of the mapping $F : \mathcal{P}_t \rightarrow (S^d)^N$ which satisfies condition (5). Take an area-regular partition $\mathcal{R} = \{R_1, \dots, R_N\}$ with

$$(8) \quad \|\mathcal{R}\| \leq B_d N^{-1/d} < \frac{r_d}{54dt}$$

as provided by Theorem B, and choose an arbitrary $x_i \in R_i$ for each $i = 1, \dots, N$. Put $\varepsilon = \frac{1}{6\sqrt{d}}$ and consider the function

$$h_\varepsilon(u) := \begin{cases} u & \text{if } u > \varepsilon, \\ \varepsilon & \text{otherwise.} \end{cases}$$

Take a mapping $U : \mathcal{P}_t \times S^d \rightarrow \mathbb{R}^{d+1}$ such that

$$U(P, y) = \frac{\nabla P(y)}{h_\varepsilon(|\nabla P(y)|)}.$$

For each $i = 1, \dots, N$ let $y_i : \mathcal{P}_t \times [0, \infty) \rightarrow S^d$ be the map satisfying the differential equation

$$(9) \quad \frac{d}{ds} y_i(P, s) = U(P, y_i(P, s))$$

with the initial condition

$$y_i(P, 0) = x_i,$$

for each $P \in \mathcal{P}_t$. Note that each mapping y_i has its values in S^d by definition of spherical gradient (3). Since the mapping $U(P, y)$ is Lipschitz continuous in both P and y , each y_i is well defined and continuous in both P and s , where the metric on \mathcal{P}_t is given by the inner product. Finally put

$$(10) \quad F(P) = (x_1(P), \dots, x_N(P)) := (y_1(P, \frac{r_d}{3t}), \dots, y_N(P, \frac{r_d}{3t})).$$

By definition the mapping F is continuous on \mathcal{P}_t . So, as explained in Section 2, to finish the proof of Theorem 1 it suffices to prove

Lemma 1. *Let $F : \mathcal{P}_t \rightarrow (S^d)^N$ be the mapping defined by (10). Then for each $P \in \partial\Omega$,*

$$\frac{1}{N} \sum_{i=1}^N P(x_i(P)) > 0,$$

where Ω is given by (4).

Proof. Fix $P \in \partial\Omega$. For the sake of simplicity we write $y_i(s)$ in place of $y_i(P, s)$. By the Newton-Leibniz formula we have

$$(11) \quad \begin{aligned} & \frac{1}{N} \sum_{i=1}^N P(x_i(P)) = \frac{1}{N} \sum_{i=1}^N P(y_i(r_d/3t)) \\ & = \frac{1}{N} \sum_{i=1}^N P(x_i) + \int_0^{r_d/3t} \frac{d}{ds} \left[\frac{1}{N} \sum_{i=1}^N P(y_i(s)) \right] ds. \end{aligned}$$

Now to prove Lemma 1, we first estimate the value

$$\left| \frac{1}{N} \sum_{i=1}^N P(x_i) \right|$$

from above, and then estimate the value

$$\frac{d}{ds} \left[\frac{1}{N} \sum_{i=1}^N P(y_i(s)) \right]$$

from below, for each $s \in [0, r_d/3t]$. We have

$$\begin{aligned} \left| \frac{1}{N} \sum_{i=1}^N P(x_i) \right| &= \left| \sum_{i=1}^N \int_{R_i} P(x_i) - P(x) d\mu_d(x) \right| \leq \sum_{i=1}^N \int_{R_i} |P(x_i) - P(x)| d\mu_d(x) \\ &\leq \frac{\|\mathcal{R}\|}{N} \sum_{i=1}^N \max_{z \in S^d: \text{dist}(z, x_i) \leq \|\mathcal{R}\|} |\nabla P(z)| \end{aligned}$$

where $\text{dist}(z, x_i)$ denotes the geodesic distance between z and x_i . Hence, for $z_i \in S^d$ such that $\text{dist}(z_i, x_i) \leq \|\mathcal{R}\|$ and

$$|\nabla P(z_i)| = \max_{z \in S^d: \text{dist}(z, x_i) \leq \|\mathcal{R}\|} |\nabla P(z)|,$$

we obtain

$$\left| \frac{1}{N} \sum_{i=1}^N P(x_i) \right| \leq \frac{\|\mathcal{R}\|}{N} \sum_{i=1}^N |\nabla P(z_i)|.$$

Consider another area-regular partition $\mathcal{R}' = \{R'_1, \dots, R'_N\}$ defined by $R'_i = R_i \cup \{z_i\}$. Clearly $\|\mathcal{R}'\| \leq 2\|\mathcal{R}\|$ and so, by (8), we get $\|\mathcal{R}'\| < r_d/(27dt)$. Applying inequality (7) to the partition \mathcal{R}' and the collection of points z_i we obtain that

$$(12) \quad \left| \frac{1}{N} \sum_{i=1}^N P(x_i) \right| \leq 3\sqrt{d} \|\mathcal{R}\| \int_{S^d} |\nabla P(x)| d\mu_d(x) < \frac{r_d}{18\sqrt{d}t}$$

for any $P \in \partial\Omega$. On the other hand, the differential equation (9) implies

$$\begin{aligned}
\frac{d}{ds} \left[\frac{1}{N} \sum_{i=1}^N P(y_i(s)) \right] &= \frac{1}{N} \sum_{i=1}^N \frac{|\nabla P(y_i(s))|^2}{h_\varepsilon(|\nabla P(y_i(s))|)} \\
&\geq \frac{1}{N} \sum_{i: |\nabla P(y_i(s))| \geq \varepsilon} |\nabla P(y_i(s))| \\
(13) \qquad \qquad \qquad &\geq \frac{1}{N} \sum_{i=1}^N |\nabla P(y_i(s))| - \varepsilon.
\end{aligned}$$

Since

$$\left| \frac{\nabla P(y)}{h_\varepsilon(|\nabla P(y)|)} \right| \leq 1$$

for each $y \in S^d$, it follows again from (9) that $\left| \frac{dy_i(s)}{ds} \right| \leq 1$. Hence we arrive at

$$\text{dist}(x_i, y_i(s)) \leq s.$$

Now for each $s \in [0, r_d/3t]$ consider the area-regular partition $\mathcal{R}'' = \{R_1'', \dots, R_N''\}$ given by $R_i'' = R_i \cup \{y_i(s)\}$. By (8) we have

$$\|\mathcal{R}''\| < \frac{r_d}{54dt} + \frac{r_d}{3t};$$

so we can apply (7) to the partition \mathcal{R}'' and the collection of points $y_i(s)$.

This and inequality (13) yield

$$\begin{aligned}
\frac{d}{ds} \left[\frac{1}{N} \sum_{i=1}^N P(y_i(s)) \right] &\geq \frac{1}{N} \sum_{i=1}^N |\nabla P(y_i(s))| - \frac{1}{6\sqrt{d}} \\
(14) \qquad \qquad \qquad &\geq \frac{1}{3\sqrt{d}} \int_{S^d} |\nabla P(x)| d\mu_d(x) - \frac{1}{6\sqrt{d}} = \frac{1}{6\sqrt{d}},
\end{aligned}$$

for each $P \in \partial\Omega$ and $s \in [0, r_d/3t]$. Finally, equation (11) and inequalities (12) and (14) imply

$$(15) \qquad \frac{1}{N} \sum_{i=1}^N P(x_i(P)) > \frac{1}{6\sqrt{d}} \frac{r_d}{3t} - \frac{r_d}{18\sqrt{d}t} = 0.$$

Lemma 1 is proved. □

References

- [1] B. BAJNOK, *Construction of spherical t -designs*, *Geom. Dedicata*, 43 (1992), 167–179.
- [2] E. BANNAI AND R.M. DAMERELL, *Tight spherical designs I*, *J. Math. Soc. Japan* 31 (1979), pp. 199–207.
- [3] E. BANNAI AND R.M. DAMERELL, *Tight spherical designs. II*, *J. London Math. Soc.* 21 (1980), pp. 13–30.
- [4] A. BONDARENKO AND M. VIAZOVSKA, *Spherical designs via Brouwer fixed point theorem*, *SIAM J. Discrete Math.*, 24 (2010), 207–217.
- [5] J. BOURGAIN AND J. LINDENSTRAUSS, *Distribution of points on spheres and approximation by zonotopes*, *Israel J. Math.*, 64 (1988), 25–31.
- [6] P. BOYVALENKOV, D. DANEV, AND S. NIKOVA, *Nonexistence of certain spherical designs of odd strengths and cardinalities*, *Discrete Comput. Geom.* 21 (1999), 143–156.
- [7] H. COHN, J. H. CONWAY, N. D. ELKIES, AND A. KUMAR, *The D_4 root system is not universally optimal*, *Experiment. Math.*, 16 (2007), 313–320.
- [8] H. COHN AND A. KUMAR, *Universally optimal distribution of points on spheres*, *J. Amer. Math. Soc.*, 20 (2007), 99–148.
- [9] X. CHEN, A. FROMMER, AND B. LANG, *Computational existence proofs for spherical t -designs*, *Numerische Mathematik*, 117 (2011), 289–305.
- [10] X. CHEN AND R. S. WOMERSLEY, *Existence of solutions to systems of underdetermined equations and spherical designs*, *SIAM Journal on Numerical Analysis* 44, 6 (2006), 2326 - 2341.

- [11] J. H. CONWAY AND N. J. A. SLOANE, *Sphere packings, lattices and groups.*, 3rd ed., Springer, New York, 1999.
- [12] P. DELSARTE, J. M. GOETHALS, AND J. J. SEIDEL, *Spherical codes and designs*, *Geom. Dedicata*, 6 (1977), 363–388.
- [13] R. H. HARDIN AND N. J. A. SLOANE, *McLaren’s Improved Snub Cube and Other New Spherical Designs in Three Dimensions*, *Discrete Comput. Geom.*, 15 (1996), 429–441.
- [14] J. KOREVAAR AND J. L. H. MEYERS, *Spherical Faraday cage for the case of equal point charges and Chebyshev-type quadrature on the sphere*, *Integral Transforms Spec. Funct.* 1 (1993), 105–117.
- [15] A. B. J. KUIJLAARS AND E. B. SAFF, *Asymptotics for minimal discrete energy on the sphere*, *Trans. Amer. Math. Soc.*, 350 (1998), 523–538.
- [16] H. N. MHASKAR, F. J. NARCOWICH, AND J. D. WARD, *Spherical Marcinkiewicz-Zygmund inequalities and positive quadrature*, *Math. Comp.*, 70 (2001), 1113–1130.
- [17] O. R. MUSIN, *The kissing number in four dimensions*, *Annals of Math.*, 68 (2008), 1–32.
- [18] DONAL O’REGAN, YEOL JE CHO AND YU QING CHEN, *Topological degree theory and applications*, Chapman & Hall/CRC, 2006.
- [19] E. B. SAFF AND A. B. J. KUIJLAARS, *Distributing many points on a sphere*, *Math. Intelligencer*, 19 (1997), 5–11.
- [20] P. D. SEYMOUR AND T. ZASLAVSKY, *Averaging sets: a generalization of mean values and spherical designs*, *Adv. Math.*, 52 (1984), 213–240.
- [21] G. WAGNER, *On averaging sets*, *Monatsh. Math.*, 111 (1991), 69–78.

- [22] V. A. YUDIN, Lower bounds for spherical designs, *Izv. Ross. Akad. Nauk. Ser. Mat.* 61 (1997), 211–233. English transl., *Izv. Math.* 61 (1997), 673–683.

Centre de Recerca Matemàtica, Campus de Bellaterra, 08193 Bellaterra (Barcelona), Spain
and

Department of Mathematical Analysis, National Taras Shevchenko University, str. Volodymyrska,
64, Kyiv, 01033, Ukraine

Email address: andriybond@gmail.com

Department of Mathematical Analysis, National Taras Shevchenko University, str. Volodymyrska,
64, Kyiv, 01033, Ukraine

Email address: danradchenko@gmail.com

Max Planck Institute for Mathematics, Vivatsgasse 7, 53111 Bonn, Germany

Email address: viazovsk@mpim-bonn.mpg.de