

# Optimal Consumption Policies in Illiquid Markets

Alessandra Cretarola

LUISS Guido Carli, Rome

*joint work with:*

Fausto Gozzi, LUISS Guido Carli, Rome

Huyên Pham, Université Paris 7, CREST-ENSAE and Institut Universitaire de France, Paris

Peter Tankov, Ecole Polytechnique, Palaiseau

GIORNATA DI FINANZA MATEMATICA, CNR-IMATI  
Milan, 8th April 2009

# Outline

- 1 Pham-Tankov model and problem formulation.
- 2 Dynamic Programming and first-order coupled nonlinear IPDE.
- 3 Regularity results:  $C^1$  regularity for the value functions.
- 4 Existence and characterization of optimal strategies:
  - ▶ feedback representation form;
  - ▶ solution of the Euler-Lagrange ODE.
- 5 Numerical illustrations.

# Outline

- 1 Pham-Tankov model and problem formulation.
- 2 Dynamic Programming and first-order coupled nonlinear IPDE.
- 3 Regularity results:  $C^1$  regularity for the value functions.
- 4 Existence and characterization of optimal strategies:
  - ▶ feedback representation form;
  - ▶ solution of the Euler-Lagrange ODE.
- 5 Numerical illustrations.

# Outline

- 1 Pham-Tankov model and problem formulation.
- 2 Dynamic Programming and first-order coupled nonlinear IPDE.
- 3 Regularity results:  $C^1$  regularity for the value functions.
- 4 Existence and characterization of optimal strategies:
  - ▶ feedback representation form;
  - ▶ solution of the Euler-Lagrange ODE.
- 5 Numerical illustrations.

# Outline

- 1 Pham-Tankov model and problem formulation.
- 2 Dynamic Programming and first-order coupled nonlinear IPDE.
- 3 Regularity results:  $C^1$  regularity for the value functions.
- 4 Existence and characterization of optimal strategies:
  - ▶ feedback representation form;
  - ▶ solution of the Euler-Lagrange ODE.
- 5 Numerical illustrations.

# Outline

- 1 Pham-Tankov model and problem formulation.
- 2 Dynamic Programming and first-order coupled nonlinear IPDE.
- 3 Regularity results:  $C^1$  regularity for the value functions.
- 4 Existence and characterization of optimal strategies:
  - ▶ feedback representation form;
  - ▶ solution of the Euler-Lagrange ODE.
- 5 Numerical illustrations.

# Outline

- 1 Pham-Tankov model and problem formulation.
- 2 Dynamic Programming and first-order coupled nonlinear IPDE.
- 3 Regularity results:  $C^1$  regularity for the value functions.
- 4 Existence and characterization of optimal strategies:
  - ▶ feedback representation form;
  - ▶ solution of the Euler-Lagrange ODE.
- 5 Numerical illustrations.

# Outline

- 1 Pham-Tankov model and problem formulation.
- 2 Dynamic Programming and first-order coupled nonlinear IPDE.
- 3 Regularity results:  $C^1$  regularity for the value functions.
- 4 Existence and characterization of optimal strategies:
  - ▶ feedback representation form;
  - ▶ solution of the Euler-Lagrange ODE.
- 5 Numerical illustrations.



# 1. The Pham-Tankov model and problem formulation

## 1.a The Pham-Tankov model of liquidity risk

### Optimal portfolio/consumption problem:

- *The investor has access to a market in which an illiquid asset (stock or fund) is traded;*
- *The asset price is observed only at random times;*
- *Discrete trading is possible only at these random times;*
- *The investor may consume continuously from her/his cash holding.*



Nonstandard mixed discrete/continuous stochastic control problem

⇒ new type of dynamic programming equation.

## 1.a The Pham-Tankov model of liquidity risk

### Optimal portfolio/consumption problem:

- *The investor has access to a market in which an **illiquid** asset (stock or fund) is traded;*
- *The asset price is observed only at random times;*
- *Discrete trading is possible only at these random times;*
- *The investor may consume continuously from her/his cash holding.*



Nonstandard **mixed discrete/continuous** stochastic control problem

⇒ new type of dynamic programming equation.

## 1.a The Pham-Tankov model of liquidity risk

### Optimal portfolio/consumption problem:

- *The investor has access to a market in which an **illiquid** asset (stock or fund) is traded;*
- *The asset price is observed only at random times;*
  - *Discrete trading is possible only at these random times;*
  - *The investor may consume continuously from her/his cash holding.*



Nonstandard **mixed discrete/continuous** stochastic control problem

⇒ new type of dynamic programming equation.

## 1.a The Pham-Tankov model of liquidity risk

### Optimal portfolio/consumption problem:

- *The investor has access to a market in which an **illiquid** asset (stock or fund) is traded;*
- *The asset price is observed only at random times;*
- *Discrete trading is possible only at these random times;*
- *The investor may consume continuously from her/his cash holding.*



Nonstandard **mixed discrete/continuous** stochastic control problem

⇒ new type of dynamic programming equation.

## 1.a The Pham-Tankov model of liquidity risk

### Optimal portfolio/consumption problem:

- *The investor has access to a market in which an **illiquid** asset (stock or fund) is traded;*
- *The asset price is observed only at random times;*
- *Discrete trading is possible only at these random times;*
- *The investor may consume continuously from her/his cash holding.*



Nonstandard **mixed discrete/continuous** stochastic control problem

⇒ new type of dynamic programming equation.

## 1.a The Pham-Tankov model of liquidity risk

### Optimal portfolio/consumption problem:

- *The investor has access to a market in which an **illiquid** asset (stock or fund) is traded;*
- *The asset price is observed only at random times;*
- *Discrete trading is possible only at these random times;*
- *The investor may consume continuously from her/his cash holding.*



Nonstandard **mixed discrete/continuous** stochastic control problem

⇒ new type of dynamic programming equation.

## 1.a The Pham-Tankov model of liquidity risk

### Optimal portfolio/consumption problem:

- *The investor has access to a market in which an **illiquid** asset (stock or fund) is traded;*
- *The asset price is observed only at random times;*
- *Discrete trading is possible only at these random times;*
- *The investor may consume continuously from her/his cash holding.*



Nonstandard **mixed discrete/continuous** stochastic control problem

⇒ new type of dynamic programming equation.



The model is first introduced in the paper:

[PT 07(a)] H. Pham and P. Tankov, (2009) **A coupled system of integrodifferential equations arising in a liquidity risk model**, *Applied Mathematics and Optimization* **59**, 147-173,

where they investigated various theoretical issues, including the viscosity characterization of the value functions.

Then, a convergent numerical algorithm to compute the value functions and several numerical illustrations are provided in the paper:

[PT 07(b)] H. Pham and P. Tankov, (2008) **A model of optimal consumption under liquidity risk with random trading times**, *Mathematical Finance* **18** (4) 613-627.

The model is first introduced in the paper:

[PT 07(a)] H. Pham and P. Tankov, (2009) **A coupled system of integrodifferential equations arising in a liquidity risk model**, *Applied Mathematics and Optimization* **59**, 147-173,

where they investigated various theoretical issues, including the viscosity characterization of the value functions.

Then, a convergent numerical algorithm to compute the value functions and several numerical illustrations are provided in the paper:

[PT 07(b)] H. Pham and P. Tankov, (2008) **A model of optimal consumption under liquidity risk with random trading times**, *Mathematical Finance* **18** (4) 613-627.

## 1.b Problem formulation

- Stock price  $S$  is observed and traded only at exogenous random times  $(\tau_k)_{k \geq 0}$  with  $\tau_0 = 0 < \tau_1 < \dots < \tau_k < \dots$ ;
- The investor may consume continuously from the bank account between two trading dates.

### *Continuous observation filtration*

$$\mathbb{G}^c = (\mathcal{G}_t)_{t \geq 0}, \quad \mathcal{G}_t = \sigma\{(\tau_k, S_{\tau_k}) : \tau_k \leq t\}$$

### *Discrete observation filtration*

$$\mathbb{G}^d = (\mathcal{G}_{\tau_k})_{k \geq 0}.$$

## 1.b Problem formulation

- Stock price  $S$  is observed and traded only at exogenous random times  $(\tau_k)_{k \geq 0}$  with  $\tau_0 = 0 < \tau_1 < \dots < \tau_k < \dots$ ;
- The investor may consume continuously from the bank account between two trading dates.

### *Continuous observation filtration*

$$\mathbb{G}^c = (\mathcal{G}_t)_{t \geq 0}, \quad \mathcal{G}_t = \sigma\{(\tau_k, S_{\tau_k}) : \tau_k \leq t\}$$

### *Discrete observation filtration*

$$\mathbb{G}^d = (\mathcal{G}_{\tau_k})_{k \geq 0}.$$

## 1.b Problem formulation

- Stock price  $S$  is observed and traded only at exogenous random times  $(\tau_k)_{k \geq 0}$  with  $\tau_0 = 0 < \tau_1 < \dots < \tau_k < \dots$ ;
- The investor may consume continuously from the bank account between two trading dates.

### *Continuous observation filtration*

$$\mathbb{G}^c = (\mathcal{G}_t)_{t \geq 0}, \quad \mathcal{G}_t = \sigma\{(\tau_k, S_{\tau_k}) : \tau_k \leq t\}$$

### *Discrete observation filtration*

$$\mathbb{G}^d = (\mathcal{G}_{\tau_k})_{k \geq 0}.$$

## 1.b Problem formulation

- Stock price  $S$  is observed and traded only at exogenous random times  $(\tau_k)_{k \geq 0}$  with  $\tau_0 = 0 < \tau_1 < \dots < \tau_k < \dots$ ;
- The investor may consume continuously from the bank account between two trading dates.

### *Continuous observation filtration*

$$\mathbb{G}^c = (\mathcal{G}_t)_{t \geq 0}, \quad \mathcal{G}_t = \sigma\{(\tau_k, S_{\tau_k}) : \tau_k \leq t\}$$

### *Discrete observation filtration*

$$\mathbb{G}^d = (\mathcal{G}_{\tau_k})_{k \geq 0}.$$

- **Control policy** : mixed discrete/continuous time process  $(\alpha, c)$  :

★  $\alpha = (\alpha_k)_{k \geq 1}$  is a real-valued  $\mathbb{G}^d$ -predictable process :

$\alpha_k$  represents the *amount of stock* invested for the period  $(\tau_{k-1}, \tau_k]$  after observing the stock price  $S_{\tau_{k-1}}$  at time  $\tau_{k-1}$

★  $c = (c_t)_{t \geq 0}$  is a nonnegative  $\mathbb{G}^c$ -predictable process :

$c_t$  represents the *consumption rate* at time  $t$  based on the observation of random arrival times and stock prices until  $t$ .

- **Control policy** : mixed discrete/continuous time process  $(\alpha, c)$  :

★  $\alpha = (\alpha_k)_{k \geq 1}$  is a real-valued  $\mathbb{G}^d$ -predictable process :

$\alpha_k$  represents the *amount of stock* invested for the period  $(\tau_{k-1}, \tau_k]$  after observing the stock price  $S_{\tau_{k-1}}$  at time  $\tau_{k-1}$

★  $c = (c_t)_{t \geq 0}$  is a nonnegative  $\mathbb{G}^c$ -predictable process :

$c_t$  represents the *consumption rate* at time  $t$  based on the observation of random arrival times and stock prices until  $t$ .



- **Wealth process** : starting from an initial capital  $x \geq 0$ , and given a strategy  $(\alpha, c)$ , the wealth  $X_k^x$  of the investor at time  $\tau_k$  is :

- **Admissible control policy** : given  $x \geq 0$ , we say that  $(\alpha, c)$  is **admissible**, and we denote  $(\alpha, c) \in \mathcal{A}(x)$ , if :

$$X_k^x \geq 0, \quad a.s. \quad \forall k \geq 1.$$

- **Wealth process** : starting from an initial capital  $x \geq 0$ , and given a strategy  $(\alpha, c)$ , the wealth  $X_k^x$  of the investor at time  $\tau_k$  is :

$$X_k^x = x - \int_0^{\tau_k} c_t dt + \sum_{i=1}^k \alpha_i \frac{S_{\tau_i} - S_{\tau_{i-1}}}{S_{\tau_{i-1}}}, \quad k \geq 1,$$

- **Admissible control policy** : given  $x \geq 0$ , we say that  $(\alpha, c)$  is **admissible**, and we denote  $(\alpha, c) \in \mathcal{A}(x)$ , if :

$$X_k^x \geq 0, \quad \text{a.s.} \quad \forall k \geq 1.$$

- **Wealth process** : starting from an initial capital  $x \geq 0$ , and given a strategy  $(\alpha, c)$ , the wealth  $X_k^x$  of the investor at time  $\tau_k$  is :

$$X_k^x = x - \int_0^{\tau_k} c_t dt + \sum_{i=1}^k \alpha_i Z_i, \quad k \geq 1,$$

where

$$Z_k = \frac{S_{\tau_k} - S_{\tau_{k-1}}}{S_{\tau_{k-1}}}$$

is the observed return process valued in  $(-1, \infty)$ .

- **Admissible control policy** : given  $x \geq 0$ , we say that  $(\alpha, c)$  is **admissible**, and we denote  $(\alpha, c) \in \mathcal{A}(x)$ , if :

$$X_k^x \geq 0, \quad \text{a.s.} \quad \forall k \geq 1.$$

- **Wealth process** : starting from an initial capital  $x \geq 0$ , and given a strategy  $(\alpha, c)$ , the wealth  $X_k^x$  of the investor at time  $\tau_k$  is :

$$X_k^x = x - \int_0^{\tau_k} c_t dt + \sum_{i=1}^k \alpha_i Z_i, \quad k \geq 1,$$

where

$$Z_k = \frac{S_{\tau_k} - S_{\tau_{k-1}}}{S_{\tau_{k-1}}}$$

is the observed return process valued in  $(-1, \infty)$ .

- **Admissible control policy** : given  $x \geq 0$ , we say that  $(\alpha, c)$  is **admissible**, and we denote  $(\alpha, c) \in \mathcal{A}(x)$ , if :

$$X_k^x \geq 0, \quad a.s. \quad \forall k \geq 1.$$

● **Assumption (1).** Conditions on  $(\tau_k, Z_k)$  :

a)  $\{\tau_k\}_{k \geq 1}$  is the sequence of jump times of a Poisson process with intensity  $\lambda$ .

b) (i) For all  $k \geq 1$ , conditionally on the interarrival time  $\tau_k - \tau_{k-1} = t$ ,  $Z_k$  is independent from  $\{\tau_i, Z_i\}_{i < k}$  and has a distribution  $p(t, dz)$ .

(ii) The support of  $p(t, dz)$  is

- ▶ either an interval with interior equal to  $(-\underline{z}, \bar{z})$ ,  $\underline{z} \in (0, 1]$  and  $\bar{z} \in (0, +\infty]$ ;
- ▶ or it is finite equal to  $\{-\underline{z}, \dots, \bar{z}\}$ ,  $\underline{z} \in (0, 1]$  and  $\bar{z} \in (0, +\infty)$ .

Go

c)  $\int zp(t, dz) \geq 0$ ,  $\forall t \geq 0$  and there exist some  $\kappa, b \in \mathbb{R}_+$  s.t.

$$\int (1+z)p(t, dz) \leq \kappa e^{bt}, \quad \forall t \geq 0.$$

d) The following continuity condition is fulfilled by the measure  $p(t, dz)$ :

$$\lim_{t \rightarrow b} \int w(z)p(t, dz) = \int w(z)p(b, dz), \quad \forall b \geq 0,$$

for all measurable functions  $w$  on  $(-\underline{z}, \bar{z})$  with at most linear growth.

● **Assumption (1).** Conditions on  $(\tau_k, Z_k)$  :

a)  $\{\tau_k\}_{k \geq 1}$  is the sequence of jump times of a Poisson process with intensity  $\lambda$ .

b) (i) For all  $k \geq 1$ , conditionally on the interarrival time  $\tau_k - \tau_{k-1} = t$ ,  $Z_k$  is independent from  $\{\tau_i, Z_i\}_{i < k}$  and has a distribution  $p(t, dz)$ .

(ii) The **support of  $p(t, dz)$**  is

- ▶ either an interval with interior equal to  $(-\underline{z}, \bar{z})$ ,  $\underline{z} \in (0, 1]$  and  $\bar{z} \in (0, +\infty]$ ;
- ▶ or it is finite equal to  $\{-\underline{z}, \dots, \bar{z}\}$ ,  $\underline{z} \in (0, 1]$  and  $\bar{z} \in (0, +\infty)$ .

◀ Go

c)  $\int zp(t, dz) \geq 0$ ,  $\forall t \geq 0$  and there exist some  $\kappa, b \in \mathbb{R}_+$  s.t.

$$\int (1+z)p(t, dz) \leq \kappa e^{bt}, \quad \forall t \geq 0.$$

d) The following continuity condition is fulfilled by the measure  $p(t, dz)$ :

$$\lim_{t \rightarrow b} \int w(z)p(t, dz) = \int w(z)p(b, dz), \quad \forall b \geq 0,$$

for all measurable functions  $w$  on  $(-\underline{z}, \bar{z})$  with at most linear growth.

● **Assumption (1).** Conditions on  $(\tau_k, Z_k)$  :

a)  $\{\tau_k\}_{k \geq 1}$  is the sequence of jump times of a Poisson process with intensity  $\lambda$ .

b) (i) For all  $k \geq 1$ , conditionally on the interarrival time  $\tau_k - \tau_{k-1} = t$ ,  $Z_k$  is independent from  $\{\tau_i, Z_i\}_{i < k}$  and has a distribution  $p(t, dz)$ .

(ii) The support of  $p(t, dz)$  is

- ▶ either an interval with interior equal to  $(-\underline{z}, \bar{z})$ ,  $\underline{z} \in (0, 1]$  and  $\bar{z} \in (0, +\infty]$ ;
- ▶ or it is finite equal to  $\{-\underline{z}, \dots, \bar{z}\}$ ,  $\underline{z} \in (0, 1]$  and  $\bar{z} \in (0, +\infty)$ .

◀ Go

c)  $\int zp(t, dz) \geq 0$ ,  $\forall t \geq 0$  and there exist some  $\kappa, b \in \mathbb{R}_+$  s.t.

$$\int (1+z)p(t, dz) \leq \kappa e^{bt}, \quad \forall t \geq 0.$$

d) The following continuity condition is fulfilled by the measure  $p(t, dz)$ :

$$\lim_{t \rightarrow b} \int w(z)p(t, dz) = \int w(z)p(b, dz), \quad \forall b \geq 0,$$

for all measurable functions  $w$  on  $(-\underline{z}, \bar{z})$  with at most linear growth.

● **Assumption (1).** Conditions on  $(\tau_k, Z_k)$  :

- a)  $\{\tau_k\}_{k \geq 1}$  is the sequence of jump times of a Poisson process with intensity  $\lambda$ .
- b) (i) For all  $k \geq 1$ , conditionally on the interarrival time  $\tau_k - \tau_{k-1} = t$ ,  $Z_k$  is independent from  $\{\tau_i, Z_i\}_{i < k}$  and has a distribution  $p(t, dz)$ .
- (ii) The **support of  $p(t, dz)$**  is
- ▶ either an interval with interior equal to  $(-\underline{z}, \bar{z})$ ,  $\underline{z} \in (0, 1]$  and  $\bar{z} \in (0, +\infty]$ ;
  - ▶ or it is finite equal to  $\{-\underline{z}, \dots, \bar{z}\}$ ,  $\underline{z} \in (0, 1]$  and  $\bar{z} \in (0, +\infty)$ .

◀ Go

- c)  $\int zp(t, dz) \geq 0$ ,  $\forall t \geq 0$  and there exist some  $\kappa, b \in \mathbb{R}_+$  s.t.

$$\int (1+z)p(t, dz) \leq \kappa e^{bt}, \quad \forall t \geq 0.$$

- d) The following continuity condition is fulfilled by the measure  $p(t, dz)$ :

$$\lim_{t \rightarrow t_0} \int w(z)p(t, dz) = \int w(z)p(t_0, dz), \quad \forall t_0 \geq 0,$$

for all measurable functions  $w$  on  $(-\underline{z}, \bar{z})$  with at most linear growth.



- **Typical Examples:**

- ★ **S** extracted from a Black-Scholes model :  $dS_t = bS_t dt + \sigma S_t dW_t$ . Then  $p(t, dz)$  is the distribution of

$$Z(t) = \exp \left[ \left( b - \frac{\sigma^2}{2} \right) t + \sigma W_t \right] - 1,$$

with support  $(-1, \infty)$ , and condition **c)** of **Assumption (1)** is clearly satisfied :  $\int (1+z)p(t, dz) = e^{bt}$ .

- ★ Return process  $Z_k$  independent of waiting times  $\tau_k - \tau_{k-1}$ , so  $p(t, dz) \equiv p(dz)$ . In particular  $p(dz)$  may be a discrete distribution with finite support in  $(-1, +\infty)$ .

- **Typical Examples:**

- ★ **S** extracted from a Black-Scholes model :  $dS_t = bS_t dt + \sigma S_t dW_t$ . Then  $p(t, dz)$  is the distribution of

$$Z(t) = \exp \left[ \left( b - \frac{\sigma^2}{2} \right) t + \sigma W_t \right] - 1,$$

with support  $(-1, \infty)$ , and condition **c)** of **Assumption (1)** is clearly satisfied :  $\int (1+z)p(t, dz) = e^{bt}$ .

- ★ Return process  $Z_k$  independent of waiting times  $\tau_k - \tau_{k-1}$ , so  $p(t, dz) \equiv p(dz)$ . In particular  $p(dz)$  may be a discrete distribution with finite support in  $(-1, +\infty)$ .

- **Optimal portfolio/consumption problem:**

## Value function

$$v(x) = \sup_{(\alpha, c) \in \mathcal{A}(x)} \mathbb{E} \left[ \int_0^{+\infty} e^{-\rho t} U(c_t) dt \right], \quad x \geq 0.$$

### Assumption (2)

- ▶ Utility function  $U : \mathbb{R}_+ \rightarrow \mathbb{R}$ ,  $U(0) = 0$ ,  $C^1$ , strictly increasing, strictly concave, satisfying the Inada conditions  $U'(0) = \infty$ ,  $U'(\infty) = 0$ , and the growth condition

$$U(w) \leq K_1 w^\gamma, \quad \gamma \in (0, 1).$$

- ▶ Discount factor  $\rho$  such that

$$\rho > b\gamma + \lambda \left( \frac{\kappa^\gamma}{\underline{z}^\gamma} - 1 \right).$$

- **Optimal portfolio/consumption problem:**

## Value function

$$v(x) = \sup_{(\alpha, c) \in \mathcal{A}(x)} \mathbb{E} \left[ \int_0^{+\infty} e^{-\rho t} U(c_t) dt \right], \quad x \geq 0.$$

### Assumption (2)

- ▶ Utility function  $U : \mathbb{R}_+ \rightarrow \mathbb{R}$ ,  $U(0) = 0$ ,  $C^1$ , strictly increasing, strictly concave, satisfying the Inada conditions  $U'(0) = \infty$ ,  $U'(\infty) = 0$ , and the growth condition

$$U(w) \leq K_1 w^\gamma, \quad \gamma \in (0, 1).$$

- ▶ Discount factor  $\rho$  such that

$$\rho > b\gamma + \lambda \left( \frac{\kappa^\gamma}{z^\gamma} - 1 \right).$$

- **Optimal portfolio/consumption problem:**

## Value function

$$v(x) = \sup_{(\alpha, c) \in \mathcal{A}(x)} \mathbb{E} \left[ \int_0^{+\infty} e^{-\rho t} U(c_t) dt \right], \quad x \geq 0.$$

### Assumption (2)

- ▶ Utility function  $U : \mathbb{R}_+ \rightarrow \mathbb{R}$ ,  $U(0) = 0$ ,  $C^1$ , strictly increasing, strictly concave, satisfying the Inada conditions  $U'(0) = \infty$ ,  $U'(\infty) = 0$ , and the growth condition

$$U(w) \leq K_1 w^\gamma, \quad \gamma \in (0, 1).$$

- ▶ Discount factor  $\rho$  such that

$$\rho > b\gamma + \lambda \left( \frac{\kappa^\gamma}{\underline{z}^\gamma} - 1 \right).$$

## 2. Dynamic Programming

- Dynamic programming is a mathematical technique for making a sequence of interrelated decisions, which can be applied to many optimization problems (including optimal control problems). Ex: optimal investment problems, consumer maximization problems in macroeconomic models, etc.
- Basic idea: consider a family of optimal control problems with different initial times and states and establish relationships among these problems via the so-called Hamilton-Jacobi-Bellman equation (HJB), which is a nonlinear first-order (in the deterministic case) or second-order (in the stochastic case) partial differential equation.  
If the HJB equation is solvable, then one can obtain an optimal feedback control by taking the maximizer/minimizer of the Hamiltonian or generalized Hamiltonian involved in the HJB equation. This is the so-called *verification technique*.
- It is required that the HJB equation admits *classical solutions*, meaning that the solutions have to be smooth enough.

- To overcome this difficulty a weaker definition of solutions was introduced.  
*Viscosity solutions*: a kind of nonsmooth solutions to partial differential equations, where the definition of solution is given in terms of sub-superdifferential of the function.  
This concept of solution usually provides a characterization of the value function of the control problem as unique solution of the HJB equation.
- **Weak point**: such a characterization is not good to use in order to construct optimal controls and regularity results for this kind of solutions are needed to go ahead.



## 2.a Dynamic Programming Principle (DPP)

Relation on the value function by considering two consecutive trading dates :

$$\begin{aligned}v(x) &= \sup_{(\alpha, c) \in \mathcal{A}(x)} \mathbb{E} \left[ \int_0^{\tau_1} e^{-\rho t} U(c_t) dt + e^{-\rho \tau_1} v(X_1^x) \right], \\ &= \sup_{(a, c) \in A_d(x)} \mathbb{E} \left[ \int_0^{\tau_1} e^{-\rho t} U(c_t) dt + e^{-\rho \tau_1} v(x - \int_0^{\tau_1} c_t dt + aZ_1) \right],\end{aligned}$$

## 2.a Dynamic Programming Principle (DPP)

Relation on the value function by considering two consecutive trading dates :

$$\begin{aligned} v(x) &= \sup_{(\alpha, c) \in \mathcal{A}(x)} \mathbb{E} \left[ \int_0^{\tau_1} e^{-\rho t} U(c_t) dt + e^{-\rho \tau_1} v(X_1^x) \right], \\ &= \sup_{(a, c) \in A_d(x)} \mathbb{E} \left[ \int_0^{\tau_1} e^{-\rho t} U(c_t) dt + e^{-\rho \tau_1} v(x - \int_0^{\tau_1} c_t dt + aZ_1) \right], \end{aligned}$$

$A_d(x)$  : pair of **deterministic** constants  $a$  and nonnegative **deterministic** processes  $c = (c_t)_{t \geq 0}$  s.t. :  $x - \int_0^{\tau_1} c_t dt + aZ_1 \geq 0$  a.s. , i.e.

$$-\frac{x}{\underline{z}} \leq a \leq \frac{x}{\underline{z}}$$

$$x - \int_0^t c_u du \geq l(a), \quad \forall t \geq 0, \text{ with } l(a) = \max(a\underline{z}, -a\bar{z}) \quad (c \in \mathcal{C}_a(x)).$$

## 2.a Dynamic Programming Principle (DPP)

Relation on the value function by considering two consecutive trading dates :

$$\begin{aligned} v(x) &= \sup_{(\alpha, c) \in \mathcal{A}(x)} \mathbb{E} \left[ \int_0^{\tau_1} e^{-\rho t} U(c_t) dt + e^{-\rho \tau_1} v(X_1^x) \right], \\ &= \sup_{(a, c) \in A_d(x)} \mathbb{E} \left[ \int_0^{\tau_1} e^{-\rho t} U(c_t) dt + e^{-\rho \tau_1} v(x - \int_0^{\tau_1} c_t dt + aZ_1) \right], \end{aligned}$$

$A_d(x)$  : pair of **deterministic** constants  $a$  and nonnegative **deterministic** processes  $c = (c_t)_{t \geq 0}$  s.t. :  $x - \int_0^{\tau_1} c_t dt + aZ_1 \geq 0$  a.s. , i.e.

$$-\frac{x}{\bar{z}} \leq a \leq \frac{x}{\underline{z}}$$

$$x - \int_0^t c_u du \geq -az, \quad \forall t \geq 0, \quad \forall z \in (-\underline{z}, \bar{z}) \quad (c \in \mathcal{C}_a(x)).$$

Under conditions **a)** and **b)** of **Assumption (1)** on  $(\tau_1, Z_1)$  **Ass (1)**, we compute the expectation :

$$v(x) = \sup_{\substack{a \in \left[-\frac{x}{\bar{z}}, \frac{x}{\bar{z}}\right] \\ c \in C_a(x)}} \int_0^{+\infty} e^{-(\rho+\lambda)s} \left[ U(c_s) + \lambda \int_0^s v\left(x - \int_0^s c_u du + az\right) p(s, dz) \right] ds.$$

- Dynamic auxiliary deterministic control problem on consumption :

$$\hat{v}(t, x, a) = \sup_{c \in C_a(t, x)} \int_t^{+\infty} e^{-(\rho+\lambda)(s-t)} \left[ U(c_s) + \lambda \int_t^s v(Y_s^{t, x} + az) p(s, dz) \right] ds,$$

where  $C_a(t, x)$  is the set of deterministic nonnegative processes

$c = (c_s)_{s \geq t}$ , s.t.

$$Y_s^{t, x} = x - \int_t^s c_u du \geq l(a), \quad s \geq t,$$

for  $(t, x, a) \in \mathcal{D} := \mathbb{R}_+ \times \mathcal{X}$ , with  $\mathcal{X} = \{(x, a) \in \mathbb{R}_+ \times A : x \geq l(a)\}$ ,  
by setting  $A = \mathbb{R}$  if  $\bar{z} < +\infty$  and  $A = \mathbb{R}_+$  if  $\bar{z} = +\infty$ .

Under conditions **a)** and **b)** of **Assumption (1)** on  $(\tau_1, Z_1)$  **Ass (1)**, we compute the expectation :

$$v(x) = \sup_{\substack{a \in \left[-\frac{x}{\bar{z}}, \frac{x}{\bar{z}}\right] \\ c \in C_a(x)}} \int_0^{+\infty} e^{-(\rho+\lambda)s} \left[ U(c_s) + \lambda \int_0^s v\left(x - \int_0^s c_u du + az\right) p(s, dz) \right] ds.$$

- **Dynamic auxiliary deterministic control problem on consumption :**

$$\hat{v}(t, x, a) = \sup_{c \in C_a(t, x)} \int_t^{+\infty} e^{-(\rho+\lambda)(s-t)} \left[ U(c_s) + \lambda \int_t^s v(Y_s^{t, x} + az) p(s, dz) \right] ds,$$

where  $C_a(t, x)$  is the set of deterministic nonnegative processes  $c = (c_s)_{s \geq t}$ , s.t.

$$Y_s^{t, x} = x - \int_t^s c_u du \geq l(a), \quad s \geq t,$$

for  $(t, x, a) \in \mathcal{D} := \mathbb{R}_+ \times \mathcal{X}$ , with  $\mathcal{X} = \{(x, a) \in \mathbb{R}_+ \times A : x \geq l(a)\}$ , by setting  $A = \mathbb{R}$  if  $\bar{z} < +\infty$  and  $A = \mathbb{R}_+$  if  $\bar{z} = +\infty$ .

## 2.b The equivalent *coupled* deterministic optimization problem

- For each  $a \in A$ , a deterministic control problem :

$$\hat{v}(t, x, a) = \sup_{c \in C_a(t, x)} \int_t^{+\infty} e^{-(\rho+\lambda)(s-t)} \left[ U(c_s) + \lambda \int v(Y_s^{t,x} + az) p(s, dz) \right] ds,$$

- A classical (scalar) maximum problem on a concave function

$$v(x) = \sup_{a \in \left[-\frac{x}{Z}, \frac{x}{Z}\right]} \hat{v}(0, x, a), \quad x \geq 0,$$

★ The difficulty comes from this coupling between  $\hat{v}$  and  $v$ .

## 2.b The equivalent *coupled* deterministic optimization problem

- For each  $a \in A$ , a deterministic control problem :

$$\hat{v}(t, x, a) = \sup_{c \in C_a(t, x)} \int_t^{+\infty} e^{-(\rho+\lambda)(s-t)} \left[ U(c_s) + \lambda \int v(Y_s^{t,x} + az) p(s, dz) \right] ds,$$

- A classical (scalar) maximum problem on a concave function

$$v(x) = \sup_{a \in \left[-\frac{x}{Z}, \frac{x}{Z}\right]} \hat{v}(0, x, a), \quad x \geq 0,$$

★ The difficulty comes from this coupling between  $\hat{v}$  and  $v$ .

## 2.b The equivalent *coupled* deterministic optimization problem

- For each  $a \in A$ , a deterministic control problem :

$$\hat{v}(t, x, a) = \sup_{c \in \mathcal{C}_a(t, x)} \int_t^{+\infty} e^{-(\rho+\lambda)(s-t)} \left[ U(c_s) + \lambda \int v(Y_s^{t,x} + az) p(s, dz) \right] ds,$$

- A classical (scalar) maximum problem on a concave function

$$v(x) = \sup_{a \in \left[-\frac{x}{Z}, \frac{x}{Z}\right]} \hat{v}(0, x, a), \quad x \geq 0,$$

★ The difficulty comes from this coupling between  $\hat{v}$  and  $v$ .



## 2.c The system of coupled nonlinear IPDE

- The Hamilton-Jacobi (HJ) equation associated to the first problem is:

$$(\rho + \lambda)\hat{v} - \frac{\partial \hat{v}}{\partial t} - \tilde{U}\left(\frac{\partial \hat{v}}{\partial x}\right) - \lambda \int \mathbf{v}(x + az)p(t, dz) = 0, \quad (t, x, a) \in \mathcal{D}, \quad (1)$$

where  $\tilde{U}(p) = \sup_{c > 0} [U(c) - cp]$ ,  $p \geq 0$  is the convex conjugate of  $U$ .

- This is coupled with the above:

$$\mathbf{v}(x) := \sup_{a \in \left[-\frac{x}{z}, \frac{x}{z}\right]} \hat{v}(0, x, a), \quad x \geq 0, \quad (2)$$

Dynamic programming  $\longrightarrow$  system of coupled nonlinear IPDE (1)-(2).

## 2.c The system of coupled nonlinear IPDE

- The Hamilton-Jacobi (HJ) equation associated to the first problem is:

$$(\rho + \lambda)\hat{v} - \frac{\partial \hat{v}}{\partial t} - \tilde{U}\left(\frac{\partial \hat{v}}{\partial x}\right) - \lambda \int v(x + az)p(t, dz) = 0, \quad (t, x, a) \in \mathcal{D}, \quad (1)$$

where  $\tilde{U}(p) = \sup_{c>0}[U(c) - cp]$ ,  $p \geq 0$  is the convex conjugate of  $U$ .

- This is coupled with the above:

$$v(x) := \sup_{a \in \left[-\frac{x}{z}, \frac{x}{z}\right]} \hat{v}(0, x, a), \quad x \geq 0, \quad (2)$$

Dynamic programming  $\longrightarrow$  system of coupled nonlinear IPDE (1)-(2).

## 2.c The system of coupled nonlinear IPDE

- The Hamilton-Jacobi (HJ) equation associated to the first problem is:

$$(\rho + \lambda)\hat{v} - \frac{\partial \hat{v}}{\partial t} - \tilde{U}\left(\frac{\partial \hat{v}}{\partial x}\right) - \lambda \int v(x + az)p(t, dz) = 0, \quad (t, x, a) \in \mathcal{D}, \quad (1)$$

where  $\tilde{U}(p) = \sup_{c > 0} [U(c) - cp]$ ,  $p \geq 0$  is the convex conjugate of  $U$ .

- This is coupled with the above:

$$v(x) := \sup_{a \in \left[-\frac{x}{Z}, \frac{x}{Z}\right]} \hat{v}(0, x, a), \quad x \geq 0, \quad (2)$$

Dynamic programming  $\longrightarrow$  system of coupled nonlinear IPDE (1)-(2).

## 2.d Some properties of $v, \hat{v}$

Under **Assumptions (1)-(2)**, the following properties hold:

- $v$  is nondecreasing, concave, continuous on  $\mathbb{R}_+$  with  $v(0) = 0$ ;
- $\hat{v}$  is concave in  $(x, a)$  and continuous on  $\mathcal{D}$ ;
- **Growth condition (G1)** : there exists some positive constant  $K$  s.t.

$$\hat{v}(t, x, a) \leq K(e^{bt}x)^\gamma, \quad \forall (t, x, a) \in \mathcal{D},$$

$$v(x) \leq Kx^\gamma, \quad \forall x \geq 0.$$

- **Boundary data (B1)**  $\longleftrightarrow$  nonnegative wealth constraint :

$$\hat{v}(t, x, a) = \lambda \int_t^{+\infty} e^{-(\rho+\lambda)(s-t)} \int v(x+az)p(s, dz) ds, \quad \forall t \geq 0, (x, a) \in \partial\mathcal{X}$$

◀ go

- ▶  $v$  is strictly increasing on  $\mathbb{R}_+$ ;
- ▶  $\hat{v}$  is strictly increasing in  $x \geq l(a)$ , given  $a \in A$ ;
- ▶ **The scaling relation for power utility**: in the case where

$$U(x) = K_1 x^\gamma, \quad 0 < \gamma < 1,$$

$$\hat{v}(t, \beta x, \beta a) = \beta^\gamma \hat{v}(t, x, a), \quad v(\beta x) = \beta^\gamma v(x).$$

## 2.d Some properties of $v, \hat{v}$

Under **Assumptions (1)-(2)**, the following properties hold:

- $v$  is nondecreasing, concave, continuous on  $\mathbb{R}_+$  with  $v(0) = 0$ ;
- $\hat{v}$  is concave in  $(x, a)$  and continuous on  $\mathcal{D}$ ;
- **Growth condition (G1)** : there exists some positive constant  $K$  s.t.

$$\hat{v}(t, x, a) \leq K(e^{bt}x)^\gamma, \quad \forall (t, x, a) \in \mathcal{D},$$

$$v(x) \leq Kx^\gamma, \quad \forall x \geq 0.$$

- **Boundary data (B1)**  $\longleftrightarrow$  nonnegative wealth constraint :

$$\hat{v}(t, x, a) = \lambda \int_t^{+\infty} e^{-(\rho+\lambda)(s-t)} \int v(x+az)p(s, dz)ds, \quad \forall t \geq 0, (x, a) \in \partial\mathcal{X}$$

◀ go

- ▶  $v$  is strictly increasing on  $\mathbb{R}_+$ ;
- ▶  $\hat{v}$  is strictly increasing in  $x \geq l(a)$ , given  $a \in A$ ;
- ▶ **The scaling relation for power utility**: in the case where

$$U(x) = K_1x^\gamma, \quad 0 < \gamma < 1,$$

$$\hat{v}(t, \beta x, \beta a) = \beta^\gamma \hat{v}(t, x, a), \quad v(\beta x) = \beta^\gamma v(x).$$

### 3. Regularity results

- In the paper [PT 07(a)] the viscosity solution approach is developed giving existence and uniqueness, but no regularity result is proved, so the *verification technique* cannot be used to obtain an optimal control in feedback form.
- In the present paper we prove that the viscosity solution is indeed *regular* in a wide class of cases that includes the examples.
- This allows to get the existence of the optimal control that we characterize both in feedback form in terms of the derivatives of the value functions and as the solution of a second-order ODE.

### 3.a Stationary case

Assume that the distribution  $p(t, dz)$  is *time independent*.

$$p(t, dz) \equiv p(dz).$$

Then the coupled HJ becomes:

$$(\rho + \lambda)\hat{v} - \tilde{U}\left(\frac{\partial \hat{v}}{\partial x}\right) - \lambda \int v(x + az)p(dz) = 0, \quad t \geq 0, x \geq l(a), \quad (3)$$

$$v(x) = \sup_{a \in \left[-\frac{x}{z}, \frac{x}{z}\right]} \hat{v}(x, a), \quad x \in \mathbb{R}_+, \quad (4)$$

#### Theorem

Suppose *Assumptions (1)-(2)* are satisfied. Then

- ★  $\forall a \geq 0$  we have  $\hat{v}(\cdot, a) \in C^2(l(a), +\infty)$  and  $\frac{\partial \hat{v}}{\partial x}(l(a)^+, a) = +\infty$ .
- ★ We have  $v \in C^1(0, +\infty)$  and  $v'(0^+) = +\infty$ .



### 3.a Stationary case

Assume that the distribution  $p(t, dz)$  is *time independent*.

$$p(t, dz) \equiv p(dz).$$

Then the coupled HJ becomes:

$$(\rho + \lambda)\hat{v} - \tilde{U}\left(\frac{\partial \hat{v}}{\partial x}\right) - \lambda \int v(x + az)p(dz) = 0, \quad t \geq 0, x \geq l(a), \quad (3)$$

$$v(x) = \sup_{a \in \left[-\frac{x}{z}, \frac{x}{z}\right]} \hat{v}(x, a), \quad x \in \mathbb{R}_+, \quad (4)$$

#### Theorem

Suppose *Assumptions (1)-(2)* are satisfied. Then

- ★  $\forall a \geq 0$  we have  $\hat{v}(\cdot, a) \in C^2(l(a), +\infty)$  and  $\frac{\partial \hat{v}}{\partial x}(l(a)^+, a) = +\infty$ .
- ★ We have  $v \in C^1(0, +\infty)$  and  $v'(0^+) = +\infty$ .

## ► Idea of the proof

First we prove that  $\hat{v}(\cdot, a) \in C^1(I(a), +\infty)$ . The other regularity properties follow by a kind of bootstrap argument based on the coupling.

Arguing by contradiction, we use:

- ★ the *viscosity characterization* of the value functions to the original control problem by means of viscosity solution to the coupled IPDE (see [PT 07(a)]).
- ★ the concavity of the function  $\hat{v}(\cdot, a)$  together with the strict convexity of the convex conjugate  $\tilde{U}$  with similar arguments to those given in:

[BCD 97] Bardi M. and Capuzzo-Dolcetta I., OPTIMAL CONTROL AND VISCOSITY SOLUTIONS OF HAMILTON-JACOBI-BELLMAN EQUATIONS. Birkhäuser Boston Inc., Boston, MA.

## ► Idea of the proof

First we prove that  $\hat{v}(\cdot, a) \in C^1(I(a), +\infty)$ . The other regularity properties follow by a kind of bootstrap argument based on the coupling.

Arguing by contradiction, we use:

- ★ the *viscosity characterization* of the value functions to the original control problem by means of viscosity solution to the coupled IPDE (see [PT 07(a)]). viscosity solutions
- ★ the concavity of the function  $\hat{v}(\cdot, a)$  together with the strict convexity of the convex conjugate  $\tilde{U}$  with similar arguments to those given in:

[BCD 97] Bardi M. and Capuzzo-Dolcetta I., *OPTIMAL CONTROL AND VISCOSITY SOLUTIONS OF HAMILTON-JACOBI-BELLMAN EQUATIONS*. Birkhäuser Boston Inc., Boston, MA.

sketch of the proof

## ► Idea of the proof

First we prove that  $\hat{v}(\cdot, a) \in C^1(I(a), +\infty)$ . The other regularity properties follow by a kind of bootstrap argument based on the coupling.

Arguing by contradiction, we use:

- ★ the *viscosity characterization* of the value functions to the original control problem by means of viscosity solution to the coupled IPDE (see [PT 07(a)]). viscosity solutions
- ★ the concavity of the function  $\hat{v}(\cdot, a)$  together with the strict convexity of the convex conjugate  $\tilde{U}$  with similar arguments to those given in:

[BCD 97] Bardi M. and Capuzzo-Dolcetta I., **OPTIMAL CONTROL AND VISCOSITY SOLUTIONS OF HAMILTON-JACOBI-BELLMAN EQUATIONS**. Birkhäuser Boston Inc., Boston, MA.

sketch of the proof

## 3.b Nonstationary case

To use the same argument as in the stationary case we need the **joint semiconcavity** in  $(t, x)$  of the function  $\hat{v}(\cdot, \cdot, a)$ . semiconcavity

Hence we need to introduce an additional assumption on the measure  $p(t, dz)$  in order to guarantee it.

**Assumption (3):** for every  $a \in A - \{0\}$ , the map

$$(t, x) \longrightarrow \lambda \int w(x + az)p(t, dz)$$

is (locally) semiconcave for  $(t, x) \in (0, +\infty) \times (l(a), +\infty)$  for all measurable continuous functions  $w$  with linear growth.

## 3.b Nonstationary case

To use the same argument as in the stationary case we need the **joint semiconcavity** in  $(t, x)$  of the function  $\hat{v}(\cdot, \cdot, a)$ . semiconcavity

Hence we need to introduce an additional assumption on the measure  $p(t, dz)$  in order to guarantee it.

**Assumption (3)**: for every  $a \in A - \{0\}$ , the map

$$(t, x) \longrightarrow \lambda \int w(x + az)p(t, dz)$$

is (locally) semiconcave for  $(t, x) \in (0, +\infty) \times (l(a), +\infty)$  for all measurable continuous functions  $w$  with linear growth.

We have the following.

## Theorem

Let *Assumptions (1)-(2)-(3)* stand in force + *local semiconcavity* of  $\hat{v}$ .

- $\forall a \geq 0$  we have  $\hat{v}(\cdot, \cdot, a) \in C^1([0, +\infty) \times (l(a), +\infty))$  and

$$\frac{\partial \hat{v}}{\partial x}(t, l(a)^+, a) = +\infty \text{ for every } t \geq 0.$$

- We have  $v \in C^1(0, +\infty)$  and  $v'(0^+) = +\infty$ .

## 4. Existence and characterization of optimal strategies



## 4.a Feedback representation form of the optimal strategies

Let Assumptions (1)-(2)-(3) stand in force from now on.

Since the viscosity solution to the coupled IPDE is *regular*, we have the following result which provides the optimal control in feedback form.

### Verification Theorem

Let  $(v, \hat{v})$  be the regular value functions solution to:

$$(\rho + \lambda)\hat{v} - \frac{\partial \hat{v}}{\partial t} - \tilde{U}\left(\frac{\partial \hat{v}}{\partial x}\right) - \lambda \int v(x + az)p(t, dz) = 0, \quad (t, x, a) \in \mathcal{D}$$
$$v(x) = \mathcal{H}\hat{v}(x) = \sup_{a \in \left[-\frac{x}{z}, \frac{x}{z}\right]} \hat{v}(0, x, a), \quad x \geq 0,$$

together with the boundary conditions (G1)-(B1).

Then there exists an optimal control policy  $(\alpha^*, c^*)$  given by:

- trading portfolio from the scalar maximum problem

$$\alpha_{k+1}^* = \arg \max_{-\frac{X_k^x}{Z} \leq a \leq \frac{X_k^x}{Z}} \hat{v}(0, X_k^x, a), \quad k \geq 0.$$

- consumption from the deterministic control problem

$$c_t^* = \hat{c} \left( t - \tau_k, Y_t^{(k)}, \alpha_{k+1}^* \right), \quad \tau_k < t \leq \tau_{k+1},$$

with

$$\hat{c}(t, x, a) = \arg \max_{c \geq 0} \left[ U(c) - c \frac{\partial \hat{v}(t, x, a)}{\partial x} \right] = I \left( \frac{\partial \hat{v}(t, x, a)}{\partial x} \right),$$

where  $X_k^x$  is the wealth investor at time  $\tau_k$ ,  $Y_t^{(k)} = X_k^x - \int_{\tau_k}^t c_s^* ds$  and  $I = (U')^{-1}$ .

## 4.b Representation of the optimal strategies as the solution of the Euler-Lagrange ODE

From the regularity results, we can deduce more properties of the optimal consumption policy.

- **Stationary case**

We get an autonomous equation for the optimal consumption policy between two trading dates.

### Proposition

Suppose that  $U \in C^2((0, +\infty))$  with  $U''(x) < 0$ , for all  $x$ . Then the wealth process  $Y$  between two trading dates is twice differentiable and satisfies the second-order ODE:

$$\frac{d^2 Y_t}{dt^2} = \frac{g'(Y_t, a) - (\rho + \lambda)U'(c_t)}{U''(c_t)}, \quad c_t = -\frac{dY_t}{dt}, \quad (5)$$

where

$$g : \mathcal{X} \longrightarrow \mathbb{R}_+, \quad g(x, a) = \lambda \int v(x + az)p(dz).$$

## Case of power utility

In this case, equation (5) takes the form

$$\frac{d^2 Y_t}{dt^2} = \frac{\rho + \lambda}{1 - \gamma} c_t - \frac{1}{K_1 \gamma (1 - \gamma)} c_t^{2-\gamma} g'(Y_t, a), \quad Y_0 = x, \quad Y_\infty = l(a). \quad (6)$$

Then we can deduce a simple *exponential lower bound* on the integrated consumption, corresponding to the solution of (6) in the case  $g \equiv 0$ :

$$Y_t \geq Y_t^0, \quad t \geq 0,$$

with

$$Y_t^0 = x - (x - l(a)) \left( 1 - e^{-\frac{(\rho + \lambda)t}{1 - \gamma}} \right). \quad (7)$$

## ● Nonstationary case

The regularity results for the optimal strategies are weaker and more difficult to prove.

As in the S.C., we can deduce an autonomous equation for the optimal wealth process between two trading dates. However, the proof is different and makes use of the Maximum Principle.

### Proposition

Let  $U \in C^2((0, +\infty))$  with  $U''(x) < 0$ , for all  $x$ . Then the optimal wealth process  $Y$  between two trading dates is twice differentiable and satisfies:

$$\frac{d^2 Y_s}{ds^2} = \frac{\frac{\partial g(s, Y_s, a)}{\partial x} - (\rho + \lambda)U'(c_s)}{U''(c_s)}, \quad c_s = -\frac{dY_s}{ds}, \quad Y_t = x. \quad (8)$$

and  $\lim_{t \rightarrow +\infty} Y_t = I(a)$ .

### Case of power utility

Equation (8) becomes:

$$\frac{d^2 Y_t}{dt^2} = \frac{\rho + \lambda}{1 - \gamma} c_t - \frac{\lambda \theta_1 c_t^{2-\gamma}}{K_1(1 - \gamma)} \int (Y_t + az)^{\gamma-1} p(t, dz), \quad Y_0 = x, \quad Y_\infty = I(a).$$

## ● Nonstationary case

The regularity results for the optimal strategies are weaker and more difficult to prove.

As in the S.C., we can deduce an autonomous equation for the optimal wealth process between two trading dates. However, the proof is different and makes use of the Maximum Principle.

### Proposition

Let  $U \in C^2((0, +\infty))$  with  $U''(x) < 0$ , for all  $x$ . Then the optimal wealth process  $Y$  between two trading dates is twice differentiable and satisfies:

$$\frac{d^2 Y_s}{ds^2} = \frac{\frac{\partial g(s, Y_s, a)}{\partial x} - (\rho + \lambda)U'(c_s)}{U''(c_s)}, \quad c_s = -\frac{dY_s}{ds}, \quad Y_t = x. \quad (8)$$

and  $\lim_{t \rightarrow +\infty} Y_t = I(a)$ .

### Case of power utility

Equation (8) becomes:

$$\frac{d^2 Y_t}{dt^2} = \frac{\rho + \lambda}{1 - \gamma} c_t - \frac{\lambda \theta_1 c_t^{2-\gamma}}{K_1(1 - \gamma)} \int (Y_t + az)^{\gamma-1} p(t, dz), \quad Y_0 = x, \quad Y_\infty = I(a).$$

## • Nonstationary case

The regularity results for the optimal strategies are weaker and more difficult to prove.

As in the S.C., we can deduce an autonomous equation for the optimal wealth process between two trading dates. However, the proof is different and makes use of the Maximum Principle.

### Proposition

Let  $U \in C^2((0, +\infty))$  with  $U''(x) < 0$ , for all  $x$ . Then the optimal wealth process  $Y$  between two trading dates is twice differentiable and satisfies:

$$\frac{d^2 Y_s}{ds^2} = \frac{\frac{\partial g(s, Y_s, a)}{\partial x} - (\rho + \lambda)U'(c_s)}{U''(c_s)}, \quad c_s = -\frac{dY_s}{ds}, \quad Y_t = x. \quad (8)$$

and  $\lim_{t \rightarrow +\infty} Y_t = l(a)$ .

### Case of power utility

Equation (8) becomes:

$$\frac{d^2 Y_t}{dt^2} = \frac{\rho + \lambda}{1 - \gamma} c_t - \frac{\lambda \theta_1 c_t^{2-\gamma}}{K_1(1 - \gamma)} \int (Y_t + az)^{\gamma-1} p(t, dz), \quad Y_0 = x, \quad Y_\infty = l(a).$$

## • Nonstationary case

The regularity results for the optimal strategies are weaker and more difficult to prove.

As in the S.C., we can deduce an autonomous equation for the optimal wealth process between two trading dates. However, the proof is different and makes use of the Maximum Principle.

### Proposition

Let  $U \in C^2((0, +\infty))$  with  $U''(x) < 0$ , for all  $x$ . Then the optimal wealth process  $Y$  between two trading dates is twice differentiable and satisfies:

$$\frac{d^2 Y_s}{ds^2} = \frac{\frac{\partial g(s, Y_s, a)}{\partial x} - (\rho + \lambda)U'(c_s)}{U''(c_s)}, \quad c_s = -\frac{dY_s}{ds}, \quad Y_t = x. \quad (8)$$

and  $\lim_{t \rightarrow +\infty} Y_t = l(a)$ .



### Case of power utility

Equation (8) becomes:

$$\frac{d^2 Y_t}{dt^2} = \frac{\rho + \lambda}{1 - \gamma} c_t - \frac{\lambda \theta_1 c_t^{2-\gamma}}{K_1(1 - \gamma)} \int (Y_t + az)^{\gamma-1} p(t, dz), \quad Y_0 = x, \quad Y_\infty = l(a).$$





## 5.a Numerical illustrations: stationary case



*Left:* typical profile of the optimal wealth process  $Y_t$  and the exponential lower bound (7). *Right:* the corresponding consumption strategies. In the presence of investment opportunities, the agent first consumes slowly but if the investment opportunity does not appear, the agent eventually “gets disappointed” and starts to consume fast.

## 5.b Numerical illustrations: nonstationary case



Optimal wealth (*left*) and consumption policy (*right*) for the probability distribution extracted from the Black-Scholes model (solid line) and from the stationary model having the same distribution as the Black-Scholes model in 3 years' time (dashed line).

Same parameter values as in [PT 07(b)]: drift  $b = 0.4$ , volatility  $\sigma = 1$ , discount factor  $\rho = 0.2$ , intensity  $\lambda = 2$  and risk aversion coefficient  $\gamma = 0.5$ . At least qualitatively, the consumption profile is similar to the one observed in the stationary model, with exponential decay.

## Viscosity solutions

In the paper [PT 07(a)] the viscosity solution approach is developed giving existence and uniqueness. Viscosity solutions for the coupled IPDE :

$$(\rho + \lambda)\hat{v} - \frac{\partial \hat{v}}{\partial t} - \tilde{U}\left(\frac{\partial \hat{v}}{\partial x}\right) - \lambda \int v(x + az)p(t, dz) = 0, \quad t \geq 0, x > l(a),$$
$$v(x) = \mathcal{H}\hat{v}(x) = \sup_{a \in \left[-\frac{x}{z}, \frac{x}{z}\right]} \hat{v}(0, x, a), \quad x \in \mathbb{R}_+,$$

Definition

## Viscosity solutions

In the paper [PT 07(a)] the viscosity solution approach is developed giving existence and uniqueness. Viscosity solutions for the coupled IPDE :

$$\begin{aligned}
 (\rho + \lambda)\hat{v} - \frac{\partial \hat{v}}{\partial t} - \tilde{U}\left(\frac{\partial \hat{v}}{\partial x}\right) - \lambda \int v(x + az)p(t, dz) &= 0, \quad t \geq 0, x > l(a), \\
 v(x) = \mathcal{H}\hat{v}(x) &= \sup_{a \in \left[-\frac{x}{z}, \frac{x}{z}\right]} \hat{v}(0, x, a), \quad x \in \mathbb{R}_+,
 \end{aligned}$$

### Definition

The pair of functions  $(v, \hat{v}) \in C_+(\mathbb{R}_+) \times C_+(\mathcal{D})$  is a viscosity *supersolution* to the above IPDE if :

- (i)  $v \geq \mathcal{H}\hat{v}$ ;
- (ii) for all  $a \in A, (\bar{t}, \bar{x}) \in \mathbb{R}_+ \times (l(a), \infty)$ ,

$$(\rho + \lambda)\hat{v}(\bar{t}, \bar{x}, a) - \frac{\partial \varphi}{\partial t}(\bar{t}, \bar{x}) - \tilde{U}\left(\frac{\partial \varphi}{\partial x}(\bar{t}, d\bar{x})\right) - \lambda \int v(\bar{x} + az)p(\bar{t}, dz) \geq 0,$$

for any test function  $\varphi \in C^1(\mathbb{R}_+ \times (l(a), +\infty))$ , which is a local *minimum* of  $(\hat{v}(\cdot, \cdot, a) - \varphi)$ .

## Viscosity solutions

In the paper [PT 07(a)] the viscosity solution approach is developed giving existence and uniqueness. Viscosity solutions for the coupled IPDE :

$$\begin{aligned}
 (\rho + \lambda)\hat{v} - \frac{\partial \hat{v}}{\partial t} - \tilde{U}\left(\frac{\partial \hat{v}}{\partial x}\right) - \lambda \int v(x + az)p(t, dz) &= 0, \quad t \geq 0, x > l(a), \\
 v(x) = \mathcal{H}\hat{v}(x) &= \sup_{a \in \left[-\frac{x}{z}, \frac{x}{z}\right]} \hat{v}(0, x, a), \quad x \in \mathbb{R}_+,
 \end{aligned}$$

### Definition

The pair of functions  $(v, \hat{v}) \in C_+(\mathbb{R}_+) \times C_+(\mathcal{D})$  is a viscosity *subsolution* to the above IPDE if :

- (i)  $v \leq \mathcal{H}\hat{v}$ ;
- (ii) for all  $a \in A, (\bar{t}, \bar{x}) \in \mathbb{R}_+ \times (l(a), \infty)$ ,

$$(\rho + \lambda)\hat{v}(\bar{t}, \bar{x}, a) - \frac{\partial \varphi}{\partial t}(\bar{t}, \bar{x}) - \tilde{U}\left(\frac{\partial \varphi}{\partial x}(\bar{t}, \bar{x})\right) - \lambda \int v(\bar{x} + az)p(\bar{t}, dz) \leq 0,$$

for any test function  $\varphi \in C^1(\mathbb{R}_+ \times (a, \infty))$ , which is a local *maximum* of  $(\hat{v}(\cdot, \cdot, a) - \varphi)$ .

## Theorem

Under **Assumptions (1)-(2)**, the pair of value functions  $(v, \hat{v})$  is the **unique viscosity solution** to the IPDE :

$$(\rho + \lambda)\hat{v} - \frac{\partial \hat{v}}{\partial t} - \tilde{U}\left(\frac{\partial \hat{v}}{\partial x}\right) - \lambda \int v(x + az)p(t, dz) = 0, \quad t \geq 0, x > l(a),$$
$$v(x) = \sup_{a \in \left[-\frac{x}{Z}, \frac{x}{Z}\right]} \hat{v}(0, x, a), \quad x \in \mathbb{R}_+,$$

satisfying the boundary conditions (G1)-(B1). BC

## Remark

*In particular for every  $a \in A$ , the value function  $\hat{v}(\cdot, \cdot, a)$  is the unique viscosity solution of the first HJ equation with the BC (G1)-(B1).*

← Return

## Theorem

Under *Assumptions (1)-(2)*, the pair of value functions  $(v, \hat{v})$  is the **unique viscosity solution** to the IPDE :

$$(\rho + \lambda)\hat{v} - \frac{\partial \hat{v}}{\partial t} - \tilde{U}\left(\frac{\partial \hat{v}}{\partial x}\right) - \lambda \int v(x + az)p(t, dz) = 0, \quad t \geq 0, x > l(a),$$
$$v(x) = \sup_{a \in \left[-\frac{x}{Z}, \frac{x}{Z}\right]} \hat{v}(0, x, a), \quad x \in \mathbb{R}_+,$$

satisfying the boundary conditions (G1)-(B1). BC

## Remark

In particular for every  $a \in A$ , the value function  $\hat{v}(\cdot, \cdot, a)$  is the unique viscosity solution of the first HJ equation with the BC (G1)-(B1).

[Return](#)

## ► Idea of the proof

First we prove that  $\hat{v}(\cdot, a) \in C^1(I(a), +\infty)$ . The other regularity properties follow by a kind of bootstrap argument based on the coupling.

Arguing by contradiction, we use:

- ★ the *viscosity characterization* of the value functions to the original control problem by means of viscosity solution to the coupled IPDE (see [PT 07(a)]).
- ★ the concavity of the function  $\hat{v}(\cdot, a)$  together with the strict convexity of the convex conjugate  $\tilde{U}$  with similar arguments to those given in:

[BCD 97] Bardi M. and Capuzzo-Dolcetta I., OPTIMAL CONTROL AND VISCOSITY SOLUTIONS OF HAMILTON-JACOBI-BELLMAN EQUATIONS. Birkhäuser Boston Inc., Boston, MA.



## ► Idea of the proof

First we prove that  $\hat{v}(\cdot, a) \in C^1(I(a), +\infty)$ . The other regularity properties follow by a kind of bootstrap argument based on the coupling.

Arguing by contradiction, we use:

★ the *viscosity characterization* of the value functions to the original control problem by means of viscosity solution to the coupled IPDE (see [PT 07(a)]). viscosity solutions

★ the concavity of the function  $\hat{v}(\cdot, a)$  together with the strict convexity of the convex conjugate  $\tilde{U}$  with similar arguments to those given in:

[BCD 97] Bardi M. and Capuzzo-Dolcetta I., *OPTIMAL CONTROL AND VISCOSITY SOLUTIONS OF HAMILTON-JACOBI-BELLMAN EQUATIONS*. Birkhäuser Boston Inc., Boston, MA.

sketch of the proof

## ► Idea of the proof

First we prove that  $\hat{v}(\cdot, a) \in C^1(I(a), +\infty)$ . The other regularity properties follow by a kind of bootstrap argument based on the coupling.

Arguing by contradiction, we use:

- ★ the *viscosity characterization* of the value functions to the original control problem by means of viscosity solution to the coupled IPDE (see [PT 07(a)]). viscosity solutions
- ★ the concavity of the function  $\hat{v}(\cdot, a)$  together with the strict convexity of the convex conjugate  $\tilde{U}$  with similar arguments to those given in:

[BCD 97] Bardi M. and Capuzzo-Dolcetta I., **OPTIMAL CONTROL AND VISCOSITY SOLUTIONS OF HAMILTON-JACOBI-BELLMAN EQUATIONS**. Birkhäuser Boston Inc., Boston, MA.

sketch of the proof

We show that  $\hat{v}$  is differentiable on  $(I(a), +\infty)$ , for any fixed  $a \in A$ .

- $D_x^+ \hat{v}(x, a) \neq \emptyset$  since  $\hat{v}$  is concave and it is enough to prove that it is always a single point, for any  $x \in (I(a), +\infty)$ . semidifferentials
- Assume by contradiction that  $D_x^+ \hat{v}(x, a) = [p_1, p_2]$ ,  $p_1 \neq p_2$ . Then there exist sequences  $x_n, y_m \in \mathbb{R}_+$  where  $\hat{v}$  is differentiable and s.t.

$$x = \lim_{n \rightarrow +\infty} x_n = \lim_{m \rightarrow +\infty} y_m, \quad p_1 = \lim_{n \rightarrow +\infty} D_x \hat{v}(x_n, a), \quad p_2 = \lim_{m \rightarrow +\infty} D_x \hat{v}(y_m, a).$$

- Since the pair  $(v, \hat{v})$  is a viscosity solution to (3)-(4), by continuity we have

$$(\rho + \lambda) \hat{v}(x, a) - \tilde{U}(p_1) - \lambda \int v(x + az) p(dz) = 0$$

$$(\rho + \lambda) \hat{v}(x, a) - \tilde{U}(p_2) - \lambda \int v(x + az) p(dz) = 0.$$

We show that  $\hat{v}$  is differentiable on  $(I(a), +\infty)$ , for any fixed  $a \in A$ .

- $D_x^+ \hat{v}(x, a) \neq \emptyset$  since  $\hat{v}$  is concave and it is enough to prove that it is always a single point, for any  $x \in (I(a), +\infty)$ . semidifferentials
- Assume by contradiction that  $D_x^+ \hat{v}(x, a) = [p_1, p_2]$ ,  $p_1 \neq p_2$ . Then there exist sequences  $x_n, y_m \in \mathbb{R}_+$  where  $\hat{v}$  is differentiable and s.t.

$$x = \lim_{n \rightarrow +\infty} x_n = \lim_{m \rightarrow +\infty} y_m, \quad p_1 = \lim_{n \rightarrow +\infty} D_x \hat{v}(x_n, a), \quad p_2 = \lim_{m \rightarrow +\infty} D_x \hat{v}(y_m, a).$$

- Since the pair  $(v, \hat{v})$  is a viscosity solution to (3)-(4), by continuity we have

$$(\rho + \lambda) \hat{v}(x, a) - \tilde{U}(p_1) - \lambda \int v(x + az) p(dz) = 0$$

$$(\rho + \lambda) \hat{v}(x, a) - \tilde{U}(p_2) - \lambda \int v(x + az) p(dz) = 0.$$

We show that  $\hat{v}$  is differentiable on  $(I(a), +\infty)$ , for any fixed  $a \in A$ .

- $D_x^+ \hat{v}(x, a) \neq \emptyset$  since  $\hat{v}$  is concave and it is enough to prove that it is always a single point, for any  $x \in (I(a), +\infty)$ . semidifferentials
- Assume by contradiction that  $D_x^+ \hat{v}(x, a) = [p_1, p_2]$ ,  $p_1 \neq p_2$ . Then there exist sequences  $x_n, y_m \in \mathbb{R}_+$  where  $\hat{v}$  is differentiable and s.t.

$$x = \lim_{n \rightarrow +\infty} x_n = \lim_{m \rightarrow +\infty} y_m, \quad p_1 = \lim_{n \rightarrow +\infty} D_x \hat{v}(x_n, a), \quad p_2 = \lim_{m \rightarrow +\infty} D_x \hat{v}(y_m, a).$$

- Since the pair  $(v, \hat{v})$  is a viscosity solution to (3)-(4), by continuity we have

$$(\rho + \lambda)\hat{v}(x, a) - \tilde{U}(p_1) - \lambda \int v(x + az)p(dz) = 0$$

$$(\rho + \lambda)\hat{v}(x, a) - \tilde{U}(p_2) - \lambda \int v(x + az)p(dz) = 0.$$

- Moreover, for  $\eta \in (0, 1)$  we have  $\bar{p} = \eta p_1 + (1 - \eta)p_2 \in (p_1, p_2)$  included in  $D_x^+ \hat{v}(x, a)$ , then by the viscosity subsolution property of  $\hat{v}$ :

$$(\rho + \lambda)\hat{v}(x, a) - \tilde{U}(\bar{p}) - \lambda \int v(x + az)p(dz) \leq 0.$$

- By strict convexity of  $\tilde{U}$ , we get

$$\tilde{U}(\bar{p}) = \tilde{U}(\eta p_1 + (1 - \eta)p_2) < \eta \tilde{U}(p_1) + (1 - \eta)\tilde{U}(p_2),$$

that is a contradiction.

We can deduce the continuity of  $\frac{\partial \hat{v}}{\partial x}$  from (3). It also follows from [CS 2004] Cannarsa P. and Sinestrari C., SEMICONCAVE FUNCTIONS, HAMILTON-JACOBI EQUATIONS AND OPTIMAL CONTROL.

← Return

- Moreover, for  $\eta \in (0, 1)$  we have  $\bar{p} = \eta p_1 + (1 - \eta)p_2 \in (p_1, p_2)$  included in  $D_x^+ \hat{v}(x, a)$ , then by the viscosity subsolution property of  $\hat{v}$ :

$$(\rho + \lambda)\hat{v}(x, a) - \tilde{U}(\bar{p}) - \lambda \int v(x + az)p(dz) \leq 0.$$

- By strict convexity of  $\tilde{U}$ , we get

$$\tilde{U}(\bar{p}) = \tilde{U}(\eta p_1 + (1 - \eta)p_2) < \eta \tilde{U}(p_1) + (1 - \eta)\tilde{U}(p_2),$$

that is a contradiction.

We can deduce the continuity of  $\frac{\partial \hat{v}}{\partial x}$  from (3). It also follows from [CS 2004] Cannarsa P. and Sinestrari C., SEMICONCAVE FUNCTIONS, HAMILTON-JACOBI EQUATIONS AND OPTIMAL CONTROL.

← Return

- Moreover, for  $\eta \in (0, 1)$  we have  $\bar{p} = \eta p_1 + (1 - \eta)p_2 \in (p_1, p_2)$  included in  $D_x^+ \hat{v}(x, a)$ , then by the viscosity subsolution property of  $\hat{v}$ :

$$(\rho + \lambda)\hat{v}(x, a) - \tilde{U}(\bar{p}) - \lambda \int v(x + az)p(dz) \leq 0.$$

- By strict convexity of  $\tilde{U}$ , we get

$$\tilde{U}(\bar{p}) = \tilde{U}(\eta p_1 + (1 - \eta)p_2) < \eta \tilde{U}(p_1) + (1 - \eta)\tilde{U}(p_2),$$

that is a contradiction.

We can deduce the continuity of  $\frac{\partial \hat{v}}{\partial x}$  from (3). It also follows from [\[CS 2004\]](#) Cannarsa P. and Sinestrari C., [SEMICONCAVE FUNCTIONS, HAMILTON-JACOBI EQUATIONS AND OPTIMAL CONTROL](#).



## ► Idea of the proof

First we prove that  $\hat{v}(\cdot, a) \in C^1(I(a), +\infty)$ . The other regularity properties follow by a kind of bootstrap argument based on the coupling.

Arguing by contradiction, we use:

- ★ the *viscosity characterization* of the value functions to the original control problem by means of viscosity solution to the coupled IPDE (see [PT 07(a)]).
- ★ the concavity of the function  $\hat{v}(\cdot, a)$  together with the strict convexity of the convex conjugate  $\tilde{U}$  with similar arguments to those given in:

[BCD 97] Bardi M. and Capuzzo-Dolcetta I., OPTIMAL CONTROL AND VISCOSITY SOLUTIONS OF HAMILTON-JACOBI-BELLMAN EQUATIONS. Birkhäuser Boston Inc., Boston, MA.

## ► Idea of the proof

First we prove that  $\hat{v}(\cdot, a) \in C^1(I(a), +\infty)$ . The other regularity properties follow by a kind of bootstrap argument based on the coupling.

Arguing by contradiction, we use:

★ the *viscosity characterization* of the value functions to the original control problem by means of viscosity solution to the coupled IPDE (see [PT 07(a)]). viscosity solutions

★ the concavity of the function  $\hat{v}(\cdot, a)$  together with the strict convexity of the convex conjugate  $\tilde{U}$  with similar arguments to those given in:

[BCD 97] Bardi M. and Capuzzo-Dolcetta I., OPTIMAL CONTROL AND VISCOSITY SOLUTIONS OF HAMILTON-JACOBI-BELLMAN EQUATIONS. Birkhäuser Boston Inc., Boston, MA.

sketch of the proof

## ► Idea of the proof

First we prove that  $\hat{v}(\cdot, a) \in C^1(I(a), +\infty)$ . The other regularity properties follow by a kind of bootstrap argument based on the coupling.

Arguing by contradiction, we use:

- ★ the *viscosity characterization* of the value functions to the original control problem by means of viscosity solution to the coupled IPDE (see [PT 07(a)]). viscosity solutions
- ★ the concavity of the function  $\hat{v}(\cdot, a)$  together with the strict convexity of the convex conjugate  $\tilde{U}$  with similar arguments to those given in:

[BCD 97] Bardi M. and Capuzzo-Dolcetta I., **OPTIMAL CONTROL AND VISCOSITY SOLUTIONS OF HAMILTON-JACOBI-BELLMAN EQUATIONS**. Birkhäuser Boston Inc., Boston, MA.

sketch of the proof

# Semidifferentials

## Definition

Let  $u$  be a continuous function on an open set  $D \subset \Omega$ . For any  $y \in D$ , the sets

$$D^- u(y) = \left\{ p \in \Omega : \liminf_{z \in D, z \rightarrow y} \frac{u(z) - u(y) - \langle p, z - y \rangle}{|z - y|} \geq 0 \right\},$$

$$D^+ u(y) = \left\{ p \in \Omega : \limsup_{z \in D, z \rightarrow y} \frac{u(z) - u(y) - \langle p, z - y \rangle}{|z - y|} \leq 0 \right\}$$

are called respectively, the (Fréchet) *subdifferential* and *superdifferential* of  $u$  at  $y$ .

◀ Return

We show that  $\hat{v}$  is differentiable on  $(I(a), +\infty)$ , for any fixed  $a \in A$ .

- $D_x^+ \hat{v}(x, a) \neq \emptyset$  since  $\hat{v}$  is concave and it is enough to prove that it is always a single point, for any  $x \in (I(a), +\infty)$ . semidifferentials
- Assume by contradiction that  $D_x^+ \hat{v}(x, a) = [p_1, p_2]$ ,  $p_1 \neq p_2$ . Then there exist sequences  $x_n, y_m \in \mathbb{R}_+$  where  $\hat{v}$  is differentiable and s.t.

$$x = \lim_{n \rightarrow +\infty} x_n = \lim_{m \rightarrow +\infty} y_m, \quad p_1 = \lim_{n \rightarrow +\infty} D_x \hat{v}(x_n, a), \quad p_2 = \lim_{m \rightarrow +\infty} D_x \hat{v}(y_m, a).$$

- Since the pair  $(v, \hat{v})$  is a viscosity solution to (3)-(4), by continuity we have

$$(\rho + \lambda) \hat{v}(x, a) - \tilde{U}(p_1) - \lambda \int v(x + az) p(dz) = 0$$

$$(\rho + \lambda) \hat{v}(x, a) - \tilde{U}(p_2) - \lambda \int v(x + az) p(dz) = 0.$$

We show that  $\hat{v}$  is differentiable on  $(I(a), +\infty)$ , for any fixed  $a \in A$ .

- $D_x^+ \hat{v}(x, a) \neq \emptyset$  since  $\hat{v}$  is concave and it is enough to prove that it is always a single point, for any  $x \in (I(a), +\infty)$ . semidifferentials
- Assume by contradiction that  $D_x^+ \hat{v}(x, a) = [p_1, p_2]$ ,  $p_1 \neq p_2$ . Then there exist sequences  $x_n, y_m \in \mathbb{R}_+$  where  $\hat{v}$  is differentiable and s.t.

$$x = \lim_{n \rightarrow +\infty} x_n = \lim_{m \rightarrow +\infty} y_m, \quad p_1 = \lim_{n \rightarrow +\infty} D_x \hat{v}(x_n, a), \quad p_2 = \lim_{m \rightarrow +\infty} D_x \hat{v}(y_m, a).$$

- Since the pair  $(v, \hat{v})$  is a viscosity solution to (3)-(4), by continuity we have

$$(\rho + \lambda) \hat{v}(x, a) - \tilde{U}(p_1) - \lambda \int v(x + az) p(dz) = 0$$

$$(\rho + \lambda) \hat{v}(x, a) - \tilde{U}(p_2) - \lambda \int v(x + az) p(dz) = 0.$$

We show that  $\hat{v}$  is differentiable on  $(l(a), +\infty)$ , for any fixed  $a \in A$ .

- $D_x^+ \hat{v}(x, a) \neq \emptyset$  since  $\hat{v}$  is concave and it is enough to prove that it is always a single point, for any  $x \in (l(a), +\infty)$ . semidifferentials
- Assume by contradiction that  $D_x^+ \hat{v}(x, a) = [p_1, p_2]$ ,  $p_1 \neq p_2$ . Then there exist sequences  $x_n, y_m \in \mathbb{R}_+$  where  $\hat{v}$  is differentiable and s.t.

$$x = \lim_{n \rightarrow +\infty} x_n = \lim_{m \rightarrow +\infty} y_m, \quad p_1 = \lim_{n \rightarrow +\infty} D_x \hat{v}(x_n, a), \quad p_2 = \lim_{m \rightarrow +\infty} D_x \hat{v}(y_m, a).$$

- Since the pair  $(v, \hat{v})$  is a viscosity solution to (3)-(4), by continuity we have

$$(\rho + \lambda)\hat{v}(x, a) - \tilde{U}(p_1) - \lambda \int v(x + az)p(dz) = 0$$

$$(\rho + \lambda)\hat{v}(x, a) - \tilde{U}(p_2) - \lambda \int v(x + az)p(dz) = 0.$$

# Semiconcavity

## Definition

We say that a function  $u : S \rightarrow \mathbb{R}$  is *semiconcave* if there exists a nondecreasing upper semicontinuous function  $\omega : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  such that  $\lim_{\rho \rightarrow 0^+} \omega(\rho) = 0$  and

$$\eta u(x_1) + (1 - \eta)u(x_2) - u(\eta x_1 + (1 - \eta)x_2) \leq \eta(1 - \eta)|x_1 - x_2|\omega(|x_1 - x_2|),$$

for any pair  $x_1, x_2$  such that the segment  $[x_1, x_2]$  is contained in  $S$  and for  $\eta \in [0, 1]$ . In particular we call *locally semiconcave* a function which is semiconcave on every compact subset of its domain of definition.

Clearly, a concave function is also semiconcave. An important example of semiconcave functions is given by the smooth ones.

## Proposition

*Let  $u \in C^1(A)$ , with  $A$  open. Then both  $u$  and  $-u$  are locally semiconcave in  $A$  with modulus equal to the modulus of continuity of  $Du$ .*



## 3.b Nonstationary case

To use the same argument as in the stationary case we need the **joint semiconcavity** in  $(t, x)$  of the function  $\hat{v}(\cdot, \cdot, a)$ . semiconcavity

Hence we need to introduce an additional assumption on the measure  $p(t, dz)$  in order to guarantee it.

**Assumption (3):** for every  $a \in A - \{0\}$ , the map

$$(t, x) \longrightarrow \lambda \int w(x + az)p(t, dz)$$

is (locally) semiconcave for  $(t, x) \in (0, +\infty) \times (l(a), +\infty)$  for all measurable continuous functions  $w$  with linear growth.

## 3.b Nonstationary case

To use the same argument as in the stationary case we need the **joint semiconcavity** in  $(t, x)$  of the function  $\hat{v}(\cdot, \cdot, a)$ . semiconcavity

Hence we need to introduce an additional assumption on the measure  $p(t, dz)$  in order to guarantee it.

**Assumption (3)**: for every  $a \in A - \{0\}$ , the map

$$(t, x) \longrightarrow \lambda \int w(x + az)p(t, dz)$$

is (locally) semiconcave for  $(t, x) \in (0, +\infty) \times (l(a), +\infty)$  for all measurable continuous functions  $w$  with linear growth.