

 Open access • Journal Article • DOI:10.1007/S11856-015-1264-7

Optimal hardy-littlewood type inequalities for polynomials and multilinear operators

— [Source link](#) 

N. Albuquerque, [Frédéric Bayart](#), [Daniel Pellegrino](#), [Juan B. Seoane-Sepúlveda](#) ...+1 more authors

Institutions: [Federal University of Paraíba](#), [Blaise Pascal University](#), [Complutense University of Madrid](#), [Spanish National Research Council](#)

Published on: 01 Feb 2016 - [Israel Journal of Mathematics](#) (The Hebrew University Magnes Press)

Topics: [Multilinear map](#) and [Hardy–Littlewood inequality](#)

Related papers:

- [Sharp generalizations of the multilinear Bohnenblust–Hille inequality](#)
- [On the Absolute Convergence of Dirichlet Series](#)
- [On bounded multilinear forms on a class of \$l_p\$ spaces](#)
- [BILINEAR FORMS BOUNDED IN SPACE \[p, q\]](#)
- [Summation of Coefficients of Polynomials on \$\ell_p\$ Spaces](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/optimal-hardy-littlewood-type-inequalities-for-polynomials-2zk0uu0cs4>

OPTIMAL HARDY–LITTLEWOOD TYPE INEQUALITIES FOR POLYNOMIALS AND MULTILINEAR OPERATORS

N. ALBUQUERQUE, F. BAYART, D. PELLEGRINO, AND J. B. SEOANE-SEPÚLVEDA

ABSTRACT. In this paper we obtain quite general and definitive forms for Hardy–Littlewood type inequalities. Moreover, when restricted to the original particular cases, our approach provides much simpler and straightforward proofs and we are able to show that in most cases the exponents involved are optimal. The technique we used is a combination of probabilistic tools and of an interpolative approach; this former technique is also employed in this paper to improve the constants for vector-valued Bohnenblust–Hille type inequalities.

1. INTRODUCTION

In 1930 Littlewood [20] has shown the following result on bilinear forms on $c_0 \times c_0$, now called Littlewood’s 4/3 inequality: for any bounded bilinear form $A : c_0 \times c_0 \rightarrow \mathbb{C}$,

$$\left(\sum_{i,j=1}^{+\infty} |A(e_i, e_j)|^{\frac{4}{3}} \right)^{\frac{3}{4}} \leq \sqrt{2} \|A\|$$

and, moreover, the exponent 4/3 is optimal. From now on, $m \geq 1$ is a positive integer, $\mathbf{p} := (p_1, \dots, p_m) \in [1, +\infty]^m$ and

$$\left| \frac{1}{\mathbf{p}} \right| := \frac{1}{p_1} + \dots + \frac{1}{p_m}.$$

For $1 \leq p < +\infty$, let us set $X_p := \ell_p$ and let us define $X_\infty = c_0$. As soon as Littlewood’s 4/3 inequality appeared, it was rapidly extended to more general frameworks. For instance:

- (Bohnenblust and Hille, [7, Theorem I], 1931 (see also [11])) There exists a constant $C = C(m) \geq 1$ such that

$$(1.1) \quad \left(\sum_{i_1, \dots, i_m=1}^{+\infty} |A(e_{i_1}, \dots, e_{i_m})|^{\frac{2m}{m+1}} \right)^{\frac{m+1}{2m}} \leq C \|A\|$$

for all continuous m -linear forms $A : c_0 \times \dots \times c_0 \rightarrow \mathbb{C}$ and the exponent $\frac{2m}{m+1}$ is optimal.

- (Hardy and Littlewood, [16], 1934 (see also [17, page 224])/Praciano-Pereira, [25, Theorems A and B], 1981) Let $\mathbf{p} \in [1, +\infty]^m$ with

$$\left| \frac{1}{\mathbf{p}} \right| \leq \frac{1}{2},$$

then there exists a constant $C > 0$ such that, for every continuous m -linear form $A : X_{p_1} \times \dots \times X_{p_m} \rightarrow \mathbb{C}$,

$$(1.2) \quad \left(\sum_{i_1, \dots, i_m=1}^{+\infty} |A(e_{i_1}, \dots, e_{i_m})|^{\frac{2m}{m+1-2|\frac{1}{\mathbf{p}}|}} \right)^{\frac{m+1-2|\frac{1}{\mathbf{p}}|}{2m}} \leq C \|A\|.$$

2010 *Mathematics Subject Classification.* 46G25, 47H60.

Key words and phrases. Absolutely summing operators, multilinear operators, Bohnenblust–Hille inequality.

D. Pellegrino and J.B. Seoane-Sepúlveda was supported by CNPq Grant 401735/2013-3 (PVE - Linha 2). N. Albuquerque was supported by CAPES.

- (Defant and Sevilla-Peris, [10, Theorem 1], 2009) If $1 \leq s \leq q \leq 2$, there exists a constant $C > 0$ such that, for every continuous m -linear mapping $A : c_0 \times \cdots \times c_0 \rightarrow \ell_s$, then

$$\left(\sum_{i_1, \dots, i_m=1}^{+\infty} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^{\frac{2m}{m+2(\frac{1}{s}-\frac{1}{q})}} \right)^{\frac{m+2(\frac{1}{s}-\frac{1}{q})}{2m}} \leq C \|A\|.$$

Very recently the previous results were generalized by the authors and by Dimant and Sevilla-Peris:

- ([1, Corollary 1.3], 2013) Let $1 \leq s \leq q \leq 2$ and $\mathbf{p} \in [1, +\infty]^m$ such that

$$(1.3) \quad \frac{1}{s} - \frac{1}{q} - \left| \frac{1}{\mathbf{p}} \right| \geq 0.$$

Then there exists a constant $C > 0$ such that, for every continuous m -linear mapping $A : X_{p_1} \times \cdots \times X_{p_m} \rightarrow X_s$, we have

$$\left(\sum_{i_1, \dots, i_m=1}^{+\infty} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^{\frac{2m}{m+2(\frac{1}{s}-\frac{1}{q}-\left|\frac{1}{\mathbf{p}}\right|)}} \right)^{\frac{m+2(\frac{1}{s}-\frac{1}{q}-\left|\frac{1}{\mathbf{p}}\right|)}{2m}} \leq C \|A\|$$

and the exponent is optimal.

- (Dimant and Sevilla-Peris, [13, Proposition 4.4], 2013) Let $\mathbf{p} \in [1, +\infty]^m$ and $s, q \in [1, +\infty]$ be such that $s \leq q$. Then there exists a constant $C > 0$ such that, for every continuous m -linear mapping $A : X_{p_1} \times \cdots \times X_{p_m} \rightarrow X_s$, we have

$$\left(\sum_{i_1, \dots, i_m=1}^{+\infty} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^{\rho} \right)^{\frac{1}{\rho}} \leq C \|A\|,$$

where ρ is given by

(i) If $s \leq q \leq 2$, and

(a) if $0 \leq \left| \frac{1}{\mathbf{p}} \right| < \frac{1}{s} - \frac{1}{q}$, then $\frac{1}{\rho} = \frac{1}{2} + \frac{1}{m} \left(\frac{1}{s} - \frac{1}{q} - \left| \frac{1}{\mathbf{p}} \right| \right)$.

(b) if $\frac{1}{s} - \frac{1}{q} \leq \left| \frac{1}{\mathbf{p}} \right| < \frac{1}{2} + \frac{1}{s} - \frac{1}{q}$, then $\frac{1}{\rho} = \frac{1}{2} + \frac{1}{s} - \frac{1}{q} - \left| \frac{1}{\mathbf{p}} \right|$.

(ii) If $s \leq 2 \leq q$, and

(a) if $0 \leq \left| \frac{1}{\mathbf{p}} \right| < \frac{1}{s} - \frac{1}{2}$, then $\frac{1}{\rho} = \frac{1}{2} + \frac{1}{m} \left(\frac{1}{s} - \frac{1}{2} - \left| \frac{1}{\mathbf{p}} \right| \right)$.

(b) if $\frac{1}{s} - \frac{1}{2} \leq \left| \frac{1}{\mathbf{p}} \right| < \frac{1}{s}$, then $\frac{1}{\rho} = \frac{1}{s} - \left| \frac{1}{\mathbf{p}} \right|$.

(iii) If $2 \leq s \leq q$ and $0 \leq \left| \frac{1}{\mathbf{p}} \right| < \frac{1}{s}$, then $\frac{1}{\rho} = \frac{1}{s} - \left| \frac{1}{\mathbf{p}} \right|$.

Moreover, the exponents in the cases (ia),(iib) and (iii) are optimal. Also, the exponent in (ib) is optimal for $\frac{1}{s} - \frac{1}{q} \leq \left| \frac{1}{\mathbf{p}} \right| < \frac{1}{2}$.

Our main intention, in this paper, is to improve the previous theorems in three directions.

- (1) We study in depth the remaining cases of the Dimant and Sevilla-Peris result. Surprisingly, we show that in case (iia), the exponent given above is optimal whereas it is not optimal in case (ib) when $\left| \frac{1}{\mathbf{p}} \right| > \frac{1}{2}$. We give a better exponent in that case and show a necessary condition on it. These two bounds coincide when $s = 1$. We can summarize this into the two following statements.

Theorem 1.1. *Let $\mathbf{p} \in [1, +\infty]^m$ and let $\rho > 0$. Assume moreover that either $q \geq 2$ or $q < 2$ and $\left| \frac{1}{\mathbf{p}} \right| < \frac{1}{2}$. Let*

$$\frac{1}{\lambda} := \frac{1}{2} + \frac{1}{s} - \frac{1}{\min\{q, 2\}} - \left| \frac{1}{\mathbf{p}} \right| > 0.$$

Then there exists $C > 0$ such that, for every continuous m -linear operator $A : X_{p_1} \times \cdots \times X_{p_m} \rightarrow X_s$, we have

$$\left(\sum_{i_1, \dots, i_m=1}^{+\infty} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^\rho \right)^{\frac{1}{\rho}} \leq C \|A\|$$

if and only if

$$\frac{m}{\rho} \leq \frac{1}{\lambda} + \frac{m-1}{\max\{\lambda, s, 2\}}.$$

The following table summarizes the optimal value of $\frac{1}{\rho}$ following the respective values of s, q, p_1, \dots, p_m :

$1 \leq s \leq q \leq 2, \lambda < 2$	$\frac{1}{2} + \frac{1}{ms} - \frac{1}{mq} - \frac{1}{m} \times \left \frac{1}{\mathbf{p}} \right $
$1 \leq s \leq q \leq 2, \lambda \geq 2, \left \frac{1}{\mathbf{p}} \right < \frac{1}{2}$	$\frac{1}{2} + \frac{1}{s} - \frac{1}{q} - \left \frac{1}{\mathbf{p}} \right $
$1 \leq s \leq 2 \leq q, \lambda < 2$	$\frac{m-1}{2m} + \frac{1}{ms} - \frac{1}{m} \times \left \frac{1}{\mathbf{p}} \right $
$1 \leq s \leq 2 \leq q, \lambda \geq 2$	$\frac{1}{s} - \left \frac{1}{\mathbf{p}} \right $
$2 \leq s \leq q$	$\frac{1}{s} - \left \frac{1}{\mathbf{p}} \right $

We note that (1.1) and (1.2) are recovered by Theorem 1.1 just by choosing $s = 1$ and $q = 2$.

When $q < 2$ and $\left| \frac{1}{\mathbf{p}} \right| > \frac{1}{2}$ (observe that this automatically implies $\lambda \geq 2$), the situation is more difficult and we get the following statement.

Theorem 1.2. Let $\mathbf{p} \in [1, +\infty]^m$, $\left| \frac{1}{\mathbf{p}} \right| > \frac{1}{2}$, $1 \leq s \leq q \leq 2$ and let $\rho > 0$. Let us consider the following property.

There exists $C > 0$ such that, for every continuous m -linear operator $A : X_{p_1} \times \cdots \times X_{p_m} \rightarrow X_s$, we have

$$\left(\sum_{i_1, \dots, i_m=1}^{+\infty} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^\rho \right)^{\frac{1}{\rho}} \leq C \|A\|.$$

(A) The property is satisfied as soon as

$$\frac{1}{\rho} \leq \frac{\left(\frac{1}{s} - \frac{1}{q} \right) \left(\frac{1}{s} - \left| \frac{1}{\mathbf{p}} \right| \right)}{\frac{1}{2} - \frac{1}{s}}.$$

(B) If the property is satisfied, then

$$\frac{1}{\rho} \leq 2 \left(1 - \left| \frac{1}{\mathbf{p}} \right| \right) \left(\frac{1}{s} - \frac{1}{q} \right).$$

In particular, if $s = 1$, then the property is satisfied if and only if

$$\frac{1}{\rho} \leq 2 \left(1 - \left| \frac{1}{\mathbf{p}} \right| \right) \left(1 - \frac{1}{q} \right).$$

(2) We give a simpler proof of the sufficient part of the Dimant and Sevilla-Peris theorem. It turns out that it is easier to prove a more general result.

Theorem 1.3. *Let $\mathbf{p} \in [1, +\infty]^m$ and $1 \leq s \leq q \leq \infty$ be such that*

$$\left| \frac{1}{\mathbf{p}} \right| < \frac{1}{2} + \frac{1}{s} - \frac{1}{\min\{q, 2\}}.$$

Let

$$\frac{1}{\lambda} := \frac{1}{2} + \frac{1}{s} - \frac{1}{\min\{q, 2\}} - \left| \frac{1}{\mathbf{p}} \right|.$$

If $\lambda > 0$ and $t_1, \dots, t_m \in [\lambda, \max\{\lambda, s, 2\}]$ are such that

$$(1.4) \quad \frac{1}{t_1} + \dots + \frac{1}{t_m} \leq \frac{1}{\lambda} + \frac{m-1}{\max\{\lambda, s, 2\}},$$

then there exists $C > 0$ satisfying, for every continuous m -linear map $A : X_{p_1} \times \dots \times X_{p_m} \rightarrow X_s$,

$$(1.5) \quad \left(\sum_{i_1=1}^{+\infty} \left(\dots \left(\sum_{i_m=1}^{+\infty} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^{t_m} \right) \dots \right)^{\frac{t_{m-1}}{t_m}} \dots \right)^{\frac{t_1}{t_2}} \frac{1}{t_1} \leq C \|A\|.$$

Moreover, the exponents are optimal except eventually if $q \leq 2$ and $\left| \frac{1}{\mathbf{p}} \right| > \frac{1}{2}$.

Remark 1.4. *The optimality in the above theorem shall be understood in a strong sense: when $\lambda < 2$, we prove that if $t_1, \dots, t_m \in [1, +\infty)$ are so that (1.5) holds then (1.4) is valid. When $\lambda \geq 2$, note that $\lambda = \max\{\lambda, s, 2\}$ and we prove that if $t = t_1 = \dots = t_m$ are in $[1, +\infty)$ and (1.5) is valid, then we have (1.4) and, as a direct consequence, $t \geq \lambda$.*

- (3) We prove similar results for m -linear mappings with arbitrary codomains which assume their cotype. For a Banach space X , let $q_X = \inf\{q \geq 2; X \text{ has cotype } q\}$.

The proof that (B) implies (A) in the theorem below appears in [13, Proposition 4.3].

Theorem 1.5. *Let $\mathbf{p} \in [2, +\infty]^m$, let X be an infinite dimensional Banach space with cotype q_X , $\left| \frac{1}{\mathbf{p}} \right| < \frac{1}{q_X}$, and let $\rho > 0$. The following assertions are equivalent:*

(A) *Every bounded m -linear operator $A : X_{p_1} \times \dots \times X_{p_m} \rightarrow X$ is such that*

$$\sum_{i_1, \dots, i_m=1}^{+\infty} \|A(e_{i_1}, \dots, e_{i_m})\|^\rho < +\infty.$$

(B) $\frac{1}{\rho} \leq \frac{1}{q_X} - \left| \frac{1}{\mathbf{p}} \right|$.

Finally, in the last section of the paper we obtain better estimates for the constants of vector-valued Bohnenblust–Hille inequalities.

We conclude this introduction by noting that our theorems can be naturally stated in the context of homogeneous polynomials. Given an m -homogeneous polynomial $P : X \rightarrow Y$, we denote its coefficients $(c_\alpha(P))$. In [10, Lemma 5], it is shown that an inequality

$$\left(\sum_{\alpha} \|c_\alpha(P)\|^\rho \right)^{\frac{1}{\rho}} \leq C \|P\|$$

holds for every m -homogeneous polynomial $P : X \rightarrow Y$ if and only if a similar inequality

$$\left(\sum_{i_1, \dots, i_m} \|T(e_{i_1}, \dots, e_{i_m})\|^\rho \right)^{\frac{1}{\rho}} \leq C' \|T\|$$

is satisfied for every m -linear mapping $A : X \times \dots \times X \rightarrow Y$, where X is a Banach sequence space.

NOTATIONS. For two positive integers n, k , we set

$$\mathcal{M}(k, n) := \{\mathbf{i} = (i_1, \dots, i_k); i_1, \dots, i_k \in \{1, \dots, n\}\}.$$

For $q \in [1, +\infty]$, q^* will denote its conjugate exponent.

2. PROOF OF THEOREM 1.3 (SUFFICIENCY)

Let $1 \leq q \leq +\infty$. We recall that a Banach space X has *cotype* q if there is a constant $\kappa > 0$ such that, no matter how we select finitely many vectors $x_1, \dots, x_n \in X$,

$$\left(\sum_{k=1}^n \|x_k\|^q \right)^{\frac{1}{q}} \leq \kappa \left(\int_I \left\| \sum_{k=1}^n r_k(t)x_k \right\|^2 dt \right)^{\frac{1}{2}}$$

where $I = [0, 1]$ and r_k denotes the k -th Rademacher function. To cover the case $q = +\infty$, the left hand side should be replaced by $\max_{1 \leq k \leq n} \|x_k\|$. The smallest of all these constants is denoted by $C_q(X)$ and named the *cotype* q constant of X .

An operator between Banach spaces $v : X \rightarrow Y$ is (r, s) -summing (with $s \leq r \leq +\infty$) if there exists $C > 0$ such that, for all $n \geq 1$ and for all vectors $x_1, \dots, x_n \in X$,

$$\left(\sum_{k=1}^n \|vx_k\|^r \right)^{\frac{1}{r}} \leq C \sup_{x^* \in B_{X^*}} \left(\sum_{k=1}^n |x^*(x_k)|^s \right)^{\frac{1}{s}}.$$

The smallest constant in this inequality is denoted by $\pi_{r,s}(v)$.

We need a cotype q version of [1, Proposition 4.1], whose proof can be found in [13, Proposition 3.1]:

Lemma 2.1. *Let X be a Banach space, let Y be a cotype q space, let $r \in [1, q]$ and let $\mathbf{p} \in [1, +\infty]^m$ with*

$$\left| \frac{1}{\mathbf{p}} \right| < \frac{1}{r} - \frac{1}{q}.$$

Define

$$\frac{1}{\lambda} := \frac{1}{r} - \left| \frac{1}{\mathbf{p}} \right|.$$

Then, for every continuous m -linear map $A : X_{p_1} \times \dots \times X_{p_m} \rightarrow X$ and every $(r, 1)$ -summing operator $v : X \rightarrow Y$, we have

$$(2.1) \quad \left(\sum_{i_k} \left(\sum_{\widehat{i_k}} \|vA(e_{i_1}, \dots, e_{i_m})\|_Y^q \right)^{\lambda/q} \right)^{1/\lambda} \leq \left(\sqrt{2}C_q(Y) \right)^{m-1} \pi_{r,1}(v) \|A\|$$

for all $k = 1, \dots, m$.

The symbol $\sum_{\widehat{i_k}}$ means that we are fixing the k -th index and that we are summing over all the remaining indices.

We shall deduce from this lemma the following theorem, which extends results of [1] and [13]:

Theorem 2.2. *Let $\mathbf{p} \in [1, +\infty]^m$, X be a Banach space, Y be a cotype q space and $1 \leq r \leq q$, with $\left| \frac{1}{\mathbf{p}} \right| < \frac{1}{r}$. Define*

$$\frac{1}{\lambda} := \frac{1}{r} - \left| \frac{1}{\mathbf{p}} \right|.$$

If $t_1, \dots, t_m \in [\lambda, \max\{\lambda, q\}]$ are such that

$$\frac{1}{t_1} + \dots + \frac{1}{t_m} \leq \frac{1}{\lambda} + \frac{m-1}{\max\{\lambda, q\}},$$

then, for every continuous m -linear map $A : X_{p_1} \times \dots \times X_{p_m} \rightarrow X$ and every $(r, 1)$ -summing operator $v : X \rightarrow Y$, we have

$$(2.2) \quad \left(\sum_{i_1=1}^{+\infty} \left(\dots \left(\sum_{i_m=1}^{+\infty} \|vA(e_{i_1}, \dots, e_{i_m})\|_Y^{t_m} \right)^{\frac{t_{m-1}}{t_m}} \dots \right)^{\frac{t_1}{t_2}} \right)^{\frac{1}{t_1}} \leq \left(\sqrt{2}C_{\max\{\lambda, q\}}(Y) \right)^{m-1} \pi_{r,1}(v) \|A\|.$$

Proof. If $\lambda < q$, from Lemma 2.1, we have (2.2) for

$$(t_1, \dots, t_m) = (\lambda, q, \dots, q).$$

Since $\lambda < q$, the mixed (ℓ_λ, ℓ_q) – norm inequality (see [1, Proposition 3.1]), we also have (2.2) for the exponents

$$(t_1, \dots, t_m) = (q, \dots, q, \lambda, q, \dots, q)$$

with λ in the k -th position, for all $k = 1, \dots, m$. Now, using a general version of Hölder's inequality (see [14, Theorem 2.1]), or interpolating

$$(\lambda, q, \dots, q), (q, \lambda, q, \dots, q), \dots, (q, \dots, q, \lambda)$$

in the sense of [1], we get (2.2) for all $(t_1, \dots, t_m) \in [\lambda, q]^m$ such that

$$\frac{1}{t_1} + \dots + \frac{1}{t_m} = \frac{1}{\lambda} + \frac{m-1}{q} = \frac{1}{\lambda} + \frac{m-1}{\max\{\lambda, q\}}.$$

If $\lambda \geq q$, for any $\varepsilon > 0$, let $q_\varepsilon = \lambda + \varepsilon$. So $\lambda < q_\varepsilon$ and this automatically implies that

$$\left| \frac{1}{\mathbf{p}} \right| < \frac{1}{r} - \frac{1}{q_\varepsilon}.$$

Since Y has cotype $q_\varepsilon > q$, we may apply Lemma 2.1 to get

$$\left(\sum_{i_1=1}^N \left(\sum_{i_2, \dots, i_m=1}^N \|vA(e_{i_1}, \dots, e_{i_m})\|^{\lambda+\varepsilon} \right)^{\frac{\lambda}{\lambda+\varepsilon}} \right)^{\frac{1}{\lambda}} \leq \left(\sqrt{2}C_{\lambda+\varepsilon}(Y) \right)^{m-1} \pi_{r,1}(v) \|A\|$$

for all positive integer N . Making $\varepsilon \rightarrow 0$, we get

$$\left(\sum_{i_1, \dots, i_m=1}^N \|vA(e_{i_1}, \dots, e_{i_m})\|^\lambda \right)^{\frac{1}{\lambda}} \leq \left(\sqrt{2}C_\lambda(Y) \right)^{m-1} \pi_{r,1}(v) \|A\|,$$

for all N and the proof is done. \square

Remark 2.3. *If we take $t_1 = \dots = t_m$, then, upon polarization, we recover exactly [13, Theorem 1.2] with a much simpler proof due to the fact that the inequality is simpler to prove for the extremal values of (t_1, \dots, t_m) .*

We are now ready for the proof of the sufficient part of Theorem 1.3. We split the proof into three cases, and we combine Theorem 2.2 with the Bennett-Carl inequalities ([3, 8]): for $1 \leq s \leq q \leq +\infty$, the inclusion map $\ell_s \hookrightarrow \ell_q$ is $(r, 1)$ -summing, where the optimal r is given by

$$\frac{1}{r} := \frac{1}{2} + \frac{1}{s} - \frac{1}{\min\{2, q\}}.$$

(i) $s \leq q \leq 2$: The Bennet-Carl-inequalities ensure that the inclusion map $\ell_s \hookrightarrow \ell_q$ is $(r, 1)$ -summing with $\frac{1}{r} = \frac{1}{2} + \frac{1}{s} - \frac{1}{q}$, so the results follow from Theorem 2.2, with t_1, \dots, t_m satisfying

$$\frac{1}{t_1} + \dots + \frac{1}{t_m} = \frac{1}{2} + \frac{1}{s} - \frac{1}{q} - \left| \frac{1}{\mathbf{p}} \right| + \frac{m-1}{\max\{\lambda, 2\}}.$$

(ii) $s \leq 2 \leq q$: Also by using Bennet-Carl inequalities, $\ell_s \hookrightarrow \ell_2$ is $(s, 1)$ -summing, thus we get (1.5) applying Theorem 2.2, with t_1, \dots, t_m satisfying

$$\frac{1}{t_1} + \dots + \frac{1}{t_m} = \frac{1}{s} - \left| \frac{1}{\mathbf{p}} \right| + \frac{m-1}{\max\{\lambda, 2\}}.$$

(iii) $2 \leq s \leq q$: Since $\ell_s \hookrightarrow \ell_s$ is $(s, 1)$ -summing, the result follows from Theorem 2.2, with $t_1 = \dots = t_m = \lambda$ and $\lambda \geq s$, since $r = s$ and

$$\frac{1}{\lambda} := \frac{1}{s} - \left| \frac{1}{\mathbf{p}} \right| \leq \frac{1}{s}.$$

Remark 2.4. *Let us set*

$$c_{qs} := \begin{cases} q, & \text{if } s \leq q \leq 2, \\ 2, & \text{if } s \leq 2 \leq q, \\ s, & \text{if } 2 \leq s \leq q. \end{cases}$$

With the above notations, a careful look at the proof shows that the constant C which appears in Theorem 1.3 is dominated by

$$\left(\sqrt{2}C_{\max\{\lambda,s,2\}}(\ell_{c_{qs}})\right)^{m-1} \pi_{r,1}(\ell_s \hookrightarrow \ell_{c_{qs}})$$

3. PROOF OF THEOREM 1.3 (OPTIMALITY)

In this section we show that the exponents in Theorem 1.3 are optimal except when $q \leq 2$ and $\left|\frac{1}{\mathbf{p}}\right| > \frac{1}{2}$. More precisely, if $(t_1, \dots, t_m) \in [1, +\infty)^m$ are such that there exists $C > 1$ satisfying, for any continuous multilinear map $X_{p_1} \times \dots \times X_{p_m} \rightarrow X_s$,

$$(3.1) \quad \left(\sum_{i_1=1}^{+\infty} \left(\dots \left(\sum_{i_m=1}^{+\infty} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^{t_m} \right)^{\frac{t_{m-1}}{t_m}} \dots \right)^{\frac{t_1}{t_2}} \right)^{\frac{1}{t_1}} \leq C \|A\|,$$

then we prove that (1.4) holds. When $\lambda \geq 2$, we will always assume that $t_1 = \dots = t_m = t$, since $\lambda = \max\{\lambda, s, 2\}$ and our inequality holds true when all the exponents are equal. We split the proof into several cases. Most of the cases are a consequence of a random construction. The main tool is the following lemma, from [1, Lemma 6.2].

Lemma 3.1. *Let $d, n \geq 1$, $q_1, \dots, q_{d+1} \in [1, +\infty]^{d+1}$ and let, for $q \geq 1$,*

$$\alpha(q) = \begin{cases} \frac{1}{2} - \frac{1}{q} & \text{if } q \geq 2 \\ 0 & \text{otherwise.} \end{cases}$$

Then there exists a d -linear mapping $A : \ell_{p_1}^n \times \dots \times \ell_{p_d}^n \rightarrow \ell_{p_{d+1}}^n$ which may be written

$$A(x^{(1)}, \dots, x^{(d)}) = \sum_{i_1, \dots, i_{d+1}=1}^n \pm x_{i_1}^{(1)} \dots x_{i_d}^{(d)} e_{i_{d+1}}$$

such that

$$\|A\| \leq C_d n^{\frac{1}{2} + \alpha(p_1) + \dots + \alpha(p_d) + \alpha(p_{d+1}^*)}.$$

3.1. Case 1: $1 \leq s \leq q \leq 2$ and $\lambda < 2$. This case has already been solved in [1, Section 6.2], using Lemma 3.1 with $d = m$ and $(q_1, \dots, q_{m+1}) = (p_1, \dots, p_m, s)$.

3.2. Case 2: $1 \leq s \leq q \leq 2$, $\lambda \geq 2$ and $\left|\frac{1}{\mathbf{p}}\right| \leq \frac{1}{2}$. This case has already been solved in [13, Proposition 4.4(ib)] using a Fourier matrix. We shall give an alternative probabilistic proof. Let $p \in [2, +\infty]$ be such that $\frac{1}{p} = \left|\frac{1}{\mathbf{p}}\right|$. By Lemma 3.1, there exists a linear map $T : \ell_p^n \rightarrow \ell_s^n$ which may be written $T(x) = \sum_{i,j} \varepsilon_{i,j} x_i e_j$ with $\varepsilon_{i,j} = \pm 1$ and such that

$$\|T\| \leq C n^{\frac{1}{2} + \frac{1}{2} - \frac{1}{p} + \frac{1}{2} - \frac{1}{s^*}} = C n^{\frac{1}{2} + \frac{1}{s} - \left|\frac{1}{\mathbf{p}}\right|}.$$

Let $A : \ell_{p_1}^n \times \dots \times \ell_{p_m}^n \rightarrow \ell_s^n$ defined by

$$A(x^{(1)}, \dots, x^{(m)}) := \sum_{i,j} \varepsilon_{i,j} x_i^{(1)} \dots x_i^{(m)} e_j.$$

By Hölder's inequality, it is plain that $\|A\| \leq \|T\| \leq C n^{\frac{1}{2} + \frac{1}{s} - \left|\frac{1}{\mathbf{p}}\right|}$. On the other hand, since $A(e_{i_1}, \dots, e_{i_m}) \neq 0$ if and only if $i_1 = \dots = i_m$, and

$$\|A(e_i, \dots, e_i)\|_{\ell_q} = n^{1/q},$$

we have

$$\left(\sum_{\mathbf{i} \in \mathcal{M}(m,n)} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^t \right)^{\frac{1}{t}} = n^{\frac{1}{q} + \frac{1}{t}}.$$

This clearly implies

$$\frac{1}{t} \leq \frac{1}{2} + \frac{1}{s} - \frac{1}{q} - \left| \frac{1}{\mathbf{p}} \right|.$$

3.3. Case 3: $1 \leq s \leq 2 \leq q$ and $\lambda < 2$. Let $p \in [0, +\infty]$ be defined by

$$\frac{1}{p} = \frac{1}{p_m} + \frac{1}{s^*}.$$

Since $\lambda < 2$, it is easy to check that $p \geq 2$ and that $p_i \geq 2$ for any $i = 1, \dots, m$. We then apply Lemma 3.1 with $d = m - 1$ and $(q_1, \dots, q_m) = (p_1, \dots, p_{m-1}, p^*)$. We get an $(m - 1)$ -linear form $T : \ell_{p_1}^n \times \dots \times \ell_{p_{m-1}}^n \rightarrow \ell_{p^*}^n$ which can be written

$$T(x^{(1)}, \dots, x^{(m-1)}) = \sum_{i_1, \dots, i_m} \varepsilon_{i_1, \dots, i_m} x_{i_1}^{(1)} \dots x_{i_{m-1}}^{(m-1)} e_{i_m}$$

and such that

$$\|T\| \leq C n^{\frac{1}{2} + \frac{m}{2} - \left| \frac{1}{\mathbf{p}} \right| - \frac{1}{s^*}} = C n^{\frac{m-1}{2} - \left| \frac{1}{\mathbf{p}} \right| + \frac{1}{s}}.$$

We then define $A : \ell_{p_1}^n \times \dots \times \ell_{p_m}^n \rightarrow \ell_s^n$ by

$$A(x^{(1)}, \dots, x^{(m)}) = \sum_{i_1, \dots, i_m} \varepsilon_{i_1, \dots, i_m} x_{i_1}^{(1)} \dots x_{i_m}^{(m)} e_{i_m}.$$

Then, for any $x^{(1)}, \dots, x^{(m)} \in B_{\ell_{p_1}^n} \times \dots \times B_{\ell_{p_m}^n}$,

$$\begin{aligned} \|A(x^{(1)}, \dots, x^{(m)})\| &= \sup_{y \in B_{\ell_s^n}} \left| \sum_{i_1, \dots, i_m} \varepsilon_{i_1, \dots, i_m} x_{i_1}^{(1)} \dots x_{i_m}^{(m)} y_{i_m} \right| \\ &\leq \sup_{z \in B_{\ell_p^n}} \left| \sum_{i_1, \dots, i_m} \varepsilon_{i_1, \dots, i_m} x_{i_1}^{(1)} \dots x_{i_{m-1}}^{(m-1)} z_{i_m} \right| \\ &\leq \|T\|. \end{aligned}$$

Moreover, given any $\mathbf{i} \in \mathcal{M}(m, n)$, $\|A(e_{i_1}, \dots, e_{i_m})\|_q = \|e_{i_m}\|_q = 1$, so that

$$\left(\sum_{i_1=1}^{+\infty} \left(\dots \left(\sum_{i_m=1}^{+\infty} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^{t_m} \right)^{\frac{t_{m-1}}{t_m}} \dots \right)^{\frac{t_1}{t_2}} \right)^{\frac{1}{t_1}} = n^{\frac{1}{t_1} + \dots + \frac{1}{t_m}}.$$

Hence, provided (3.1) is satisfied, (t_1, \dots, t_m) has to satisfy

$$\frac{1}{t_1} + \dots + \frac{1}{t_m} \leq \frac{m-1}{2} + \frac{1}{s} - \left| \frac{1}{\mathbf{p}} \right|.$$

3.4. Case 4 and Case 5: $1 \leq s \leq 2 \leq q$ and $\lambda \geq 2$, $2 \leq s \leq q$. These cases have a deterministic proof, as noted in [13, Proposition 4.4 (ii), (iii)], considering $A : \ell_{p_1}^n \times \dots \times \ell_{p_m}^n \rightarrow \ell_s^n$ given by

$$A(x^{(1)}, \dots, x^{(m)}) := \sum_{i=1}^n x_i^{(1)} \dots x_i^{(m)} e_i.$$

3.5. The proof of Theorem 1.1. From Theorem 1.3, by choosing $t_1 = \dots = t_m$ we conclude that provided

$$\left| \frac{1}{\mathbf{p}} \right| < \frac{1}{2} + \frac{1}{s} - \frac{1}{\min\{q, 2\}},$$

the best exponent ρ in Theorem 1.1 satisfies

$$\frac{m}{\rho} = \frac{1}{\lambda} + \frac{m-1}{\max\{\lambda, s, 2\}}.$$

To conclude the proof, it remains to prove that, whenever

$$\left| \frac{1}{\mathbf{p}} \right| \geq \frac{1}{2} + \frac{1}{s} - \frac{1}{\min\{q, 2\}},$$

we cannot find an exponent $\rho > 0$ such that (1.1) is satisfied for all m -linear operators $A : X_{p_1} \times \cdots \times X_{p_m} \rightarrow X_s$. In fact, everything has already been done before: if $q \leq 2$, then we have just to follow the lines of Case 2 and if $q \geq 2$, then we may consider the m -linear mapping of Cases 4 and 5.

4. THE CASE $1 \leq s \leq q \leq 2$, $\lambda \geq 2$ AND $\left|\frac{1}{\mathbf{p}}\right| > \frac{1}{2}$

4.1. A reformulation of the Hardy-Littlewood type inequalities. We shall improve in this section the bound given by Theorem 1.1. We shall proceed by interpolation. To do this, we need a reformulation of the result of this theorem, as Villanueva and Perez-Garcia reformulated the Bohnenblust-Hille inequality in [24]. The proof is a combination of [24, Corollary 3.20] and Proposition 2.2 and will be omitted.

Theorem 4.1. *Let $1 \leq p_1, \dots, p_m \leq +\infty$, $1 \leq s \leq q \leq \infty$ and let $\rho > 0$. The following assertions are equivalent.*

(A) *There exists $C > 0$ such that, for every continuous m -linear mapping $A : X_{p_1} \times \cdots \times X_{p_m} \rightarrow X_s$, we have*

$$\left(\sum_{i_1, \dots, i_m} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^\rho \right)^{1/\rho} \leq C \|A\|.$$

(B) *There exists $C > 0$ such that, for any $n \geq 1$, for any Banach spaces Y_1, \dots, Y_m , for any continuous m -linear mapping $S : Y_1 \times \cdots \times Y_m \rightarrow X_s$, the induced operator*

$$\begin{aligned} T : \ell_{p_1^*, w}^n(Y_1) \times \cdots \times \ell_{p_m^*, w}^n(Y_m) &\rightarrow \ell_\rho^m(X_s) \\ (x^{(1)}, \dots, x^{(m)}) &\mapsto (S(x_{i_1}^{(1)}, \dots, x_{i_m}^{(m)}))_{\mathbf{i} \in \mathcal{M}(m, n)} \end{aligned}$$

satisfies $\|T\| \leq C \|S\|$.

We recall that, for any $p \in [1, +\infty]$ and any Banach space Y ,

$$\ell_{p, w}^n(Y) = \left\{ (x_j)_{j=1}^n \subset Y; \|(x_j)\|_{w, p} := \sup_{\varphi \in B_{Y^*}} \left(\sum_{j=1}^n |\varphi(x_j)|^p \right)^{1/p} < +\infty \right\}$$

with the appropriate modifications for $p = \infty$.

4.2. Proof of the sufficient condition. We now prove our better upper bound in the case $1 \leq s \leq q \leq 2$, $\left|\frac{1}{\mathbf{p}}\right| > \frac{1}{2}$ (namely we prove the first part of Theorem 1.2). Let $n \geq 1$, let Y_1, \dots, Y_m be Banach spaces and let $S : Y_1 \times \cdots \times Y_m \rightarrow X_s$ be bounded. Let $n \geq 1$ and let T be the operator induced by S on $\mathcal{Y} = \ell_{p_1^*, w}^n(Y_1) \times \cdots \times \ell_{p_m^*, w}^n(Y_m)$, defined by

$$T(x^{(1)}, \dots, x^{(m)}) = (S(x_{i_1}^{(1)}, \dots, x_{i_m}^{(m)})).$$

Then T is bounded as an operator from \mathcal{Y} into $\ell_\infty^m(X_s)$ (this is trivial). T is also bounded as an operator from \mathcal{Y} into $\ell_\rho^m(X_s)$ with $\frac{1}{\rho} = \frac{1}{s} - \left|\frac{1}{\mathbf{p}}\right|$ (this is Theorem 1.1 for $1 \leq s \leq 2$ and $q \geq 2$). We can interpolate between these two extreme situations. Hence, let $q \in [s, 2]$ and let $\theta \in [0, 1]$ be such that

$$\frac{1}{q} = \frac{1-\theta}{s} + \frac{\theta}{2} \iff \theta = \frac{\frac{1}{s} - \frac{1}{q}}{\frac{1}{s} - \frac{1}{2}}.$$

By [4, Theorem 4.4.1], T is bounded as an operator from \mathcal{Y} into $\ell_t^m(X_q)$ where

$$\frac{1}{t} = \frac{1-\theta}{\infty} + \frac{\theta}{\rho} = \frac{\left(\frac{1}{s} - \frac{1}{q}\right) \left(\frac{1}{s} - \left|\frac{1}{\mathbf{p}}\right|\right)}{\frac{1}{s} - \frac{1}{2}}.$$

Remark 4.2. *It is easy to check that, for $1 \leq s \leq q \leq 2$ and $\left|\frac{1}{\mathbf{p}}\right| \geq \frac{1}{2}$, then the bound $\frac{\left(\frac{1}{s} - \frac{1}{q}\right) \left(\frac{1}{s} - \left|\frac{1}{\mathbf{p}}\right|\right)}{\frac{1}{s} - \frac{1}{2}}$ is always better (namely larger) than the bound $\frac{1}{2} + \frac{1}{s} - \frac{1}{q} - \left|\frac{1}{\mathbf{p}}\right|$ obtained in Theorem 1.1.*

4.3. The necessary condition. We now prove the second part of Theorem 1.2. It also uses a probabilistic device for linear maps when the two spaces do not need to have the same dimension. The forthcoming lemma can be found in [3, Proposition 3.2].

Lemma 4.3. *Let $n, d \geq 1$, $1 \leq p, s \leq 2$. There exists $T : \ell_p^d \rightarrow \ell_s^n$, $T(x) = \sum_{i,j} \pm x_j e_i$ such that*

$$\|T\| \leq C_{p,s} \max \left(d^{1/s}, n^{1-\frac{1}{p}} d^{\frac{1}{s}-\frac{1}{2}} \right).$$

Coming back to the proof of Theorem 1.2, we first observe that we may always assume that $\left| \frac{1}{\mathbf{p}} \right| < 1$. Otherwise, we can consider the m -linear map $A : X_{p_1} \times \cdots \times X_{p_m} \rightarrow X_s$ defined by

$$A(x^{(1)}, \dots, x^{(m)}) = \sum_{i \geq 1} x_i^{(1)} \cdots x_i^{(m)} e_0$$

and observe that it is bounded whereas it has infinitely many coefficients equal to 1. We then define $p \in [1, 2]$ by $\frac{1}{p} = \left| \frac{1}{\mathbf{p}} \right|$ and we consider $T : \ell_p^d \rightarrow \ell_s^n$, $T(x) = \sum_{i,j} \varepsilon_{i,j} x_j e_i$ the map given by Lemma 4.3. We then define

$$\begin{aligned} A : \ell_{p_1}^d \times \cdots \times \ell_{p_m}^d &\rightarrow \ell_s^n \\ (x^{(1)}, \dots, x^{(m)}) &\mapsto \sum_{i,j} \varepsilon_{i,j} x_j^{(1)} \cdots x_j^{(m)} e_i \end{aligned}$$

and we observe that, by Hölder's inequality, $\|A\| \leq \|T\|$. Furthermore,

$$\left(\sum_{i_1, \dots, i_m} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^t \right)^{1/t} = n^{1/t} d^{1/q}.$$

Taking $d^{1/2} = n^{1-\frac{1}{p}}$ (this is the optimal relation between d and n), we get that if

$$\left(\sum_{i_1, \dots, i_m} \|A(e_{i_1}, \dots, e_{i_m})\|_{\ell_q}^t \right)^{1/t} \leq C \|A\|,$$

then it is necessary that

$$\frac{1}{t} \leq 2 \left(1 - \frac{1}{p} \right) \left(\frac{1}{s} - \frac{1}{q} \right).$$

Remark 4.4. *This last condition is optimal when $s = 1$ or when $\left| \frac{1}{\mathbf{p}} \right| = \frac{1}{2}$ (with, in fact, the same proof as in Case 2 above). When $1 < s < 2$, another necessary condition is*

$$\frac{1}{t} \leq \frac{1}{s} - \left| \frac{1}{\mathbf{p}} \right|$$

(see Case 4 or Case 5 above).

5. OPTIMAL ESTIMATES UNDER COTYPE ASSUMPTIONS

For a Banach space X , let $q_X := \inf\{q \geq 2; X \text{ has cotype } q\}$. For scalar-valued multilinear operators it is easy to observe that summability in multiple indexes behaves in a quite different way than summability in just one index. For instance, for any bounded bilinear form $A : c_0 \times c_0 \rightarrow \mathbb{C}$,

$$\left(\sum_{i,j=1}^{+\infty} |A(e_i, e_j)|^{\frac{4}{3}} \right)^{\frac{3}{4}} \leq \sqrt{2} \|A\|$$

and the exponent $4/3$ is optimal. But, if we sum diagonally ($i = j$) the exponent $4/3$ can be reduced to 1 since

$$\sum_{i=1}^{+\infty} |A(e_i, e_i)| \leq \|A\|$$

for any bounded bilinear form $A : c_0 \times c_0 \rightarrow \mathbb{C}$. Now we prove Theorem 1.5 which shows that when replacing the scalar field by infinite-dimensional spaces the situation is quite different.

Proof. (A) \Rightarrow (B). From a deep result of Maurey and Pisier ([22] and [12, Section 14]), ℓ_{q_X} is finitely representable in X , which means that, for any $n \geq 1$, one may find unit vectors $z_1, \dots, z_n \in X$ such that, for any $a_1, \dots, a_n \in \mathbb{C}$,

$$\sum_{i=1}^n \|a_i z_i\|_X \leq 2 \left(\sum_{i=1}^n |a_i|^{q_X} \right)^{1/q_X}.$$

We then consider the m -linear map $A : \ell_{p_1}^n \times \dots \times \ell_{p_m}^n \rightarrow X$ defined by

$$A(x^{(1)}, \dots, x^{(m)}) := \sum_{i=1}^n x_i^{(1)} \dots x_i^{(m)} z_i.$$

Then, for any $(x^{(1)}, \dots, x^{(m)})$ belonging to $B_{\ell_{p_1}^n} \times \dots \times B_{\ell_{p_m}^n}$,

$$\begin{aligned} \left\| A(x^{(1)}, \dots, x^{(m)}) \right\| &\leq 2 \left(\sum_{i=1}^n |x_i^{(1)}|^{q_X} \dots |x_i^{(m)}|^{q_X} \right)^{1/q_X} \\ &\leq 2n^{\frac{1}{q_X} - \left| \frac{1}{\mathbf{p}} \right|} \end{aligned}$$

where the last inequality follows from Hölder's inequality applied to the exponents

$$\frac{p_1}{q_X}, \dots, \frac{p_m}{q_X}, \left(1 - q_X \left| \frac{1}{\mathbf{p}} \right| \right)^{-1}.$$

On the other hand,

$$\left(\sum_{i=1}^n \|A(e_i, \dots, e_i)\|^\rho \right)^{1/\rho} = n^{\frac{1}{\rho}}$$

and we obtain (3).

(B) \Rightarrow (A). This implication is proved in [13, Proposition 4.3]. \square

If X does not have cotype q_X , the condition remains necessary. But now we just have the following sufficient condition:

$$\frac{m}{\rho} < \frac{1}{q_X} - \left| \frac{1}{\mathbf{p}} \right|.$$

Of course, it would be nice to determine what happens in this case. A look at [12, page 304] shows that the situation does not look simple.

As a consequence of the previous result we conclude that under certain circumstances the concepts of absolutely summing multilinear operator and multiple summing multilinear operator (see [6, 21, 23]) are precisely the same.

Corollary 5.1. *Let $p \in [2, +\infty]$, let X be an infinite dimensional Banach space with cotype $q_X < \frac{p}{m}$ and let $\rho > 0$. The following assertions are equivalent:*

- (A) *Every bounded m -linear operator $A : X_p \times \dots \times X_p \rightarrow X$ is absolutely $(\rho; p^*)$ -summing.*
- (B) *Every bounded m -linear operator $A : X_p \times \dots \times X_p \rightarrow X$ is multiple $(\rho; p^*)$ -summing.*
- (C) $\frac{1}{\rho} \leq \frac{1}{q_X} - \frac{m}{p}$.

We stress the equivalence between (A) and (B) is not true, in general. For instance, every bounded bilinear operator $A : \ell_2 \times \ell_2 \rightarrow \ell_2$ is absolutely $(1; 1)$ -summing but this is no longer true for multiple summability.

6. CONSTANTS OF VECTOR-VALUED BOHNENBLUST–HILLE INEQUALITIES

A particular case of our main result is the following vector-valued Bohnenblust–Hille inequality (see [10, Lemma 3] and also [26, Section 2.2]):

Theorem 6.1. *Let X be a Banach space, Y a cotype q space and $v : X \rightarrow Y$ an $(r, 1)$ -summing operator with $1 \leq r \leq q$. Then, for all m -linear operators $T : c_0 \times \cdots \times c_0 \rightarrow X$,*

$$\left(\sum_{i_1, \dots, i_m=1}^{+\infty} \|vT(e_{i_1}, \dots, e_{i_m})\|_Y^{\frac{qrm}{q+(m-1)r}} \right)^{\frac{q+(m-1)r}{qrm}} \leq C_{Y,m} \pi_{r,1}(v) \|T\|$$

with $C_{Y,m} = (\sqrt{2}C_q(Y))^{m-1}$.

In this section, in Theorem 6.2, we improve the above estimate for $C_{Y,m}$. The proof of Theorem 6.2 follows almost word by word the proof of [2, Proposition 3.1] using [9, Lemma 2.2] and Kahane's inequality instead of the Khinchine inequality. We present the proof for the sake of completeness. We need the following inequality due to Kahane:

Kahane's Inequality. *Let $0 < p, q < +\infty$. Then there is a constant $K_{p,q} > 0$ for which*

$$\left(\int_I \left\| \sum_{k=1}^n r_k(t)x_k \right\|^q dt \right)^{\frac{1}{q}} \leq K_{p,q} \left(\int_I \left\| \sum_{k=1}^n r_k(t)x_k \right\|^p dt \right)^{\frac{1}{p}},$$

regardless of the choice of a Banach space X and of finitely many vectors $x_1, \dots, x_n \in X$.

Theorem 6.2. *For all m and all $1 \leq k < m$,*

$$C_{Y,m} \leq \left(C_q(Y) K_{\frac{qrk}{q+(k-1)r}, 2} \right)^{m-k} C_{Y,k}.$$

Proof. Let $\rho := \frac{qrm}{q+(m-1)r}$ and to simplify notation let us write

$$vTe_i = vT(e_{i_1}, \dots, e_{i_m}).$$

Let us make use of [2, Remark 2.2] with $m \geq 2$, $1 \leq k \leq m-1$ and $s = \frac{qrk}{q+(k-1)r}$. So we have

$$(6.1) \quad \left(\sum_{\mathbf{i}} \|vTe_{\mathbf{i}}\|_Y^\rho \right)^{\frac{1}{\rho}} \leq \prod_{S \in P_k(m)} \left(\sum_{\mathbf{i}_S} \left(\sum_{\mathbf{i}_{\hat{S}}} \|vT(e_{\mathbf{i}_S}, e_{\mathbf{i}_{\hat{S}}})\|_Y^q \right)^{\frac{s}{q}} \right)^{\frac{1}{s} \binom{m}{k}},$$

where $P_k(m)$ denotes the set of all subsets of $\{1, \dots, m\}$ with cardinality k . For sake of clarity, we shall assume that $S = \{1, \dots, k\}$. By the multilinear cotype inequality (see [9, Lemma 2.2]) and the Kahane inequality, we have

$$\begin{aligned} & \left(\sum_{\mathbf{i}_{\hat{S}}} \|vT(e_{\mathbf{i}_S}, e_{\mathbf{i}_{\hat{S}}})\|_Y^q \right)^{\frac{s}{q}} \\ & \leq (C_q(Y)K_{s,2})^{s(m-k)} \int_{I^{m-k}} \left\| \sum_{\mathbf{i}_{\hat{S}}} r_{\mathbf{i}_{\hat{S}}}(t_{\hat{S}}) vT(e_{\mathbf{i}_S}, e_{\mathbf{i}_{\hat{S}}}) \right\|_Y^s dt_{\hat{S}} \\ & = (C_q(Y)K_{s,2})^{s(m-k)} \int_{I^{m-k}} \left\| vT \left(e_{\mathbf{i}_S}, \sum_{\mathbf{i}_{\hat{S}}} r_{\mathbf{i}_{\hat{S}}}(t_{\hat{S}}) e_{\mathbf{i}_{\hat{S}}} \right) \right\|_Y^s dt_{\hat{S}} \\ & = (C_q(Y)K_{s,2})^{s(m-k)} \int_{I^{m-k}} \left\| v \left(T \left(e_{i_1}, \dots, e_{i_k}, \sum_{i_{k+1}} r_{k+1}(t_{k+1}) e_{k+1}, \dots, \sum_{i_m} r_m(t_m) e_m \right) \right) \right\|_Y^s dt_{k+1} \dots dt_m \end{aligned}$$

But for a fixed choice of $(t_{k+1}, \dots, t_m) \in I^{m-k} = [0, 1]^{m-k}$, we know, by Theorem 6.1, that

$$\sum_{i_1, \dots, i_k} \left\| v \left(T \left(e_{i_1}, \dots, e_{i_k}, \sum_{i_{k+1}} r_{k+1}(t_{k+1})e_{k+1}, \dots, \sum_{i_m} r_m(t_m)e_m \right) \right) \right\|_Y^s \leq (C_{Y,k} \pi_{r,1}(v) \|T\|)^s.$$

Thus,

$$\begin{aligned} & \sum_{\mathbf{i}_S} \left(\sum_{\mathbf{i}_{\hat{S}}} \|vT(e_{\mathbf{i}_S}, e_{\mathbf{i}_{\hat{S}}})\|_Y^q \right)^{\frac{s}{q}} \\ & \leq (C_q(Y)K_{s,2})^{s(m-k)} \cdot \sum_{i_1, \dots, i_k} \left\| v \left(T \left(e_{i_1}, \dots, e_{i_k}, \sum_{i_{k+1}} r_{k+1}(t_{k+1})e_{k+1}, \dots, \sum_{i_m} r_m(t_m)e_m \right) \right) \right\|_Y^s \\ (6.2) \quad & \leq \left((C_q(Y)K_{s,2})^{m-k} \pi_{r,1}(v) C_{Y,k} \|T\| \right)^s, \end{aligned}$$

namely

$$\left(\sum_{\mathbf{i}_S} \left(\sum_{\mathbf{i}_{\hat{S}}} \|vT(e_{\mathbf{i}_S}, e_{\mathbf{i}_{\hat{S}}})\|_Y^q \right)^{\frac{s}{q}} \right)^{\frac{1}{s}} \leq (C_q(Y)K_{s,2})^{m-k} \pi_{r,1}(v) C_{Y,k} \|T\|.$$

From (6.1) we conclude that

$$\left(\sum_{\mathbf{i}} \|vT e_{\mathbf{i}}\|_Y^\rho \right)^{\frac{1}{\rho}} \leq (C_q(Y)K_{s,2})^{m-k} C_{Y,k} \pi_{r,1}(v) \|T\|.$$

□

When m is even, the case $k = \frac{m}{2}$ recovers the constants from [26].

Corollary 6.3. *For all m ,*

$$C_{Y,m} \leq C_q(Y)^{m-1} \prod_{k=1}^{m-1} K_{\frac{qrk}{q+(k-1)r}, 2}.$$

7. OTHER EXPONENTS

From now on $1 \leq r \leq q$ and $(q_1, \dots, q_m) \in [r, q]^m$ so that

$$\frac{1}{q_1} + \dots + \frac{1}{q_m} = \frac{q + (m-1)r}{qr} = \frac{1}{r} + \frac{m-1}{q}$$

are called vector-valued Bohnenblust–Hille exponents. From Theorem 2.2 we have:

Theorem 7.1 (Multiple exponent vector-valued Bohnenblust–Hille inequality). *Let X be a Banach space and Y a cotype q space with $1 \leq r \leq q$. If $(q_1, \dots, q_m) \in [r, q]^m$ are vector-valued Bohnenblust–Hille exponents, then there exists $C_{Y,q_1, \dots, q_m} \geq 1$ such that, for all m -linear operators $T : c_0 \times \dots \times c_0 \rightarrow X$ and every $(r, 1)$ -summing operator $v : X \rightarrow Y$, we have*

$$(7.1) \quad \left(\sum_{i_1=1}^{+\infty} \dots \left(\sum_{i_m=1}^{+\infty} \|vT e_{\mathbf{i}}\|_Y^{q_m} \right)^{\frac{q_m-1}{q_m}} \dots \right)^{\frac{1}{q_1}} \leq C_{Y,q_1, \dots, q_m} \pi_{r,1}(v) \|T\|,$$

with $C_{Y,q_1, \dots, q_m} = (\sqrt{2}C_q(Y))^{m-1}$.

Our final result gives better estimates for the constants C_{Y,q_1, \dots, q_m} :

Theorem 7.2. *If (q_1, \dots, q_m) is a vector-valued Bohnenblust–Hille exponent, then*

$$C_{Y,q_1, \dots, q_m} \leq \prod_{k=1}^m \left(\left(C_q(Y) K_{\frac{kqr}{q+(k-1)r}, 2} \right)^{m-k} C_{Y,k} \right)^{\theta_k}$$

with

$$(7.2) \quad \theta_m = m \left(\frac{1}{r} - \frac{1}{q} \right)^{-1} \left(\frac{1}{q_m} - \frac{1}{q} \right)$$

and

$$(7.3) \quad \theta_k = k \left(\frac{1}{r} - \frac{1}{q} \right)^{-1} \left(\frac{1}{q_k} - \frac{1}{q_{k+1}} \right), \quad \text{for } k = 1, \dots, m-1.$$

Proof. It suffices to consider $q_i \leq q_j$ whenever $i < j$. For each $k = 1, \dots, m$, define

$$s_k = \frac{kqr}{q + (k-1)r}.$$

From the proof of Theorem 6.2 we have (7.1) for each exponent $(s_k, k \text{ times}, s_k, q, \dots, q)$. More precisely, from (6.2) we have

$$\left(\sum_{i_1, \dots, i_k} \left(\sum_{i_{k+1}, \dots, i_m} \|vTe_i\|_Y^q \right)^{\frac{s_k}{q}} \right)^{\frac{1}{s_k}} \leq (C_q(Y)K_{s_k,2})^{m-k} C_{Y,k} \pi_{r,1}(v) \|T\|.$$

Consequently, for each $k = 1, \dots, m$ we have

$$C_{Y, s_k, k \text{ times}, s_k, q, \dots, q} \leq (C_q(Y)K_{s_k,2})^{m-k} C_{Y,k}.$$

Since every vector-valued Bohnenblust–Hille exponent (q_1, \dots, q_m) with $q_1 \leq \dots \leq q_m$ is obtained by interpolation of $\alpha_1, \dots, \alpha_m$ with $\alpha_k = (s_k, k \text{ times}, s_k, q, \dots, q)$, and $\theta_1, \dots, \theta_m$ as in (7.2) and (7.3), we conclude that

$$C_{Y, q_1, \dots, q_m} \leq \prod_{k=1}^m \left(C_{Y, s_k, k \text{ times}, s_k, q, \dots, q} \right)^{\theta_k} \leq \prod_{k=1}^m \left((C_q(Y)K_{s_k,2})^{m-k} C_{Y,k} \right)^{\theta_k}$$

□

A particular case of Kahane’s inequality is Khintchine’s inequality: if (ε_i) is a sequence of independent Rademacher variables, then, for any $p \in [1, 2]$, there exists a constant $A_{\mathbb{R},p}$ such that, for any $n \geq 1$ and any $a_1, \dots, a_n \in \mathbb{R}$,

$$\left(\sum_{i=1}^n |a_i|^2 \right)^{\frac{1}{2}} \leq A_{\mathbb{R},p} \left(\int_{\Omega} \left| \sum_{i=1}^n a_i \varepsilon_i(\omega) \right|^p d\omega \right)^{\frac{1}{p}}.$$

It has a complex counterpart: for any $p \in [1, 2]$, there exists a constant $A_{\mathbb{C},p}$ such that, for any $n \geq 1$ and any $a_1, \dots, a_n \in \mathbb{C}$,

$$\left(\sum_{i=1}^n |a_i|^2 \right)^{\frac{1}{2}} \leq A_{\mathbb{C},p} \left(\int_{\mathbb{T}^n} \left| \sum_{i=1}^n a_i z_i \right|^p dz \right)^{\frac{1}{p}}.$$

The best constants $A_{\mathbb{R},p}$ and $A_{\mathbb{C},p}$ are known (see [15] and [18]):

- $A_{\mathbb{R},p} = \begin{cases} 2^{\frac{1}{p}-\frac{1}{2}}, & \text{if } 0 < p \leq p_0 \approx 1.847 \\ \frac{1}{\sqrt{2}} \left(\frac{\Gamma(\frac{1+p}{2})}{\sqrt{\pi}} \right)^{-\frac{1}{p}}, & \text{if } p > p_0; \end{cases}$
- $A_{\mathbb{C},p} = \Gamma\left(\frac{1+p}{2}\right)^{-\frac{1}{p}}, \text{ if } 1 < p \leq 2.$

Taking $X = Y = \mathbb{K}$ and $r = 1$ we obtain estimates for the constants of the scalar-valued Bohnenblust–Hille inequality with multiple exponents:

Corollary 7.3. *If $(q_1, \dots, q_m) \in [1, 2]^m$ are so that*

$$\frac{1}{q_1} + \dots + \frac{1}{q_m} = \frac{m+1}{2},$$

then

$$\left(\sum_{i_1=1}^{+\infty} \dots \left(\sum_{i_m=1}^{+\infty} |T(e_{i_1}, \dots, e_{i_m})|^{q_m} \right)^{\frac{q_{m-1}}{q_m}} \dots \right)^{\frac{1}{q_1}} \leq C_{\mathbb{K}, m}^{2m\left(\frac{1}{q_m} - \frac{1}{2}\right)} \left(\prod_{k=1}^{m-1} \left(A_{\mathbb{K}, \frac{2k}{k+1}}^{m-k} C_{\mathbb{K}, k} \right)^{2k\left(\frac{1}{q_k} - \frac{1}{q_{k+1}}\right)} \right) \|T\|$$

for all m -linear operators $T : c_0 \times \dots \times c_0 \rightarrow \mathbb{K}$. In particular, for complex scalars, the left hand side of the above inequality can be replaced by

$$\left(\prod_{j=1}^m \Gamma\left(2 - \frac{1}{j}\right)^{\frac{j}{2-2j}} \right)^{2m\left(\frac{1}{q_m} - \frac{1}{2}\right)} \left(\prod_{k=1}^{m-1} \left(\Gamma\left(\frac{3k+1}{2k+2}\right)^{\left(\frac{-k-1}{2k}\right)(m-k)} \prod_{j=1}^k \Gamma\left(2 - \frac{1}{j}\right)^{\frac{j}{2-2j}} \right)^{2k\left(\frac{1}{q_k} - \frac{1}{q_{k+1}}\right)} \right) \|T\|.$$

Acknowledgement. The authors thank the referee for important comments and suggestions.

REFERENCES

- [1] N. Albuquerque, F. Bayart, D. Pellegrino, and J.B. Seoane-Sepúlveda, *Sharp generalizations of the multilinear Bohnenblust–Hille inequality*, J. Funct. Anal., **266** (2014), 3726–3740.
- [2] F. Bayart, D. Pellegrino, and J.B. Seoane-Sepúlveda, *The Bohr radius of the n -dimensional polydisk is equivalent to $\sqrt{(\log n)/n}$* , arXiv:1310.2834v2 [math.FA], 15 Oct 2013.
- [3] G. Bennett, *Schur multipliers*, Duke Math. J, **44** (1977), 603–639.
- [4] J. Bergh and J. Löfström, *Interpolation spaces. An introduction*, Springer-Verlag, Berlin, 1976. Grundlehren der Mathematischen Wissenschaften, No. 223.
- [5] H.P. Boas, *Majorant series*, J. Korean Math. Soc., **37** (2000), 321–337.
- [6] G. Botelho, *Cotype and absolutely summing multilinear mappings and homogeneous polynomials*, Proc. Roy. Irish Acad. Sect. A **97** (1997), 145–153.
- [7] H. Bohnenblust and E. Hille, *On the absolute convergence of Dirichlet series*, Ann. of Math. (2) **32**, (1931), no.3, 600–622.
- [8] B. Carl, *Absolut- $(p, 1)$ -summierende identische Operatoren von l_u in l_v* , Math. Nachr. **63** (1974), 353–360.
- [9] A. Defant, D. Popa, and U. Schwarting, *Coordinatewise multiple summing operators in Banach spaces*, J. Funct. Anal. **259** (2010), no.1, 220–242.
- [10] A. Defant and P. Sevilla-Peris, *A new multilinear insight on Littlewood’s 4/3-inequality*, J. Funct. Anal. **256**(5) (2009) 1642–1664.
- [11] A. Defant and P. Sevilla-Peris, *The Bohnenblust–Hille cycle of ideas from a modern point of view*. Funct. Approx. Comment. Math. **50** (2014), no. 1, 55–127.
- [12] J. Diestel, H. Jarchow, and A. Tonge, *Absolutely summing operators*, Cambridge Studies in Advanced Mathematics 43, Cambridge University Press, Cambridge, 1995.
- [13] V. Dimant and P. Sevilla-Peris, *Summation of coefficients of polynomials on ℓ_p spaces*, arXiv:1309.6063v1 [math.FA] 24 Sep 2013.
- [14] J.F. Fournier, *Sobolev’s inequality and Littlewood’s inequality*, Ann. Mat. Pura Appl. **148** (1987), 51–76.
- [15] U. Haagerup, *The best constants in the Khintchine inequality*, Studia Math. **70** (1981), no. 3, 231–283 (1982).
- [16] G. Hardy and J. Littlewood. *Bilinear forms bounded in space $[p, q]$* . Q. J. Math. **5** (1934), 241–254.
- [17] G. Hardy, J. Littlewood and G. Polya, *Inequalities*, Cambridge Mathematical Library, Second Edition, 1951.
- [18] H. König and S. Kwapien, *Best Khintchine type inequalities for sums of independent, rotationally invariant random vectors*, Positivity **5** (2001), no. 2, 115–152.
- [19] S. Kwapien, *Some remarks on (p, q) -absolutely summing operators in ℓ_p -spaces*, Studia Math., 29:327–337, 1968.
- [20] J. E. Littlewood, *On bounded bilinear forms in an infinite number of variables*, Q. J. Math. **1** (1930), 164–174.
- [21] M.C. Matos, *Fully absolutely summing mappings and Hilbert Schmidt operators*, Collect. Math. **54** (2003) 111–136.
- [22] B. Maurey and G. Pisier, *Séries de variables aleatoires vectorielles indépendantes et propriétés géométriques des espaces de Banach*, Studia Math. **58** (1976), 45–90.
- [23] D. Pérez-García and I. Villanueva, *Multiple summing operators on Banach spaces*, J. Math. Anal. Appl. **285** (2003), 86–96.
- [24] D. Pérez-García and I. Villanueva, *Multiple summing operators on $C(K)$ spaces*, Ark. Math. **42** (2003), 153–171.
- [25] T. Praciano-Pereira, *On bounded multilinear forms on a class of ℓ_p spaces*. J. Math. Anal. Appl. **81** (2): 561–568, 1981.
- [26] U. C. Schwarting, *Vector Valued Bohnenblust–Hille Inequalities*, Thesis, (2013).

DEPARTAMENTO DE MATEMÁTICA,
UNIVERSIDADE FEDERAL DA PARAÍBA,
58.051-900 - JOÃO PESSOA, BRAZIL.
E-mail address: `ngalbqrrq@gmail.com`

LABORATOIRE DE MATHÉMATIQUES,
UNIVERSITÉ BLAISE PASCAL CAMPUS DES CÉZEAUX,
F-63177 AUBIERE CEDEX, FRANCE.
E-mail address: `Frederic.Bayart@math.univ-bpclermont.fr`

DEPARTAMENTO DE MATEMÁTICA,
UNIVERSIDADE FEDERAL DA PARAÍBA,
58.051-900 - JOÃO PESSOA, BRAZIL.
E-mail address: `dmpellegrino@gmail.com` and `pellegrino@pq.cnpq.br`

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO,
FACULTAD DE CIENCIAS MATEMÁTICAS,
PLAZA DE CIENCIAS 3,
UNIVERSIDAD COMPLUTENSE DE MADRID,
MADRID, 28040, SPAIN.

AND

INSTITUTO DE CIENCIAS MATEMÁTICAS – ICMAT
CALLE NICOLÁS CABRERA 13–15,
MADRID, 28049, SPAIN.
E-mail address: `jseoane@mat.ucm.es`