

Research Article

Optimization of the Two Fishermen's Profits Exploiting Three Competing Species Where Prices Depend on Harvest

Imane Agmour, Meriem Bentounsi, Naceur Achtaich, and Youssef El Foutayeni

Analysis, Modeling and Simulation Laboratory, Hassan II University, Casablanca, Morocco

Correspondence should be addressed to Youssef El Foutayeni; foutayeni@gmail.com

Received 21 August 2017; Accepted 13 November 2017; Published 26 December 2017

Academic Editor: Abid A. Lashari

Copyright © 2017 Imane Agmour et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Bioeconomic modeling of the exploitation of biological resources such as fisheries has gained importance in recent years. In this work we propose to define and study a bioeconomic equilibrium model for two fishermen who catch three species taking into consideration the fact that the prices of fish populations vary according to the quantity harvested; these species compete with each other for space or food; the natural growth of each species is modeled using a logistic law. The main purpose of this work is to define the fishing effort that maximizes the profit of each fisherman, but all of them have to respect two constraints: the first one is the sustainable management of the resources and the second one is the preservation of the biodiversity. The existence of the steady states and their stability are studied using eigenvalue analysis. The problem of determining the equilibrium point that maximizes the profit of each fisherman leads to Nash equilibrium problem; to solve this problem we transform it into a linear complementarity problem (LCP); then we prove that the obtained problem (LCP) admits a unique solution that represents the Nash equilibrium point of our problem. We close our paper with some numerical simulations.

1. Introduction

Overfishing leads to resource destruction, that is why there is an increasing need for the bioeconomic modeling tool that evaluates the biological and economic effects of different harvesting strategies directed at extracting the long-term maximum sustainable production while avoiding the risk of recruitment overfishing. The techniques and issues associated with the bioeconomic modeling for the exploitation of marine resources have been discussed in detail by Clark and Munro [1, 2]. Clark and Munro [1] demonstrated that, with the aid of optimal control theory, fisheries economics can without difficulty be cast in a capital-theoretic framework yielding results that are both general and readily comprehensible. Chaudhuri [3] discussed the problem of combined harvesting of two competing fish species, each of which obeys the law of logistic growth; it is shown that the open-access fishery may possess a bioeconomic equilibrium which drives one species to extinction. In this context, Chaudhuri [4] considered the problem of dynamic optimization of the exploitation policy connected with the combined harvesting of two competing fish species, each of which obeys the

logistic growth law. Models on the combined harvesting of a two-species prey-predator fishery have been discussed by Chaudhuri and Ray [5]. Kar and Chaudhuri [6] studied the problem of harvesting two competing species in the presence of a predator species which feeds on both the competing species; a combined harvesting effort is devoted to the exploitation of the first two (prey) species while the third (predator) species is not harvested. Mchich et al. [7] proposed a specific stock-effort dynamic model; the stock corresponds to two fish populations growing and moving between two fishing zones, on which they are harvested by two different fleets; the effort represents the number of fishing vessels of the two fleets which operate on the two fishing zones; the bioeconomic model is a set of four ordinary differential equations governing the stocks and the fishing efforts in the two fishing areas; fish migration, as well as vessels displacements, between the two zones is assumed to take place at a faster time scale than the variation of the stocks and the changes of fleets sizes, respectively; the vessels movements between the two fishing areas are assumed to be stock dependent, that is, the larger the stock density is in a zone, the more the vessels tend to remain in it.

Many mathematical models have been developed to describe the dynamics of fisheries; we can refer, for example, to El Foutayeni et al. [8] who in their work have built a bioeconomic equilibrium model for several fishermen who catch two fish species; in this work, the authors have showed that the problem of determining the equilibrium point that maximizes the profit of each fisherman is solved by using linear complementarity problem. El Foutayeni et al. [9] have also defined a bioeconomic equilibrium model for “ n ” fishermen who catch three species; these species compete with each other for space or food; the natural growth of each species is modeled using a logistic law; the objective of their work is to calculate the fishing effort that maximizes the profit of each fisherman at biological equilibrium by using the generalized Nash equilibrium problem.

Most bioeconomic models do not take into account the variational of the price of fish population. Usually, the existing models consider that the prices of the fish populations are constants. In this context, El Foutayeni and Khaladi [10, 11] have presented a bioeconomic model of fish populations taking into consideration the fact that the prices of fish populations vary according to the quantity harvested. But in these articles they assumed the existence of a single fisherman.

This paper is situated in this general context; in this work we present a bioeconomic model for three species which compete with each other for space or food and each of which obeys the law logistic growth. These species are caught by two fishermen. We will assume that the price of the fish population increases with decreasing harvest and the price of the fish population decreases with the increase of the harvest, but the minimum price is equal to a fixed positive constant. The aim of this paper consists in determining the fishing effort strategy adopted by each fisherman to maximize its income under two assumptions; the first one is the sustainable management of the resources, and the second one is the preservation of the biodiversity.

The paper is structured as follows. In Section 2, we give a description of the biological model of fish populations; we will define the mathematical model and study the stability of the equilibrium of our system. In Section 3, we give the bioeconomic model of the fish populations taking into consideration the fact that the prices of fish populations vary according to the quantity harvested; in this section we prove that the resolution of bioeconomic equilibrium model of the three fish populations is equivalent to solving a Nash equilibrium problem and then we show that the latter problem is equivalent to a linear complementarity problem, then we prove that the obtained problem (LCP) admits a unique solution that represents the Nash equilibrium of our problem. Some numerical simulations are given in Section 4 to illustrate the results. Finally, in Section 5 we give a conclusion.

2. The Biological Model of Fish Populations

The aim of this section is to define a biological model of three marine species that compete with each other for space or food and whose natural growth of each is obtained by means of a

logistic law. We study the existence of the steady states and their stability using eigenvalue analysis and Routh-Hurwitz stability criterion.

2.1. The Mathematical Model and Hypotheses. The evolution of the biomass of the first species is given by the following mathematical equation:

$$\begin{aligned} \dot{x}_1(t) = & r_1 x_1(t) \left(1 - \frac{x_1(t)}{K_1} \right) - c_{12} x_1(t) x_2(t) \\ & - c_{13} x_1(t) x_3(t), \end{aligned} \quad (1)$$

where $x_1(t)$ is the biomass of population 1; r_1 is the intrinsic growth rate of species 1; K_1 is the carrying capacity for species 1; c_{12} is the coefficient of competition between species 2 and species 1; and c_{13} is the coefficient of competition between species 3 and species 1.

The evolution of the biomass of the second population is given by the following mathematical equation:

$$\begin{aligned} \dot{x}_2(t) = & r_2 x_2(t) \left(1 - \frac{x_2(t)}{K_2} \right) - c_{21} x_1(t) x_2(t) \\ & - c_{23} x_2(t) x_3(t), \end{aligned} \quad (2)$$

where $x_2(t)$ is the biomass of population 2; r_2 is the intrinsic growth rate of species 2; K_2 is the carrying capacity for species 2; c_{21} is the coefficient of competition between species 1 and species 2; and c_{23} is the coefficient of competition between species 3 and species 2.

The evolution of the biomass of the third species is given by the following mathematical equation:

$$\begin{aligned} \dot{x}_3(t) = & r_3 x_3(t) \left(1 - \frac{x_3(t)}{K_3} \right) - c_{31} x_1(t) x_3(t) \\ & - c_{32} x_2(t) x_3(t), \end{aligned} \quad (3)$$

where $x_3(t)$ is the biomass of population 3; r_3 is the intrinsic growth rate 3; K_3 is the carrying capacity for the species of species 3; c_{31} is the coefficient of competition between species 1 and species 3; and c_{32} is the coefficient of competition between species 2 and species 3.

It is interesting to note that to assure the existence of the three species and their stability we should assume that

$$r_i > c_{ij} K_j, \quad \forall i, j = 1, 2, 3, \text{ with } i \neq j. \quad (4)$$

The evolution of the biomass of fish populations is modeled by the following equations:

$$\begin{aligned} \dot{x}_1(t) = & r_1 x_1(t) \left(1 - \frac{x_1(t)}{K_1} \right) - c_{12} x_1(t) x_2(t) \\ & - c_{13} x_1(t) x_3(t), \\ \dot{x}_2(t) = & r_2 x_2(t) \left(1 - \frac{x_2(t)}{K_2} \right) - c_{21} x_1(t) x_2(t) \\ & - c_{23} x_2(t) x_3(t), \\ \dot{x}_3(t) = & r_3 x_3(t) \left(1 - \frac{x_3(t)}{K_3} \right) - c_{31} x_1(t) x_3(t) \\ & - c_{32} x_2(t) x_3(t). \end{aligned} \quad (5)$$

Let $x(t) = (x_1(t), x_2(t), x_3(t))$ be the solution of system (5). Then all the solutions of the system (5) are nonnegative. To demonstrate that, we must recall that by [12] the system of equation

$$\dot{x} = f(x_1, x_2, \dots, x_n) \quad \text{with } x(t = 0) = x_0 \quad (6)$$

is a positive system if and only if

$$\begin{aligned} \dot{x}_i = f_i(x_1 \geq 0, \dots, x_i = 0, \dots, x_n \geq 0) &\geq 0; \\ \forall i \in [1 \dots n]. \end{aligned} \quad (7)$$

In our case, for $x_1 = 0, x_2, x_3 \geq 0$, we have $dx_1/dt = 0 \geq 0$. By the same, for $x_2 = 0, x_1, x_3 \geq 0$, we have $dx_2/dt = 0 \geq 0$. Also for $x_3 = 0, x_1, x_2 \geq 0$, we have $dx_3/dt = 0 \geq 0$. Therefore, all the solutions of system (5) are nonnegative.

Theorem 1. All the solutions of system (5) which start in \mathbb{R}_+^3 are uniformly bounded.

Proof. We define the function

$$W = x_1 + x_2 + x_3. \quad (8)$$

Therefore, the time derivative along a solution of (5) is

$$\begin{aligned} \frac{dW}{dt} &= r_1 x_1 \left(1 - \frac{x_1}{K_1}\right) + r_2 x_2 \left(1 - \frac{x_2}{K_2}\right) \\ &+ r_3 x_3 \left(1 - \frac{x_3}{K_3}\right) - c_{12} x_1 x_2 - c_{13} x_1 x_3 \\ &- c_{21} x_1 x_2 - c_{23} x_2 x_3 - c_{31} x_1 x_3 - c_{32} x_2 x_3. \end{aligned} \quad (9)$$

For each $\vartheta > 0$, we have

$$\begin{aligned} \frac{dW}{dt} + \vartheta W &= r_1 x_1 \left(1 - \frac{x_1}{K_1}\right) + r_2 x_2 \left(1 - \frac{x_2}{K_2}\right) \\ &+ r_3 x_3 \left(1 - \frac{x_3}{K_3}\right) - c_{12} x_1 x_2 - c_{13} x_1 x_3 \\ &- c_{21} x_1 x_2 - c_{23} x_2 x_3 - c_{31} x_1 x_3 \\ &- c_{32} x_2 x_3 + \vartheta x_1 + \vartheta x_2 + \vartheta x_3 \\ &\leq r_1 x_1 \left(1 - \frac{x_1}{K_1}\right) + r_2 x_2 \left(1 - \frac{x_2}{K_2}\right) \\ &+ r_3 x_3 \left(1 - \frac{x_3}{K_3}\right) + \vartheta x_1 + \vartheta x_2 + \vartheta x_3 \\ &= x_1 \left[r_1 \left(1 - \frac{x_1}{K_1}\right) + \vartheta \right] \\ &+ x_2 \left[r_2 \left(1 - \frac{x_2}{K_2}\right) + \vartheta \right] \\ &+ x_3 \left[r_3 \left(1 - \frac{x_3}{K_3}\right) + \vartheta \right] \end{aligned}$$

$$\begin{aligned} &= x_1 (r_1 + \vartheta) - \frac{r_1}{K_1} x_1^2 + x_2 (r_2 + \vartheta) \\ &- \frac{r_2}{K_2} x_2^2 + x_3 (r_3 + \vartheta) - \frac{r_3}{K_3} x_3^2. \end{aligned} \quad (10)$$

We can easily show that

$$\begin{aligned} -\frac{r_1}{K_1} x_1^2 + x_1 (r_1 + \vartheta) - \frac{K_1}{4r_1} (r_1 + \vartheta)^2 &\leq 0, \\ -\frac{r_2}{K_2} x_2^2 + x_2 (r_2 + \vartheta) - \frac{K_2}{4r_2} (r_2 + \vartheta)^2 &\leq 0, \\ -\frac{r_3}{K_3} x_3^2 + x_3 (r_3 + \vartheta) - \frac{K_3}{4r_3} (r_3 + \vartheta)^2 &\leq 0. \end{aligned} \quad (11)$$

Then

$$\begin{aligned} -\frac{r_1}{K_1} x_1^2 + x_1 (r_1 + \vartheta) &\leq \frac{K_1}{4r_1} (r_1 + \vartheta)^2, \\ -\frac{r_2}{K_2} x_2^2 + x_2 (r_2 + \vartheta) &\leq \frac{K_2}{4r_2} (r_2 + \vartheta)^2, \\ -\frac{r_3}{K_3} x_3^2 + x_3 (r_3 + \vartheta) &\leq \frac{K_3}{4r_3} (r_3 + \vartheta)^2. \end{aligned} \quad (12)$$

Therefore, we can deduce that

$$\begin{aligned} \frac{dW}{dt} + \vartheta W &\leq \frac{K_1}{4r_1} (r_1 + \vartheta)^2 + \frac{K_2}{4r_2} (r_2 + \vartheta)^2 \\ &+ \frac{K_3}{4r_3} (r_3 + \vartheta)^2. \end{aligned} \quad (13)$$

So the right-hand side is positive; therefore it is bounded for all $(x_1, x_2, x_3) \in \mathbb{R}_+^3$. Therefore we find a $\theta > 0$ with $dW/dt + \vartheta W < \theta$. Using the theory of differential inequality [13], we obtain

$$\begin{aligned} 0 &\leq W(x_1, x_2, x_3) \\ &\leq \frac{\theta}{\vartheta} + \left[W(x_1(0), x_2(0), x_3(0)) - \frac{\theta}{\vartheta} \right] e^{-\vartheta t} \end{aligned} \quad (14)$$

which, upon letting $t \rightarrow \infty$, yields $0 \leq W \leq \theta/\vartheta$.

Then, we have

$$B = \left\{ (x_1, x_2, x_3) \in \mathbb{R}_+^3 : W < \frac{\theta}{\vartheta} + \varepsilon, \text{ for any } \varepsilon > 0 \right\}, \quad (15)$$

where B is the region in which all the solutions of system of (5) that start in \mathbb{R}_+^3 are confined. \square

2.2. The Steady States of the System. The steady states of the system of (5) are obtained by solving the system of equations

$$\begin{aligned}
 r_1 x_1 \left(1 - \frac{x_1}{K_1} \right) - c_{12} x_1 x_2 - c_{13} x_1 x_3 &= 0, \\
 r_2 x_2 \left(1 - \frac{x_2}{K_2} \right) - c_{21} x_1 x_2 - c_{23} x_2 x_3 &= 0, \\
 r_3 x_3 \left(1 - \frac{x_3}{K_3} \right) - c_{31} x_1 x_3 - c_{32} x_2 x_3 &= 0.
 \end{aligned}
 \tag{16}$$

This system of equations has eight solutions

$$P_1(0, 0, 0), P_2(K_1, 0, 0), P_3(0, K_2, 0), P_4(0, 0, K_3), P_5(x_1^{(5)}, x_2^{(5)}, 0), \text{ where}$$

$$\begin{aligned}
 x_1^{(5)} &= K_1 r_2 \frac{r_1 - c_{12} K_2}{r_1 r_2 - c_{12} c_{21} K_2 K_1}, \\
 x_2^{(5)} &= K_2 r_1 \frac{r_2 - c_{21} K_1}{r_1 r_2 - c_{12} c_{21} K_2 K_1},
 \end{aligned}
 \tag{17}$$

$P_6(x_1^{(6)}, 0, x_3^{(6)})$, where

$$\begin{aligned}
 x_1^{(6)} &= K_1 r_3 \frac{r_1 - c_{13} K_3}{r_1 r_3 - c_{13} c_{31} K_3 K_1}, \\
 x_3^{(6)} &= K_3 r_1 \frac{r_3 - c_{31} K_1}{r_1 r_3 - c_{13} c_{31} K_3 K_1},
 \end{aligned}
 \tag{18}$$

$P_7(0, x_2^{(7)}, x_3^{(7)})$, where

$$\begin{aligned}
 x_2^{(7)} &= K_2 r_3 \frac{r_2 - c_{23} K_3}{r_3 r_2 - c_{32} c_{23} K_2 K_3}, \\
 x_3^{(7)} &= K_3 r_2 \frac{r_3 - c_{32} K_2}{r_3 r_2 - c_{32} c_{23} K_2 K_3},
 \end{aligned}
 \tag{19}$$

and $P_8(x_1^*, x_2^*, x_3^*)$, where

$$\begin{aligned}
 x_1^* &= \frac{K_1 (r_1 r_2 r_3 - r_1 c_{23} c_{32} K_2 K_3 + r_3 c_{12} c_{23} K_2 K_3 - r_2 r_3 c_{12} K_2 - r_2 r_3 c_{13} K_3 + r_2 c_{13} c_{32} K_2 K_3)}{\Delta}, \\
 x_2^* &= \frac{K_2 (r_1 r_2 r_3 - r_2 c_{13} c_{31} K_1 K_3 + r_1 c_{23} c_{31} K_1 K_3 - r_1 r_3 c_{21} K_1 - r_1 r_3 c_{23} K_3 + r_3 c_{13} c_{21} K_1 K_3)}{\Delta}, \\
 x_3^* &= \frac{K_3 (r_1 r_2 r_3 - r_3 c_{12} c_{21} K_1 K_2 - r_1 r_2 c_{31} K_1 + r_1 c_{21} c_{32} K_1 K_2 + r_2 c_{12} c_{31} K_1 K_2 - r_1 r_2 c_{32} K_2)}{\Delta}, \\
 \Delta &= r_1 r_2 r_3 - r_1 c_{23} c_{32} K_2 K_3 - r_2 c_{13} c_{31} K_1 K_3 - r_3 c_{12} c_{21} K_1 K_2 + c_{12} c_{23} c_{31} K_1 K_2 K_3 + c_{13} c_{21} c_{32} K_1 K_2 K_3.
 \end{aligned}
 \tag{20}$$

The system of (16) has several solutions, but only one of them can give the coexistence of the biomass of the three species; this solution is the point $P_8(x_1^*, x_2^*, x_3^*)$.

2.3. *The Stability of the Steady States.* The variational matrix of system (5) is

$$J = \begin{bmatrix} J_{11} & -c_{12} x_1 & -c_{13} x_1 \\ -c_{21} x_2 & J_{22} & -c_{23} x_2 \\ -c_{31} x_3 & -c_{32} x_3 & J_{33} \end{bmatrix}, \tag{21}$$

where

$$\begin{aligned}
 J_{11} &= r_1 \left(1 - \frac{2}{K_1} x_1 \right) - c_{12} x_2 - c_{13} x_3, \\
 J_{22} &= r_2 \left(1 - \frac{2}{K_2} x_2 \right) - c_{21} x_1 - c_{23} x_3, \\
 J_{33} &= r_3 \left(1 - \frac{2}{K_3} x_3 \right) - c_{31} x_1 - c_{32} x_2.
 \end{aligned}
 \tag{22}$$

Proposition 2. *The point $P_1(0, 0, 0)$ is unstable.*

Proof. The variational matrix of system (5) at the steady state $P_1(0, 0, 0)$ is

$$J_1 = \begin{bmatrix} r_1 & 0 & 0 \\ 0 & r_2 & 0 \\ 0 & 0 & r_3 \end{bmatrix}. \tag{23}$$

The eigenvalues of J_1 are

$$\begin{aligned}
 \lambda_1 &= r_1 > 0, \\
 \lambda_2 &= r_2 > 0, \\
 \lambda_3 &= r_3 > 0,
 \end{aligned}
 \tag{24}$$

then, the point $P_1(0, 0, 0)$ is unstable. \square

Let $r_1 = 2, r_2 = 1, r_3 = 3, c_{12} = 0.009, c_{21} = 0.007, c_{13} = 0.008, c_{23} = 0.001, c_{31} = 0.002, c_{32} = 0.001, K_1 = 70, K_2 = 50, K_3 = 40$ in appropriate units. Figure 1 shows the dynamical behaviors and phase space trajectory of the three marine species against time, beginning with the initial values $x(0) = 0.01, y(0) = 0.01, z(0) = 0.01$. By Figure 1 we find that the steady state point P_1 is unstable, and more precisely this point tends to the point P_8 .

Proposition 3. *The point $P_2(K_1, 0, 0)$ is unstable if the conditions of existence given by (4) hold; if not, it is stable.*




FIGURE 1: Dynamical behaviors and phase space trajectories of the three marine species.

Proof. The variational matrix of system (5) at the steady state $P_2(K_1, 0, 0)$ is

$$J_2 = \begin{bmatrix} -r_1 & -c_{12}K_1 & -c_{13}K_1 \\ 0 & r_2 - c_{21}K_1 & 0 \\ 0 & 0 & r_3 - c_{31}K_1 \end{bmatrix}. \tag{25}$$

The eigenvalues of J_2 are

$$\begin{aligned} \lambda_1 &= -r_1 < 0, \\ \lambda_2 &= r_2 - c_{21}K_1, \\ \lambda_3 &= r_3 - c_{31}K_1 \end{aligned} \tag{26}$$

if

$$\begin{aligned} r_2 &> c_{21}K_1 \\ r_3 &> c_{31}K_1, \end{aligned} \tag{27}$$

then, the point $P_2(K_1, 0, 0)$ is unstable; if not, it is stable. \square

Let $r_1 = 2, r_2 = 1, r_3 = 3, c_{12} = 0.009, c_{21} = 0.007, c_{13} = 0.008, c_{23} = 0.001, c_{31} = 0.002, c_{32} = 0.001, K_1 = 70, K_2 = 50, K_3 = 40$ in appropriate units. Figure 2 shows the dynamical behaviors and phase space trajectory of the three marine species against time, beginning with the initial values $x(0) =$

$70, y(0) = 0.01, z(0) = 0.01$. By Figure 2 we can see that the steady state point P_2 is unstable, and more precisely this point tends to the point P_8 too.

Proposition 4. *The point $P_3(0, K_2, 0)$ is unstable if the conditions of existence given by (4) hold; if not, it is stable.*

Proof. The variational matrix of system (5) at the steady state $P_3(0, K_2, 0)$ is

$$J_3 = \begin{bmatrix} r_1 - c_{12}K_2 & 0 & 0 \\ -c_{21}K_2 & -r_2 & -c_{23}K_2 \\ 0 & 0 & r_3 - c_{32}K_2 \end{bmatrix}. \tag{28}$$

The eigenvalues of J_3 are

$$\begin{aligned} \lambda_1 &= r_1 - c_{12}K_2, \\ \lambda_2 &= -r_2 < 0, \\ \lambda_3 &= r_3 - c_{32}K_2 \end{aligned} \tag{29}$$

if

$$\begin{aligned} r_1 &> c_{12}K_2, \\ r_3 &> c_{32}K_2; \end{aligned} \tag{30}$$

therefore, the point $P_3(0, K_2, 0)$ is unstable; if not, it is stable. \square




FIGURE 2: Dynamical behaviors and phase space trajectories of the three marine species.

Let $r_1 = 2, r_2 = 1, r_3 = 3, c_{12} = 0.009, c_{21} = 0.007, c_{13} = 0.008, c_{23} = 0.001, c_{31} = 0.002, c_{32} = 0.001, K_1 = 70, K_2 = 50, K_3 = 40$ in appropriate units. Figure 3 shows the dynamical behaviors and phase space trajectory of the three marine species against time, beginning with the initial values $x(0) = 0.01, y(0) = 50, z(0) = 0.01$. By Figure 3 we can see that the steady state point P_3 is also unstable and tends to the point P_8 .

Proposition 5. *The point $P_4(0, 0, K_3)$ is unstable if the conditions of existence given by (4) hold; if not, it is stable.*

Proof. The variational matrix of system (5) at the steady state $P_4(0, 0, K_3)$ is

$$J_4 = \begin{bmatrix} r_1 - c_{13}K_3 & 0 & 0 \\ 0 & r_2 - c_{23}K_3 & 0 \\ -c_{13}K_3 & -c_{32}K_3 & -r_3 \end{bmatrix}. \tag{31}$$

The eigenvalues of J_4 are

$$\begin{aligned} \lambda_1 &= r_1 - c_{13}K_3, \\ \lambda_2 &= r_2 - c_{23}K_3, \\ \lambda_3 &= -r_3 < 0 \end{aligned} \tag{32}$$

if

$$\begin{aligned} r_1 &> c_{13}K_3, \\ r_2 &> c_{23}K_3, \end{aligned} \tag{33}$$

then, the point $P_4(0, 0, K_3)$ is unstable; if not, it is stable. \square

Let $r_1 = 2, r_2 = 1, r_3 = 3, c_{12} = 0.009, c_{21} = 0.007, c_{13} = 0.008, c_{23} = 0.001, c_{31} = 0.002, c_{32} = 0.001, K_1 = 70, K_2 = 50, K_3 = 40$ in appropriate units. Figure 4 indicates the dynamical behaviors and phase space trajectory of the three marine species against time, beginning with the initial values $x(0) = 0.01, y(0) = 0.01, z(0) = 40$. Following Figure 4 we can see that the steady state point P_3 is unstable and also tends to the point P_8 .

Proposition 6. *The point $P_5(x_1^{(5)}, x_2^{(5)}, 0)$ is unstable.*

Proof. The variational matrix of system (5) at the steady state $P_5(x_1^{(5)}, x_2^{(5)}, 0)$ is

$$J_5 = \begin{bmatrix} -\frac{r_1}{K_1}x_1^{(5)} & -c_{12}x_1^{(5)} & -c_{13}x_1^{(5)} \\ -c_{21}x_2^{(5)} & -\frac{r_2}{K_2}x_2^{(5)} & -c_{23}x_2^{(5)} \\ 0 & 0 & r_3 - c_{31}x_1^{(5)} - c_{32}x_2^{(5)} \end{bmatrix}. \tag{34}$$




FIGURE 3: Dynamical behaviors and phase space trajectories of the three marine species.

The eigenvalues of J_5 are

$$\begin{aligned} \lambda_1 &= -\frac{1}{2K_1K_2} (M - \sqrt{N}), \\ \lambda_2 &= -\frac{1}{2K_1K_2} (M + \sqrt{N}), \\ \lambda_3 &= r_3 - c_{31}x_1^{(5)} - c_{32}x_2^{(5)}, \end{aligned} \tag{35}$$

where

$$\begin{aligned} M &= r_1x_1^{(5)}K_2 + K_1r_2x_2^{(5)}, \\ N &= [r_1x_1^{(5)}K_2 - K_1r_2x_2^{(5)}]^2 + 4K_1^2K_2^2c_{21}x_2^{(5)}c_{12}x_1^{(5)}. \end{aligned} \tag{36}$$

If

$$\begin{aligned} r_1 &> c_{12}K_2, \\ r_2 &> c_{21}K_1 \end{aligned} \tag{37}$$

then, $\lambda_3 > 0$; if not, then $\lambda_1 > 0$. Therefore, the point $P_5(x_1^{(5)}, x_2^{(5)}, 0)$ is unstable in all cases. \square

Let $r_1 = 2, r_2 = 1, r_3 = 3, c_{12} = 0.009, c_{21} = 0.007, c_{13} = 0.008, c_{23} = 0.001, c_{31} = 0.002, c_{32} = 0.001, K_1 = 70, K_2 = 50, K_3 = 40$ in appropriate units. Figure 5 represents the

dynamical behaviors and phase space trajectory of the three marine species against time, beginning with the initial values $x(0) = 60, y(0) = 28, z(0) = 0.01$. Following Figure 5 we can deduce that the steady state point P_5 is unstable and also tends to the point P_8 .

Proposition 7. *The point $P_6(x_1^{(6)}, 0, x_3^{(6)})$ is unstable.*

Proof. The variational matrix of system (5) at the steady state $P_6(x_1^{(6)}, 0, x_3^{(6)})$ is

$$J_6 = \begin{bmatrix} -\frac{r_1}{k_1}x_1^{(6)} & -c_{12}x_1^{(6)} & -c_{13}x_1^{(6)} \\ 0 & r_2 - c_{21}x_1^{(6)} - c_{23}x_3^{(6)} & 0 \\ -c_{31}x_3^{(6)} & -c_{32}x_3^{(6)} & -\frac{r_3}{K_3}x_3^{(6)} \end{bmatrix}. \tag{38}$$

The eigenvalues of J_6 are

$$\begin{aligned} \lambda_1 &= -\frac{1}{2K_1K_3} (G - \sqrt{L}), \\ \lambda_2 &= r_2 - c_{21}x_1^{(6)} - c_{23}x_3^{(6)}, \\ \lambda_3 &= -\frac{1}{2K_1K_3} (G + \sqrt{L}), \end{aligned} \tag{39}$$




FIGURE 4: Dynamical behaviors and phase space trajectories of the three marine species.

where

$$G = r_1 x_1^{(6)} K_3 + K_1 r_3 x_3^{(6)}, \tag{40}$$

$$L = [r_1 x_1^{(6)} K_3 - K_1 r_3 x_3^{(6)}]^2 + 4K_1^2 K_3^2 c_{31} x_3^{(6)} c_{13} x_1^{(6)}.$$

If

$$r_1 > c_{13} K_3, \tag{41}$$

$$r_3 > c_{31} K_1$$

then, $\lambda_2 > 0$; if not, then $\lambda_1 > 0$; therefore, point $P_6(x_1^{(6)}, 0, x_3^{(6)})$ is unstable. \square

Let $r_1 = 2, r_2 = 1, r_3 = 3, c_{12} = 0.009, c_{21} = 0.007, c_{13} = 0.008, c_{23} = 0.001, c_{31} = 0.002, c_{32} = 0.001, K_1 = 70, K_2 = 50, K_3 = 40$ in appropriate units. Figure 6 indicates the dynamical behaviors and phase space trajectory of the three marine species against time, beginning with the initial values $x(0) = 59, y(0) = 0.01, z(0) = 38$. Following Figure 6 we can deduce that the steady state point P_6 is unstable and also tends to the point P_8 .

Proposition 8. *The point $P_7(0, x_2^{(7)}, x_3^{(7)})$ is unstable.*

Proof. The variational matrix of system (5) at the steady state $P_7(0, x_2^{(7)}, x_3^{(7)})$ is

$$J_7 = \begin{bmatrix} r_1 - c_{12}x_2^{(7)} - c_{13}x_3^{(7)} & 0 & 0 \\ -c_{21}x_2^{(7)} & -\frac{r_2}{K_2}x_2^{(7)} & -c_{23}x_2^{(7)} \\ -c_{31}x_3^{(7)} & -c_{32}x_3^{(7)} & -\frac{r_3}{K_3}x_3^{(7)} \end{bmatrix}. \tag{42}$$

The eigenvalues of J_7 are

$$\begin{aligned} \lambda_1 &= r_1 - c_{12}x_2^{(7)} - c_{13}x_3^{(7)}, \\ \lambda_2 &= -\frac{1}{2K_2K_3} (R - \sqrt{S}), \\ \lambda_3 &= -\frac{1}{2K_2K_3} (R + \sqrt{S}), \end{aligned} \tag{43}$$

where

$$R = r_2 x_2^{(7)} K_3 + K_2 r_3 x_3^{(7)}, \tag{44}$$

$$S = [r_2 x_2^{(7)} K_3 + K_2 r_3 x_3^{(7)}]^2 + 4K_2^2 K_3^2 c_{23} x_2^{(7)} c_{32} x_3^{(7)}.$$

If

$$r_2 > c_{23} K_3, \tag{45}$$

$$r_3 > c_{32} K_2$$




FIGURE 5: Dynamical behaviors and phase space trajectories of the three marine species.

then, $\lambda_1 > 0$; if not, then $\lambda_2 > 0$. Therefore, point $P_7(0, x_2^{(7)}, x_3^{(7)})$ is unstable. \square

Let $r_1 = 2, r_2 = 1, r_3 = 3, c_{12} = 0.009, c_{21} = 0.007, c_{13} = 0.008, c_{23} = 0.001, c_{31} = 0.002, c_{32} = 0.001, K_1 = 70, K_2 = 50, K_3 = 40$ in appropriate units. Figure 7 shows the dynamical behaviors and phase space trajectory of the three marine species against time, beginning with the initial values $x(0) = 0.01, y(0) = 48, z(0) = 39$. By Figure 7 we can conclude that the steady state point P_7 is unstable and also tends to point P_8 .

Theorem 9. *The point $P_8(x_1^*, x_2^*, x_3^*)$ is locally asymptotically stable.*

Proof. We proof this theorem by using Routh-Hurwitz stability criterion.

The variational matrix of system (5) in the steady state $P_8(x_1^*, x_2^*, x_3^*)$ is

$$J_8 = \begin{bmatrix} J_{11} & -c_{12}x_1^* & -c_{13}x_1^* \\ -c_{21}x_2^* & J_{22} & -c_{23}x_2^* \\ -c_{31}x_3^* & -c_{32}x_3^* & J_{33} \end{bmatrix}, \quad (46)$$

where

$$J_{11} = r_1 \left(1 - \frac{2}{K_1} x_1^* \right) - c_{12}x_2^* - c_{13}x_3^*,$$

$$J_{22} = r_2 \left(1 - \frac{2}{K_2} x_2^* \right) - c_{21}x_1^* - c_{23}x_3^*,$$

$$J_{33} = r_3 \left(1 - \frac{2}{K_3} x_3^* \right) - c_{31}x_1^* - c_{32}x_2^*.$$

(47)

Using the fact that by (16) we have

$$r_1 \left(1 - \frac{2}{K_1} x_1^* \right) - c_{12}x_2^* - c_{13}x_3^* = -\frac{r_1}{K_1} x_1^*,$$

$$r_2 \left(1 - \frac{2}{K_2} x_2^* \right) - c_{21}x_1^* - c_{23}x_3^* = -\frac{r_2}{K_2} x_2^*, \quad (48)$$

$$r_3 \left(1 - \frac{2}{K_3} x_3^* \right) - c_{31}x_1^* - c_{32}x_2^* = -\frac{r_3}{K_3} x_3^*$$

then

$$J_8 = \begin{bmatrix} -\frac{r_1}{K_1} x_1^* & -c_{12}x_1^* & -c_{13}x_1^* \\ -c_{21}x_2^* & -\frac{r_2}{K_2} x_2^* & -c_{23}x_2^* \\ -c_{31}x_3^* & -c_{32}x_3^* & -\frac{r_3}{K_3} x_3^* \end{bmatrix}. \quad (49)$$

The characteristic polynomial of the variational matrix is

$$P(\lambda) = a_0\lambda^3 + a_1\lambda^2 + a_2\lambda + a_3, \quad (50)$$




FIGURE 6: Dynamical behaviors and phase space trajectories of the three marine species.

where

$$\begin{aligned}
 a_0 &= 1, \\
 a_1 &= \frac{r_1}{K_1}x_1^* + \frac{r_2}{K_2}x_2^* + \frac{r_3}{K_3}x_3^*, \\
 a_2 &= \frac{r_1}{K_1}x_1^* \frac{r_2}{K_2}x_2^* + \frac{r_1}{K_1}x_1^* \frac{r_3}{K_3}x_3^* + \frac{r_2}{K_2}x_2^* \frac{r_3}{K_3}x_3^* \\
 &\quad - c_{23}x_2^*c_{32}x_3^* - c_{12}x_1^*c_{21}x_2^* - c_{13}x_1^*c_{31}x_3^*, \\
 a_3 &= \frac{r_1}{K_1}x_1^* \frac{r_2}{K_2}x_2^* \frac{r_3}{K_3}x_3^* + c_{12}x_1^*c_{23}x_2^*c_{31}x_3^* \\
 &\quad + c_{13}x_3^*c_{32}x_2^*c_{21}x_1^* - c_{12}x_1^*c_{21}x_2^* \frac{r_3}{K_3}x_3^* \\
 &\quad - c_{23}x_2^*c_{32}x_3^* \frac{r_1}{K_1}x_1^* - c_{13}x_3^*c_{31}x_1^* \frac{r_2}{K_2}x_2^*;
 \end{aligned} \tag{51}$$

we have $a_i > 0, \forall i = 0, 1, 2, 3$. In fact,

- (i) $a_0 = 1 > 0$,
- (ii) $a_1 = (r_1/K_1)x_1^* + (r_2/K_2)x_2^* + (r_3/K_3)x_3^* > 0$,
- (iii) using the fact that by (4) we have

$$\begin{aligned}
 r_1r_2 &> c_{12}K_2c_{21}K_1, \\
 r_2r_3 &> c_{23}K_3c_{32}K_2, \\
 r_1r_3 &> c_{13}K_3c_{31}K_1
 \end{aligned} \tag{52}$$

so

$$\begin{aligned}
 a_2 &= \frac{r_1}{K_1}x_1^* \frac{r_2}{K_2}x_2^* + \frac{r_1}{K_1}x_1^* \frac{r_3}{K_3}x_3^* + \frac{r_2}{K_2}x_2^* \frac{r_3}{K_3}x_3^* \\
 &\quad - c_{23}x_2^*c_{32}x_3^* - c_{12}x_1^*c_{21}x_2^* - c_{13}x_1^*c_{31}x_3^* > 0, \\
 a_3 &= \frac{r_1}{K_1}x_1^* \frac{r_2}{K_2}x_2^* \frac{r_3}{K_3}x_3^* + c_{12}x_1^*c_{23}x_2^*c_{31}x_3^* \\
 &\quad + c_{13}x_3^*c_{32}x_2^*c_{21}x_1^* - c_{12}x_1^*c_{21}x_2^* \frac{r_3}{K_3}x_3^* \\
 &\quad - c_{23}x_2^*c_{32}x_3^* \frac{r_1}{K_1}x_1^* - c_{13}x_3^*c_{31}x_1^* \frac{r_2}{K_2}x_2^* > 0,
 \end{aligned} \tag{53}$$

(iv)

$$\begin{aligned}
 a_1a_2 - a_0a_3 &= \left(\frac{r_1}{K_1}x_1^* + \frac{r_2}{K_2}x_2^* + \frac{r_3}{K_3}x_3^* \right) \\
 &\quad \cdot \left(\frac{r_1}{K_1}x_1^* \frac{r_2}{K_2}x_2^* + \frac{r_1}{K_1}x_1^* \frac{r_3}{K_3}x_3^* + \frac{r_2}{K_2}x_2^* \frac{r_3}{K_3}x_3^* \right)
 \end{aligned}$$




FIGURE 7: Dynamical behaviors and phase space trajectories of the three marine species.

$$\begin{aligned}
 & -c_{23}x_2^*c_{32}x_3^* - c_{12}x_1^*c_{21}x_2^* - c_{13}x_1^*c_{31}x_3^* \\
 & - \left(\frac{r_1}{K_1}x_1^* \frac{r_2}{K_2}x_2^* \frac{r_3}{K_3}x_3^* + c_{12}x_1^*c_{23}x_2^*c_{31}x_3^* \right. \\
 & \left. + c_{13}x_3^*c_{32}x_2^*c_{21}x_1^* - c_{12}x_1^*c_{21}x_2^*r_3 \frac{x_3^*}{K_3} \right. \\
 & \left. - c_{23}x_2^*c_{32}x_3^* \frac{r_1}{K_1}x_1^* - c_{13}x_3^*c_{31}x_1^* \frac{r_2}{K_2}x_2^* \right) = \frac{r_1}{K_1}x_1^* \\
 & \cdot \frac{r_2}{K_2}x_2^* \frac{r_3}{K_3}x_3^* - c_{12}x_1^*c_{23}x_2^*c_{31}x_3^* + \frac{r_1}{K_1}x_1^* \frac{r_2}{K_2}x_2^* \\
 & \cdot \frac{r_3}{K_3}x_3^* - c_{13}x_3^*c_{32}x_2^*c_{21}x_1^* + \frac{r_1^2}{K_1^2}x_1^{*2} \frac{r_2}{K_2}x_2^* - \frac{r_1}{K_1} \\
 & \cdot x_1^{*2} c_{12}c_{21}x_2^* + \frac{r_2^2}{K_2^2}x_2^{*2} \frac{r_3}{K_3}x_3^* - \frac{r_2}{K_2}x_2^{*2} c_{32}c_{23}x_3^* \\
 & + \frac{r_1^2}{K_1^2}x_1^{*2} \frac{r_3}{K_3}x_3^* - \frac{r_1}{K_1}x_1^{*2} c_{13}c_{31}x_3^* + \frac{r_3^2}{K_3^2}x_3^{*2} \frac{r_1}{K_1}x_1^* \\
 & - \frac{r_3}{K_3}x_3^{*2} c_{13}c_{31}x_1^* + \frac{r_2^2}{K_2^2}x_2^{*2} \frac{r_1}{K_1}x_1^* - \frac{r_2}{K_2}x_2^{*2} c_{12}c_{21}x_1^* \\
 & + \frac{r_3^2}{K_3^2}x_3^{*2} \frac{r_2}{K_2}x_2^* - \frac{r_3}{K_3}x_3^{*2} c_{23}c_{32}x_2^*.
 \end{aligned} \tag{54}$$

From (4) we deduce that

$$a_1a_2 - a_0a_3 > 0. \tag{55}$$

Then, using the Routh-Hurwitz stability criterion we conclude that the steady state point $P_8(x_1^*, x_2^*, x_3^*)$ is locally asymptotically stable. \square

Let $r_1 = 2, r_2 = 1, r_3 = 3, c_{12} = 0.009, c_{21} = 0.007, c_{13} = 0.008, c_{23} = 0.001, c_{31} = 0.002, c_{32} = 0.001, K_1 = 70, K_2 = 50, K_3 = 40$ in appropriate units. Figure 8 shows the dynamical behaviors and phase space trajectory of the three marine species against time, beginning with the initial values $x(0) = 49, y(0) = 30, z(0) = 38$. By Figure 8 one can see that the steady state point P_8 is locally asymptotically stable.

More precisely, beginning with different initial values we can confirm that the three marine species tend to point P_8 , and according to the phase space trajectories given by




FIGURE 8: Dynamical behaviors and phase space trajectories of the three marine species.

Figures 1–7 we can confirm that the steady state point P_8 is a global attractor.

3. Bioeconomic Model of Fishery

The main purpose of this section is to define and study a bioeconomic equilibrium model for two fishermen who catch three fish populations.

More specifically, this bioeconomic model includes three parts: a biological part connecting the catch to the biomass stock, an exploitation part connecting the catch to the fishing effort, and an economic part connecting the fishing effort to the profit.

So, introducing the fishing by reducing the rate of fish population growth by the amount

$$H_{ij} = q_j E_{ij} x_j, \tag{56}$$

where H_{ij} is the catches of fish population j by the fisherman i , E_{ij} is the fishing effort to exploit a fish population j by the fisherman i , and q_j is the catchability coefficient of fish population j , the model for the evolution of fish populations is given by the following mathematical system of equations:

$$\dot{x}_1 = r_1 x_1 \left(1 - \frac{x_1}{K_1} \right) - c_{12} x_1 x_2 - c_{13} x_1 x_3 - q_1 E_1 x_1,$$

$$\dot{x}_2 = r_2 x_2 \left(1 - \frac{x_2}{K_2} \right) - c_{21} x_1 x_2 - c_{23} x_2 x_3 - q_2 E_2 x_2,$$

$$\dot{x}_3 = r_3 x_3 \left(1 - \frac{x_3}{K_3} \right) - c_{31} x_1 x_3 - c_{32} x_2 x_3 - q_3 E_3 x_3. \tag{57}$$

On one hand, we denote by $H_j = H_{1j} + H_{2j}$ the total catches of species j by all fishermen; on the other hand, we denote by $E_j = E_{1j} + E_{2j}$ the total fishing effort dedicated to species j by all fishermen, and we denote by $E^{(i)} = (E_{i1}, E_{i2}, E_{i3})^T$ the vector fishing effort which must be provided by the fisherman i to catch the three species.

In what follows of this paper, the product of two vectors $\alpha \in \mathbb{R}^3$ and $\beta \in \mathbb{R}^3$ is the vector noted by $\alpha\beta$ or $\beta\alpha$ and is defined by

$$\alpha\beta := (\alpha_1 \beta_1, \alpha_2 \beta_2, \alpha_3 \beta_3)^T \in \mathbb{R}^3. \tag{58}$$

The scalar product is noted by $\alpha^T \beta$. The division of the vector $\alpha \in \mathbb{R}^3$ and the not null vector $\beta \in \mathbb{R}^3$ (i.e., $\beta_i \neq 0, \forall i = 1, 2, 3$) is the vector noted by α/β and is defined by

$$\frac{\alpha}{\beta} := \left(\frac{\alpha_1}{\beta_1}, \frac{\alpha_2}{\beta_2}, \frac{\alpha_3}{\beta_3} \right)^T \in \mathbb{R}^3. \tag{59}$$

The product of the vector $\alpha \in \mathbb{R}^3$ and the matrix $A \in \mathbb{R}^{3 \times 3}$ is noted by αA and is defined by

$$\alpha A := \text{diag}(\alpha) \cdot A \in \mathbb{R}^{3 \times 3}. \tag{60}$$

Now we give the expression of biomass as a function of fishing effort.

The biomasses at biological equilibrium are the solutions of the system

$$\begin{aligned} r_1 \left(1 - \frac{x_1}{K_1}\right) &= c_{12}x_2 + c_{13}x_3 + q_1E_1, \\ r_2 \left(1 - \frac{x_2}{K_2}\right) &= c_{21}x_1 + c_{23}x_3 + q_2E_2, \\ r_3 \left(1 - \frac{x_3}{K_3}\right) &= c_{31}x_1 + c_{32}x_2 + q_3E_3. \end{aligned} \tag{61}$$

The solutions of this system are given by

$$\begin{aligned} x_1 &= a_{11}E_1 + a_{12}E_2 + a_{13}E_3 + x_1^*, \\ x_2 &= a_{21}E_1 + a_{22}E_2 + a_{23}E_3 + x_2^*, \\ x_3 &= a_{31}E_1 + a_{32}E_2 + a_{33}E_3 + x_3^*, \end{aligned} \tag{62}$$

where

$$\begin{aligned} a_{11} &= \frac{K_1(c_{32}K_2K_3c_{23}q_1 - r_3r_2q_1)}{\Delta}, \\ a_{12} &= \frac{K_1(-c_{32}K_2q_2c_{13}K_3 + K_2q_2c_{12}r_3)}{\Delta}, \\ a_{13} &= \frac{K_1(-K_2K_3c_{23}c_{12}q_3 + q_3r_2c_{13}K_3)}{\Delta}, \\ a_{21} &= \frac{K_2(-K_3c_{23}q_1K_1c_{31} + K_1c_{21}r_3q_1)}{\Delta}, \\ a_{22} &= \frac{K_2(q_2c_{13}K_1K_3c_{31} - q_2r_1r_3)}{\Delta}, \\ a_{23} &= \frac{K_2(K_3c_{23}r_1q_3 - K_1c_{21}q_3c_{13}K_3)}{\Delta}, \\ a_{31} &= \frac{K_3(-q_1K_1c_{32}K_2c_{21} + q_1K_1r_2c_{31})}{\Delta}, \\ a_{32} &= \frac{K_3(r_1c_{32}K_2q_2 - c_{12}K_1K_2q_2c_{31})}{\Delta}, \\ a_{33} &= \frac{K_3(c_{12}K_1K_2c_{21}q_3 - r_1r_2q_3)}{\Delta} \end{aligned} \tag{63}$$

or in matrix form $X = -AE + X^*$, where $A = (-a_{ij})_{1 \leq i, j \leq 3}$, $E = (E_1, E_2, E_3)^T$, and $X^* = (x_1^*, x_2^*, x_3^*)^T$.

3.1. Expression of the Total Revenue. It is interesting to note that there exist many different variables that affect the fish price; in this paper, we will consider that the price of the fish population depends on the quantity harvested; specifically we

assumed that the price of the marine species increases with the decreasing harvest and the price of the marine species decreases with the increase of the harvest, but the minimum price is equal to a fixed positive constant. More precisely, the price of marine species j exploited by the fisherman i is given by $p_{ij} = a_j/H_{ij} + p_{0j}$, where a_j and p_{0j} are given positive parameters for all $j = 1, 2, 3$. Under these more realistic assumptions the total revenue of the fisherman i is

$$\begin{aligned} (\text{TR})_i &= p_{i1}H_{i1} + p_{i2}H_{i2} + p_{i3}H_{i3} \\ &= \left(\frac{a_1}{H_{i1}} + p_{01}\right)H_{i1} + \left(\frac{a_2}{H_{i2}} + p_{02}\right)H_{i2} \\ &\quad + \left(\frac{a_3}{H_{i3}} + p_{03}\right)H_{i3} \\ &= p_{01}H_{i1} + p_{02}H_{i2} + p_{03}H_{i3} + \sum_{k=1}^3 a_k \\ &= p_{01}q_1E_{i1}x_1 + p_{02}q_2E_{i2}x_2 + p_{03}q_3E_{i3}x_3 \\ &\quad + \sum_{k=1}^3 a_k = \langle P_0, qE^{(i)}X \rangle + \sum_{k=1}^3 a_k \\ &= \langle P_0, qE^{(i)}(-AE + X^*) \rangle + \sum_{k=1}^3 a_k \end{aligned} \tag{64}$$

so,

$$\begin{aligned} (\text{TR})_i &= \langle E^{(i)}, -P_0qAE^{(i)} \rangle \\ &\quad + \langle E^{(i)}, P_0qxX^* - P_0qAE^{(j)} \rangle + \sum_{k=1}^3 a_k, \end{aligned} \tag{65}$$

where $P_0 = \text{diag}(p_0)$.

3.2. Expression of the Total Effort Cost. In accordance with many standard fisheries models, we consider that expression of the total effort cost is

$$(\text{TC})_i = \langle c^{(i)}, E^{(i)} \rangle, \tag{66}$$

where $c^{(i)}$ is the constant cost per unit of harvesting and $E^{(i)}$ is the total effort of the fisherman i .

3.3. Expression of the Profit. The profit of each fisherman π_i is equal to total revenue $(\text{TR})_i$; minus total cost $(\text{TC})_i$; it is represented by the following function:

$$\begin{aligned} \pi_i(E^{(i)}) &= (\text{TR})_i - (\text{TC})_i \\ &= \langle E^{(i)}, -P_0qAE^{(i)} \rangle \end{aligned}$$

$$\begin{aligned}
 & + \langle E^{(i)}, P_0qX^* - P_0qAE^{(j)} \rangle + \sum_{k=1}^3 a_k \\
 & - \langle c^{(i)}, E^{(i)} \rangle \\
 = & \langle E^{(i)}, -P_0qAE^{(i)} \rangle \\
 & + \langle E^{(i)}, P_0qX^* - c^{(i)} - P_0qAE^{(j)} \rangle + \sum_{k=1}^3 a_k.
 \end{aligned} \tag{67}$$

3.4. *Constraints of the Model.* The biological model has a meaning if and only if the biomass of all the marine species are strictly positive, then we have

$$X = -AE + X^* \geq X_0 > 0. \tag{68}$$

In other words, for the fisherman i ,

$$AE^{(i)} \leq -AE^{(j)} + X^*. \tag{69}$$

3.5. *Nash Equilibrium Problem.* The problem of determining the fishing effort that maximizes the profit of each fisherman leads to a Nash equilibrium problem. By definition a Nash equilibrium exists when there is no unilateral profitable deviation from any of the fishermen involved. In other words, no fisherman would take a different action as long as every other fisherman remains the same. This problem can be translated into the following two mathematical problems.

The first fisherman must solve the problem (P_1) :

$$\begin{aligned}
 \max \quad & \pi_1(E^{(1)}) \\
 = & \langle E^{(1)}, -P_0qAE^{(1)} + P_0qX^* - c^{(1)} - P_0qAE^{(2)} \rangle \\
 & + \sum_{k=1}^3 a_k
 \end{aligned} \tag{P_1}$$

subject to $AE^{(1)} \leq -AE^{(2)} + X^*$

$E^{(1)} \geq 0$

$E^{(2)}$ given

and the second fisherman must solve the problem (P_2) :

$$\begin{aligned}
 \max \quad & \pi_2(E^{(2)}) \\
 = & \langle E^{(2)}, -P_0qAE^{(2)} + P_0qX^* - c^{(2)} - P_0qAE^{(1)} \rangle \\
 & + \sum_{k=1}^3 a_k
 \end{aligned} \tag{P_2}$$

subject to $AE^{(2)} \leq -AE^{(1)} + X^*$

$E^{(2)} \geq 0$

$E^{(1)}$ given.

The point $(E^{(1)}, E^{(2)})$ is called Nash equilibrium point if and only if $E^{(1)}$ is a solution of problem (P_1) for $E^{(2)}$ given, and $E^{(2)}$ is solution of problem (P_2) for $E^{(1)}$ given.

The essential conditions of Karush-Kuhn-Tucker applied to the problem (P_1) confirm that if $E^{(1)}$ is a solution of the problem (P_1) then there exist constants $u^{(1)} \in \mathbb{R}_+^3$, $v^{(1)} \in \mathbb{R}_+^3$, and $\lambda^{(1)} \in \mathbb{R}_+^3$ such that

$$\begin{aligned}
 & 2P_0qAE^{(1)} + c^{(1)} - P_0qX^* + P_0qAE^{(2)} - u^{(1)} \\
 & + A^T \lambda^{(1)} = 0, \\
 & AE^{(1)} + v^{(1)} = -AE^{(2)} + X^*, \\
 & \langle u^{(1)}, E^{(1)} \rangle = \langle \lambda^{(1)}, v^{(1)} \rangle = 0.
 \end{aligned} \tag{KKT1}$$

In the same way, the essential conditions of Karush-Kuhn-Tucker applied to the problem (P_2) confirm that if $E^{(2)}$ is a solution of the problem (P_2) then there exist constants $u^{(2)} \in \mathbb{R}_+^3$, $v^{(2)} \in \mathbb{R}_+^3$, and $\lambda^{(2)} \in \mathbb{R}_+^3$ such that

$$\begin{aligned}
 & 2P_0qAE^{(2)} + c^{(2)} - P_0qX^* + P_0qAE^{(1)} - u^{(2)} \\
 & + A^T \lambda^{(2)} = 0, \\
 & AE^{(2)} + v^{(2)} = -AE^{(1)} + X^*, \\
 & \langle u^{(2)}, E^{(2)} \rangle = \langle \lambda^{(2)}, v^{(2)} \rangle = 0;
 \end{aligned} \tag{KKT2}$$

we remark that (KKT1) and (KKT2) lead to the following system:

$$\begin{aligned}
 u^{(1)} &= 2P_0qAE^{(1)} + c^{(1)} - P_0qX^* \\
 & + P_0qAE^{(2)} + A^T \lambda^{(1)}, \\
 u^{(2)} &= 2P_0qAE^{(2)} + c^{(2)} - P_0qX^* \\
 & + P_0qAE^{(1)} + A^T \lambda^{(2)}, \\
 v^{(1)} &= -AE^{(1)} - AE^{(2)} + X^*, \\
 v^{(2)} &= -AE^{(1)} - AE^{(2)} + X^*, \\
 \langle u^{(i)}, E^{(i)} \rangle &= \langle \lambda^{(i)}, v^{(i)} \rangle = 0 \quad \forall i = 1, 2,
 \end{aligned} \tag{70}$$

$$E^{(i)}, u^{(i)}, \lambda^{(i)}, v^{(i)} \geq 0 \quad \forall i = 1, 2.$$

To maintain the biodiversity of species, it is natural to assume that all biomasses remain strictly positive; that is, $x_j > 0$ for all $j = 1, 2, 3$; therefore $v^{(1)} = v^{(2)} > 0$.

As the scalar product of $(\lambda^{(i)})_{i=1,2}$ and $(v^{(i)})_{i=1,2}$ is zero, $\lambda^{(i)} = 0$ for all $i = 1, 2$. So, we denote $v = v^{(1)} = v^{(2)}$. Then we have the following expressions:

$$\begin{aligned}
 u^{(1)} &= 2P_0qAE^{(1)} + P_0qAE^{(2)} + c^{(1)} - P_0qX^*, \\
 u^{(2)} &= 2P_0qAE^{(2)} + P_0qAE^{(1)} + c^{(2)} - P_0qX^*, \\
 v &= -AE^{(1)} - AE^{(2)} + X^*,
 \end{aligned} \tag{71}$$

$$\langle u^{(i)}, E^{(i)} \rangle = 0 \quad \forall i = 1, 2,$$

$$E^{(i)}, u^{(i)}, v \geq 0 \quad \forall i = 1, 2.$$

Thus

$$\begin{pmatrix} u^{(1)} \\ u^{(2)} \\ v \end{pmatrix} = \begin{bmatrix} 2P_0qA & P_0qA & A^T \\ P_0qA & 2P_0qA & 0 \\ -A & -A & 0 \end{bmatrix} \begin{pmatrix} E^{(1)} \\ E^{(2)} \\ 0 \end{pmatrix} + \begin{pmatrix} c^{(1)} - P_0qX^* \\ c^{(2)} - P_0qX^* \\ X^* \end{pmatrix}. \tag{72}$$

3.6. Linear Complementarity Problem. We denote

$$\begin{aligned} z &= \begin{pmatrix} E^{(1)} \\ E^{(2)} \\ 0 \end{pmatrix}, \\ w &= \begin{pmatrix} u^{(1)} \\ u^{(2)} \\ v \end{pmatrix}, \\ M &= \begin{bmatrix} 2P_0qA & P_0qA & A^T \\ P_0qA & 2P_0qA & 0 \\ -A & -A & 0 \end{bmatrix}, \\ b &= \begin{pmatrix} c^{(1)} - P_0qX^* \\ c^{(2)} - P_0qX^* \\ X^* \end{pmatrix}. \end{aligned} \tag{73}$$

The Nash equilibrium problem is equivalent to the Linear Complementarity Problem $LCP(M, b)$. Find vectors $z, w \in \mathbb{R}^6$ such that

$$\begin{aligned} w &= Mz + b \geq 0, \\ z, w &\geq 0, \\ z^T w &= 0. \end{aligned} \tag{74}$$

The following proposition confirms that $LCP(M, b)$ has a unique solution.

Proposition 10. *The matrix*

$$M = \begin{bmatrix} 2P_0qA & P_0qA & A^T \\ P_0qA & 2P_0qA & 0 \\ -A & -A & 0 \end{bmatrix} \tag{75}$$

is P -matrix.

Proof. We have $a_{ii} < 0$ for all $i = 1, 2, 3$ and $\Delta > 0$ so, if we note by $(M_i)_{i=1, \dots, 9}$ the submatrix of M , we obtain

$$\begin{aligned} \det(M_1) &= -2p_{01}q_1a_{11} > 0, \\ \det(M_2) &= 4p_{01}q_1p_{02}q_2q_2q_1K_1r_3q_2K_2\Delta > 0, \\ \det(M_3) &= 8p_{01}q_1p_{02}q_2p_3q_3q_3K_3q_1K_1q_2K_2\Delta^2 > 0, \end{aligned}$$

$$\begin{aligned} \det(M_4) &= -12a_{11}p_{01}^2q_1^2p_{02}q_2p_{03}q_3q_3K_3q_1K_1q_2K_2\Delta^2 \\ &> 0, \end{aligned}$$

$$\begin{aligned} \det(M_5) &= 18p_{01}^2q_1^2p_{02}^2q_2^2p_{03}q_3q_1K_1r_3q_2K_2q_3K_3q_1K_1q_2K_2\Delta^3 \\ &> 0, \end{aligned}$$

$$\begin{aligned} \det(M_6) &= 27p_{01}^2q_1^2p_{02}^2q_2^2p_{03}^2q_3^2(q_3K_3q_1K_1q_2K_2\Delta^2)^2 \\ &> 0, \end{aligned}$$

$$\begin{aligned} \det(M_7) &= -9p_{01}q_1p_{02}^2q_2^2p_{03}^2q_3^2a_{11}(q_3K_3q_1K_1q_2K_2\Delta^2)^2 \\ &> 0, \end{aligned}$$

$$\begin{aligned} \det(M_8) &= 3p_{01}q_1p_{02}q_2p_{03}^2q_3^2q_1K_1r_3q_2K_2\Delta(q_3K_3q_1K_1q_2K_2\Delta^2)^2 \\ &> 0, \end{aligned}$$

$$\det(M) = p_{01}q_1p_{02}q_2p_{03}q_3(q_3K_3q_1K_1q_2K_2\Delta^2)^2 > 0. \tag{76}$$

Then, the matrix M is P -matrix and therefore the linear complementarity problem $LCP(M, b)$ admits one and only one solution. \square

The unique solution of $LCP(M, b)$ represents the Nash equilibrium point of our problem and it is given by

$$\begin{aligned} E^{(1)} &= \frac{1}{3}A^{-1} \left(X^* - \frac{c^{(1)}}{P_0q} \right), \\ E^{(2)} &= \frac{1}{3}A^{-1} \left(X^* - \frac{c^{(2)}}{P_0q} \right), \end{aligned} \tag{77}$$

where

$$A^{-1} = \begin{bmatrix} \frac{r_1}{K_1q_1} & \frac{c_{12}}{q_1} & \frac{c_{13}}{q_1} \\ \frac{c_{21}}{q_2} & \frac{r_2}{K_2q_2} & \frac{c_{23}}{q_2} \\ \frac{c_{31}}{q_3} & \frac{c_{32}}{q_3} & \frac{r_3}{K_3q_3} \end{bmatrix}. \tag{78}$$

Then, the fishing effort that maximizes the profit of the first fisherman for catching the first species is

$$\begin{aligned} E_{11} &= \frac{1}{3} \left[\frac{r_1}{K_1q_1} \left(x_1^* - \frac{c_1}{P_{01}q_1} \right) + \frac{c_{12}}{q_1} \left(x_2^* - \frac{c_1}{P_{02}q_2} \right) \right. \\ &\quad \left. + \frac{c_{13}}{q_1} \left(x_3^* - \frac{c_1}{P_{03}q_3} \right) \right]; \end{aligned} \tag{79}$$

the fishing effort that maximizes the profit of the first fisherman for catching the second species is

$$E_{12} = \frac{1}{3} \left[\frac{r_2}{K_2 q_2} \left(x_2^* - \frac{c_1}{P_{02} q_2} \right) + \frac{c_{21}}{q_2} \left(x_1^* - \frac{c_1}{P_{01} q_1} \right) + \frac{c_{23}}{q_2} \left(x_3^* - \frac{c_1}{P_{03} q_3} \right) \right]; \tag{80}$$

the fishing effort that maximizes the profit of the first fisherman for catching the third species is

$$E_{13} = \frac{1}{3} \left[\frac{r_3}{K_3 q_3} \left(x_3^* - \frac{c_1}{P_{03} q_3} \right) + \frac{c_{31}}{q_3} \left(x_1^* - \frac{c_1}{P_{01} q_1} \right) + \frac{c_{32}}{q_3} \left(x_2^* - \frac{c_1}{P_{02} q_2} \right) \right]. \tag{81}$$

The fishing effort that maximizes the profit of the second fisherman for catching the first species is

$$E_{21} = \frac{1}{3} \left[\frac{r_1}{K_1 q_1} \left(x_1^* - \frac{c_2}{P_{01} q_1} \right) + \frac{c_{12}}{q_1} \left(x_2^* - \frac{c_2}{P_{02} q_2} \right) + \frac{c_{13}}{q_1} \left(x_3^* - \frac{c_2}{P_{03} q_3} \right) \right]; \tag{82}$$

the fishing effort that maximizes the profit of the second fisherman for catching the second species is

$$E_{22} = \frac{1}{3} \left[\frac{r_2}{K_2 q_2} \left(x_2^* - \frac{c_2}{P_{02} q_2} \right) + \frac{c_{21}}{q_2} \left(x_1^* - \frac{c_2}{P_{01} q_1} \right) + \frac{c_{23}}{q_2} \left(x_3^* - \frac{c_2}{P_{03} q_3} \right) \right]; \tag{83}$$

the fishing effort that maximizes the profit of the second fisherman for catching the third species is

$$E_{23} = \frac{1}{3} \left[\frac{r_3}{K_3 q_3} \left(x_3^* - \frac{c_2}{P_{03} q_3} \right) + \frac{c_{31}}{q_3} \left(x_1^* - \frac{c_2}{P_{01} q_1} \right) + \frac{c_{32}}{q_3} \left(x_2^* - \frac{c_2}{P_{02} q_2} \right) \right]. \tag{84}$$

4. Numerical Simulations and Discussion of the Results

In this section, we take as case of study two fishermen who catch three fish species competing with each other for space or food. In order to assure the existence and stability of the locally asymptotically stable state of the three fish populations, we consider the parameters of the model system (5) as shown in Table 1.

Let us consider the economic parameters such as that shown in Table 2.

Using the parameters cited in Tables 1 and 2, thereafter we will see how changes in the minimum prices can affect effort fishing, catches, and profit.

TABLE 1: Characteristics of the three marine species.

Species 1	Species 2	Species 3
$r_1 = 0,5$	$r_2 = 0,3$	$r_3 = 0,2$
$K_1 = 1000$	$K_2 = 700$	$K_3 = 600$
$c_{12} = 2 \cdot 10^{-4}$	$c_{21} = 10^{-5}$	$c_{31} = 10^{-4}$
$c_{13} = 3 \cdot 10^{-4}$	$c_{23} = 2 \cdot 10^{-5}$	$c_{32} = 10^{-4}$

TABLE 2: Economic parameters of the model.

Species 1	Species 2	Species 3
$a_1 = 0,1$	$a_2 = 0,2$	$a_3 = 0,3$
$p_{01} = 1$	$p_{02} = 2$	$p_{03} = 3$
$q_1 = 0,1$	$q_2 = 0,02$	$q_3 = 0,004$
$c_1 = 0,1$	$c_1 = 0,1$	$c_1 = 0,1$
$c_2 = 0,2$	$c_2 = 0,2$	$c_2 = 0,2$

TABLE 3: The influence of the price on the fishing effort.

p_{01}	p_{02}	p_{03}	E_1	E_2
1	2	3	17,0451	16,5151
11	17	23	17,5943	17,5314
16	27	48	17,6383	17,6073
31	47	78	17,6552	17,6363
51	70	108	17,6627	17,6492
84	101	273	17,6734	17,6677
106	133	327	17,6749	17,6702
340	378	427	17,6769	17,6736
574	577	606	17,6783	17,6760
808	811	914	17,6794	17,6778
917	956	981	17,6795	17,6781
1000	1079	1090	17,6797	17,6784

TABLE 4: The influence of the price on the catches.

p_{01}	p_{02}	p_{03}	H_1	H_2
1	2	3	245,0957	234,4651
11	17	23	246,4411	245,9382
16	27	48	246,5725	246,2429
31	47	78	246,6298	246,3781
51	70	108	246,6552	246,5718
84	101	273	246,6865	246,5974
106	133	327	246,6923	246,6334
340	378	427	246,7020	246,6334
574	577	606	246,7063	246,6582
808	811	914	246,7095	246,6775
917	956	981	246,7101	246,6804
1000	1079	1090	246,7107	246,6839

4.1. The Influence of the Price on the Fishing Effort, Catches, and Profit. By Tables 3, 4, and 5 we will discover how changes in the price can affect the fishing effort, catches, and profit.

According to Tables 3, 4, and 5, one can remark that an increase in the price leads to an increase in fishing effort, catches, and profit. But it is clear that when the price

TABLE 5: The influence of the price on the profits.

P_{01}	P_{02}	P_{03}	π_1	π_2
1	2	3	282	269
11	17	23	2959	2942
16	27	48	4513	4500
31	47	78	8479	8465
51	70	108	13584	13567
84	101	273	23130	23118
106	133	327	29111	29099
340	378	427	85598	85572
574	577	606	142017	141987
808	811	914	200558	200531
917	956	981	227721	227692
1000	1079	1090	249295	249266

level increases significantly, that is, when it varies in a large amplitude interval, the fishing effort and the catches increase by varying in an interval of small amplitude. More precisely, when the price is between 1 and 1090, the fishing effort varies between 16,51 and 17,68, and the catches vary between 234,4 and 246,7. This is justified by the need for conservation of marine species even if the price increases significantly.

From Table 5 one can see that the level of profit increases, which allows fishermen to have highest returns through more reasonable catches, taking into account the conservation of biodiversity.

These results allow us to deduce that our model is pertinent since it allows us to determine the fishing effort that maximizes the profit of each fisherman without being obliged to make more catches that lead to the overexploitation of these marine species.

Let us add that when the price tends to infinity, the fishing efforts of the two fishermen are equal and they do not exceed 18, as well as the catches which do not exceed 250; contrariwise the profit is always increasing thanks to the increase of the price. Then we can deduce the effect of the price change on the fishing effort, catches, and profit.

It is very interesting to note that if the price tends to infinity and the fishing effort is superior to 18, then the catches and the profit decrease.

5. Conclusions


In this paper, we have developed a bioeconomic model for three species catches by two fishermen. In one hand, we have assumed that the evolution of these species is described by a density dependent model taking into account the competition between the species which compete with each other for space or food. The natural growth of each species is modeled using a logistic law. On the other hand, we have assumed that the prices of these species vary according to the quantity harvested. In this work we have calculated fishing effort that maximizes the income of each fisherman at biological equilibrium by using the Nash equilibrium problem. The existence of the steady states and their stability are studied using eigenvalue analysis and Routh-Hurwitz criterion.

Conflicts of Interest

The authors declare that there are no conflicts of interest.

References

- [1] C. W. Clark and G. R. Munro, "The economics of fishing and modern capital theory: a simplified approach," *Journal of Environmental Economics and Management*, vol. 2, no. 2, pp. 92–106, 1975.
- [2] C. W. Clark, *Bioeconomic modelling and fisheries management*, John Wiley and Sons, New York, NY, USA, 1985.
- [3] K. S. Chaudhuri, "A bioeconomic model of harvesting a multi-species fishery," *Ecological Modelling*, vol. 32, no. 4, pp. 267–279, 1986.
- [4] K. Chaudhuri, "Dynamic optimization of combined harvesting of a two-species fishery," *Ecological Modelling*, vol. 41, no. 1-2, pp. 17–25, 1988.
- [5] K. S. Chaudhuri and S. S. Ray, "On the combined harvesting of a prey-predator system," *Journal of Biological Systems*, vol. 4, no. 3, pp. 373–389, 1996.
- [6] T. K. Kar and K. S. Chaudhuri, "Harvesting in a two-prey one-predator fishery: A bioeconomic model," *Anziam Journal*, vol. 45, no. 3, pp. 443–456, 2004.
- [7] R. Mchich, P. M. Auger, R. Bravo De La Parra, and N. Raissi, "Dynamics of a fishery on two fishing zones with fish stock dependent migrations: Aggregation and control," *Ecological Modelling*, vol. 158, no. 1-2, pp. 51–62, 2002.
- [8] Y. El Foutayeni, M. Khaladi, and A. Zegzouti, *Profit maximization of fishermen exploiting two fish species in competition*, vol. 15 of *AMO Advanced Modeling and Optimisation*, 2013, 457–469.
- [9] Y. El Foutayeni, M. Khaladi, and A. Zegzouti, *A generalized Nash equilibrium for a bioeconomic problem of fishing*, *Studia Informatica Universalis-HERMANN*, 2012, 186–204.
- [10] Y. EL Foutayeni and M. Khaladi, "A bio-economic model of fishery where prices depend on harvest," *Advanced Modeling and Optimization. The Electronic International Journal*, vol. 14, no. 3, pp. 543–555, 2012.
- [11] Y. El Foutayeni and M. Khaladi, "A generalized bio-economic model for competing multiple-fish populations where prices depend on harvest," *Advanced Modeling and Optimization. The Electronic International Journal*, vol. 14, no. 3, pp. 531–542, 2012.
- [12] Mailleret Ludovic, *Stabilisation Globale de Systèmes Dynamiques Positifs Mal Connus*, Applications en Biologie. Ecologie, Environnement, Université Nice Sophia Antipolis, 2004.
- [13] G. Birkhoff and G.-C. Rota, *Ordinary Differential Equations*, Ginn, 1982.



Hindawi

Submit your manuscripts at
<https://www.hindawi.com>

