

 Open access • Journal Article • DOI:10.1126/SCIENCE.1169659

Origin and Radiation of the Earliest Vascular Land Plants — [Source link](#)

[Philippe Steemans](#), [Alain Le Hérissé](#), [John Melvin](#), [Merrell A. Miller](#) ...+3 more authors

Published on: 17 Apr 2009 - [Science](#) (American Association for the Advancement of Science)

Topics: [Vascular plant](#)

Related papers:

- [Early Middle Ordovician evidence for land plants in Argentina \(eastern Gondwana\)](#)
- [Fragments of the earliest land plants](#)
- [The origin and early evolution of plants on land](#)
- [The Microfossil Record of Early Land Plants](#)
- [New evidence for land plants from the lower Middle Ordovician of Saudi Arabia](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/origin-and-radiation-of-the-earliest-vascular-land-plants-nzveov99hf>

Origin and Radiation of the Earliest Vascular Land Plants

Philippe Steemans,^{1*} Alain Le Hérisse,² John Melvin,³ Merrell A. Miller,³ Florentin Paris,⁴ Jacques Verniers,⁵ Charles H. Wellman^{6*†}

The earliest land plants (embryophytes) evolved from charophycean green algal ancestors and probably possessed bryophyte-like anatomy and physiology (1–3). Plant megafossils are rare, presumably because they lacked fossilizable tissues, although there is a large microfossil record of dispersed spores, termed cryptospores because they occur in unusual configurations such as dyads and tetrads (3–6). The oldest uncontroversial cryptospore record is Mid-Ordovician (Darrivilian) in age (Fig. 1A). Similar cryptospore assemblages have been reported throughout the globe, suggesting that the earliest terrestrial vegetation consisted of ecological generalists and was cosmopolitan. The cryptospore record shows little spatial or temporal variation over about 30 million years. Vascular plants are considered to have originated and adaptively radiated from the Early Silurian on the basis of major changes in spore assemblages (from cryptospore to hilate/trilete spore dominated) and the appearance of vascular plant megafossils.

We examined palynological preparations from 37 core samples from Ordovician-Silurian strata penetrated in the Qusaiba-1 corehole, northern Saudi Arabia (7). They yielded abundant, well-preserved, and thermally immature marine palynomorphs (chitinozoans and acritarchs) that were used to determine the age of the cores (7). Subsidiary land plant spores were present in all samples. The upper part of the Qasim Formation is a shallowing upward succession of offshore shelf sandy silt and mudstones topped by a sandstone. The oldest samples from this unit are no younger than Mid-Katian in age, and succeeding samples from the uppermost Qasim Formation are Late Katian in age (Fig. 1A) on the basis of chitinozoan and acritarch biostratigraphy. The Qasim Formation was unconformably overlain by glacial-marine sediments of the Sarah Formation. These sediments are characterized by Himantian acritarch species with reworked earlier Ordovician forms. The Qalibah Formation (Qusaiba Member) overlies the Sarah Formation and consists of marine shelf deposits of Early Silurian age.

The ancestral condition among embryophytes may have been meiotically produced spores dis-

persed in tetrad or dyad configurations (cryptospores). Dissociation of these units into individual hilate spores (from dyads) or trilete spores (from tetrads) is a more derived condition. Cryptospores were the most abundant plant spores recovered from the Late Ordo-

although rare specimens have been recorded in Hirnantian deposits from Turkey (northern Gondwana). Ornamented hilate spores and trilete spores have not been reported from strata older than Wenlock.

Trilete spores are generally considered to derive from the vascular plant lineage (7) because extant and fossil basal vascular plant groups (nonseed plants) nearly all produce dissociated single spores of trilete or the more derived monolete form. A small proportion of extant bryophytes produce trilete spores, but in these cases the trilete condition is likely non-functional or convergent (3, 7). Also, the earliest reported vascular plant megafossils are more-or-less coincident with the appearance and diversification of hilate/trilete spores. Thus, hilate/trilete spores from the Late Ordovician may represent an earlier emergence and diversification of the vascular plant lineage in Gondwana. Hilate/trilete spores found elsewhere suggests that vascular plants may have subsequently migrated out of Gondwana and colonized other continents where they secondarily diversified.




Fig. 1. (A) Ordovician to Lower Devonian time scale. Black bar represents the studied interval. Stratigraphic position of 1, the oldest cryptospores; 2, the oldest mesofossils of sporangia containing cryptospores; 3, the previous oldest laevigate trilete spores from Turkey; 4, the previous oldest ornamented trilete spores; 5, the oldest megafossils of the vascular plant lineage; and 6, the oldest vascular plant megafossils with preserved conducting tissues. (B to I) Fossil trilete spores from the Qusaiba-1 corehole. Descriptions of spores provided in (7). Scale bars indicate 10 μ m.

vician sediments of the Qusaiba-1 corehole and are similar to forms in other coeval assemblages found worldwide. Surprisingly, all samples also contained plant spores naturally dissociated from polyads: one species of hilate spore and seven species of trilete spores (Fig. 1, B to I). Two trilete spore species were unornamented, whereas the other five had an ornament of verrucae and/or muri, indicating high species diversity. Neither hilate spores nor trilete spores have been reported from strata of this age (3–6). The oldest unornamented hilate spores are from the Llandovery,

References and Notes

- D. Edwards, C. H. Wellman, in *Plants Invade the Land*, P. G. Gensel, D. Edwards, Eds. (Columbia Univ. Press, New York, 2001), chap. 2.
- C. H. Wellman, P. L. Osterloff, U. Mohiuddin, *Nature* **425**, 282 (2003).
- J. Gray, *Philos. Trans. R. Soc. London Ser. B* **309**, 167 (1985).
- J. B. Richardson, in *Palynology: Principles and Applications*, J. Jansonius, D. C. McGregor, Eds. (AASP-The Palynological Society, Dallas, TX, 1996), vol. 2, chap. 18A.
- P. Steemans, *Rev. Palaeobot. Palynol.* **113**, 189 (2000).
- P. K. Strother, in *Phanerozoic Terrestrial Ecosystems*, R. A. Gastaldo, W. A. DiMichele, Eds. (Palaeontological Society Papers No. 6, Pittsburgh, PA, 2000), pp. 3–20.
- Materials and methods, age dating, spore descriptions, and discussion of trilete spore affinities are available as supporting material on Science Online.
- P.S. is a National Fund for Scientific Research Research Associate. Saudi Aramco provided material for study and publication. Work was undertaken as part of the Commission Internationale de Microflora du Paléozoïque/Saudi Aramco joint project "Understanding the Palaeozoic palynology of Saudi Arabia."

Supporting Online Material

www.sciencemag.org/cgi/content/full/324/5925/353/DC1

Materials and Methods

SOM Text

References

11 December 2008; accepted 17 February 2009
10.1126/science.1169659

¹Palaeobotany, B-18, University of Liège, 4000 Liège, Belgium.
²Université de Brest, UMR 6538 du CNRS, Institut Universitaire Européen de la Mer, 6 Avenue le Gorgeu, 29238 Brest-cedex 3, France.
³Saudi Aramco, Dhahran, 31311, Saudi Arabia.
⁴Université de Rennes 1, UMR 6118 du CNRS, 35042 Rennes-cedex, France.
⁵Research Unit Palaeontology, Ghent University, Krijgslaan 281/58, B-9000 Ghent, Belgium.
⁶Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK.

*These authors contributed equally to this work.

†To whom correspondence should be addressed. E-mail: c.wellman@sheffield.ac.uk