

ORIGINAL RESEARCH PAPER

Origin and spatial distribution of metals in agricultural soils

Gh.A. Mohammadpour¹, A.R. Karbassi^{2,*}, A. Baghvand²

¹Department of Environmental Science, Graduate School of the Environment and Energy, Science and Research Branch, Islamic Azad University, Tehran, Iran

²Graduate Faculty of Environment, University of Tehran, P.O. Box 14135, Tehran, Iran

Received 16 December 2015; revised 5 January 2016; accepted 30 January 2016; available online 1 March 2016

ABSTRACT: Presence of toxic metals in agricultural soils can impose adverse health impact on consumers. The main purpose of this study was to determine spatial distribution of elements Fe, Sb, Mn in agriculture soils and crops of Hamedan Province in Iran. Soil samples (0-20 cm depth) were collected from an area of 2831 km². Iron, Antimony and Manganese in samples of soil and agricultural crops were extracted and their amount was determined using atomic absorption spectrometer. The spatial distribution map of the studied elements was developed using Kriging method. The main concentration of Fe, Sb and Mn in the soil of the study area is about 3.8%, 2.5 and 403 mg/kg, respectively. According to chemical partitioning studies, the anthropogenic share of Fe, Sb and Mn is about 28.51%, 34.83% and 30.35%, respectively. Results of comparison of heavy metals pollution intensity in the agricultural soil with geo-accumulation index and also pollution index, illustrated that iron and manganese are classified in the Non-polluted class and antimony is in the moderately polluted class. Analysis of zoning map of pollution index showed that Fe, Sb and Mn are of geological sources. In fact, these metals are naturally found in soil. However, anthropogenic activities have led to more accumulation of these metals in the soil. The obtained health risk for metals in agricultural crops is indicative of safe value for consumers.

KEYWORDS: Agricultural yields; Health index ratio (HRI); Heavy metals; Soil; Zoning

INTRODUCTION

The levels of toxic metals is increasing in the agricultural soils due to over utilization of various chemicals for better yields. The pollutants can include metals, organic wastes and other organic and inorganic substances (Shetty and Rajkumar, 2009). The term "heavy metals" is very general but generally refers to any metallic element that has a relatively high density and is toxic even at low concentrations (Uba *et al.*, 2009). Due to the adverse health impacts of metals, intensive research has been carried out to determine heavy metals in foods and other common food

✉ *Corresponding Author Email: akarbasi@ut.ac.ir

Tel.: +98 2161 113 139; Fax: +98 2166 407 719

Note. Discussion period for this manuscript open until June 1, 2016 on GJESM website at the "Show Article".

contaminants. The adverse health effects of heavy metals could mainly be due to their interference with normal body biochemistry through various metabolic processes (Okunola *et al.*, 2011). Some of the heavy metals are extremely persistent in the environment (Taghinia Hejabi *et al.*, 2011). They are not easily biodegradable and thus their accumulation reaches to critical levels (Khan *et al.*, 2009). Metals have a high degree of toxicity that can be dangerous for both the human and the environment (Singanan *et al.*, 2008). The soil pollution by heavy metals has received ample attention in the recent decades (Rafiei *et al.*, 2010). Since soil is a thin part of the earth surface system that comes into direct contact with man (Kargar *et al.*, 2012), therefore, it can transfer pollutants to food chain and

on the other hand it can be polluted by man's activities. For instance, soil properties and quality can be adversely affected by the agricultural and industrial activities (Nasrabadi et al., 2010). Thus it is vital to preserve soil conditions for sustainable development (Ghaderi et al., 2012). For these reasons, more attention has been paid to the geochemistry of soil and also the intensity of pollutants (Soffianian et al., 2014; Karbassi et al., 2008). Since metals contents of soil can find their way into the agricultural crops, more awareness is built up in societies in connection health agricultural products (Akoto et al., 2008). Wastewater irrigation, solid waste disposal, sludge applications, vehicular exhaust and industrial activities are the major sources of metals in soil and an increased metal uptake by food crops grown on such contaminated soils is often observed (Nasrabadi et al., 2011). In general, farmers prefer to use wastewater for their beneficial nutrients and they ignore the presence of toxic metals (Chen et al., 2005; Singh et al., 2004). Although low concentrations of these metals are naturally found in soils, human activities have elevated their concentrations (Karbassi et al., 2015). Thus it is very important to assess soil pollution and take the necessary remediation measures (Romic et al., 2007). Mining, industries, road traffic, waste disposal, and agricultural use of fertilizers and chemicals are amongst human activities that can lead to heavy metal contamination of the soil (Karbassi et al., 2016). On the other hand, main natural factors contributing to metal contamination of the soil include volcanoes, fires in forests, and chemical composition of parent materials (Lado et al., 2008). Heavy metal accumulation in soils and plants is of increasing concern because of the potential human health risks (Karbassi et al., 2011). Heavy metal accumulation in plants depends on many factors that has brought out the subject of transfer factors of the metals (Rattan et al., 2005). The transfer factors of metals from soil to plant vary from soil to soil (Karbassi, 1998). The texture of soil, organic content of soil and also type of plant are among the main factors for transfer of metals from soil to plants (Kabata-Pendias and Pendias, 1984). As a general rule, there should be a balance amongst uptake of various materials by plants to ensure health products for consumers as well as for the plants (Iyengar and Nair, 2000; Turkdogan et al., 2003). Thus, it will be wise to determine the elemental contents of soil before beginning new crop production. Thus, it is essential

to set up a guideline to balance amongst heavy metals transfer in soil, the accumulation in vegetables and the health of consumers. Thus, it will be useful to assess health risk of metals in many ways (Raj Shakya and Malla Khwaounjoo, 2013). There are several studies in different countries for interpolation and determination of spatial distribution of heavy metals concentrations in soil (Biati et al., 2010; Karbassi et al., 2014). Lado et al. (2008) carried out modeling the distribution of eight critical metals (As, Cd, Cr, Cu, Hg, Ni and Zn) in European topsoil. They used regression-Kriging method and accuracy of predictions was evaluated by cross validation method. Spatial distribution of Cr, As, Cu, Pb, Cd and Hg in Changxing of Zhejiang Province in China was investigated by geo-statistics and log normal Kriging and ordinary Kriging for mapping (Juang et al., 2001). Juang et al. (2001) and Rodriguez et al. (2009) applied geo-statistics in their studies in order to prepare distribution map of heavy metals. Preparation of map to show the spatial distribution of metal contents in soils can help decision makers to select suitable areas for various land use. Hence, due to importance of Hamedan City as one of the main centers of agricultural activities in Iran, it is of utmost importance to determine the pollution of topsoil and agricultural crops of this area. Thus in the present investigation has tried to bring out the sources of Fe, Mn and Sb in agricultural soils of Hamedan. Subsequently the transfer factor of metals from soil to crops was determined. Finally the health risk of metals as a result of consumption of various agricultural products was computed. This study has been performed in Hamadan Province located in west of Iran in 2014.

MATERIALS AND METHODS

Hamedan Province occupies an area of 2831 km² (34° 35' 34" N, 20° 49' 13" E). Samples were taken from depth of 0-20 cm in accordance with systematic method. A total of 58 compound samples of topsoil and crops of wheat, barley, corn, alfalfa and potatoes were collected. Sampling was conducted in late September after harvest. Fig. 1 shows the location map of the study area and sampling points.

Soil samples were transferred to the laboratory for testing and measuring soil physical and chemical properties and were passed through 63-micron sieve. Phosphorus was measured by Olsen P extracting solution (0.5 M NaHCO₃, pH 8.5); total nitrogen by




Fig. 1: Location map of the study area along with the sampling points

Kjeldahl digestion; pH was analyzed by glass electrode using a 1:1 soil: water ratio; and EC and salinity was measured by conductivity meter in a soil-water extract (1:2 soil: water ratio) (USDA, 2004; ASTM, 2000; Dewis and Freitas.,1984). For measuring soil-lime titration method was used (Gee and Bauder, 1986). Sodium and potassium were measured by a flame detector (AOAC, 2005). Soil texture was done in accordance with Gee and Bauder (1986). Total C were measured as described by Allison (1965) and Supaphol *et al.*, (2006). Cation exchange capacity was measured according to standard methods (APHA, 2005; Aparna *et al.*, 2010). Total organic carbon is an alternative analytical method for measuring hydrocarbons using the wet oxidation technique as previously reported by Nelson and Sommers (1975). Measurement of calcium and magnesium of soil was carried out by EDTA solution using complex metric (titrations) (AOAC, 2005; Gee and Bauder, 1986).

For digestion of soil samples, at first they were air dried and passed through a 63-micron sieve. Then 2g of soil was poured into a capped container and 15 mL of 4N nitric acid was added. Then the flasks were left for 12 h in a hot water bath at 80°C. Subsequently, the samples were filtered and concentrations of heavy metals were measured by atomic absorption spectrometer (Varian AA-400 model; available in laboratory of Environment Protection Organization of Hamedan Province). The reagents used include distilled water, 0.01M calcium chloride, 0.05M diethylene tri-amine penta-acetic acid (DTPA) (Black, 1965).The dry ash extraction method was used for digestion of plant samples (Shaw, 1989). Then concentration of iron, manganese and antimony of

plant samples in the considered extraction was determined by atomic absorption spectrometer. The instrument was calibrated with diluted solution prepared from a known stock solution of each element. The accuracy and precision of the overall procedure have been determined and are estimated to be around 5% for most elements. The quality assurance of the analytical results was controlled with the use of certified reference marine organism IAEA-407 provided by International Atomic Energy Agency (Mzoughi and Chouba, 2012). Soil to plant metal transfer was computed as translocation factor (TF), which was calculated by using the equation $TF = C_{Plant}/C_{Soil}$ where, C_{Plant} is the concentration of heavy metals in plants and soil is the concentration of heavy metals in soil (Mahmood and Malik, 2014). Daily intake of vegetables in adult was calculated by data obtained during the survey though a questionnaire. DIM was calculated by the following equation: $DIM = C_{metal} \times C_{factor} \times D_{food\ intake} / B_{average\ weight}$ (Chary *et al.*, 2008).

Where C_{metal} , C_{factor} , $D_{food\ intake}$ and $B_{average\ weight}$ represent the heavy metal concentrations in plants (mg/kg), conversion factor: 0.085 (Chary *et al.*, 2008), daily intake of vegetables and average body weight, respectively. In the present research work vegetables grown at the soils were collected from the study area and their metal concentration was used to calculate the health risk index (HRI). Value of HRI depends upon the daily intake of metals (DIM) and oral reference dose (RfD). RfD is an estimated per day exposure of metal to the human body that has no hazardous effect during life time (US-EPA IRIS, 2006). The health risk index for Sb, Mn and Fe by consumption

of contaminated vegetables was calculated by following equation:

$$HRI = \frac{DIM}{Rfd} \quad (\text{Akoto et al., 2014}). \quad (1)$$

Where DIM represents the daily intake of metals and Rfd represents reference oral dose. Rfd value for Sb, Mn and Fe is 0.0004, 0.14 and 7000 (mg/kg bw/day) respectively (WHO, 1993; EPA, 2007). To quantify the degree of heavy metal pollution in soil, Igeo was calculated according to Muller and is given in Eq. 1 (Praveena et al., 2008). The results were interpreted using Igeo classes given in Table 1.

$$I_g = \text{Log}_2 \left(\frac{C_n}{1.5 B_n} \right) \quad (2)$$

Where Cn is the concentration of the examined metal in the soil, Bn is the geochemical background value of a given metal in the soil (Turekian and Wedepohl, 1961) and the factor 1.5 is used to account the possible variations in the background values.

To assess the intensity of metal contamination in Hamedan soils, the pollution index (Karbassi et al., 2008) was calculated using:

$$I_p = \text{Log}_2 \left(\frac{C_n}{B_n} \right) \quad (3)$$

Where, Cn is the total elemental content in soils and Bn is geochemical background of element.

Chemical fractionation method suggests that heavy metals form five bonds with soil and sediments including loosely bond, sulfide bond, organometallic bond, most resistant bond and within lattice bond among which loosely bond, sulfide bond and organometallic bond indicate anthropogenic elements of the environment and the other two bonds, namely, most resistant bond and within lattice bond indicate natural part of heavy metals in soil (Spencer and

Macleod, 2002; Karbassi, Shankar., 2005). Five-stage chemical fractionation method was applied in this study in order to specify the anthropogenic part of heavy metals and the natural part of heavy metals. To understand the relationship amongst various metals and environmental indicators, Multi Variable Statistical Program (MVSP) was used. This analytical software is frequently used by various researchers (Karbassi et al., 2004 and 2008). Highest similarities are clustered/linked first. Two variables are connected only if they are highly correlated. After two variables are clustered, their correlations with all the other variables are averaged. The results of clustering are displayed in the form of a Dendrogram (Karbassi et al., 2007). Kriging interpolation method was applied in Arc GIS 9.3 software to show changes in concentration distribution of the investigated heavy metals (Johnston et al., 2001).

RESULTS AND DISCUSSION

Investigation of physicochemical parameters of soil showed that the parameters of electrical conductivity (EC), sodium, calcium carbonate (CaCO₃) and salinity have coefficients of variation above 50% which indicates the large variations of concentration of these variables in soil of Hamedan City. The rest of soil physicochemical parameters have coefficients of variation less than 50% which indicates its low variation in soil of Hamedan City. The average acidity of soil is 7.72 which indicates the soil of the study area is of alkaline nature. The dominant soil texture in the area is silty clay loam soil in accordance with USDA soil taxonomy. A summary of statistical status of some physicochemical properties of the soil of study area is presented in Table 2.

The results of Kolmogorov-Smirnov test showed that all data of soil and harvested crops are normal. A summary of statistical status of concentration of heavy metals in samples is presented in Table 3. The average concentration of heavy metals in soil of Hamedan City is as follows: iron (3.77 %), manganese (403.38 mg/kg) and antimony (2.44 mg/kg). It was concluded that among the investigated metals, concentration of manganese in soil is above the standard level. Table 4 compares concentrations of iron and antimony in the soil of study area with mean concentrations of Europe and the mean concentration of the world is soil As it is seen, mean concentrations of iron and antimony in this study are more than those of Europe but mean concentration of iron is lower than that of the world

Table 1: Igeo classes in relation to soil quality (Serbaji et al., 2012)

Igeo	Igeo class	Soil quality
0-0	0	Unpolluted
0-1	1	Unpolluted to moderately polluted
1-2	2	Moderately polluted
2-3	3	Moderately to highly polluted
3-4	4	Highly polluted
4-5	5	Highly to very highly polluted
5-6	>5	Very highly polluted

Table 2: Physio-chemical characteristics of the soils used in the study

Factors	Unit	Minimum	Maximum	Mean	SD*	Coefficient of variation
pH	-	6.80	8.90	7.7224	0.52249	0.068
Organic	%	0.90	2.00	1.4200	0.25782	0.182
CEC	Cmole/kg	9.40	39.40	23.1112	7.72925	0.334
clay	%	8.50	54.30	27.2512	12.16787	0.447
sand	%	8.90	69.40	30.9724	14.25832	0.460
silt	%	7.90	77.40	46.0547	17.81106	0.387
EC	(ds/m)	0.05	3.70	1.6533	1.14063	0.691
CaCO ₃	%	1.00	43.00	12.3793	6.75856	0.546
N	%	0.01	0.09	.0526	0.02283	0.457
P	(mg/kg)	3.40	39.40	21.9807	8.21287	0.374
Ca	(mg/kg)	4.40	49.40	22.9752	10.42741	0.454
mg	(mg/kg)	69.40	199.50	143.6884	29.98786	0.209
Na	(mg/kg)	8.50	71.60	32.2821	17.76203	0.550
K	(mg/kg)	10.40	79.40	28.6617	12.34805	0.431
Salinity	(ds/m)	0.10	2.24	0.7386	0.40293	0.545

*Standard Deviation

Table 3: Mean concentration of heavy metals in the soils

Heavy metals	Minimum	Maximum	Mean	Std. Deviation	Standard level
Fe (%)	2.20	5.00	3.769	0.63057	1000 (WHO, 1996)
Mn (mg/kg)	124.00	768.00	403.379	167.056	80 (Tsafe et al., 2012)
Sb (mg/kg)	1.00	5.50	2.439	0.86814	2-10 (USEPA, 1983)

Table 4: Comparison of the concentrations of elements in the studied area with the values from the region, Europe, and the world (Facchinelli et al., 2001; Franco-Uria et al., 2009)

Heavy metal	Europe mean	Area mean	World median	Area median
Fe (%)	2.17	3.769	4	3.9
Mn (mg/kg)	NA*	403.379	NA*	427
Sb (mg/kg)	1.04	2.439	0.5	2.35

*Not Announced

Table 5: Mean concentration heavy metals in the crops of Hamedan region

Crops	Fe (%)	Mn (mg/kg)	Sb (mg/kg)
Wheat	0.6227	46.0258	0.0269
Barley	0.379	44.3914	0.0222
Potato	0.2398	42.0538	0.0416
Alfalfa	0.2888	41.0107	0.0259
Corn	0.3218	45.6891	0.0209

and finally mean concentration of antimony is higher than that of the world. In this study, concentration of iron is much lower than the amount reported by in Jordan (5370.6 mg/kg), Malaysia (1790 mg/kg), England (2600 mg/kg), Spain (19300 mg/kg), and Canada (25660 mg/kg). The average concentration of iron in Earth's crust is 41000 mg/kg (Rasmussen et al., 2001; De Miguel et al., 1997; Ramlan and Badri, 1989; Jiries, 2003; Schwar et al., 1988; Karbassi et al., 2005; Niencheski, 2002).also concentration of manganese in this study was more than reported by Jordan (144.6 mg/

kg), Malaysia (153 mg/kg), Spain (362 mg/kg). Table 5 shows average concentration of iron, manganese and antimony in agricultural crops harvested in Hamedan City. International levels of iron, manganese and antimony in the agricultural crops are 5, 6.61 and 0.003 mg/kg, respectively (FAO/WHO, 2011; McGrath et al., 1997; Lone et al., 2008). In this study, concentration of manganese (in all crops) is higher than the standard level. According to the studies carried out by Tsafe et al. (2012) in Nigeria, concentration of iron and manganese in soil are 0.0142% and 195.25 mg/

Table 6: Transfer factor (TF) of heavy metals in agriculture crops grown at Hamedan City

Factor	Range	Minimum	Maximum	Mean	SD
Mn	644.00	124.00	768.00	403.3793	167.05696
Fe	2.80	2.20	5.00	3.7690	0.63057
Sb	4.50	1.00	5.50	2.4397	0.86814

Table 7: DIM and HRI of heavy metals in adults and children

Heavy metal	Heavy metals in crops	DIM in adults	DIM in children	HRI in adults	HRI in children
Mn	43.834	0.023	0.0264	0.164	0.189
Fe	0.37	1.94×10^{-4}	2.232×10^{-4}	2.774×10^{-4}	3.187×10^{-4}
Sb	0.0275	1.44×10^{-5}	2.466×10^{-5}	0.036	0.0617

Table 8: Muller’s geochemical index, Karbassi modified geochemical accumulation index and contamination factor of studied soils (Karbassi et al., 2008)

Heavy metals	Background concentration of heavy metal ¹	I _{poll}	I _{geo}
Mn	850	0	0
Fe	3800	0	0
Sb	0.6	2.025	1.439

¹Manta et al., 2002; McBride et al., 1997; Istvan, 1997




Fig. 2: Cluster analysis amongst heavy metals and physico-chemical parameters in Hamedan soil

kg, respectively and the said concentrations in vegetables are 0.0044% and 54.05 mg/kg, respectively (Tsafe et al., 2012). Concentrations of iron and manganese in vegetables might be very negligible (Anthony et al., 2007; Zhuang et al., 2009).

The ability of a metal species to migrate from the soil in to the plant parts is called as transfer factor (TF). Table 6 shows the transfer factor from soil to agricultural crops in Hamedan region. According to the studies carried out by Mahmood and Malik (2014), the transfer factor for manganese is less than the standard level. That is in accordance with the result of present study. The degree of toxicity of heavy metals to human being depends upon their daily intake (Anita

et al., 2010). The population will pose no risk, if the ratio is less than 1 and if the ratio is equal or greater than 1 then population will experience health risk (Sajjad et al., 2009). Table 7, presents the daily intake and health risk of the studied metals in children and adults in Hamedan region. Given the daily intake of metals from agricultural products as well as considering the fact that health risk index is less than 1 for all metals, hence it will be safe for children and adults to consume the agricultural products. In a study performed by Mahmood and Malik (2014), HRI was found to be maximum for *Spinacia oleracea* (2.42 mg/kg) and *Brassica campestris* (1.22 mg/kg).

Intensity of soil pollution with heavy metals using

Table 9: Chemical bonds of heavy metals in Hamedan City soils

Heavy metals	Total concentration	Fractional Steps					Anthropogenic portion	Lithogenic portion
		Step 1	Step 2	Step 3	Step 4	Step 5		
Mn (mg/kg)	403.38	27.43	36.02	58.98	278.76	2.194	122.43	280.95
Fe (%)	3.77	0.37	0.67	0.035	2.691	0.0043	1.075	2.695
Sb (mg/kg)	2.44	0.18	0.29	0.38	1.586	0.0043	0.85	1.590




Fig. 3: Anthropogenic and lithogenic portion of Fe, Mn and Sb in the agricultural soil of Hamedan area




Fig. 4: Spatial distribution of iron in the soil of Hamedan region

two indices of Muller’s geo-chemical index (I_{geo}) and Karbassi’s modified geochemical accumulation index (I_{poll}) is presented in Table 8, Results of comparison of heavy metals pollution in agricultural soil with Muller’s index and Karbassi’s modified index showed that iron and manganese fall within “no pollution- class” and antimony is classified as moderately polluted.

Fig. 2 shows the relationship amongst studied parameter. It should be pointed out that correlation

coefficients obtained on the basis of Pearson formulae are used in cluster analysis. Hence, the similar parameters are linked together. The confidence level is set as 0.05 and Student’s Test is indicative of meaningful values over 0.45. These is not any meaningful relationship amongst parameters within Hamedan soil. This could be probably be due to human influence on soil chemistry by using various compost and fertilizers. There is only a rather acceptable relationship between




Fig. 5: Spatial distribution of manganese in the soils of Hamedan region




Fig. 6: Spatial distribution of antimony in soil of Hamedan region

Fe and Sb that is indicative of the similar behavior in soil.

Table 9, shows the chemical fractionation of heavy metals in soil of Hamedan City. The results show that the highest concentrations of iron, manganese and antimony in the soil are related to the most resistant bonds and the lowest one is related to within lattice bonds. Fig. 3 shows the natural source and anthropogenic source of iron, manganese and antimony in soil. According to these curves, concentrations of all three metals are mostly affected by the parent materials.

In this study, Kriging method is applied on the studied elements to find out their spatial distribution (Figs. 4 to 6). According to the results, it was found that

the highest concentration of iron in the soil of study area (4.25 %) is detected in the north, southwest and partly in the west of Hamedan City (Fig. 4). By overlaying various maps such as zone, geological and land use of the study area, it was found that this part of Hamedan City has agricultural and pastures use and its bed rock is composed of igneous rocks, shale rocks, sandstone and sedimentary rocks. In fact, it might be concluded that the iron in soil of Hamedan City is mostly of geological source. Anthropogenic sources of iron in soil include industrial activities such as steel industries, dust and wastes of iron mines, iron sulfate used in herbicides and chemical fertilizers (De Ves *et al.*, 2005). Except

for use of herbicides and chemical fertilizers, these mentioned industries do not exist in the study area. Mico *et al.* (2006) applied multi-factor analysis in assessment of heavy metals sources in agricultural soils and concluded that iron concentration is controlled by the bed rock.

The highest concentration of manganese (about 457 mg/kg) in the soil of the study area is detected in the west of Hamedan City (Fig. 5). By overlaying the resulted zoning and geological and land use maps of the study area, it was found that this part of Hamedan City has mainly agricultural use and its bedrock is composed of igneous and metamorphic rocks. In fact, it may be said that manganese in soil of Hamedan City is mainly of geological source, however anthropogenic activities related to use of fertilizers and sewage in the agricultural lands may have led to increased amount of this heavy metal in the soil.

The highest concentration of antimony (about 3.2 mg/kg) in the soil of study area is detected in the southwest of Hamedan City (Fig. 6). By overlaying the zoning, geological and land use maps of the study area, it was found that the areas with high Sb concentrations coincide with cultivation pattern of the area. Its bedrock is composed of igneous, sandstone and shale rocks. In fact, it may be said that the antimony found in soil of Hamedan City is mainly of geological source, however the anthropogenic activities related to use of fertilizer and sewage in agricultural lands may have led to increased amount of this heavy metal in soil of Hamedan City. According to studies, concentration of antimony is naturally high in bedrocks of shale and igneous rocks (De Vos *et al.*, 2005). The greatest concentration of Sb ranges from 5.0 to 9.8 mg/kg and was seen in the form of two spots on igneous stones and shale bedrocks in Hamedan province. Overlaying of the distribution maps of Fe and Sb, land use maps, and geological strata of the studied area revealed that the distribution pattern of the elements did not conform to the land use pattern of the area. On the other hand, since the concentration of each element is naturally high in its bedrock (De Vos *et al.*, 2005), the most important factor affecting the concentration of the mentioned elements in soil must have been the geology (Soffianian *et al.*, 2014). As mentioned earlier, cluster analysis showed a rather good relationship between Fe and Sb.

The most important findings of visual adaptation of spatial distribution maps of heavy metals and land use

and geological maps of the area suggests that the main cause of high concentration of iron, antimony and manganese in the study area is the parent material of soil. However, the possible increase in concentration of these elements in agricultural land is not far from expectation due to presence of heavy metals in chemical fertilizers, pesticides and sewage. Therefore, given the hazards of heavy metals intake especially in the existing agricultural lands, it is recommended to prevent further distribution of heavy metals by managing the use of chemical fertilizers and pesticides and non-utilization of sewage for irrigation of lands as well as to avoid cultivation of plants possessing high potential of heavy metals intake in the polluted areas.

ACKNOWLEDGEMENT

Authors sincerely thank Dr. Nasrabadi, T. for his assistance in sample collection and analysis. Also thank to Dr. Tajziehchi, for her assistance in many parts of present investigation.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interest regarding the publication of this manuscript.

REFERENCES

- Akoto, O.; Ephraim, J.H.; Darko, G., (2008). Heavy Metals Pollution in Surface Soils in the Vicinity of Abundant Railway Servicing Workshop in Kumasi, Ghana, *Int. J. Environ. Res.*, 2(4): 359-364 (6 pages).
- Akoto, O.; Bismark Eshun, F.; Darko, G.; Adei, E., (2014). Concentrations and Health Risk Assessments of Heavy Metals in Fish from the Fosu Lagoon, *Int. J. Environ. Res.*, 8(2):403-410 (8 pages).
- Allison, L.E., (1965). Organic carbon. In: Black, C.A. (Ed.). *Methods of soil analysis. Part 2, Agronomy monograph*, ASA, Madison., 2(1):1367– 1389 (23 pages).
- Alhashemi, A.H.; Karbassi, A.R.; Kiabi, B.K.; Monavari, S.M.; Sekhvatjou, M.S. (2012). Bioaccumulation of trace elements in different tissues of three commonly available fish species regarding their gender, gonadosomatic index, and condition factor in a wetland ecosystem. *Environ. Monit. Assess.* 184 (4): 1865-1878 (14 pages)
- Anita, S.; Rajesh, K. S., (2010). Madhoolika Agrawal and Fiona M. Marshall; Risk assessment of heavy metal toxicity through contaminated vegetables from waste water irrigated area of Varanasi, India; *Tropical ecologist: Int. Soc. Trop. Ecol.*, 51(2): 375-387 (13 pages).
- Anthony, G.K.; Balwant, S.; Naveen, P. B., (2007). Heavy metal tolerance in common fern species. *Austral. J. Bot.*, 55, 63–73 (11 pages)
- AOAC, (2005). *Methods of analysis.*, Association of Official Analytical Chemists. Washington D.C.
- Aparna, C.; Saritha, P.; Himabindu, V.; Bhandari, A.; Anjaneyulu, Y., (2010). Evaluation of bioremediation effectiveness on

- sediments Contaminated with industrial wastes. *Int. J. Environ. Sci.*,1(4): 607-620 (**14 pages**).
- APHA., (2005). Standard methods for the examination of water and wastewater, 21st Ed., American Public Health Association, WWA and WEF, Washington~ D.C.
- ASTM., (2000). Standard test method for moisture, Ash and organic matter of peat and other organic soils, ASTM D2974-00.
- Biati, A.; Karbassi, A.R.; Hassani, A.H.; Monavari, S.M.; Moattar, F., (2010). Role of metal species in flocculation rate during estuarine mixing. *Int. J. Environ. Sci. Tech.*, 7(2): 327-336 (**10 pages**).
- Black, C.A., (1965). Methods of soil analysis, Part 2. Second Ed, Agron. Monogr. 9, ASA, Madison, WI.
- Chary, N.S.; Kamala, C.T.; Raj, D.S.S., (2008). Assessing risk of heavy metals from consuming food grown on sewage irrigated soils and food chain transfer. *Ecotox. Environ. Safe.*, 6(9): 513–524 (**12 pages**).
- Chen, Y.; Wang, C.; Wang, Z., (2005). Residues and source identification of per-sistent organic pollutants in farmland soils irrigated by effluents from biological treatment plants. *Environ. Int.*, 3(1): 778-783 (**6 pages**).
- De Miguel, E., (1997). Origin and patterns of distribution of trace elements in street dust : Unleaded petrol and urban lead. *Atmos. Environ.*, 31(17), 2733-2740 (**8 pages**).
- De Vos, W.; Batista, M.J.; Demetriades, A.; Duris , M.J.; Lexa, J.; Lis, J.; Sina, K.; Connor, P.J., (2005). Metallogenic Mineral Provinces and World Class Ore Deposits in Europe”, In: *Geochemical Atlas of Europe. Part 1 Background Information, Methodology and Maps*, Geological Survey of Finland. [Online] availability.
- Dewis, J.; Freitas, H., (1984). Physical and chemical methods of soil and water analysis. *FAO Soil Bulletin 10*, Oxford and IBH Publishing CO. PVT LTD. New Delhi.
- Facchinelli, A.; Sacchi, E.; mallen, L., (2001). Multivariate statistical and GIS-based approach to identify heavy metal sources in soils. *Environ. Pollut.*, 1(14): 313-324 (**12 pages**).
- FAO/WHO., (2011). Joint FAO/WHO food standards program codex committee on contaminants in foods, 5(1): 64-89 (**26 pages**).
- Franco-Uria, A.; Lopez-Mateo, C.; Roca , E.; Fernandez-Marcos, ML., (2009). Source identification of heavy metals in pastureland by multivariate analysis in NW Spain. *J Hazard Mater.*, 16(5): 1008–1015 (**8 pages**).
- Ghaderi, A.A., Abdulji, M.A., Karbassi, A.R., Nasrabadi, T., Khajeh, M., (2012). Evaluating the effects of fertilizers on bioavailable metallic pollution of soils. Case study: Sistan farms, Iran. *Int. J. Environ. Res.*, 6(2): 565-570 (**6 pages**).
- Gee, G.W.; Bauder, J.W ., (1986). Particle-size analysis. In: Klute, A. (Ed.), *Methods of soil analysis*. American Society of Agronomy. Agron. Monogr., 9(1) :383– 411(**29 pages**).
- Istvan, P.; Benton, J., (1997). Trace elements. Lucie Press. Boca Raton, Florida.
- Iyengar, V.; Nair, P., (2000). Global outlook on nutrition and the environment: meeting the challenges of the next millennium. *Sci. Total Environ.*, 24(9), 331-346 (**16 pages**).
- Jiries, A., (2003). Vehicular contamination of dust in Amman, Jordan. *Environmentalist*, 23(1): 205-210 (**6 pages**).
- Johnston, K.; VerHoef, J.M.; Krivoruchko, K.; Lucas, N., (2001). *Using ArcGIS geostatistical analyst*. ESRI Press, New York.
- Juang, K.W.; Lee, D.Y.; Ellsworth, T.R.,(2001). Using rank-order geostatistics for spatial interpolation of highly skewed data in heavy metal contaminated site. *J. Environ. Qual.* 30(1): 894-903 (**10 pages**).
- Kabata-Pendias, A.; Pendias, H., (1984). Trace elements in soils and plants. CRC, Press Boca Raton, FL.
- Karbassi, A.R., (1998). Geochemistry of Ni, Zn, Cu, Pb, Co, Cd, V, Mn, Fe, Al and Ca in sediments of North Western part of the Persian Gulf. *Int. J. Environ. Stud.*, 54(3), 205-212 (**8 pages**).
- Karbassi, A.R.; Monavari, S.M.; Bidhendi, G.R.N.; Nouri, J.; Nematpour, K., (2008). Metal pollution assessment of sediment and water in the Shur River. *Environ. Monit. Assess.* 147 (1-3): 107-116 (**10 pages**).
- Karbassi, A.R.; Nouri, J.; Ayaz, G.O., (2007). Flocculation of trace metals during mixing of Talar River water with Caspian seawater. *Int. J. Environ. Res.* 1 (1): 66-73 (**8 pages**).
- Karbassi, A.R.; Shankar, R., (2005). Geochemistry of two sediment cores from the west coast of India. *Int. J. Environ. Sci. Tech.*, 1(4), 307-316 (**10 pages**).
- Karbassi, A.R.; Amirnezhad, R., (2004). Geochemistry of heavy metals and sedimentation rate in a bay adjacent to the Caspian sea. *Int. J. Environ. Sci. Tech.*,1 (3), 199-206 (**8 pages**).
- Karbassi, A.R.; Torabi, F.; Ghazban, F.; Ardestani, M.,(2011). Association of trace metals with various sedimentary phases in dam reservoirs. *I.J.E.S.T.*,8(4), 841-852 (**12 pages**).
- Karbassi, A.R.; Heidari, M.; Vaezi, A.R.; Valikhani Samani, A.R., Fakhraee, M.; Heidari, F., (2014). Effect of pH and salinity on flocculation process of heavy metals during mixing of Aras River water with Caspian Sea water. *Environ. Earth Sci.*, 72(2): 457-465 (**9 pages**).
- Karbassi, A.R.; Nouri, J.; Nabi Bidhendi, G.R.; Ayaz, Gh.,(2008). Behavior of Cu, Zn, Pb, Ni and Mn during mixing of freshwater with the Caspian Sea water. *Desalination.*, 229(1): 118-124 (**7 pages**).
- Karbassi, A.R.; Monavari, S.M.; Nabi Bidhendi, Gh.R.; Nouri, J.; Nematpour, K., (2008). Metal pollution assessment of sediment and water in the Shur River. *Environ. Monit. Assess.*, 14(7): 107–116 (**10 pages**).
- Karbassi, A.R.; Tajziehchi, S.; Farhang Adib, F., (2016). Role of estuarine natural processes in removal of trace metals under emergency situations. *Global J. Environ. Sci. Manage.*, 2(1), 31-38 (**8 pages**).
- Karbassi, A.R.; Tajziehchi, S.; Afshar, S., (2015). An investigation on heavy metals in soils around oil field area. *Global J. Environ. Sci. Manage.*, 1(4), 275-282 (**8 pages**).
- Kargar, M.; Khorasani, N.A.; Karami, M.; Rafiee, G.H.; Naseh, R., (2012). An investigation on As, Cd, Mo and Cu contents of soils surrounding the Meyduk Tailings Dam. *Int. J. Environ. Res.*, 6(1):173-184 (**12 pages**).
- Khan, S.; Farooq, R.; Shahbaz, S.; Aziz Khan, M.; Sadique, M.,(2009). Health risk assessment of heavy metals for population via consumption of vegetables. *World Appl. Sci. J.*, 6 (12): 1602-1606 (**5 pages**).
- Lado, L.R.; Hengl, T.; Reuter, H.I., (2008). Heavy metals in European soils: A geostatistical analysis of the FOREGS Geochemical database. *Geoderma.*,14(8):189-199 (**11pages**).
- Lone, M.I.; He, Z-l.; Stoffella, P.J.; Yang, X-e., (2008). Phytoremediation of heavy metal polluted soils and water:

- progresses and perspectives. J. Zhejiang University Sci., 9 (3): 210–220 (11 pages).
- Mahmood, A.; Malik, R.N., (2014). Human health risk assessment of heavy metals via consumption of contaminated vegetables collected from different irrigation sources in Lahore, Pakistan, Arabian J. Chem.,7(1): 91-99 (9 pages).
- Manta, D.S.; Angelone, M.; Bellanca, A.; Neri,R.; Sprovieri, M., (2002). Heavy metals in urban soils: a case study from the city of Palermo, Sicily, Italy. Sci. Total Environ., 30(2):229–243(15 pages).
- McBride, M.; Sauve, S.; Hendershole, W., (1997). Solubility control of Cu, Zn, Cd and Pb in contaminated soils, Eur. J. Soil Sci.,48(2), 337-346 (10 pages).
- McGrath S.P., Shen Z.G., Zhao F.J., (1997). Heavy metal uptake and chemical changes in the rhizosphere of *Thlaspi caerulescens* and *Thlaspi ochroleucum* grown in contaminated soils. Plant Soil, 188: 153-159 (7 pages).
- Mico, C.; Recatala, L.; Peris, M.; Sanchez, J., (2006). Assessing heavy metal sources in agricultural soils of an European Mediterranean area by multivariate analysis. Chemosphere., 6(5): 863-872(10 pages).
- Mzoughi, N.; Chouba, L.,(2012). Heavy Metals and PAH Assessment based on mussel caging in the north coast of Tunisia (Mediterranean Sea), Int. J. Environ. Res., 6(1):109-118 (10 pages).
- Nasrabadi, T.; Nabi Bidhendi, G.; Karbassi, A.R.; Mehrdadi, N.,(2010). Partitioning of metals in sediments of the Haraz River: Southern Caspian Sea basin., Environ. Earth Sci. J., 59(5): 1111-1117 (7 pages).
- Nasrabadi, T.; Nabi Bidhendi, G.R.; Karbassi, A.R.; Grathwohl, P.; Mehrdadi, N., (2011).Impact of major organophosphate pesticides used in agriculture to surface water and sediment quality: Southern Caspian Sea basin, Haraz River, Environ. Earth Sci., 63(4): 873-883 (11 pages).
- Nelson, D.W.; Sommers, L.E ., (1975). A rapid and accurate method for estimating organic carbon in soil. Proceedings of the Indiana Academy of Science., 8(4): 456-462 (7 pages).
- Niencheski Hax, L.F., (2002). Lithium as a normalizer for the assessment of anthropogenic metal contamination of sediments of the southern area of Patos Lagoon. Aquatic Ecosystem Health & Management., 5(4), 473-483 (11 pages).
- Okunola, O.J.; Alhassan, Y.; Yapbella, G.G.; Uzairu, A.; Tsafe, A.I.; Embassey, E.; Abechi, S., (2011). Risk assessment of using mobile phone recharge cards in Nigeria. J. Environ. Chem. Ecotox., 3(4): 80-85 (6 pages).
- Praveena, S.M.; Ahmed, A.; Radojevic, M.; Abdullah, M.H.; Aris, A.Z., (2008). Heavy metals in mangrove surface sediment of Mengkabong Lagoon, Sabah: Multivariate and geo-accumulation index approaches, Int. J. Environ. Res., 2(2): 139-148 (10 pages).
- Rafiei,B.; Bakhtiari Nejad, M .; Hashemi, M.; Khodaei, A.S., (2010). Distribution of heavy metals around the Dashkasan Au Mine. Int. J. Environ. Res., 4(4): 647-654 (8 pages).
- Raj Shakya, P.; Malla Khwaounjoo,N., (2013). Heavy metal contamination in green leafy vegetables collected from different market sites of Kathmandu and their associated health risks, Sci. World., 11(11): 37-42 (6 pages).
- Ramlan, M.N.; Badri, M.A., (1989). Heavy metals in tropical city street dust and road side soils: A case of Kuala Lumpur, Malaysia. Environ. Tech. Lett., 10(1): 435-444 (10 pages).
- Rasmussen,P.E., Subramanian,K.S., Jessiman,B.J., (2001). A multi-element profile of house dust in relation to exterior dust and soils in the city of Ottawa. Canada. Sci. Total Environ., 2(67):125-140 (16 pages).
- Rattan, R.K.; Datta, S.P.; Chhonkar, P.K.; Suribabu, K.; Singh, A.K., (2005). Long-term impact of irrigation with sewage effluents on heavy metal content in soils, crops and groundwater: A case study. Agric. Ecosyst. Environ., 109(3): 310-322 (13 pages).
- Rodriguez, L.; Ruiz, E.; Azcarate ,J. A.; Rincon, J., (2009). Heavy metal distribution and chemical speciation in tailing and soils around a Pb-Zn mine in Spain. J. Environ. Manage., 9(1):1106-1116 (11 pages).
- Romic, M.; Hengl, T., Romic, D.; Husnjak, S ., (2007). Representing soil pollution by heavy metals using continuous limitation scores. Comput. Geosci., 3(3): 1316–1326 (11 pages).
- Sajjad, K.; Robina, F.; Shagufta, S.; Mohammad Aziz, K.; Maria, S., (2009). Health risk assessment of heavy metals for population via consumption of vegetables. World Appl. Sci. J., 6 (12): 1602-1606 (5 pages).
- Schwar, M.J.R., (1988). Baseline metal in dust concentrations in Greater London. Sci. Total Environ., 6(8):25-43 (19 pages).
- Serbaji, M.M.; Azri, C.; Medhioub, K., (2012). Anthropogenic contributions to heavy metal distributions in the surface and sub-surface sediments of the northern coast of Sfax, Tunisia, Int. J. Environ. Res., 6(3): 613-626 (14 pages).
- Shaw, A.J., (1989). Heavy metal tolerance in plants: Evolutionary aspects. CRC Press, Inc. Florida.,2(1): 299-305 (6 pages).
- Shetty, R.; Rajkumar, Sh., (2009). Biosorption of Cu (II) by metal resistant *pseudomonas* sp., Int. J. Environ. Res., 3(1):121-128 (8 pages).
- Singanam, M.; Singanam, V.; Abebaw, A., (2008). Biosorption of Cr (III) from aqueous solutions using indigenous biomaterial. Int. J. Environ. Res., 2(2): 177-182 (6 pages).
- Singh, K.P.; Mohan, D.; Sinha, S.; Dalwani, R.,(2004). Impact assessment of treated/ untreated wastewater toxicants discharged by sewage treatment plants on health, agricultural, and environmental quality in the wastewater disposal area. Chemosphere.,5(5): 227-255 (29 pages).
- Soffianian, A.; Sadat Madani, E.; Arabi, M., (2014). Risk assessment of heavy metal soil pollution through principal components analysis and false color composition in Hamedan Province, Iran, Environ. Syst. Res., 3: 3. doi:10.1186/2193-2697-3-3
- Spencer, K.L.; Macleod, C.L., (2002). Distribution and partitioning of heavy metals in stuarine sediment cores and implications for the use of sediment quality standard. J. Hydrol. Earth Syst. Sci., 6 (6) : 989-998 (10 pages).
- Supaphol, S.; Panichsakpatana, S.; Trakulnaleamsai, S.; Tungkananuruk, N.; Roughjanajirapa, P.; Gerard, O.; Donnell, A., (2006). The selection of mixed microbial inocula in environmental biotechnology: Example using petroleum contaminated tropical soils. J.Microbiol. Methods., 6(5): 432– 441 (10 pages).
- Taghinia Hejabi, A.; Basavarajappa, H.T.; Karbassi, A.R.; Monavari, S.M., (2011). Heavy metal pollution in water and sediments in the Kabini River, Karnataka, India. Environ. Monit. Assess.,182(4): 1-13 (13 pages).

- Tsafe, A.I.; Hassan, L.G.; Sahabi, D.M.; Alhassan, Y.; Bala, B.M., (2012). Evaluation of heavy metals uptake and risk assessment of vegetables grown in Yargalma of northern Nigeria. *J. Basic. Appl. Sci. Res.*, 2(7): 6708-6714 (7 pages).
- Turekian, K.K.; Wedepohl, K.H., (1961). Distribution of the elements in some major units of the earth's crust. *Geol. Soc. Am. Bull.*, 7(2): 75-92 (18 pages).
- Turkdogan, M.K.; Fevzi, K.; Kazim, K.; Ilyas, T.; Ismail, U., (2003). Heavy metals in soil, vegetables and fruits in the endemic upper gastrointestinal cancer region of Turkey. *Environ. Toxic. Pharmacol.*, 1(3):175-179 (5 pages).
- Uba, S.; Uzairu, A.; Okunola, O.J., (2009). Content of heavy metals in *lumbricus terrestris* associated soils in dump sites. *Int. J. Environ. Res.*, 3(3): 353-358 (6 pages).
- USDA., (2004). Soil survey laboratory methods manual. United States Department of Agriculture, USA.
- US-EPA IRIS., (2006). Integrated risk information system. United States, Environmental Protection Agency.
- USEPA., (1983). Hazardous waste land treatment. Office of Solid Waste and Emergency Response. SW-874, 273.

AUTHOR (S) BIOSKETCHES

Mohammadpour, G.A., Ph.D. Candidate, Department of Environmental Science, Graduate School of the Environment and Energy, Science and Research Branch, Islamic Azad University, Tehran, Iran. Email: g.mohammadpor@gmail.com

Karbassi, A.R., Ph.D., Associate Professor, Graduate Faculty of Environment, University of Tehran, P.O. Box 14135, Tehran, Iran. Email: akarbasi@ut.ac.ir

Baghvand, A., Ph.D., Associate Professor, Graduate Faculty of Environment, University of Tehran, P.O. Box 14135, Tehran, Iran. Email: baghvand@ut.ac.ir

How to cite this article:

Mohammadpour, G.A.; Karbassi, A.R.; Baghvand, A. (2016). Origin and spatial distribution of metals in agricultural soil. Global J. Environ. Sci. Manage., 2(2): 145-156.

DOI: [10.7508/gjesm.2016.02.005](https://doi.org/10.7508/gjesm.2016.02.005)

URL: http://gjesm.net/article_17689_1931.html

