

RESEARCH

Open Access



Orlicz-Garling sequence spaces of difference operator and their domination in Orlicz-Lorentz spaces

Charu Sharma¹, Syed Abdul Mohiuddine^{2*} , Kuldip Raj¹ and Ali H. Alkhaldi³

*Correspondence: mohiuddine@gmail.com
²Operator Theory and Applications Research Group, Department of Mathematics, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia
Full list of author information is available at the end of the article

Abstract

We introduce new classes of generalized Orlicz-Garling sequences and Orlicz-Lorentz sequences by using a sequence of Orlicz functions and difference operator. We show that the Orlicz-Garling sequence space admits a unique 1-subsymmetric basis and a 1-dominated block basic sequence in $g(\mathcal{M}, \Delta^{(m)}, \nu, p)$. We also make an effort to prove that every symmetric normalized block Orlicz-Garling sequence dominates an Orlicz-Lorentz sequence. Finally, we study some geometric properties of these spaces and establish some inclusion relations between spaces.

MSC: 46B25; 46B45; 41A65

Keywords: Banach space; Lorentz sequence space; Garling sequence space; Orlicz function; Subsymmetric bases; Symmetric bases

1 Introduction and preliminaries

Let c_{00} be the space of all scalar sequences, that is, elements of $\mathbb{R}^{\mathbb{N}}$ or $\mathbb{C}^{\mathbb{N}}$ with finitely many nonzero entries. Let X and Y be infinite-dimensional Banach spaces. If $(x_k)_{k=1}^{\infty}$ is a basis of the Banach space X and $x = \sum_{k=1}^{\infty} a_k x_k$, then we denote $\text{supp } x = \{k \in \mathbb{N} : a_k \neq 0\}$, that is, the set of indices corresponding to the nonzero entries of x , which is called the support of x . When the basis $(x_k)_{k=1}^{\infty}$ is clear from context, we write $x = (a_k)_{k=1}^{\infty}$ instead of $x = \sum_{k=1}^{\infty} a_k x_k$.

Now, suppose that $(x_k)_{k=1}^{\infty}$ and $(y_k)_{k=1}^{\infty}$ are sequences in X and Y , respectively. If there exists $C \in [1, \infty)$ such that

$$\left\| \sum_{k=1}^{\infty} a_k x_k \right\|_X \leq C \left\| \sum_{k=1}^{\infty} a_k y_k \right\|_Y$$

for all finitely supported sequences $(a_k)_{k=1}^{\infty} \in c_{00}$, then $(y_k)_{k=1}^{\infty}$ C -dominates $(x_k)_{k=1}^{\infty}$, and we can write $(x_k)_{k=1}^{\infty} \lesssim_C (y_k)_{k=1}^{\infty}$. In case C does not matter, we will simply say that $(y_k)_{k=1}^{\infty}$ dominates $(x_k)_{k=1}^{\infty}$ and $(x_k)_{k=1}^{\infty} \lesssim (y_k)_{k=1}^{\infty}$. If $(x_k)_{k=1}^{\infty} \lesssim_C (y_k)_{k=1}^{\infty}$ and $(y_k)_{k=1}^{\infty} \lesssim_C (x_k)_{k=1}^{\infty}$, then $(x_k)_{k=1}^{\infty}$ and $(y_k)_{k=1}^{\infty}$ are C -equivalent, and we can write it as $(x_k)_{k=1}^{\infty} \approx_C (y_k)_{k=1}^{\infty}$.

A basic sequence $(x_k)_{k=1}^{\infty}$ is called subsymmetric just in case it is unconditional and equivalent to each of its subsequences. It is called symmetric whenever it is unconditional and equivalent to each of its permutations. Lindenstrauss and Tzafriri [1] studied the uniform boundedness which means that if $(x_k)_{k=1}^{\infty}$ is a subsymmetric basic sequence, then

there is a uniform constant $C \geq 1$ such that if $(x_{k_n})_{n=1}^\infty$ is any subsequence and $(\varepsilon_n)_{n=1}^\infty$ is any sequence of signs, then $(x_k)_{k=1}^\infty$ is C -equivalent to $(\varepsilon_n x_{k_n})_{n=1}^\infty$. In this case, we say that $(x_k)_{k=1}^\infty$ is C -subsymmetric. Similarly, if $(x_k)_{k=1}^\infty$ is symmetric, then there is $C \geq 1$ such that $(x_k)_{k=1}^\infty$ is C -equivalent to each $(\varepsilon_k x_{\varrho(k)})_{k=1}^\infty$, where ϱ is a permutation of \mathbb{N} , and in this case, we say that $(x_k)_{k=1}^\infty$ is C -symmetric. Note that C -symmetry implies C -subsymmetry, which in turn implies C -unconditionality. In this paper, we use standard facts and notation from Banach spaces and approximation theory. For necessary background, see [1, 2], and references therein. We denote by \mathbb{F} the real or complex field. The canonical basis of \mathbb{F} is denoted by $(e_k)_{k=1}^\infty$, that is, $e_k = (\delta_{k,n})_{n=1}^\infty$, where $\delta_{k,n} = 1$ if $n = k$ and $\delta_{k,n} = 0$ if $n \neq k$.

An Orlicz function M is a function that is continuous, nondecreasing, and convex with $M(0) = 0$, $M(x) > 0$ for $x > 0$ and $M(x) \rightarrow \infty$ as $x \rightarrow \infty$. An Orlicz function M is said to satisfy Δ_2 -condition if there exists $R > 0$ such that $M(2u) \leq RM(u)$, $u \geq 0$.

The study of Orlicz sequence spaces was initiated with a certain specific purpose in Banach space theory. Lindenstrauss and Tzafriri [3] used the idea of an Orlicz function to define the sequence space

$$\ell_M = \left\{ x \in w : \sum_{k=1}^\infty M\left(\frac{|x_k|}{\rho}\right) < \infty \text{ for some } \rho > 0 \right\},$$

which is called an Orlicz sequence space. The space ℓ_M is a Banach space with the norm

$$\|x\| = \inf \left\{ \rho > 0 : \sum_{k=1}^\infty M\left(\frac{|x_k|}{\rho}\right) \leq 1 \right\}.$$

It is shown in [3] that every Orlicz sequence space ℓ_M contains a subspace isomorphic to ℓ_p ($p \geq 1$).

A sequence $\mathcal{M} = (M_k)$ of Orlicz functions is called a *Musielak-Orlicz function* (see [4, 5]). For more detail about sequence spaces, see [6–26], and references therein.

The notion of a difference operator in the sequence spaces was first introduced by Kizmaz [27]. The idea of difference sequence spaces of Kizmaz was further generalized by Et and Çolak [28]. Later, this concept was studied by Bektaş et al. [29] and Et et al. [30]. Now, the difference matrix $\Delta = \delta_{nk}$ defined by

$$\delta_{nk} = \begin{cases} (-1)^{n-k} & (n - 1 \leq k \leq n), \\ 0 & (0 \leq k < n - 1 \text{ or } n > k). \end{cases}$$

The difference operator of order m is defined by $\Delta^m : w \rightarrow w$, $(\Delta^1 x)_k = (x_k - x_{k-1})$ and $\Delta^m x = (\Delta^1 x)_k \circ (\Delta^{m-1} x)_k$ for $m \geq 2$.

The triangle matrix $\Delta^{(m)} = \delta_{nk}^{(m)}$ defined by

$$\delta_{nk}^{(m)} = \begin{cases} (-1)^{n-k} \binom{m}{n-k} & (\max\{0, n - m\} \leq k \leq n), \\ 0 & (0 \leq k < \max\{0, n - m\} \text{ or } n > k) \end{cases}$$

for all $k, n \in \mathbb{N}$ and any fixed $m \in \mathbb{N}$.

The Lorentz sequence space was introduced by Lorentz [31, 32]. This space plays an important role in the theory of Banach spaces, whereas Garling [33] studied, for $v = (k^{-1/2})_{k=1}^\infty$,

the canonical unit vectors of $g(v, 1)$ from a subsymmetric basic sequence that is not symmetric. Pujara [34] studied $g(v, 2)$ for $v = (k^{-1/2})_{k=1}^\infty$ and showed that these spaces are uniformly convex and that their canonical basis is subsymmetric but not symmetric. The space from [35] can be viewed as being constructed from variations of $g(w, 1)$ for various choices of weights and gives a canonical basis that is 1-greedy and subsymmetric but not symmetric. Both the Garling sequence spaces and the Lorentz sequence spaces are defined by taking the completion of c_{00} under the norms $\|\cdot\|_g$ or $\|\cdot\|_d$, respectively. The only difference between these norms is that $\|\cdot\|_g$ is defined by taking a certain supremum over subsequences instead of permutations of sequences as is the case for $\|\cdot\|_d$. Wallis [36] generalized the construction of Garling for each $1 \leq p < \infty$ and normalized nonincreasing sequence of positive numbers that exhibited complementably homogeneous Banach Garling space related to the Lorentz sequence space. For more detail about these spaces, see [37–39], and references therein.

Let us consider the set of weights

$$V = \{(v_k)_{k=1}^\infty \in c_0 \setminus l_1 : 1 = v_1 \geq v_2 > \dots \geq v_k \geq v_{k+1} \geq \dots > 0\}.$$

Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions, and let $v = (v_k)_{k=1}^\infty \in V$ be a weight, that is, a sequence of positive scalars. Then the Orlicz-Garling sequence space is defined as the Banach space consisting of all scalar sequences $A = (a_k)_{k=1}^\infty$ such that

$$\|A\|_{g(\mathcal{M}, \Delta^{(m)}, v, p)} = \inf \left\{ \rho > 0 : \sup_{\psi \in \mathcal{O}} \left(\sum_{k=1}^\infty \left(M_k \left(\frac{|\Delta^{(m)} a_{\psi(k)}|^p v_k}{\rho} \right) \right) \right)^{\frac{1}{p}} \leq 1 \text{ for some } \rho > 0 \right\},$$

where \mathcal{O} denotes the set of all increasing functions from \mathbb{N} to \mathbb{N} . We will assume that v is normalized, that is, $v_1 = 1$. We consider the normed space $g(\mathcal{M}, \Delta^{(m)}, v, p) = \{A = (a_k)_{k=1}^\infty : \|A\|_{g(\mathcal{M}, \Delta^{(m)}, v, p)} < \infty\}$.

Let Π denote the set of permutations on \mathbb{N} , and let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions. Then, for any $1 \leq p < \infty$ and $v \in V$, we define the Orlicz-Lorentz sequence space $d(\mathcal{M}, \Delta^{(m)}, v, p)$ as the completion of c_{00} under the norm $\|\cdot\|_{d(\mathcal{M}, \Delta^{(m)}, v, p)}$ defined by

$$\|A\|_{d(\mathcal{M}, \Delta^{(m)}, v, p)} = \inf \left\{ \rho > 0 : \sup_{\varrho \in \Pi} \left(\sum_{k=1}^\infty \left(M_k \left(\frac{|\Delta^{(m)} a_{\varrho(k)}|^p v_k}{\rho} \right) \right) \right)^{\frac{1}{p}} \leq 1 \text{ for some } \rho > 0 \right\}.$$

We consider the normed space $d(\mathcal{M}, \Delta^{(m)}, v, p) = \{A = (a_k)_{k=1}^\infty : \|A\|_{d(\mathcal{M}, \Delta^{(m)}, v, p)} < \infty\}$.

The main aim of this paper is to introduce and study some difference Orlicz-Garling sequence spaces and Orlicz-Lorentz sequence spaces. Using the originally developed Orlicz-Lorentz sequence space, we show that the Orlicz-Garling sequence space admits a unique 1-subsymmetric basis. Finally, we discuss some additional geometric properties of $g(\mathcal{M}, \Delta^{(m)}, v, p)$ and also establish some inclusion relations between these spaces.

2 Main results on $g(\mathcal{M}, \Delta^{(m)}, v, p)$

Given a function ψ , we denote by $U(\psi)$ its range. Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions, and let $v = (v_k)_{k=1}^\infty \in V$ be a weight. Let \mathcal{O}_A be the set of increasing functions from an integer interval $[1, \dots, u] \cap \mathbb{N}$ into \mathbb{N} . For a given $\psi \in \mathcal{O}_A$, we denote by $u(\psi)$ the largest integer in its domain; a function in \mathcal{O}_A is univocally determined by its range. For $A = (a_k)_{k=1}^\infty$, we have

$$\|A\|_{g(\mathcal{M}, \Delta^{(m)}, v, p)} = \inf \left\{ \rho > 0 : \sup_{\psi \in \mathcal{O}_A} \sum_{k=1}^{u(\psi)} \left(M_k \left(\frac{|\Delta^{(m)} a_{\psi(k)}|^p v_k}{\rho} \right) \right) \leq 1 \right. \\ \left. \text{for some } \rho > 0 \right\}. \tag{2.1}$$

A Banach space with an unconditional basis is said to have a unique unconditional if any two seminormalized unconditional bases of X are equivalent.

Now we define the weak Orlicz-Lorentz sequence space $d_\infty(\mathcal{M}, \Delta^{(m)}, v, p)$ for a sequence of Orlicz functions $\mathcal{M} = (M_k)$. For $v \in V$ and $1 \leq p < \infty$, it consists of all sequences $A = (a_k)_{k=1}^\infty \in c_0$ such that

$$\|A\|_{d_\infty(\mathcal{M}, \Delta^{(m)}, v, p)} = \inf \left\{ \rho > 0 : \sup_k \left(\sum_{n=1}^k \left(M_k \left(\frac{|\Delta^{(m)} a_k^*| v_n}{\rho} \right) \right) \right)^{\frac{1}{p}} \leq 1 \text{ for some } \rho > 0 \right\},$$

where $(a_k^*)_{k=1}^\infty$ denotes the decreasing rearrangement of A .

Let $(x_k)_{k=1}^\infty$ be a basis for a Banach space X , and let $D = \sum_{k=1}^\infty b_k x_k$ be a vector in X . For any $A = \sum_{k=1}^\infty a_k x_k \in X$, it can be written as $A < D$ if

- (i) $\text{supp}(A) \subseteq \text{supp}(D)$,
- (ii) $a_k = b_k$ for all $k \in \text{supp}(A)$, and
- (iii) $\|A\| = \|D\|$.

We can say that D is minimal in X if it is minimal in X equipped with the partial order $<$, that is, if $A < D$ implies $A = D$. If D is supported with respect to a basis $(x_k)_{k=1}^\infty$, then there exists a minimal $A < D$.

A block basic sequence of a basic sequence $(x_k)_{k=1}^\infty$ in a Banach space X is a sequence $(y_k)_{k=1}^\infty$ of nonzero vectors of the form

$$y_k = \sum_{j=p_k}^{p_{k+1}-1} a_j x_j$$

for some increasing sequence of integers $(p_k)_{k=1}^\infty$ with $p_1 = 1$ and some $(a_k)_{k=1}^\infty \in \mathbb{F}^\mathbb{N}$.

Theorem 2.1 *Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions, and let $v = (v_k)_{k=1}^\infty \in V$ be a weight. For $1 \leq p < \infty$, we have $l_p \subsetneq d(\mathcal{M}, \Delta^{(m)}, v, p) \subseteq g(\mathcal{M}, \Delta^{(m)}, v, p) \subseteq d_\infty(\mathcal{M}, \Delta^{(m)}, v, p)$, with norm inclusions.*

Proof It is clear that $l_p \subsetneq d(\mathcal{M}, \Delta^{(m)}, v, p)$. Let $\psi \in \mathcal{O}_A$, $A = (a_k)_{k=1}^\infty \in \mathbb{F}^\mathbb{N}$, and $\varrho \in \Pi$ extending ψ . Then we have

$$\begin{aligned} \sum_{k=1}^{t(\psi)} \left(M_k \left(\frac{|\Delta^{(m)} a_{\psi(k)}|^p v_k}{\rho} \right) \right) &= \sum_{k=1}^{t(\psi)} \left(M_k \left(\frac{|\Delta^{(m)} a_{\varrho(k)}|^p v_k}{\rho} \right) \right) \\ &\leq \sum_{k=1}^\infty \left(M_k \left(\frac{|\Delta^{(m)} a_{\varrho(k)}|^p v_k}{\rho} \right) \right) \\ &\leq \|A\|_{d(\mathcal{M}, \Delta^{(m)}, v, p)}. \end{aligned}$$

Taking the supremum on ψ , we get $\|A\|_{g(\mathcal{M}, \Delta^{(m)}, v, p)} \leq \|A\|_{d(\mathcal{M}, \Delta^{(m)}, v, p)}$. Let $A = (a_k)_{k=1}^\infty \in c_{00}$. Given $t(\psi) = k$ and $|a_{\psi(n)}| \geq |a_k^*|$ for $1 \leq n \leq t(\psi) = k$, there exists $\psi \in \mathcal{O}_A$ such that, for a given $k \in \mathbb{N}$, we have

$$\left(\sum_{n=1}^k \left(M_k \left(\frac{|\Delta^{(m)} a_k^*| v_n}{\rho} \right) \right) \right)^{\frac{1}{p}} \leq \left(\sum_{n=1}^k \left(M_k \left(\frac{|\Delta^{(m)} a_{\varrho(n)}|^p v_n}{\rho} \right) \right) \right)^{\frac{1}{p}} \leq \|A\|_{d(\mathcal{M}, \Delta^{(m)}, v, p)}.$$

Therefore, $\|A\|_{d_\infty(\mathcal{M}, \Delta^{(m)}, v, p)} \leq \|A\|_{g(\mathcal{M}, \Delta^{(m)}, v, p)}$, which concludes the proof. □

Let us define some linear functions as follows:

- (i) For a given sequence of signs $\varepsilon = (\varepsilon_k)_{k=1}^\infty$, we define the linear mapping $T_\varepsilon : \mathbb{F}^\mathbb{N} \rightarrow \mathbb{F}^\mathbb{N}$ by $T_\varepsilon((a_k)_{k=1}^\infty) = (\varepsilon_k a_k)_{k=1}^\infty$.
- (ii) The coordinate projection on $B \subseteq \mathbb{N}$ is defined by $P_B : \mathbb{F}^\mathbb{N} \rightarrow \mathbb{F}^\mathbb{N}$, $P_B((a_k)_{k=1}^\infty) = (\mu_k a_k)_{k=1}^\infty$, where

$$\mu_k = \begin{cases} 1 & \text{if } k \in B, \\ 0 & \text{if } k \notin B. \end{cases}$$

- (iii) Given $\psi \in \mathcal{O}$, we define the linear mapping $S_\psi : \mathbb{F}^\mathbb{N} \rightarrow \mathbb{F}^\mathbb{N}$ by $S_\psi((a_k)_{k=1}^\infty) = (a_{\psi(k)})_{k=1}^\infty$, and
- (iv) we define the linear function $R_\psi : \mathbb{F}^\mathbb{N} \rightarrow \mathbb{F}^\mathbb{N}$ by $R_\psi((a_k)_{k=1}^\infty) = (b_k)_{k=1}^\infty$, where

$$b_k = \begin{cases} a_n & \text{if } k = \psi(n), \\ 0 & \text{if } k \notin \psi(n). \end{cases}$$

Remark 2.2 Given $k \in \mathbb{N}$ and $\psi \in \mathcal{O}$, we have $R_\psi \circ S_\psi = Id_{\mathbb{F}^\mathbb{N}}$ and $S_\psi(e_k) = e_{\psi(k)}$.

Theorem 2.3 *Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions, let $v = (v_k)_{k=1}^\infty \in V$ be a weight, and let $1 \leq p < \infty$. Let $\varepsilon = (\varepsilon_k)_{k=1}^\infty$ be a sequence of signs. Let B be a subset of \mathbb{N} , and let ψ be a linear map in \mathcal{O} . Then we have the following:*

- (i) T_ε and P_B are norm-one operators from $g(\mathcal{M}, \Delta^{(m)}, v, p)$ into $g(\mathcal{M}, \Delta^{(m)}, v, p)$.
- (ii) S_ψ and R_ψ are norm-one operators from $g(\mathcal{M}, \Delta^{(m)}, v, p)$ into $g(\mathcal{M}, \Delta^{(m)}, v, p)$.
- (iii) S_ψ is an isometry from $g(\mathcal{M}, \Delta^{(m)}, v, p)$ into $g(\mathcal{M}, \Delta^{(m)}, v, p)$.
- (iv) The standard unit vectors form a 1-subsymmetric basic sequence in $g(\mathcal{M}, \Delta^{(m)}, v, p)$.

Proof The proof of (i) is clear since it is a consequence of (ii) and Remark 2.2. (iv) is a consequence of (i), (iii), and Remark 2.2. Therefore, we only prove (ii). Let $A = (a_k)_{k=1}^\infty \in \mathbb{F}^N$. It is clear that $\|T_\varepsilon(A)\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)} = \|A\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)}$ and $\|P_B(A)\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)} = \|A\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)}$. Since $\psi \circ \varphi \in \mathcal{O}$, we have

$$\begin{aligned} \|S_\psi(A)\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)} &= \inf \left\{ \rho > 0 : \sup_{\varphi \in \mathcal{O}} \sum_{k=1}^\infty \left(M_k \left(\frac{|\Delta^{(m)} a_{\psi(\varphi(k))}|^p \nu_k}{\rho} \right) \right) \leq 1 \text{ for some } \rho > 0 \right\} \\ &\leq \|A\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)}. \end{aligned}$$

Let us take $\varphi \in \mathcal{O}$ and $R_\psi(A) = (b_k)_{k=1}^\infty$. As we know, $\psi^{-1} \circ \varphi$ is an increasing function from a set $B \subseteq \mathbb{N}$ to \mathbb{N} . Let $\phi : L \rightarrow B$ be increasing and bijective, where $L = \{k \in \mathbb{N} : k \leq |B|\}$. Since $\xi : \psi^{-1} \circ \varphi \circ \phi \in \mathcal{O} \cup \mathcal{O}_A$ and $k \leq \phi(k)$ for all $k \in L$, we have

$$\begin{aligned} \sum_{k=1}^\infty \left(M_k \left(\frac{|\Delta^{(m)} b_{\psi(\varphi(k))}|^p \nu_k}{\rho} \right) \right) &= \sum_{k \in B} \left(M_k \left(\frac{|\Delta^{(m)} a_{\psi^{-1} \circ \varphi(k)}|^p \nu_k}{\rho} \right) \right) \\ &= \sum_{k \in L} \left(M_k \left(\frac{|\Delta^{(m)} a_{\xi(k)}|^p \nu_{\phi(k)}}{\rho} \right) \right) \\ &\leq \sum_{k \in L} \left(M_k \left(\frac{|\Delta^{(m)} a_{\xi(k)}|^p \nu_k}{\rho} \right) \right) \leq \|A\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)}. \end{aligned}$$

Taking the supremum on φ , we get $\|R_\psi(A)\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)} \leq \|A\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)}$. □

Theorem 2.4 *Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions, let $\nu = (\nu_k)_{k=1}^\infty \in V$ be a weight, and let $1 \leq p < \infty$. Suppose $A \in c_{00}$ is minimal in $g(\mathcal{M}, \Delta^{(m)}, \nu, p)$ with respect to the canonical basis. Then*

$$\begin{aligned} \|A\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)} &= \inf \left\{ \rho > 0 : \left(\sum_{k=1}^{u(\varphi)} \left(M_k \left(\frac{|\Delta^{(m)} a_{\varphi(k)}|^p \nu_k}{\rho} \right) \right) \right)^{\frac{1}{p}} \leq 1, \right. \\ &\quad \left. \text{for some } \rho > 0 \right\}, \end{aligned} \tag{2.2}$$

where $\varphi \in \mathcal{O}_A$ is determined by $U(\varphi) = \text{supp}(A)$.

Proof For given $\psi \in \mathcal{O}_A$, let φ be the function in \mathcal{O}_A determined by $U(\varphi) = U(\psi) \cap \text{supp}(A)$. Letting ϕ be the inverse function of ψ restricted to $U(\varphi)$, we have

$$\begin{aligned} \sum_{k=1}^{u(\psi)} \left(M_k \left(\frac{|\Delta^{(m)} a_{\psi(k)}|^p \nu_k}{\rho} \right) \right) &= \sum_{k \in U(\varphi)} \left(M_k \left(\frac{|\Delta^{(m)} a_k|^p \nu_{\phi(k)}}{\rho} \right) \right) \\ &\leq \sum_{k \in U(\varphi)} \left(M_k \left(\frac{|\Delta^{(m)} a_k|^p \nu_{\varphi^{-1}(k)}}{\rho} \right) \right) \\ &= \sum_{k \in u(\varphi)} \left(M_k \left(\frac{|\Delta^{(m)} a_{\varphi(k)}|^p \nu_k}{\rho} \right) \right). \end{aligned}$$

Therefore, we can restrict it to $\{\psi \in \mathcal{O}_A : U(\psi) \subseteq \text{supp}(A)\}$, so that supremum (2.1) is attained. Let $\varphi \in \mathcal{O}_A$ with $U(\varphi) \subseteq \text{supp}(A)$, and let H be the projection of A onto $U(\varphi)$. Then, we have

$$\begin{aligned} \|A\|_g^{\mathcal{M}, \Delta^{(m)}, p} &= \sum_{k \in u(\varphi)} \left(M_k \left(\frac{|\Delta^{(m)} a_{\varphi(k)}|^p v_k}{\rho} \right) \right) \\ &\leq \|H\|_g^{\mathcal{M}, \Delta^{(m)}, p}. \end{aligned}$$

Thus, $H = A$ and $U(\varphi) = \text{supp}(A)$. □

Theorem 2.5 *Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions, let $v = (v_k)_{k=1}^\infty \in V$ be a weight, and let $1 \leq p < \infty$. Every normalized block basic sequence of the canonical basis of $g(\mathcal{M}, \Delta^{(m)}, v, p)$ is 1-dominated by the canonical basis of l_p .*

Proof Let a normalized block basic sequence be $y_k = \sum_{j=p_k}^{p_{k+1}-1} a_j g_j$, $k \in \mathbb{N}$. Let us define $k = k(j) \in \mathbb{N}$ by $p_k \leq j \leq p_{k+1} - 1$ for $j \in \mathbb{N}$. Let $(c_j)_{j=1}^\infty = \sum_{k=1}^\infty b_k y_k$ and $(b_k)_{k=1}^\infty \in c_{00}$. Let $\psi \in \mathcal{O}$, and let for each $k \in \mathbb{N}$, $L_k = \{j \in \mathbb{N} : p_k \leq \psi(j) \leq p_{k+1} - 1\}$ be an increasing interval. Then, for some $s_k, u_k \in \mathbb{N} \cup \{0\}$, $L_k = \{j \in \mathbb{N} : 1 + s_k \leq i \leq s_k + u_k\}$. We obtain

$$\begin{aligned} &\sum_{k=1}^\infty \left(M_k \left(\frac{|\Delta^{(m)} c_{\psi(k)}|^p v_k}{\rho} \right) \right) \\ &\leq \sum_{k=1}^\infty \left(M_k \left(\frac{|\Delta^{(m)} b_k|^p}{\rho} \right) \right) \sum_{j \in L_k} \left(M_k \left(\frac{|\Delta^{(m)} a_{\psi(j)}|^p v_j}{\rho} \right) \right) \\ &= \sum_{k=1}^\infty \left(M_k \left(\frac{|\Delta^{(m)} b_k|^p}{\rho} \right) \right) \sum_{i=1}^{u_k} \left(M_k \left(\frac{|\Delta^{(m)} a_{\psi(i+s_k)}|^p v_{i+s_k}}{\rho} \right) \right) \\ &\leq \sum_{k=1}^\infty \left(M_k \left(\frac{|\Delta^{(m)} b_k|^p}{\rho} \right) \right) \sum_{i=1}^{u_k} \left(M_k \left(\frac{|\Delta^{(m)} a_{\psi(i+s_k)}|^p v_i}{\rho} \right) \right) \\ &\leq \sum_{k=1}^\infty \left(M_k \left(\frac{|\Delta^{(m)} b_k|^p}{\rho} \right) \right) \|y_k\|_g^{\mathcal{M}, \Delta^{(m)}, p} \\ &= \sum_{k=1}^\infty \left(M_k \left(\frac{|\Delta^{(m)} b_k|^p}{\rho} \right) \right). \end{aligned}$$

Taking the supremum on ψ , we get the following result. □

Theorem 2.6 *Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions, let $v = (v_k)_{k=1}^\infty \in V$ be a weight, and let $1 \leq p < \infty$. Let $(g_k)_{k=1}^\infty$ denote the canonical basis of $g(\mathcal{M}, \Delta^{(m)}, v, p)$. Suppose that $y_k = \sum_{j=p_k}^{p_{k+1}-1} a_j g_j$, $k \in \mathbb{N}$, forms a block basic sequence in $g(\mathcal{M}, \Delta^{(m)}, v, p)$. Then there exists another block basic sequence formed by $\hat{y}_k = \sum_{q=\hat{p}_k}^{\hat{p}_{k+1}-1} a_{j_q} g_q$, $k \in \mathbb{N}$, such that*

- (i) $a_{j_q} \neq 0$ for $q \in \mathbb{N}$;
- (ii) $\hat{p}_1 = 1$;
- (iii) $(j_q)_{q=\hat{p}_k}^{\hat{p}_{k+1}-1}$ is a subsequence of $(j)_{j=p_k}^{p_{k+1}-1}$ for each $k \in \mathbb{N}$;

(iv)

$$\begin{aligned} \|y_k\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)} &= \|\hat{y}_k\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)} \\ &= \inf \left\{ \rho > 0 : \left(\sum_{q=\hat{p}_k}^{\hat{p}_{k+1}-1} \left(M_k \left(\frac{|\Delta^{(m)} a_{jq}|^p \nu_{q-\hat{p}_{k+1}}}{\rho} \right) \right) \right)^{\frac{1}{p}} \leq 1 \right. \\ &\quad \left. \text{for some } \rho > 0 \text{ and for each } k \in \mathbb{N} \right\}; \end{aligned}$$

(v) $(\hat{y}_k)_{k=1}^\infty$ is 1-dominated by $(y_k)_{k=1}^\infty$.

Proof Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions, let $\nu = (\nu_k)_{k=1}^\infty \in V$ be a weight, and let $1 \leq p < \infty$. Now, for each $k \in \mathbb{N}$, we have $B_k \subseteq \{p_k, \dots, p_{k+1} - 1\}$ such that if $m(j) = \#\{u \in B_k : u \leq j\}$ for each $p_k \leq j \leq p_{k+1} - 1$, then

$$\begin{aligned} \|y_k\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)} &= \inf \left\{ \rho > 0 : \left(\sum_{q=\hat{p}_k}^{\hat{p}_{k+1}-1} \left(M_k \left(\frac{|\Delta^{(m)} a_{jq}|^p \nu_{q-\hat{p}_{k+1}}}{\rho} \right) \right) \right)^{\frac{1}{p}} \leq 1 \text{ for some } \rho > 0 \right\}. \end{aligned}$$

We suppose that $a_j \neq 0$ whenever $j \in B_k$ for some $k \in \mathbb{N}$. Therefore, for each $p_k \leq j \leq p_{k+1} - 1$, we define

$$\tilde{a}_j = \begin{cases} a_j & \text{if } j \in B_k, \\ 0 & \text{if } j \notin B_k. \end{cases}$$

Let $\tilde{y}_k = \sum_{j=p_k}^{p_{k+1}-1} \tilde{a}_j g_j$ in $g(\mathcal{M}, \Delta^{(m)} \nu, p)$ for each $k \in \mathbb{N}$. Then $(\tilde{y}_k)_{k=1}^\infty$ is a normalized block sequence in $g(\mathcal{M}, \Delta^{(m)}, \nu, p)$, which is 1-dominated by $(y_k)_{k=1}^\infty$.

By defining the sequences $(a_{jq})_{q=1}^\infty$ and $(\hat{p}_k)_{k=1}^\infty$ we shall construct the block basic sequence $(\hat{y}_k)_{k=1}^\infty$. Suppose $(\tilde{a}_{jq})_{q=1}^\infty$ is the subsequence that consists precisely of the nonzero values of $(\tilde{a}_j)_{j=1}^\infty$. It is clear that (i) holds and there exist $1 = \hat{p}_1 < \hat{p}_2 < \dots \in \mathbb{N}$ such that (ii) and (iii) also hold. Now, we find that $\|y_k\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)} = \|\tilde{y}_k\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)} = \|\hat{y}_k\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)}$, which yields (iv). Finally, we have $(\hat{y}_k)_{k=1}^\infty \approx_1 (\tilde{y}_k)_{k=1}^\infty \lesssim_1 (y_k)_{k=1}^\infty$ and get (v). \square

Theorem 2.7 *Suppose $\mathcal{M} = (M_k)$ is a sequence of Orlicz functions, $\nu = (\nu_k)_{k=1}^\infty \in V$ is a weight, and $1 \leq p < \infty$. Let $(d_k)_{k=1}^\infty$ denote the canonical basis of $d(\mathcal{M}, \Delta^{(m)}, \nu, p)$ and $(g_k)_{k=1}^\infty$ denote the canonical basis of $g(\mathcal{M}, \Delta^{(m)}, \nu, p)$. Then every symmetric normalized block sequence of $(g_k)_{k=1}^\infty$ dominates $(d_k)_{k=1}^\infty$.*

Proof Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions, let $\nu = (\nu_k)_{k=1}^\infty \in V$ be a weight, and let $1 \leq p < \infty$. Take any symmetric normalized block sequence $(y_k)_{k=1}^\infty$ of $(g_k)_{k=1}^\infty$ in

$g(\mathcal{M}, \Delta^{(m)}, v, p)$. For any finite sequence $(b_k)_{k=1}^K$ for $K \in \mathbb{N}$, we have

$$\begin{aligned} & \left\| \sum_{k=1}^K b_k d_k \right\|_{d(\mathcal{M}, \Delta^{(m)}, v, p)} \\ &= \inf \left\{ \rho > 0 : \left(\sum_{k=1}^K \left(M_k \left(\frac{|\Delta^{(m)} b_{\varrho(k)}|^p v_k}{\rho} \right) \right) \right)^{\frac{1}{p}} \leq 1 \text{ for some } \rho > 0 \right\} \end{aligned}$$

for some permutation ϱ of $\{1, \dots, K\}$. However, due to symmetry, we can find that $C \in [1, \infty)$ such that $(y_k)_{k=1}^\infty$ is C -equivalent to all its subsequences and C -dominates $(g_k)_{k=1}^\infty$. Therefore,

$$\begin{aligned} \left\| \sum_{k=1}^K b_k d_k \right\|_{d(\mathcal{M}, \Delta^{(m)}, v, p)} &= \left(\sum_{k=1}^K \left(M_k \left(\frac{|\Delta^{(m)} b_{\varrho(k)}|^p v_k}{\rho} \right) \right) \right)^{\frac{1}{p}} \\ &\leq C^2 \left\| \sum_{k=1}^K b_k d_k \right\|_{g(\mathcal{M}, \Delta^{(m)}, v, p)}. \end{aligned}$$

It gives that $(y_k)_{k=1}^\infty$ C^2 -dominates $(d_k)_{k=1}^\infty$ □

Theorem 2.8 *Let $\mathcal{M} = (M_k)$ be a sequence of Orlicz functions, let $v = (v_k)_{k=1}^\infty \in V$ be a weight, and let $1 \leq p < \infty$. Let $(g_k)_{k=1}^\infty$ denote the canonical basis of $g(\mathcal{M}, \Delta^{(m)}, v, p)$. Suppose $y_k = \sum_{j=\hat{p}_k}^{\hat{p}_{k+1}-1} a_j g_j$ forms a normalized block basic sequence and satisfies $\lim_{j \rightarrow \infty} a_j = 0$. Let $(\hat{y}_k)_{k=1}^\infty$ be the sequence from Theorem 2.6 corresponding to $(y_k)_{k=1}^\infty$. Then for each $\varepsilon > 0$, there exists a subsequence $(y_{k_n})_{n=1}^\infty$ that is $(1 + \varepsilon)$ -equivalent to the canonical basis $(a_k)_{k=1}^\infty$ such that $(\hat{y}_{k_n})_{n=1}^\infty$ is $(1 + \varepsilon)$ -complemented in $g(\mathcal{M}, \Delta^{(m)}, v, p)$.*

Proof The first part of the proof follows that of Lemma 1 in [38]. Let $\varepsilon > 0$, and let $\hat{y}_k = \sum_{q=\hat{p}_k}^{\hat{p}_{k+1}-1} a_{j_q} g_q$ form a normalized block basic sequence as defined in Theorem 2.6, and let $\psi \in (0, 1)$ be such that $\psi^{-1/p} \leq \psi^1 < 1 + \psi$. We claim that there exists a block sequence of the form $z_k = \sum_{q=l_k}^{l_{k+1}-1} b_q g_q$ for $n \in \mathbb{N}$ and a subsequence $(\hat{y}_{k_n})_{k=1}^\infty$ satisfying the following two properties for every $n \in \mathbb{N}$:

- (i) The coefficients of z_k are the same as the coefficients of \hat{y}_{k_n} , that is, $l_{n+1} - l_n = \hat{p}_{k_{n+1}} - \hat{p}_{k_n}$ and $b_{l_n+q} = \hat{a}_{j_{\hat{p}_{k_n}+1}}$ for $q = 0, \dots, l_{n+1} - l_n - 1$, and
- (ii) $\sum_{q=l_n}^{l_{n+1}-1} (M_q(\frac{|\Delta^{(m)} b_q|^p v_q}{\rho})) \geq \psi$.

Let $l_1 = 1$ and define l_n for some $n \in \mathbb{N}$. Since $v_k \rightarrow 0$, we have $i \in \mathbb{N}$ such that

$$\sum_{q=i+1}^{i+l_n} M(|v_q|) < \frac{1 - \psi}{2}.$$

Since $a_{j_q} \rightarrow 0$, we have $\hat{p}_{k_{n+1}} - \hat{p}_{k_n} \rightarrow \infty$. Therefore, for any $h \in \mathbb{N}$ such that $\hat{p}_{h+1} - \hat{p}_h > i + l_n$, we have $M(|a_{j_q}|^p) < \frac{1-\psi}{2^i}$ for all $q \geq \hat{p}_h$. Let $l_{n+1} = \hat{p}_{h+1} - \hat{p}_h + l_n$ and $b_{l_n+q} = a_{j_{\hat{p}_h-q}}$ for all $q = 0, \dots, l_{n+1} - l_n - 1$. Since $\hat{y}_{k_n} = \hat{y}_h$ satisfies (i), by property (iv) of Theorem 2.6 we have

$$1 = \|\hat{y}_h\|_{g(\mathcal{M}, \Delta^{(m)}, v, p)}^p = \inf \left\{ \rho > 0 : \sum_{q=\hat{p}_h}^{\hat{p}_{h+1}-1} \left(M_q \left(\frac{|\Delta^{(m)} a_{j_q}|^p v_{q-\hat{p}_h+1}}{\rho} \right) \right) \leq 1 \text{ for some } \rho > 0 \right\},$$

and thus we have

$$\begin{aligned}
 \sum_{q=l_n}^{\hat{l}_{n+1}-1} \left(M_q \left(\frac{|\Delta^{(m)} b_q|^p v_q}{\rho} \right) \right) &= \sum_{q=\hat{p}_h}^{\hat{p}_{h+1}-1} \left(M_q \left(\frac{|\Delta^{(m)} a_{j_q}|^p v_{q+l_n-\hat{p}_h}}{\rho} \right) \right) \\
 &= \sum_{q=\hat{p}_h}^{\hat{p}_{h+1}-1} \left(M_q \left(\frac{|\Delta^{(m)} a_{j_q}|^p v_{q-\hat{p}_h+1}}{\rho} \right) \right) \\
 &\quad - \sum_{q=\hat{p}_h}^{\hat{p}_{h+1}-1} \left(M_q \left(\frac{|\Delta^{(m)} \hat{a}_q|^p (v_{q-\hat{p}_h+1} - v_{q+l_n-\hat{p}_h})}{\rho} \right) \right) \\
 &= 1 - \sum_{q=\hat{p}_h}^{\hat{p}_{h+1}-1} \left(M_q \left(\frac{|\Delta^{(m)} a_{j_q}|^p (v_{q-\hat{p}_h+1} - v_{q+l_n-\hat{p}_h})}{\rho} \right) \right) \\
 &\quad - \sum_{q=\hat{p}_h+i}^{\hat{p}_{h+1}-1} \left(M_q \left(\frac{|\Delta^{(m)} a_{j_q}|^p (v_{q-\hat{p}_h+1} - v_{q+l_n-\hat{p}_h})}{\rho} \right) \right). \tag{2.3}
 \end{aligned}$$

Since $(v_q)_{q=1}^\infty$ is nonincreasing with $0 < v_q \leq 1$ for $q \in \mathbb{N}$ and $l_n \geq 1$, we obtain $0 \leq v_{q-\hat{p}_h+1} - v_{q+l_n-\hat{p}_h} < 1$ for $q \in \mathbb{N}$. Since there are i values between \hat{p}_h and $\hat{p}_h + i - 1$ and also $M(|a_{j_q}|^p) < \frac{1-\psi}{2i}$ for all $q \geq \hat{p}_h$, we have

$$\begin{aligned}
 &\sum_{q=\hat{p}_h}^{\hat{p}_{h+1}-1} \left(M_q \left(\frac{|\Delta^{(m)} a_{j_q}|^p (v_{q-\hat{p}_h+1} - v_{q+l_n-\hat{p}_h})}{\rho} \right) \right) \\
 &< \sum_{q=\hat{p}_h}^{\hat{p}_{h+1}-1} \left(M_q \left(\frac{|\Delta^{(m)} a_{j_q}|^p}{\rho} \right) \right) < \frac{1-\psi}{2i} \sum_{q=\hat{p}_h}^{\hat{p}_{h+1}-1} M(1) = \frac{1-\psi}{2}. \tag{2.4}
 \end{aligned}$$

Since $M(|a_{j_q}|^p) < \frac{1-\psi}{2i} < 1$ for all $q \geq \hat{p}_h$, $\sum_{q=i+1}^{i+l_n} M(|v_q|) < \frac{1-\psi}{2}$, and $\hat{p}_{h+1} + l_n - 2 \geq \hat{p}_{h+1} - 1$, we have

$$\begin{aligned}
 &\sum_{q=\hat{p}_h+i}^{\hat{p}_{h+1}-1} \left(M_q \left(\frac{|\Delta^{(m)} a_{j_q}|^p (v_{q-\hat{p}_h+1} - v_{q+l_n-\hat{p}_h})}{\rho} \right) \right) \\
 &< \sum_{q=\hat{p}_h+i}^{\hat{p}_{h+1}-1} \left(M_q \left(\frac{v_{q-\hat{p}_h+1} - v_{q+l_n-\hat{p}_h}}{\rho} \right) \right) \\
 &< \sum_{q=\hat{p}_h+i}^{\hat{p}_{h+1}+l_n-1} \left(M_q \left(\frac{v_{q-\hat{p}_h+1}}{\rho} \right) \right) < \frac{1-\psi}{2}. \tag{2.5}
 \end{aligned}$$

Combining (2.3), (2.4), and (2.5), we obtain (ii). This completes the proof of our claim.

Let us prove that a block basic sequence $(z_k)_{k=1}$ in $g(\mathcal{M}, \Delta^{(m)}, v, p)$ dominates the canonical basis $(A_k)_{k=1}^\infty$ for l_p . For $K \in \mathbb{N}$, let $(c_k)_{k=1}^\infty$ be any finite sequence of scalars. Then we

have

$$\begin{aligned} \psi \sum_{k=1}^K \left(M_k \left(\frac{|\Delta^{(m)} c_k|^p}{\rho} \right) \right) &\leq \sum_{k=1}^K \left(M_k \left(\frac{|\Delta^{(m)} c_k|^p}{\rho} \right) \right) \sum_{q=l_k}^{l_{k+1}-1} \left(M_q \left(\frac{|\Delta^{(m)} b_q|^p v_q}{\rho} \right) \right) \\ &\leq \left\| \sum_{k=1}^K c_k z_k \right\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)}. \end{aligned}$$

Therefore,

$$\left(\sum_{k=1}^K \left(M_k \left(\frac{|\Delta^{(m)} c_k|^p}{\rho} \right) \right) \right)^{\frac{1}{p}} \leq \psi^{1/p} \left\| \sum_{k=1}^K c_k z_k \right\|_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)}.$$

Thus, $(A_k)_{k=1}^\infty$ is $\psi^{-1/p}$ -dominated by $(z_k)_{k=1}^\infty$. On the other hand, $(z_k)_{k=1}^\infty$ is 1-equivalent to $(\hat{y}_k)_{k=1}^\infty$ by (i) with 1-subsymmetry of $(g_k)_{k=1}^\infty$. By property (v) in Theorem 2.6 the sequence $(\hat{y}_k)_{k=1}^\infty$ is 1-dominated by $(y_k)_{k=1}^\infty$, and by Theorem 2.5 the sequence $(y_k)_{k=1}^\infty$ is 1-dominated by $(A_k)_{k=1}^\infty$. Now, we have

$$(A_k)_{k=1}^\infty \lesssim_{\psi^{-1/p}} (z_k)_{k=1}^\infty \approx_1 (y_k)_{k=1}^\infty \lesssim_1 (A_k)_{k=1}^\infty, \tag{2.6}$$

and thus $(y_k)_{k=1}^\infty$ is $\psi^{-1/p}$ -equivalent and hence $(1 + \varepsilon)$ -equivalent to $(A_k)_{k=1}^\infty$.

To show that $(\hat{y}_k)_{k=1}^\infty$ is $(1 + \varepsilon)$ -complement in $g(\mathcal{M}, \Delta^{(m)}, \nu, p)$, we will follow the proof of Lemma 15 in [39]. Let us define the sequence $\tilde{z}_n = \sum_{q=l_n}^{l_{n+1}-1} (M_q(\frac{|\Delta^{(m)} b_q|^p v_q}{\rho}))$ for $n \in \mathbb{N}$. Let P extend to a linear operator on $(g_j)_{j=1}^\infty$. For each $n \in \mathbb{N}$, we obtain $P\tilde{z}_n = P \sum_{j=l_n}^{l_{n+1}-1} (M_j(\frac{|\Delta^{(m)} b_j|^p v_j}{\rho})) = \hat{z}_n$, so that P is the projection onto $(\tilde{z}_n)_{n=1}^\infty$. Also by Theorem 2.5, $(\tilde{z}_n)_{n=1}^\infty$ is 1-dominated by $(A_n)_{n=1}^\infty$. Since (ii) holds for each $(c_j)_{j=1}^\infty \in c_{00}$, we get

$$\begin{aligned} \left\| P \sum_{j=1}^\infty c_j g_j \right\|_{\mathcal{M}, \Delta^{(m)}, p} &= \left\| P \sum_{n=1}^\infty \sum_{j=l_n}^{l_{n+1}-1} c_j g_j \right\|_{\mathcal{M}, \Delta^{(m)}, p} \\ &\leq \psi^{-p} \sum_{n=1}^\infty \left| \sum_{j=l_n}^{l_{n+1}-1} \left(M_j \left(\frac{c_j |\Delta^{(m)} b_j|^{p-1} v_j}{\rho} \right) \right) \right|^p. \end{aligned}$$

Then by the Hölder’s inequality with $\frac{p}{p-1}$ conjugate to p and by

$$\Delta^m(c_j) |\Delta^m b_j|^{p-1} v_j = (\Delta^m(c_j) v_j^{1/p}) (|\Delta^m b_j|^{p-1} v_j^{(p-1)/p})$$

with $\sum_{j=l_n}^{l_{n+1}-1} (M_j(\frac{|\Delta^{(m)} b_j|^p v_j}{\rho})) \leq \|z_n\|_{\mathcal{M}, \Delta^{(m)}, p} = 1$ we have

$$\begin{aligned} \psi^{-p} \sum_{n=1}^\infty \left| \sum_{j=l_n}^{l_{n+1}-1} \left(M_j \left(\frac{\Delta^m(c_j) |\Delta^m b_j|^{p-1} v_j}{\rho} \right) \right) \right|^p &\leq \psi^{-p} \sum_{n=1}^\infty \sum_{j=l_n}^{l_{n+1}-1} \left(M_j \left(\frac{|\Delta^{(m)} c_j|^p v_j}{\rho} \right) \right) \\ &\leq \psi^{-p} \left\| \sum_{j=1}^\infty c_j g_j \right\|_{g, \mathcal{M}, \Delta^{(m)}, p}. \end{aligned}$$

Therefore, P is bounded by ψ^{-1} , and thus there is a ψ^{-1} -bounded continuous extension \tilde{P} in $\mathcal{L}(g(\mathcal{M}, \Delta^{(m)}, \nu, p))$, where \tilde{P} is the projection onto $(\tilde{z}_n)_{n=1}^\infty$. Now define some operators $\Psi_1, \Theta_1, \Psi_2, \Theta_2 \in \mathcal{L}(g(\mathcal{M}, \Delta^{(m)}, \nu, p))$ such that

$$\hat{y}_{k_n} \xrightarrow{\Psi_1} z_n \xrightarrow{\Psi_2} \tilde{z}_n \quad \text{and} \quad \tilde{z}_n \xrightarrow{\Theta_2} z_n \xrightarrow{\Theta_1} \hat{y}_{k_n} \quad (2.7)$$

for all $n \in \mathbb{N}$. Note that $\Psi_2\Theta_2 = \Theta_2\Psi_2 = I_{g(\mathcal{M}, \Delta^{(m)}, \nu, p)}$. By the 1-unconditionality of $(g_k)_{k=1}^\infty$ these are all norm-one operators, and all these satisfy the mappings in (2.7). Let us form the operator

$$\Theta_1\Theta_2\tilde{P}\Psi_1\Psi_2 \in \mathcal{L}(g(\mathcal{M}, \Delta^{(m)}, \nu, p)).$$

Therefore it is a projection onto $(\hat{y}_{k_n})_{n=1}^\infty$. This gives the desired result. \square

Acknowledgements

The authors extend their appreciation to the Deanship of Scientific Research at King Khalid University for funding this work through research groups program under grant number R.G.P. 1/13/38.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

The authors contributed equally and significantly in writing this paper. All authors read and approved the final manuscript.

Author details

¹School of Mathematics, Shri Mata Vaishno Devi University, Katra, India. ²Operator Theory and Applications Research Group, Department of Mathematics, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia. ³Department of Mathematics, College of Science, King Khalid University, Abha, Saudi Arabia.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 21 November 2017 Accepted: 25 January 2018 Published online: 07 February 2018

References

- Lindenstrauss, J., Tzafriri, L.: Classical Banach Spaces I: Sequence Spaces (1977). ISBN 3-540-08072-4
- Albiac, F., Kalton, N.J.: Topics in Banach Space Theory, 2nd revised and updated edn. Graduate Texts in Mathematics, vol. 233. Springer, Berlin (2016)
- Lindenstrauss, J., Tzafriri, L.: On Orlicz sequence spaces. *Isr. J. Math.* **10**, 379–390 (1971)
- Maligranda, L.: Orlicz Spaces and Interpolation. *Seminars in Mathematics*, vol. 5. Polish Academy of Science, Warsaw (1989)
- Musielak, J.: Orlicz Spaces and Modular Spaces. *Lecture Notes in Mathematics*, vol. 1034. Springer, Berlin (1983)
- Alotaibi, A., Raj, K., Mohiuddine, S.A.: Some generalized difference sequence spaces defined by a sequence of moduli in n -normed spaces. *J. Funct. Spaces* **2015**, Article ID 413850 (2015)
- Basarir, M.: Some new sequence space generated by Orlicz functions. *South East Asian J. Math. Math. Sci.* **1**(3), 65–73 (2003)
- Basarir, M.: On the generalized Riesz B -difference sequence spaces. *Filomat* **24**(4), 35–52 (2010)
- Basarir, M.: On lacunary strong σ -convergent with respect to a sequence of ϕ -functions. *Fasc. Math.* **43**, 19–32 (2010)
- Basarir, M., Altundag, S.: Some difference sequence spaces defined by a sequence of ϕ -functions. *Rend. Circ. Mat. Palermo* **57**(1), 149–160 (2008)
- Basarir, M., Altundag, S.: Some generalized difference sequence spaces defined by a sequence of Orlicz functions. *Fasc. Math.* **41**, 17–30 (2009)
- Basarir, M., Kara, E.E.: On some difference sequence spaces of weighted means and compact operators. *Ann. Funct. Anal.* **2**(2), 116–131 (2011)
- Basarir, M., Kara, E.E.: On the B -difference sequence space derived by weighted mean and compact operators. *J. Math. Anal. Appl.* **391**(1), 67–81 (2012)
- Basarir, M., Kara, E.E.: On the m th order difference sequence space of generalized weighted mean and compact operators. *Acta Math. Sci.* **33B**(3), 797–813 (2013)
- Basarir, M., Ozturk, M., Kara, E.E.: Some topological and geometric properties of generalized Euler sequence space. *Math. Slovaca* **60**(3), 385–398 (2010)
- Basarir, M., Ozturk, M.: On some generalized B_m -difference Riesz sequence spaces and uniform Opial property. *J. Inequal. Appl.* **2011**, Article ID 485730 (2011)

17. Esi, A.: On ∇ -statistical convergence in random 2-normed space. *Rev. Anal. Numér. Théor. Approx.* **43**, 175–187 (2014)
18. Konca, S., Basarir, M.: Generalized difference sequence spaces associated with a multiplier sequence on a real n -normed space. *J. Inequal. Appl.* **2013**, 335 (2013)
19. Mohiuddine, S.A., Hazarika, B.: Some classes of ideal convergent sequences and generalized difference matrix operator. *Filomat* **31**(6), 1827–1834 (2017)
20. Mohiuddine, S.A., Hazarika, B.: On strongly almost generalized difference lacunary ideal convergent sequences of fuzzy numbers. *J. Comput. Anal. Appl.* **23**(5), 925–936 (2017)
21. Mohiuddine, S.A., Raj, K.: Vector valued Orlicz-Lorentz sequence spaces and their operator ideals. *J. Nonlinear Sci. Appl.* **10**, 338–353 (2017)
22. Mohiuddine, S.A., Raj, K., Alotaibi, A.: Generalized spaces of double sequences for Orlicz functions and bounded-regular matrices over n -normed spaces. *J. Inequal. Appl.* **2014**, 332 (2014)
23. Mursaleen, M., Noman, A.K.: On some imbedding relations between certain sequence spaces. *Ukr. Math. J.* **63**, 564–579 (2011)
24. Raj, K., Sharma, C.: Applications of strongly convergent sequences to Fourier series by means of modulus functions. *Acta Math. Hung.* **150**, 396–411 (2016)
25. Raj, K., Kiliçman, A.: On certain generalized paranormed spaces. *J. Inequal. Appl.* **2015**, 37 (2015)
26. Raj, K., Jamwal, S.: On some generalized statistical convergent sequence spaces. *Kuwait J. Sci. Eng.* **42**, 86–104 (2015)
27. Kizmaz, H.: On certain sequence spaces. *Can. Math. Bull.* **24**, 169–176 (1981)
28. Et, M., Çolak, R.: On generalized difference sequence spaces. *Soochow J. Math.* **21**, 377–386 (1995)
29. Bektaş, Ç.A., Et, M., Çolak, R.: Generalized difference sequence spaces and their dual spaces. *J. Math. Anal. Appl.* **292**, 423–432 (2004)
30. Et, M., Esi, A.: On Köthe-Toeplitz duals of generalized difference sequence spaces. *Bull. Malays. Math. Sci. Soc.* **23**, 25–32 (2000)
31. Lorentz, G.G.: Some new functional spaces. *Ann. Math.* **51**, 37–55 (1950)
32. Lorentz, G.G.: On the theory of spaces Λ . *Pac. J. Math.* **1**, 411–429 (1951)
33. Garling, D.J.H.: Symmetric bases of locally convex spaces. *Stud. Math.* **30**, 163–181 (1968)
34. Pujara, L.R.: A uniformly convex Banach space with a Schauder basis which is subsymmetric but not symmetric. *Proc. Am. Math. Soc.* **54**, 207–210 (1976)
35. Dilworth, S.J., Odell, E., Schlumprecht, T., Zsák, A.: Renormings and symmetry properties of 1-greedy bases. *J. Approx. Theory* **163**, 1049–1075 (2011)
36. Wallis, B.: Garling sequence spaces (2016) arXiv:1612.01145v1 [math.FA]
37. Altshuler, Z.: Uniform convexity in Lorentz sequence spaces. *Isr. J. Math.* **20**, 260–274 (1975)
38. Altshuler, Z., Casazza, P.G., Lin, B.L.: On symmetric basic sequences in Lorentz sequence spaces. *Isr. J. Math.* **15**, 140–155 (1973)
39. Casazza, P.G., Lin, B.L.: On symmetric basic sequences in Lorentz sequence spaces II. *Isr. J. Math.* **17**, 191–218 (1974)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)
