

Aplikace matematiky

Otakar Jaroš

Orthoexponential polynomials and the Legendre polynomials

Aplikace matematiky, Vol. 23 (1978), No. 6, 467–471

Persistent URL: <http://dml.cz/dmlcz/103772>

Terms of use:

© Institute of Mathematics AS CR, 1978

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ORTHOEXPONENTIAL POLYNOMIALS AND THE LEGENDRE POLYNOMIALS

OTAKAR JAROCH

(Received April 5, 1977)

Orthogonal exponential polynomials or *orthoexponential polynomials* have interesting applications in Automatic Control – Laning-Battin [8], Electrical Circuits Theory – Čížek [2], Tuttle [10], Hydrometeorology – Dmitriyev [4], and as a means for the inversion of Laplace transforms – Jaroch [5]. A very simple connection will be shown to exist between the orthoexponential and the Legendre polynomials.

In this paper, the letters k, m, n denote nonnegative integers, t and x are real variables, δ_{mn} is the Kronecker delta function and, in Definition II only, $[n/2]$ denotes the integer part of $n/2$. As the notation and standardization of both the orthoexponential and the Legendre polynomials vary, definitions of the Legendre, Jacobi, and orthoexponential polynomials as used in this paper are re-stated here.

Definition I [6]. The functions $\varphi_n(t) = \sum_{k=1}^n b_{nk} e^{-kt}$, $n = 1, 2, 3, \dots$, are called *orthoexponential polynomials* if the coefficients b_{nk} are so chosen that for all m, n the following conditions are satisfied: $\varphi_n(0) = 1$ and $\int_0^\infty \varphi_m(t) \varphi_n(t) dt = \delta_{mn} \|\varphi_n\|^2$.

The coefficients of orthoexponential polynomials are $b_{nk} = (-1)^{n+k} \binom{n}{k} \binom{n+k-1}{k-1}$; $\varphi_1(t) = e^{-t}$. For all n we have $\varphi_n(+\infty) = 0$. Orthoexponential polynomials are bounded and $|\varphi_n(t)| \leq 1$ for all nonnegative t . Their norm is $\|\varphi_n\| = (2n)^{-1/2}$ and orthonormal exponential polynomials are therefore $t \mapsto (2n)^{1/2} \varphi_n(t)$.

Definition II [9]. The polynomials $P_n(x) = \sum_{k=0}^{[n/2]} a_{nk} x^{n-2k}$, $n = 0, 1, 2, \dots$, are called *Legendre polynomials* if the coefficients a_{nk} are so chosen that for all m, n the following conditions are satisfied: $P_n(1) = 1$ and $\int_{-1}^1 P_m(x) P_n(x) dx = \delta_{mn} \|P_n\|^2$.

The coefficients of the Legendre polynomials are $a_{nk} = (-1)^k 2^{-n} \binom{n}{k} \binom{2n-2k}{n}$; $P_0(x) \equiv 1$. Their norm is $\|P_n\| = (n + \frac{1}{2})^{-1/2}$, and $|P_n(x)| \leq 1$ if only $-1 \leq x \leq 1$.

The notation $P_n^{(\alpha,\beta)}(x)$ is used for the Jacobi polynomials, orthogonal over $(-1, 1)$ with respect to the weight function $(1-x)^\alpha(1+x)^\beta$, $\alpha > -1$, $\beta > -1$, and standardized with $P_n^{(\alpha,\beta)}(1) = \binom{n+\alpha}{n}$. Legendre polynomials are a special case of the Jacobi polynomials for $\alpha = \beta = 0$. Many properties of the Jacobi polynomials originate from their close connection with the hypergeometric function. The following relation will be used in this paper:

$$(1) \quad (1+x)P_{n-1}^{(0,1)}(x) = P_n^{(0,0)}(x) + P_{n-1}^{(0,0)}(x), \quad n \geq 1,$$

where $P_n^{(0,0)}(x) \equiv P_n(x)$ are the Legendre polynomials [1], [9].

It is known [6] that orthoexponential polynomials can be expressed in terms of the Jacobi polynomials $P_n^{(0,1)}(x)$. However, no explicit mention has been found so far of a simple relation connecting the orthoexponential and the Legendre polynomials as expressed in the following two theorems.

Theorem 1. *Let $\varphi_n(t)$ be the orthoexponential polynomials and $P_n(x)$ the Legendre polynomials in accordance with Definitions I and II; $n = 1, 2, 3, \dots$. Then, for arbitrary t ,*

$$(2) \quad \varphi_n(t) = \frac{1}{2}[P_n(2e^{-t} - 1) + P_{n-1}(2e^{-t} - 1)].$$

Proof. In Eq. (2), orthoexponential polynomials are expressed, in accordance with Definitions I and II, as linear combinations of exponential functions e^{-t} , e^{-2t} , ..., e^{-nt} . The standardization $\varphi_n(0) = 1$, $\varphi_n(+\infty) = 0$, is satisfied at the same time. It remains to prove that $\int_0^\infty \varphi_m(t) \varphi_n(t) dt = \delta_{mn}(2n)^{-1}$. Substituting $x = 2e^{-t} - 1$ in Eq. (1) we have

$$2e^{-t}P_{n-1}^{(0,1)}(2e^{-t} - 1) = P_n(2e^{-t} - 1) + P_{n-1}(2e^{-t} - 1)$$

and Eq. (2) is therefore equivalent to

$$(3) \quad \varphi_n(t) = e^{-t}P_{n-1}^{(0,1)}(2e^{-t} - 1).$$

The norm of the Jacobi polynomials is known [1] [9] and for the case under consideration we have $\|P_n^{(0,1)}\| = 2^{1/2}(n+1)^{-1/2}$. As a consequence,

$$\int_0^\infty \varphi_m(t) \varphi_n(t) dt = \int_0^\infty e^{-2t}P_{m-1}^{(0,1)}(2e^{-t} - 1)P_{n-1}^{(0,1)}(2e^{-t} - 1) dt$$

and, if the substitution $x = 2e^{-t} - 1$ is used,

$$\int_0^\infty \varphi_m(t) \varphi_n(t) dt = \frac{1}{4} \int_{-1}^{+1} (1+x)P_{m-1}^{(0,1)}(x)P_{n-1}^{(0,1)}(x) dx = \delta_{mn}(2n)^{-1}.$$

Q.e.d.

Remark. Jacobi polynomials are defined by Courant and Hilbert [3] [6] as polynomials orthogonal over $(0, 1)$ with respect to the weight function $x^{q-1}(1-x)^{p-q}$, $q > 0$, $p - q > -1$, and denoted by $G_n(p, q, x)$. The standardization is $G_n(p, q, 0) = 1$ for all n . In this case Eq. (3) becomes

$$(4) \quad \varphi_n(t) = (-1)^{n-1} n e^{-t} G_{n-1}(2, 2, e^{-t})$$

and Eq. (2) assumes the form

$$(5) \quad \varphi_n(t) = (-1)^n 2^{-1} [G_n(1, 1, e^{-t}) - G_{n-1}(1, 1, e^{-t})].$$

Example 1. The value of $\varphi_{10}(1)$ is to be computed from Eq. (2). For $t = 1$ we have $2e^{-t} - 1 = -0.264241$ and thus $\varphi_{10}(1) = \frac{1}{2}[P_{10}(-0.264241) + P_9(-0.264241)] = \frac{1}{2}(0.237018 - 0.148194)$. The result is $\varphi_{10}(1) = 0.044412$. A calculation of values of the orthoexponential polynomials based on Definition I involves, for large n , computations with prohibitively large numbers. In this example, if Definition I is used for the computation of $\varphi_{10}(1)$ then the largest coefficient involved is $b_{10,7} = -960960$.

Theorem 2. Let $\varphi_n(t)$ be the orthoexponential polynomials and $P_n(x)$ the Legendre polynomials in accordance with Definitions I and II; $n = 1, 2, 3, \dots$. Then, for arbitrary t ,

$$(6) \quad P_n(2e^{-t} - 1) = 2 \varphi_n(t) - 2 \varphi_{n-1}(t) + \dots + (-1)^{n-1} 2 \varphi_1(t) + (-1)^n.$$

Proof. For $n = 1$ we have $\varphi_1(t) = \frac{1}{2}[P_1(2e^{-t} - 1) + 1]$ as a consequence of Theorem 1 and thus $P_1(2e^{-t} - 1) = 2 \varphi_1(t) - 1$. For $n = 2$ we find $\varphi_2(t) = \frac{1}{2}[P_2(2e^{-t} - 1) + P_1(2e^{-t} - 1)]$ or $\varphi_2(t) = \frac{1}{2}[P_2(2e^{-t} - 1) + 2 \varphi_1(t) - 1]$ and that too agrees with Eq. (6). The proof proceeds by induction. Let us assume that Eq. (6) is valid for a certain integer $n > 1$. It follows from Theorem 1 that $\varphi_{n+1}(t) = \frac{1}{2}[P_{n+1}(2e^{-t} - 1) + P_n(2e^{-t} - 1)]$ or $P_{n+1}(2e^{-t} - 1) = 2 \varphi_{n+1}(t) - P_n(2e^{-t} - 1)$. Substituting for $P_n(2e^{-t} - 1)$ from Eq. (6) we have $P_{n+1}(2e^{-t} - 1) = 2 \varphi_{n+1}(t) - 2 \varphi_n(t) + \dots + (-1)^n 2 \varphi_1(t) + (-1)^{n+1}$. Q.e.d.

Remark. For the Jacobi polynomials in the Courant and Hilbert notation [3] an analogous relation follows from Eq. (5):

$$(7) \quad G_n(1, 1, e^{-t}) = 1 - 2 \varphi_1(t) + 2 \varphi_2(t) - 2 \varphi_3(t) + \dots + (-1)^n 2 \varphi_n(t).$$

Example 2. The series $1 - 2 \varphi_1(t) + 2 \varphi_2(t) - 2 \varphi_3(t) + \dots + (-1)^n 2 \varphi_n(t) + \dots = 0$ is convergent for any positive t . The n -th partial sum of this series is $(-1)^n \cdot P_n(2e^{-t} - 1)$ as a consequence of Theorem 2. Laplace's theorem on the asymptotics of the Legendre polynomials [9] states that $P_n(x) = O(n^{-1/2})$ for $n \rightarrow \infty$ if only $-1 < x < 1$. Thus, the n -th partial sum of the series under consideration is $(-1)^n \cdot$

$P_n(2e^{-t} - 1) = o(1)$ and the series therefore converges. Accordingly, the series $2 \sum_{n=1}^{\infty} (-1)^{n-1} \varphi_n(t) = 1$ converges for any positive t ; for $t = 0$ the series is divergent.

CONCLUSION

The theorems proved in this paper are useful if a formulation is sought, for orthoexponential polynomials, of some of the many known results in the theory of classical orthogonal polynomials. Furthermore, Theorem 1 may be convenient for the computation of values of orthoexponential polynomials if the values of the Legendre polynomials are available. In addition to the recurrence formula

$$(8) \quad \begin{aligned} (n+1)(2n-1)\varphi_{n+1} &= \\ &= [(4n^2-1)e^{-t} - 2n^2]2\varphi_n - (n-1)(2n+1)\varphi_{n-1}, \end{aligned}$$

$n = 1, 2, 3, \dots$; $\varphi_1 = e^{-t}$, $\varphi_0 \equiv 0$ (see Jaroch-Novotný [7]) we have here another relation, namely $\varphi_n(t) = \frac{1}{2}[P_n(2e^{-t} - 1) + P_{n-1}(2e^{-t} - 1)]$, which is useful when an effective algorithm is required for the computation of values of orthoexponential polynomials.

References

- [1] *H. Bateman, A. Erdélyi*: Higher Transcendental Functions, Vol. 2, McGraw-Hill, New York 1953.
- [2] *V. Čížek*: Methods of Time Domain Synthesis. Research Report Z-44, Czechoslovak Academy of Sciences, Institute of Radioelectronics, Praha, 1960 (in Czech).
- [3] *R. Courant, D. Hilbert*: Methoden der mathematischen Physik, Vol. 1, Berlin, 1931 (Russian translation: GITTL, 1951).
- [4] *A. A. Dmitriyev*: Orthogonal Exponential Functions in Hydrometeorology. Gidrometeoizdat, Leningrad, 1973 (in Russian).
- [5] *O. Jaroch*: A Method of Numerical Inversion of Laplace Transforms. Práce ČVUT, Series VI, No. 1, Part I, pp. 332—339. Czech Technical University, Prague 1961 (in Czech).
- [6] *O. Jaroch*: Approximation by Exponential Functions. Aplikace matematiky, Vol. 7, No. 4, pp. 249—264, 1962 (in Czech).
- [7] *O. Jaroch, J. Novotný*: Recurrence Relations for Orthogonal Exponential Polynomials and their Derivatives. Acta Polytechnica — Práce ČVUT, Vol. IV (1973), pp. 39—42 (in Czech).
- [8] *J. H. Laning, R. H. Battin*: Random Processes in Automatic Control. McGraw-Hill, New York, 1956.
- [9] *G. Szegő*: Orthogonal Polynomials. American Mathematical Society, New York, 1959.
- [10] *D. F. Tuttle*: Network Synthesis for Prescribed Transient Response. Massachusetts Institute of Technology, 1949, DSc. Thesis.

Souhrn

O SOUVISLOSTI ORTOEXPONENCIÁLNÍCH MNOHOČLENŮ S LEGENDREOVÝMI MNOHOČLENY

OTAKAR JAROCH

Vyjádření ortoexponenciálních mnohočlenů $\varphi_n(t) = \sum_{k=1}^n b_{nk} e^{-kt}$ podle definice I je pro velká n nevhodné pro numerický výpočet funkčních hodnot, protože k výsledku přicházíme odčítáním velkých čísel. V tomto případě je užitečný rekurentní vzorec (8) nebo v tomto článku dokázaná věta 1, totiž $\varphi_n(t) = \frac{1}{2}[P_n(2e^{-t} - 1) + P_{n-1}(2e^{-t} - 1)]$, kde $P_n(x)$ jsou Legendreovy mnohočleny. Věty, ukazující na souvislost ortoexponenciálních mnohočlenů s Legendreovými mnohočleny jsou užitečné jednak pro numerické výpočty jednak proto, že umožňují jednoduchým způsobem přenést na ortoexponenciální mnohočleny některé výsledky z teorie klasických ortogonálních mnohočlenů. Ortoexponenciální mnohočleny jsme definovali se standardizací $\varphi_n(0) = 1$ pro všechna n , protože při této volbě zvláště vyniknou souvislosti s klasickou teorií ortogonálních mnohočlenů.

Author's address: Doc. Dr. Otakar Jarocho, Katedra matematiky, Vysoká škola chemickotechnologická, Suchbátarova 5, 166 28 Praha 6 - Dejvice.