

Svatopluk Poljak; Vojtěch Rödl  
Orthogonal partitions and covering of graphs

*Czechoslovak Mathematical Journal*, Vol. 30 (1980), No. 3, 475,476–477,478–485

Persistent URL: <http://dml.cz/dmlcz/101696>

## Terms of use:

© Institute of Mathematics AS CR, 1980

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## ORTHOGONAL PARTITIONS AND COVERING OF GRAPHS

SVATOPLUK POLJAK and VOJTECH RÖDL, Praha

(Received December 7, 1978)

## INTRODUCTION

We investigate two notions related to the notion of chromatic number. For a given graph  $G$  one may define  $\bar{\omega}(G)$  as the minimum number of stable sets (i.e. subsets of the vertex set with no two vertices joined by an edge) such that every pair of non-adjacent vertices is contained in at least one of them, and  $\dim G$  as the minimal  $d$  such that  $G$  is contained in a product of  $d$  complete graphs as an induced subgraph.

Determination of both  $\bar{\omega}$  and  $\dim$  is an NP-complete problem. (See [15], [10].) In [8] and [12],  $\dim$  of some special graphs was investigated. The main purpose of this paper is to prove some analogous statements concerning the notion of  $\bar{\omega}$  and extend some results of [8] and [12].

The first two sections are of technical character. In § 1 we present some remarks concerning the notion of orthogonal array (which is in our sense more general than that defined in [5]).

In § 2 we introduce the notion of orthogonal  $q$ -partitions and investigate the problem of maximal number of pairwise orthogonal  $q$ -partitions on a given set. For  $q = 2$  we are able to give an exact formula for this number. For  $q \geq 3$  we give only some bounds.

In § 4 we show that  $\bar{\omega}$  of a product of two graphs with large  $\bar{\omega}$  is large, too. To prove an analogous statement concerning chromatic number is an open (and probably difficult) problem.

In last paragraphs using the results of the first two sections we give a formula for  $\bar{\omega}(nK_2)$  and some bounds for  $\bar{\omega}(nK_q)$ ,  $\dim(nK_q)$  and  $\dim\binom{n}{q}$ .

We state now three well known theorems which will be used throughout our paper.

**Theorem 0.1.** (see [5]) *Let  $q$  be a power of a prime. Then there exist  $q - 1$  pairwise orthogonal latin squares of order  $q$ .*

**Theorem 0.2.** (see [2]) *Let  $A_1, A_2, \dots, A_n$  be a system of subsets of  $1, 2, \dots, t$  such that:*

- (i)  $A_i \not\subset A_j$ ;
- (ii)  $A_i \cap A_j \neq \emptyset$ ;
- (iii)  $|A_i| \leq \frac{1}{2}t$ .

Then

$$n \leq \binom{t-1}{\lfloor \frac{t}{2} \rfloor - 1}.$$

**Theorem 0.3.** For every integer  $n$  there exists a prime  $p$  such that

$$n \leq p \leq 2n.$$

### 1. ORTHOGONAL ARRAYS

Let  $q$  be a positive integer. We say that two vectors  $\mathbf{u} = (u_1, u_2, \dots, u_{t_1})$  and  $\mathbf{v} = (v_1, v_2, \dots, v_{t_2})$  are orthogonal if for every  $k, k' = 1, 2, \dots, q$  there exists  $j = 1, 2, \dots, t$  so that  $u_j = k$  and  $v_j = k'$ .

For two vectors  $\mathbf{u} = (u_1, u_2, \dots, u_t)$  and  $\mathbf{v} = (v_1, v_2, \dots, v_t)$  we put  $(\mathbf{u}, \mathbf{v}) = (u_1, u_2, \dots, u_t, v_1, v_2, \dots, v_t)$ .

Let  $\pi$  be a permutation of the set  $\{1, 2, \dots, q\}$ . We identify  $\pi$  with the vector  $(\pi(1), \pi(2), \dots, \pi(q))$ . If  $\mathbf{u} = (u_j)$ ,  $u_j \in \{1, 2, \dots, q\}$  is another vector then we put

$$\pi(\mathbf{u}) = (\pi(u_1), \pi(u_2), \dots, \pi(u_t)).$$

Obviously the following holds

$$\pi(\mathbf{u}, \mathbf{v}) = (\pi(\mathbf{u}), \pi(\mathbf{v})).$$

We say that a vector  $\mathbf{w}$  has the permutation property if

$$\mathbf{w} = (\pi_1, \pi_2, \dots, \pi_r)$$

where  $\pi_s$ ,  $s = 1, 2, \dots, r$ , are permutations of  $\{1, 2, \dots, q\}$  for some  $q$  and  $r$ .

The orthogonal array  $\mathbf{OA}(q, n, t)$  is an  $n \times t$  matrix  $(a_{ij})$  where  $a_{ij} \in \{1, 2, \dots, q\}$  for every  $i, j$ ,  $i = 1, 2, \dots, n$ ,  $j = 1, 2, \dots, t$ , such that every two row-vectors are orthogonal. The orthogonal array  $(\mathbf{OA})$  has the permutation property if each row of the corresponding matrix has this property.

Let  $t = q^2$ . Then the existence of  $\mathbf{OA}(q, n, q^2)$  is equivalent with the existence of  $n - 2$  pairwise orthogonal latin squares of order  $q$ . (See e.g. [1, 5].) In [3], the maximal number of pairwise orthogonal latin squares is estimated. For a given  $q$  there exist at most  $q - 1$  pairwise orthogonal latin squares, and if  $q$  is a power of a prime then such a system exists.

**Theorem 1.1.** *Let  $q$  be a power of a prime. Then*

- (i) *There exists  $\mathbf{OA}(q, q + 1, q^2)$ .*
- (ii) *There exists  $\mathbf{OA}(q, q, q^2)$  which has the permutation property.*

*Proof.* By Theorem 0.1 and the relation between  $\mathbf{OA}$ 's we have (i). It remains to prove (ii).

Take  $\mathbf{OA}(q, q + 1, q^2)$ . In each row there are exactly  $q$  entries equal to  $k$  for each  $k = 1, 2, \dots, q$ . We can suppose that the last row is of the form  $(1, 1, \dots, 1, 2, 2, \dots, 2, \dots, q, q, \dots, q)$ . (This follows from the obvious fact that if we rearrange columns of  $\mathbf{OA}$  we obtain again  $\mathbf{OA}$ .) The first  $q$  rows obviously form an  $\mathbf{OA}$  with the permutation property. So for  $i = 1, 2, \dots, q$  the  $i$ -th row is of the form  $(\pi_{i1}, \pi_{i2}, \dots, \pi_{iq})$ .

The following theorem as well as its proof is a simple generalization of [9] (see [5], Theorem 13.2.1).

**Theorem 1.2.** *The existence of  $\mathbf{OA}(q, n_1, t_1)$  and  $\mathbf{OA}(q, n_2, t_2)$  implies the existence of  $\mathbf{OA}(q_1q_2, n, t_1t_2)$ .*

**Theorem 1.3.** *The existence of  $\mathbf{OA}(q, n_1, t_1)$  and  $\mathbf{OA}(q, n_2, t_2)$  implies the existence of  $\mathbf{OA}(q, n_1n_2, t_1 + t_2)$ .*

*Moreover, if the first two  $\mathbf{OA}$ 's have the permutation property then the last one has the property as well.*

*Proof.* Denote the row-vectors of  $\mathbf{OA}(q, n_1, t_1)$  and  $\mathbf{OA}(q, n_2, t_2)$  by  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{t_1}$  and  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{t_2}$ , respectively. Then  $\mathbf{w}_{i,i'} = (\mathbf{u}_i, \mathbf{v}_{i'})$ ,  $i = 1, 2, \dots, t_1$ ,  $i' = 1, 2, \dots, t_2$  are pairwise orthogonal vectors and hence  $\mathbf{OA}(q, n_1n_2, t_1 + t_2)$  exists. Moreover, if  $\mathbf{u}_i, \mathbf{v}_{i'}$  have the permutation property, then  $\mathbf{w}_{i,i'}$  has the permutation property, too.

The following theorem gives a better estimate for the case that we start the construction with  $q$   $\mathbf{OA}$ 's.

**Theorem 1.4.** *Let  $q$  be a power of a prime. Let  $\mathbf{OA}(q, n_j, t_j)$ ,  $j = 1, 2, \dots, q$ , exist. Then there exists  $\mathbf{OA}(q, qn_1n_2 \dots n_q, t_1 + \dots + t_q)$ . Moreover, if the first  $q$   $\mathbf{OA}$ 's have the permutation property then the last one has the property as well.*

*Proof.* Denote by  $\mathbf{v}_j^i$  the  $i$ -th row-vector of  $\mathbf{OA}(q, n_j, t_j)$ . Take  $\mathbf{OA}(q, q, q^2)$  with the permutation property (the existence follows from Theorem 1.1) and denote its  $i$ -th row by  $\pi_i = (\pi_{i1}, \pi_{i2}, \dots, \pi_{iq})$ . The vectors

$$\mathbf{v}_{i_0, i_1, i_2, \dots, i_q} = (\pi_{i_01}(\mathbf{u}_{i_1}^1), \pi_{i_02}(\mathbf{u}_{i_2}^2), \dots, \pi_{i_0q}(\mathbf{u}_{i_q}^q))$$

for  $i_0 \in \{1, 2, \dots, q\}$  and  $i_j \in \{1, 2, \dots, n_j\}$ , for each  $j \in \{1, 2, \dots, q\}$ , form  $\mathbf{OA}(q, qn_1n_2 \dots n_q, t_1 + t_2 + \dots + t_q)$ . If the vectors  $\mathbf{u}_j^i$ ,  $j \in \{1, 2, \dots, q\}$ , have the permutation property then the vectors  $\mathbf{v}_{i_0, i_1, \dots, i_q}$  have the property as well.

## 2. ORTHOGONAL PARTITIONS

**Definition.** Let  $\mathcal{A} = (A_1, A_2, \dots, A_q)$ ,  $\mathcal{B} = (B_1, B_2, \dots, B_q)$  be two partitions into  $q$  parts ( $q$ -partitions) of a set with  $t$  elements. We say that  $\mathcal{A}$  and  $\mathcal{B}$  are *orthogonal* if  $A_i \cap B_{i'} \neq \emptyset$  for every  $i, i' \in \{1, 2, \dots, q\}$ .

We denote by  $f(t, q)$  the maximal size of a system of pairwise orthogonal  $q$ -partitions of the  $t$ -element set.

**Theorem 2.1.**

$$f(t, 2) = \binom{t-1}{\lfloor \frac{t}{2} \rfloor - 1}.$$

*Proof.* Let  $\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_n$  be a system of pairwise orthogonal 2-partitions. For every  $i = 1, \dots, n$  choose the partition class  $A_i$  with cardinality  $\leq t/2$ . The sets  $A_1, A_2, \dots, A_n$  have the following properties:

- (i) they form an antichain in the partial order of subsets of  $\{1, \dots, t\}$ ;
- (ii)  $A_i \cap A_j \neq \emptyset$  for every  $i \neq j$ ;
- (iii)  $|A_i| \leq \frac{t}{2}$  for every  $i$ .

Hence, by Erdős Ko Rado theorem [2], we get

$$f(t, 2) \leq \binom{t-1}{\lfloor \frac{t}{2} \rfloor - 1}.$$

To prove  $\geq$  consider the 2-partitions of  $\{1, \dots, t\}$  defined as follows: To every subset with cardinality  $\lfloor \frac{t}{2} \rfloor$  containing 1, assign a 2-partition consisting of this set and its complement.

For  $q > 2$  we are far from being able to find an exact formula for  $f(t, q)$ . We give here some estimates.

**Theorem 2.2.**

$$f\left(\binom{q}{2}t, q\right) \geq f(t, 2).$$

*Proof.* Suppose that on the set  $X = \{1, 2, \dots, t\}$  there exists a system of  $n$  pairwise orthogonal 2-partitions. Consider  $\binom{q}{2}$  copies  $X^{kj}$  ( $1 \leq k < j \leq q$ ) of the set  $X$  and on each set  $X^{kj}$  a system of pairwise orthogonal 2-partitions  $\mathcal{A}_i^{kj} = (A_i^{kj}, B_i^{kj})$ ,  $i = 1, \dots, n$ . For every  $i$  define a  $q$ -partition  $\mathcal{C}_i = (C_{i1}, C_{i2}, \dots, C_{iq})$  of the set  $Y =$

$= \bigcup_{k < j} X^{kj}$  by formula

$$C_{ik} = \bigcup_{j=1}^{k-1} A_i^{jk} \cup \bigcup_{j=k+1}^q B_i^{kj}, \quad k = 1, \dots, q.$$

It is easy to verify that  $\mathcal{C}_i, i = 1, \dots, n$ , is a system of pairwise orthogonal  $q$ -partitions.

**Theorem 2.3.**  $OA(q, n, t)$  exists if and only if  $f(t, q) \geq n$ .

*Proof.* The proof follows from the fact that each  $q$ -partition  $\mathcal{A}$  of the set  $\{1, 2, \dots, t\}$  is in a 1-1 correspondence with the vector  $\mathbf{u} = (u_1, u_2, \dots, u_t)$  defined as follows:  $u_j = k$  iff  $j \in A_k$  where  $A_k$  is the  $k$ -th class of  $\mathcal{A}$ .

Theorems 1.1-1.4 and Theorem 2.3 imply immediately

**Theorem 2.4.** Let  $t, t_1, t_2, n, n_1, n_2, q, q_1, q_2$  be positive integers. Then

- (i)  $f(t_1, q_1) \geq n, f(t_2, q_2) \geq n$  implies  $f(t_1 t_2, q_1 q_2) \geq n$ ;
- (ii)  $f(t_1, q) \geq n_1, f(t_2, q) \geq n_2$  implies  $f(t_1 + t_2, q) \geq n_1 n_2$ .

**Theorem 2.5.** Let  $q$  be a power of a prime. Then

- (i)  $f(q^2, q) = q + 1$ ;
- (ii)  $f(t_j, q) \geq n_j$  for  $j = 1, \dots, q$  implies  $f(t_1 + \dots + t_q, q) \geq q n_1 n_2 \dots n_q$ .

An easy computation gives the following

**Corollary 2.6.** For  $q$  a power of a prime and  $t$  of the form

$$t = \alpha_j q^j + \alpha_{j-1} q^{j-1} + \dots + \alpha_1 q + \alpha_0$$

it is

$$f(t, q) \geq \prod_{i=2}^j [(q+1)^{q^{i-2}} q^{(q^{i-1}-1)/(q-1)}]^{z_i}$$

and hence e.g.  $f(t, q) \geq (q+1)^{\lceil t/q^2 \rceil}$ .

**Remark 2.7.** For  $q = 3, 4$  Theorem 2.2 gives a better result than Corollary 2.6. Nevertheless, we have not been able to generalize the construction given in 2.2 to improve the estimates given in 2.6 for  $q > 3$ .

**Theorem 2.8.** Let  $t, p, q$  be positive integers,  $p \leq q$ . Then

$$f(t, q) \leq f(t, p) f(q, p).$$

Particularly,  $f(t, q) \leq f(t, q-1)$ .

*Proof.* Let  $\mathfrak{S}$  and  $\mathfrak{T}$  be systems of orthogonal  $q$ -partitions and  $p$ -partitions on the  $t$ -set and  $q$ -set, respectively. To every pair  $\mathcal{A}, \mathcal{B}, \mathcal{A} = (A_1, \dots, A_q), \mathcal{B} = (B_1, \dots, B_p)$ ,

$\mathcal{A} \in \mathfrak{S}, \mathcal{B} \in \mathfrak{T}$ , assign  $\mathcal{C}(\mathcal{A}, \mathcal{B}) = (C_1, \dots, C_p)$  defined as follows:

$$C_i = \cup\{A_j | j \in B_i\}, \quad i = 1, \dots, p.$$

$\{\mathcal{C}(\mathcal{A}, \mathcal{B}) | \mathcal{A} \in \mathfrak{S}, \mathcal{B} \in \mathfrak{T}\}$  obviously forms a system of pairwise orthogonal  $p$ -partitions.

**Corollary 2.9.**

$$f(t, q) \geq (2q)^{t/4q^2}$$

for every integers  $t, q, t \geq 8q^3$ .

*Proof.* Immediately follows from 2.6 and 0.3.

**Proposition 2.10.** *Let  $t > q$  be positive integers. Then*

$$f(t, q) \leq \binom{t-1}{\left[ \frac{t}{q} - 1 \right]}.$$

*Proof.* Let  $\mathfrak{S}$  be a maximal system of pairwise orthogonal  $q$ -partitions of a set  $\{1, \dots, t\}$ . From every  $\mathcal{A}_j \in \mathfrak{S}, j = 1, \dots, f(t, q)$  choose one of its partition classes  $A^j \in \mathcal{A}_j$  with  $|A^j| \leq t/q$ . The system  $A^j$  has the following properties:

$$|A^j| \leq t/q, \quad A^i \not\subset A^j \quad \text{and} \quad A^i \cap A^j \neq \emptyset \quad \text{for every} \quad i \neq j.$$

Thus the Erdős Ko Rado theorem yields

$$|\mathfrak{S}| \leq \binom{t-1}{\left[ \frac{t}{q} - 1 \right]}.$$

### 3. $\bar{\omega}(nK_q)$

We shall now give estimates of the number  $\bar{\omega}$ , defined in the introduction, for the graphs  $nK_q$ . (The graph  $nK_q$  consists of  $n$  disjoint copies of the complete graph  $K_q$ .) The notion of  $\bar{\omega}$  was in fact studied in a "complementary form" in several papers, e.g. [4], [15]. The number  $\omega(G)$  is defined in [6] as the minimal size of a set  $S$  such that  $G$  is isomorphic to an intersection graph of a system of subsets of the set  $S$ .

**Proposition 3.1.**  $\bar{\omega}(G) = \omega(\bar{G})$  where  $\bar{G}$  denotes the complement of  $G$ . Hence  $\bar{\omega}(G)$  equals to the minimal size of a set  $S$  such that to every vertex of  $G$  one can assign a subset of  $S$  so that two vertices are adjacent iff the corresponding subsets are disjoint. (This assignment is called a disjoint representation of  $G$ .)

The proof immediately follows from Proposition 1, Chapter 17, § 4 in [0].

**Theorem 3.2.**

- (i)  $\bar{\omega}(nK_q) \leq t$  iff  $\mathbf{OA}(q, n, t)$  exists;
- (ii)  $\bar{\omega}(nK_q) = \text{Min} \{t \mid \text{there exists at least } n \text{ orthogonal } q\text{-partitions of the set } \{1, 2, \dots, t\}\}$ .

Proof. We shall write  $nK_q = K_q^1 + \dots + K_q^n$  where  $V(K_q^i) = \{x_1^i, \dots, x_q^i\}$ . Let  $A_1, \dots, A_t$  be a system of stable sets so that every pair of nonadjacent vertices of  $nK_q$  is contained in at least one of  $A_i$ 's. Without loss of generality we may suppose  $A_j$  to be maximal, i.e.  $|A_j| = n$  for each  $j$ . To every  $j \in \{1, \dots, t\}$  assign a column vector  $(u_{ij})_{1 \leq i \leq n}$  so that

$$(*) \quad u_{ij} = k \quad \text{iff} \quad x_k^i \in A_j.$$

The above defined system of column vectors forms an  $\mathbf{OA}(q, n, t)$ . On the other hand, if an  $\mathbf{OA}(q, n, t)$  exists then the formula (\*) defines a system of  $A_j$ 's with the required properties.

(ii) follows from (i) and Theorem 2.3.

The proof of the following theorem is easy and therefore it is omitted.

**Theorem 3.3.** Let  $A_1, \dots, A_{q,r}$  be a system of stable sets of  $nK_q$  such that every pair of nonadjacent vertices is contained in at least one of them. Then the sets  $A_{(s-1)q+1}, A_{(s-1)q+2}, \dots, A_{s,q}$  form a partition of  $V(nK_q)$  for every  $s = 1, \dots, r$  iff the corresponding  $\mathbf{OA}$  (constructed in the proof of the above theorem) has the permutation property.

Theorem 3.2 and the results of § 2 imply

**Corollary 3.4.**

- (i)  $\bar{\omega}(nK_2) = \text{Min} \left\{ t / \left( \binom{t-1}{\lfloor \frac{t}{2} \rfloor} - 1 \right) \geq n \right\},$
- (ii)  $\frac{\log n}{\frac{1}{q} \log q + \left(1 - \frac{1}{q}\right) \log \frac{q}{q-1}} \leq \bar{\omega}(nK_q) \leq \frac{q^2 \log n}{\log q} (1 + o(1))$  for  $q \geq 3$ .

4. PRODUCT

By a categorical product of two graphs  $G, H$  we mean the following graph  $G \times H$ :

$$V(G \times H) = V(G) \times V(H);$$

$$E(G \times H) = \{ \{ \langle x_1, y_1 \rangle, \langle x_2, y_2 \rangle \} \mid \{x_1, x_2\} \in E(G), \{y_1, y_2\} \in E(H) \}.$$



**Theorem 4.1.**

$$\bar{\omega}(G) + \bar{\omega}(H) \geq \bar{\omega}(G \times H) > \frac{1}{2}[\log_2 \bar{\omega}(G) + \log_2 \bar{\omega}(H)].$$

*Proof.* Let  $C_1, C_2, \dots, C_l$  and  $D_1, D_2, \dots, D_k$  be systems of stable sets covering all pairs of nonadjacent vertices of  $G$  and  $H$ , respectively. Then  $C_i \times V(H), V(G) \times D_j, i = 1, \dots, l, j = 1, \dots, k$ , is a system covering all pairs of nonadjacent vertices of  $G \times H$ . This proves the first inequality.

For the proof of the second inequality we shall need the following three propositions.

Let  $G$  be a graph. Define an equivalence  $\sim$  on the set of its vertices as follows:  $x \sim y$  iff  $x$  and  $y$  have the same neighbourhoods. From each equivalence class choose one point and consider the subgraph induced on this set. Denote this graph by  $m(G)$ .

**Proposition 4.2.**  $|V(m(G))| \geq \sqrt{(2\bar{\omega}(G))}$  for every graph  $G$ .

*Proof.* It follows from the well known theorem of Erdős, Goodman and Posa [4] which states that  $\omega(G) \leq |V(G)|^2/4$ , and from the fact that  $\bar{\omega}(m(G)) = \bar{\omega}(G)$ .

**Proposition 4.3.**

$$m(G) \times m(H) = m(G \times H).$$

*Proof.* It suffices to realize that a neighbourhood of a vertex  $\langle x, y \rangle$  of the product is the product of neighbourhoods of vertices  $x$  and  $y$ .

**Proposition 4.4.**  $\bar{\omega}(F) \geq \log_2 |V(F)|$  for  $F = m(F)$ .

*Proof.* Consider the disjoint representation  $\mathcal{S}$  of the graph  $F$ . From  $F = m(F)$  it follows that  $\mathcal{S}$  is a system of distinct sets and hence  $|V(F)| \leq 2^{\bar{\omega}(F)}$ .

Now we can prove the second inequality of 4.1:

$$\begin{aligned} \bar{\omega}(G \times H) &= \bar{\omega}(m(G \times H)) \geq \log_2 (|V(m(G \times H))|) = \\ &= \log_2 (V(m(G)) \cdot V(m(H))) = \log_2 |V(m(G))| + \log_2 |V(m(H))| > \\ &> \frac{1}{2}[\log_2 \bar{\omega}(G) + \log_2 \bar{\omega}(H)]. \end{aligned}$$

**Remark 4.5.** We have just proved:

If we denote  $f(r) = \min \{\bar{\omega}(G \times H) / \bar{\omega}(G) = \bar{\omega}(H) = r\}$  then  $f(r) \rightarrow \infty$  with  $r \rightarrow \infty$ .

An analogous statement for the chromatic number is not known (see [14]).

## 5. DIMENSION

The notion of dimension of graphs, defined in the introduction, was introduced in [11] and [8]. For a survey of recent results concerning this notion see [10].

The following proposition is shown in [11].

**Proposition 5.1.** *The dimension of  $G$  equals the minimal number of equivalences  $E_1, \dots, E_n$  such that*

$$1) E(G) = \bigcup_{i=1}^n E_i;$$

$$2) \bigcap_{i=1}^n E_i = 0.$$

In [8], dimension of the graphs  $nK_2$  was studied and it was shown that  $\dim nK_2 = \log_2^+ n + 1$ . We give here some estimates of  $\dim nK_q$  for  $q \geq 3$ .

**Proposition 5.2.** *If there exists  $\mathbf{OA}(q, n, qr)$  with the permutation property then  $\dim nK_q \leq r$ .*

Proof follows immediately from Theorem 3.3 and Proposition 5.1.

**Theorem 5.3.**

(i)  $\dim(qK_q) = q$  for  $q$  a power of a prime;

(ii)  $\dim(n_1 n_2 K_q) \leq \dim(n_1 K_q) + \dim(n_2 K_q)$ ;

(iii)  $\dim(nK_{q_1 q_2}) \leq \dim nK_{q_1} \cdot \dim nK_{q_2}$ .

Proof. According to (ii) of 1.1, there exists  $\mathbf{OA}(q, q, q^2)$  with the permutation property and hence according to 5.2,  $\dim qK_q \leq q$ . The second inequality follows from the easy fact that  $\dim(K_q + x) = q$ . ( $K_q + x$  is the disjoint union of  $K_q$  and a single vertex  $x$ .)

(ii) As  $n_1 n_2 K_q$  is an induced subgraph of  $n_1 K_q \times n_2 K_q$  we have  $\dim(n_1 n_2 K_q) \leq \dim(n_1 K_q \times n_2 K_q) \leq \dim n_1 K_q + \dim n_2 K_q$ .

(iii) It is easily seen that  $\dim G = r$  if one can assign to each vertex  $x$  of  $G$  a vector  $\mathbf{v}(x)$  with  $r$  coordinates so that  $x = y$  implies  $\mathbf{v}(x) = \mathbf{v}(y)$  and  $(x, y) \in E(G)$  iff  $v_i(x) \neq v_i(y)$  for every  $i = 1, \dots, r$ . ( $v_i(x)$  denotes the  $i$ -th coordinate of  $\mathbf{v}(x)$ .) We shall call such an assignment an encoding.

Denote by  $x_j^i$  and  $x_k^i$  the  $j$ -th and the  $k$ -th vertex of the  $i$ -th copy of  $K_{q_2}$  in the graph  $nK_{q_2}$  and  $nK_{q_1}$ , respectively. Put  $nK_{q_1 q_2} = K_{q_1 q_2}^1 + \dots + K_{q_1 q_2}^n$ , where  $V(K_{q_1 q_2}^i) = \{x_{1,1}^i, \dots, x_{j,k}^i, \dots, x_{q_1, q_2}^i\}$ . Let  $\mathbf{u}$  and  $\mathbf{v}$  be an encoding of  $nK_{q_1}$  and  $nK_{q_2}$ , respectively. Define a mapping  $\mathbf{w}$  as follows:  $\mathbf{w}_{\alpha, \beta}(x_{j,k}^i) = \langle \mathbf{u}_\alpha(x_j^i), \mathbf{v}_\beta(x_k^i) \rangle$ . It can be verified easily that  $\mathbf{w}$  is an encoding.

**Theorem 5.4.**

$$\dim nK_q \leq \frac{q \log n}{\log q} (1 + o(1)) \text{ for } q \geq 3.$$

Proof follows from 1.4, 5.2 and 0.3.

Let  $n > q$  be positive integers. Define the graph  $\binom{n}{q}$  whose vertices are  $q$ -point subsets of the set  $\{1, \dots, n\}$  with two vertices adjacent if and only if they are disjoint subsets.

The graphs  $\binom{n}{q}$  have interesting properties. It is not difficult to see that they are universal, i.e. every finite graph is an induced subgraph of some  $\binom{n}{q}$ . Kneser conjectured that the chromatic number of  $\binom{2n+k}{n}$  is  $k+2$ . Lovasz proved this in [7] using methods of algebraic topology. In [13] we studied  $\bar{\omega}$  of  $\binom{n}{q}$  and proved that  $\bar{\omega}\left(\binom{n}{q}\right) = n$  for  $q \leq \frac{1}{2}n$ .

In [12] the dimension of Kneser graphs was studied. It was proved that

$$(*) \quad \log_2^+ \log_2^+ n - o(1) \leq \dim \binom{n}{q} \leq (q-1)q^2 \log_2^+ \log_2^+ n.$$

Using 2.9 and [12, Remark 3.7] we obtain a slight improvement of (\*):

$$\dim \binom{n}{q} \leq \frac{4(q-1)q^2}{1 + \log_2 q} \log_2^+ \log_2^+ n.$$

( $\log_2^+ n$  denotes the smallest integer not less than  $\log_2 n$ ).

## 6. PROBLEMS

There are many open problems left. We shall mention only three of them.

- 1) Is  $nK_q$  an induced subgraph of a product of  $\dim nK_q$  copies of  $K_q$ ?  
(I.e., is the condition in Prop. 5.2 also necessary?)
- 2) Find a better estimates for  $f(n, q)$ !
- 3) Let  $G$  be an arbitrary graph. Is it true that

$$\bar{\omega}(nG) \geq \bar{\omega}(nK_{\chi(G)})?$$

## References

- [0] C. Berge: Graphs and Hypergraphs, North. Holland Publ. Company 1973.
- [1] R. C. Bose: On the application of the properties of Galois fields to the construction of hyper-Graeco-Latin squares, Sankhya 3 (1938), 323–338.
- [2] P. Erdős, Chao Ko and R. Rado: Intersection theorems for systems of finite sets, Quart. J. Math. Oxford S 12 (1961), 313–320.
- [3] P. Erdős, S. Chowla and E. G. Straus: On the maximal number of pairwise orthogonal Latin squares of a given order, Canad. J. Math. 12 (1960), 204–208.

- [4] *P. Erdős, A. W. Goodman and L. Pósa*: The representation of a graph by intersections, *Canad. J. Math.* 18 (1966), 106—112.
- [5] *M. Hall*: *Combinatorial Theory*, Blaisdell Publishing Company, Waltham (Massachusetts), Toronto, London, 1967.
- [6] *F. Harary*: *Graph Theory*, Addison-Wesley 1969.
- [7] *L. Lovasz*: Kneser's conjecture, chromatic number and homotopy, *J. Comb. Th. A* 25, 3 (1978), 319—325.
- [8] *L. Lovasz, J. Nešetřil and A. Pultr*: On a product dimension of graphs, to appear in *J. Comb. Th. B*.
- [9] *H. F. Mac Neish*: Euler squares, *Ann. Math.*, 23 (1922), 221—227.
- [10] *J. Nešetřil, A. Pultr*: Product and other representation of graphs and related characteristics, to appear in *Proc. Conf. Algebraic Methods in Graph Theory, Szeged 1978*.
- [11] *J. Nešetřil, V. Rödl*: A simple proof of the Galvin Ramsey property of the class of all finite graphs and the dimension of graphs, *Discrete Math.* 23 (1978), 49—55.
- [12] *S. Poljak, A. Pultr and V. Rödl*: On the dimension of the Kneser graphs, to appear in *Proc. Conf. Algebraic Methods in Graph Theory, Szeged 1978*.
- [13] *S. Poljak, V. Rödl*: Set systems determined by intersections, to appear.
- [14] *S. Poljak, V. Rödl*: On arc chromatic number of digraphs, to appear.
- [15] *S. Poljak, V. Rödl and D. Turzik*: Complexity of covering of edges by complete graphs, to appear.

*Authors' addresses*: S. Poljak, 118 00 Praha 9, Sokolovská 83, ČSSR (Matematicko-fyzikální fakulta UK); V. Rödl, 110 00 Praha 1, Husova 5, ČSSR (Fakulta jaderná a fyzikálně inženýrská ČVUT).