

Pacific Journal of Mathematics

OSCILLATORY PROPERTIES OF A DELAY DIFFERENTIAL EQUATION OF EVEN ORDER

RAYMOND D. TERRY

OSCILLATORY PROPERTIES OF A DELAY DIFFERENTIAL EQUATION OF EVEN ORDER

RAYMOND D. TERRY

A classification of nonoscillatory solutions according to the sign properties of their derivatives is introduced for a general nonlinear delay differential equation of order $2n$. It is seen that there are n types of positive solutions of this equation. An intermediate Riccati transformation is employed to obtain integral criteria for the nonexistence of such solutions and for the oscillation of all solutions. Analysis of the Taylor Remainder gives rise to a summability condition which is used to investigate the asymptotic behavior of a class of solutions. The major results are then shown to be special cases of a more general result based on the direct method of Lyapunov.

The purpose of this paper is to discuss the oscillatory and nonoscillatory behavior of solutions of the nonlinear delay differential equation of order $2n$:

$$(1.1) \quad D^n[r(t)D^n y(t)] + y_\tau(t)f(t, y_\tau(t)) = 0,$$

where $y_\tau(t) = y[t - \tau(t)]$, $0 \leq \tau(t) \leq T$ and $0 < m \leq r(t) \leq M$. Throughout the first three sections $f(t, u)$ is assumed to satisfy the following three hypotheses:

- (i) $f(t, u)$ is a continuous real-valued function on $[0, \infty) \times R$, $R = (-\infty, \infty)$;
- (ii) for each fixed $t \in [0, \infty)$, $f(t, u) < f(t, v)$ for $0 < u < v$; and
- (iii) for each fixed $t \in [0, \infty)$, $f(t, u) > 0$ and $f(t, u) = f(t, -u)$ for $u \neq 0$.

In section four, these assumptions on $f(t, u)$ will be replaced by others as indicated there.

1. A solution $y(t)$ of (1.1) is said to be oscillatory on $[0, \infty)$ if for each $t_0 > 0$, there exists a $T_0 > t_0$ such that $y(T_0) = 0$; it is called nonoscillatory otherwise. Following Kiguradze [4], we say that a solution $y(t)$ is of type A_j if for sufficiently large t the derivatives $D^k y(t) > 0$, $k = 0, 1, \dots, 2j + 1$ and $(-1)^{k+1} D^k y(t) > 0$, $k = 2j + 2, \dots, 2n - 1$. In an analogous manner we say that a solution $y(t)$ is of type B_j if for sufficiently large t , $y_k(t) > 0$ for $k = 0, \dots, 2j + 1$ and $(-1)^{k+1} y_k(t) > 0$, $k = 2j + 2, \dots, 2n - 1$ where

$$y_k(t) = D^k y(t), \quad k = 0, \dots, n - 1$$

and

$$y_k(t) = D^{k-n}[r(t)D^n y(t)], \quad k = n, \dots, 2n - 1.$$

When $r(t) \equiv 1$, these definitions reduce to those of types A_j ($0 \leq j \leq n - 1$) respectively. In [4] Kiguradze proved a fundamental lemma which we state here as follows.

LEMMA 1.1. *Let $u(t)$ be a continuous nonnegative function on $(0, \infty)$ having continuous derivatives up to order $2n$ inclusive which do not change sign on this interval. If $D^{2n}u(t) \leq 0$, then there exists an integer $p = 0, \dots, n - 1$ such that*

$$\begin{aligned} D^k u(t) &\geq 0, & k &= 0, \dots, l, \\ (-1)^{k+l} D^k u(t) &\geq 0, & k &= l + 1, \dots, 2n - 1, \end{aligned}$$

where $l = 2p + 1$. Furthermore, $0 \leq D^l u(t) \leq l! t^{-l} u(t)$.

In view of this result, all nonoscillatory solutions of (1.1) with $r(t) \equiv 1$ are of type A_j for some $j = 0, \dots, n - 1$. In the general case we argue as follows. Suppose $y(t)$ is a nonoscillatory solution of (1.1) which we may assume to be nonnegative because of (iii). First, no two successive y_k , $k \geq 1$ can ultimately be negative. Suppose $k \geq n + 1$ and y_k and y_{k+1} are negative for $t > a$. Then y_k is a negative decreasing function on $[a, \infty)$ and there exists a constant $C_k > 0$ such that $y_k(t) < -C_k$ for $t > a$. Thus

$$y_{k-1}(t) - y_{k-1}(a) = \int_a^t y_k(s) ds < -C_k(t - a),$$

which implies that y_{k-1} is eventually negative. Next, if y_n and y_{n+1} are eventually negative, say for $t > a$, then

$$[M y_{n-1}(s)]_a^t = M \int_a^t D^n y(s) ds < \int_a^t y_n(s) ds < -C_n(t - a),$$

which implies that y_{n-1} is eventually negative. Since $r(t) > 0$ and $y_n(t) < 0$, $D^n y(t) < 0$. Using the negativity of $D^n y$ and $D^{n-1} y$ we may show as in the first part that y_{n-k} is eventually negative for any $k = 2, \dots, n$ which contradicts the positivity of $y(t)$.

Using the same technique as above, it follows that if any two consecutive y_k , $k \geq 1$ are ultimately positive, then all the preceding y_k are eventually positive. We conclude that a positive solution of (1.1) is necessarily of type B_j for some $j = 0, \dots, n - 1$. Thus the nonexistence of nonoscillatory solutions of type B_j ($0 \leq j \leq n - 1$) will imply that all solutions are necessarily oscillatory.

In section two integral criteria are given for the nonexistence of solutions of type B_j as well as for the oscillation of all solutions

of (1.1). Section three provides a necessary and sufficient condition for the existence of a nonoscillatory solution of (1.1) with $r(t) \equiv 1$ having prescribed asymptotic form. Recently Yoshizawa [11, 12] has applied the direct method of Lyapunov to study the oscillatory behavior of solutions of certain nonlinear second order ordinary differential equations. In section four his technique is employed to investigate the nonoscillatory behavior of (1.1). For recent related studies see the papers of Burkowski [2], Gollwitzer [3], Staikos and Petsoulas [7], and Wong [9].

2. In this section integral criteria for the nonexistence of solutions of types $B_j(j = 0, \dots, n - 1)$ are derived. To obtain these results we shall first prove several lemmas.

LEMMA 2.1. *Let $y(t)$ be a solution of (1.1) of type B_j where either (i) n is even and $j \leq (n - 2)/2$ or (ii) n is odd and $j \leq (n - 3)/2$. Then for all sufficiently large t ,*

$$ty_k(t) \leq 2(2j + 2 - k)y_{k-1}(t), \quad k = 1, \dots, 2j + 1.$$

LEMMA 2.2. *Let $y(t)$ be a B_j -solution of (1.1) where n is odd and $j = (n - 1)/2$. Then for all sufficiently large t ,*

$$ty_n(t) \leq 2My_{n-1}(t)$$

and

$$ty_k(t) \leq 2[Mm^{-1} + (n - k)]y_{k-1}(t), \quad k=1, \dots, n - 1.$$

LEMMA 2.3. *Let $y(t)$ be a B_j -solution of (1.1) where either (i) n is even and $j \geq n/2$ or (ii) n is odd and $j \geq (n + 1)/2$. Then for all sufficiently large t ,*

$$(a) \quad ty_k(t) \leq 2(2j + 2 - k)y_{k-1}(t), \quad k = n + 1, \dots, 2j + 1.$$

$$(b) \quad ty_n(t) \leq 2M(2j - n + 2)y_{n-1}(t), \text{ and}$$

$$(c) \quad ty_k(t) \leq 2[M(2j - n + 2)m^{-1} + (n - k)]y_{k-1}(t),$$

$$k = 1, \dots, n - 1.$$

The proof of each of the three lemmas is elementary using only integration by parts and the definition of a B_j -solution. For brevity, we will prove only Lemma 2.3.

Proof. Suppose $y(t)$ is a solution of type B_j . Then there is a $T_0 > 0$ such that $y_k(t) > 0$, ($0 \leq k \leq 2j + 1$) and $y_{2j+2}(t) < 0$ for $t \geq T_0$. Hence, $y_\tau(t) > 0$ for $t - \tau(t) \geq T_0$, i.e., for $t \geq T_0 + T = T_1$. Thus, $y_{2j+1}(t)$ is a decreasing function of t for $t \geq T_1$. Consequently,

$$y_{2j}(t) \geq [y_{2j}(s)]_{T_1}^t = \int_{T_1}^t y_{2j+1}(s)ds \geq (t - T_1)y_{2j+1}(t).$$

Since $(t - T_1) \geq t/2$ for $t \geq 2T_1$, we have $y_{2j}(t) \geq ty_{2j+1}(t)/2$ which proves (a) for the case $k = 2j + 1$.

We proceed inductively and suppose that for some integer k , $n + 1 < k \leq 2j + 1$,

$$(2.1) \quad (t - T_1)y_k(t) \leq (2j + 2 - k)y_{k-1}(t).$$

An integration of (2.1) yields

$$[(s - T_1)y_{k-1}(s)]_{T_1}^t - \int_{T_1}^t y_{k-1}(s)ds = \int_{T_1}^t (s - T_1)y_k(s)ds$$

so that

$$\begin{aligned} (t - T_1)y_{k-1}(t) &\leq [2j + 2 - (k - 1)] \int_{T_1}^t y_{k-1}(s)ds \\ &\leq [2j + 2 - (k - 1)]y_{k-2}(t), \end{aligned}$$

which proves (a). Specifically, for $k = n + 1$, (2.1) becomes

$$(t - T_1)y_{n+1}(t) \leq (2j - n + 1)y_n(t).$$

Integrating this by parts results in

$$\begin{aligned} (t - T_1)y_n(t) - \int_{T_1}^t y_n(s)ds &= \int_{T_1}^t (s - T_1)y_{n+1}(s)ds \\ &\leq (2j - n + 1) \int_{T_1}^t y_n(s)ds. \end{aligned}$$

Thus, for $t \geq T_1$ we obtain

$$(2.2) \quad \begin{aligned} (t - T_1)y_n(t) &\leq (2j - n + 2) \int_{T_1}^t y_n(s)ds \\ &\leq (2j - n + 2)My_{n-1}(t). \end{aligned}$$

As in (a) the estimate for $t \geq 2T_1$ becomes

$$ty_n(t) \leq 2(2j - n + 2)My_{n-1}(t),$$

which proves (b).

Now integration of (2.2) by parts establishes the anchor for an inductive proof of (c).

$$\begin{aligned} [m(s - T_1)y_{n-1}(s)]_{T_1}^t - m \int_{T_1}^t y_{n-1}(s)ds &= m \int_{T_1}^t (s - T_1)D^n y(s) \\ &\leq \int_{T_1}^t (s - T_1)y_n(s)ds \\ &\leq (2j - n + 2)M \int_{T_1}^t y_{n-1}(s)ds. \end{aligned}$$

Thus, for $t \geq 2T_1$, we have

$$\begin{aligned} mty_{n-1}(t)/2 &\leq m(t - T_1)y_{n-1}(t) \leq [(2j - n + 2)M + m] \int_{T_1}^t y_{n-1}(s)ds \\ &\leq [(2j - n + 2)M + m]y_{n-2}(t), \end{aligned}$$

which proves (c) in the case $k = n - 1$. Now assume inductively that

$$(t - T_1)y_k(t) \leq [M(2j - n + 2)m^{-1} + (n - k)]y_{k-1}(t)$$

for some $k, 1 < k \leq n - 1$. Then

$$\begin{aligned} (t - T_1)y_{k-1}(t) - \int_{T_1}^t y_{k-1}(s)ds \\ = \int_{T_1}^t (s - T_1)y_k(s)ds \\ \leq [M(2j - n + 2)m^{-1} + (n - k)] \int_{T_1}^t y_{k-1}(s)ds \end{aligned}$$

so that for $t \geq T_1$

$$(t - T_1)y_{k-1}(t) \leq [M(2j - n + 2)m^{-1} + (n - k + 1)]y_{k-2}(t)$$

and for $t \leq 2T_1$

$$ty_{k-1}(t) \leq 2[M(2j - n + 2)m^{-1} + (n - k + 1)]y_{k-2}(t)$$

which proves (c).

We remark that if $y(t)$ is a B_j -solution of (1.1) on $[0, \infty)$ with $r(t) \equiv 1$ and $\tau(t) \equiv 0$, we may take $T_1 = 0$ and $m = M = 1$. The above proof will yield

$$ty_{2j+2-k} \leq ky_{2j+1-k}$$

for $k = 1, \dots, 2j + 1$, which is another form of Kiguradze's lemma [4].

LEMMA 2.4. *Let $y(t)$ be a solution of (1.1) of type B_j . Then $y_{2j}(t - \tau(t)) \sim y_{2j}(t)$.*

Proof. Let $y(t)$ be a B_j -solution of (1.1). For $j \neq (n - 1)/2$ and $t \geq T_1$, $y_\tau(t)$ and $y_k(t)$, $k = 0, \dots, 2j + 1$ are all positive while $y_{2j+2}(t) < 0$. Since $\tau(t) \geq 0$ and $y_{2j}(t)$ is an increasing function on (T_1, ∞) , $y_{2j}(t - \tau(t)) < y_{2j}(t)$, so that with the help of Lemma 2.1 or 2.3, we have

$$\left| \frac{y_{2j}(t - \tau(t))}{y_{2j}(t)} - 1 \right| = \tau(t) \frac{y_{2j+1}(s)}{y_{2j}(t)} < \frac{2T}{s} \frac{y_{2j}(s)}{y_{2j}(t)} < \frac{2T}{s},$$

where $t - \tau(t) \leq s \leq t$. Since s tends to infinity with t , the lemma follows in this case. The case $j = (n - 1)/2$ follows in a similar manner using Lemma 2.2 and the estimate $mDy_{2j}(s) \leq y_{2j+1}(s)$.

We remark that this lemma is analogous to one proved by Bradley [1] for the linear equation

$$y''(t) + p(t)y_\tau(t) = 0 .$$

His proof can be modified to yield Lemma 2.4. We note that Lemmas 2.1, 2.2, and 2.3 are valid for unbounded t , provided $\lim_{t \rightarrow \infty} (t - \tau(t)) = +\infty$. A weaker version of Lemma 2.4 is also true for unbounded $\tau(t)$, provided $0 \leq \tau(t) \leq \mu t$, where μ can be specified. In general, if $y(t)$ is a B_j -solution of (1.1) described by Lemmas 2.1 or 2.3, then we may take $\mu < 1/2$; otherwise the stronger estimate $\mu < m/(M + m)$ is required. The conclusion of Lemma 2.4 is changed to read: There are constants $k_j > 0$ and $t_j > 0$ such that $y_{2j}(t - \tau(t)) > k_j y_{2j}(t)$ for $t > t_j$. With these lemmas we can now give criteria for the non-existence of solutions of type $B_j(0 \leq j \leq n - 1)$.

THEOREM 2.5. *Suppose that for all constants $C > 0$ and some $j = 0, \dots, (n - 1)$,*

$$(2.3) \quad \int^\infty t^{2j} f(t, Ct^{2j}) dt = +\infty .$$

Then (1.1) has no solutions of type B_j .

Proof. Let $y(t)$ be a positive solution of type B_j and let $w(t) = y_{2n-1}(t)/y_{2j}(t)$. Then (1.1) shows that

$$w'(t) + y_{2n-1}(t) D y_{2j}(t) y_{2j}^{-2}(t) + y_\tau(t) f(t, y_\tau(t)) y_{2j}^{-1}(t) = 0 .$$

For $j \neq (n - 1)/2$, $D y_{2j}(t) = y_{2j+1}(t) > 0$; if n is odd and $j = (n - 1)/2$, $D y_{2j}(t) = y_{2j+1}(t)/r(t) > 0$ since $r(t) > 0$. Since $y_{2n-1}(t)$ and $y_{2j}(t)$ are positive for $t > T_1$, this reduces to

$$(2.4) \quad w'(t) + y_\tau(t) f(t, y_\tau(t)) y_{2j}^{-1}(t) < 0, t \geq T_1 .$$

There are three cases to consider

- (i) n is even and $j \leq (n - 2)/2$ or n is odd and $j \leq (n - 3)/2$;
- (ii) n is odd and $j = (n - 1)/2$; and
- (iii) n is even and $j \geq n/2$ or n is odd and $j \geq (n + 1)/2$.

Applying Lemmas 2.1, 2.2, and 2.3 to cases (i), (ii), and (iii) respectively, we obtain

$$t^{2j} y_{2j}(t) \leq M_j y(t) ,$$

where

$$2^{2j} (2j + 1)! , \quad \text{case (i),}$$

$$M_j = 2^{2j} \prod_{k=1}^{2j} [Mm^{-1} + (n - k)] , \quad \text{case (ii),}$$

$$2^{2j}(2j - n + 2)! M \prod_{k=1}^{n-1} [(2j - n + 2)Mm^{-1} + (n - k)] , \quad \text{case (iii).}$$

Letting $N_j = M_j^{-1}$, we have $y(t) \geq N_j t^{2j} y_{2j}(t)$ for $t \geq 2T_1$, so for $t \geq 2T_1 + T = T_2$ we have

$$y_-(t) \geq N_j(t - \tau(t))^{2j} y_{2j}(t - \tau(t)) \geq N_j(t - T)^{2j} y_{2j}(t - \tau(t))$$

because $0 \leq \tau(t) \leq T$. Since $Dy_{2j}(t) > 0$ for $t \geq T_1$, there is a constant C_0 such that $y_{2j}(t) \geq C_0$ for $t \geq T_1$. Hence, $y_{2j}(t - \tau(t)) \geq C_0$ for $t \geq T_1 + T$. Moreover, $(t - T) \geq t/2$ for $t \geq 2T$. Because $T_2 = 2T_1 + T > \max(T_1 + T, 2T)$, both estimates hold for $t \geq T_2$. Combining the above estimates with (2.4) via hypothesis (ii) and using Lemma 2.4, we arrive at

$$w'(t) + NKt^{2j} f(t, Ct^{2j}) \leq 0, \quad t \geq T_3$$

for some $T_3 \geq T_2$, where $N = N_j 2^{-2j}$, $C = 2^{-2j} N_j C_0$ and $K = K_j > 0$ is the constant from Lemma 2.4. An integration of this together with (2.3) shows that $w(t)$ is eventually negative, which is absurd.

Since the divergence of $t^{2j} f(t, Ct^{2j})$ implies that of $t^{2(j+1)} f(t, Ct^{2(j+1)})$ the conclusion of Theorem 2.5 may be strengthened to exclude solutions of type B_k where $j \leq k \leq n - 1$. The theorem may also be restated as follows.

THEOREM 2.6. *Suppose (2.3) holds for all constants $C > 0$ and for some $j = 0, \dots, n - 1$. Then either (1.1) is oscillatory or else $y(t)y_{2j}(t) < 0$ for t sufficiently large.*

For $j = n - 1$ and $r(t) \equiv 1$, Theorem 2.6 reduces to the alternative that either (1.1) is oscillatory or else $y(t)D^{2n-2}y(t) < 0$ for t sufficiently large, which is essentially Theorem 3.1 of Ladas [5].

Moreover, in view of the fact that all positive solutions are of types B_j for some $j(0 \leq j \leq n - 1)$, we can immediately restate Theorem 2.5 as a criterion of oscillation.

COROLLARY 2.7. *If for all constants $C > 0$*

$$\int^{\infty} f(t, C)dt = +\infty ,$$

then all solutions of (1.1) are oscillatory.

COROLLARY 2.8. *Suppose $p(t)$ is continuous and eventually positive and that*

$$\int^{\infty} p(t)dt = +\infty .$$

Then all solutions of the equation

$$(2.5) \quad D^n[r(t)D^n y(t)] + p(t)y_\tau^{2\lambda+1}(t) = 0, \lambda > 0$$

are oscillatory.

The conclusion of Corollary 2.8 is true in the case $\lambda = 0$ in (2.5). In this instance, (2.5) is not a special case of (1.1) since $f(t, u) = p(t)$ does not satisfy hypothesis (ii) of section one. To permit this extension, we may suppose that $y(t)$ is a B_j -solution of (2.5) and let $w(t) = y_{z_{n-1}}(t)/y_{z_j}(t)$. Equation (2.4) becomes

$$w'(t) + p(t)y_\tau(t)y_{z_j}^{-1}(t) < 0, t \geq T_1.$$

Applying Lemmas 2.1-2.4, which are independent of hypothesis (ii), produces the same contradiction as in the proof of Theorem 2.5. Thus, if $\lambda = 0$ and

$$\int^\infty t^{2j} p(t) dt = \infty$$

for some $j = 0, \dots, n-1$, (2.5) has no B_k -solutions for $k = j, \dots, n-1$. Corollary 2.8 then follows by specifying $j = 0$.

NOTE. When $n = 2$ and $j = 1$, Lemma 2.3 reduces to Lemma 2.1 of Terry and Wong [8]. Similarly, letting $n = 2$ and $j = 0, 1$ in Lemma 2.4, we obtain Lemma 2.2 (a), (b) of [8]. Moreover, Theorem 2.5, Corollary 2.7, and Corollary 2.8 here are, respectively, the analogues of Theorem 2.8, Theorem 2.4, and Corollary 2.5 of [8].

3. In this section an asymptotic result is established for the equation

$$(3.1) \quad D^{2n}y(t) + f(t, y_\tau(t))y_\tau(t) = 0,$$

where $f(t, u)$ satisfies the three conditions of section one.

LEMMA 3.1. *Let $y(t)$ be a solution of (3.1) which is eventually positive. Then*

$$D^{2n-1}y(t) \sim (2n-1)! t^{1-2n}y(t).$$

Proof. Suppose $y(t)$ is a solution of (3.1) such that $y(t) > 0$ for $t \geq T_1$. Then $y_\tau(t) > 0$ for $t - \tau(t) \geq T_1$, i.e., for $t \geq T_1 + T = T^*$. By Taylor's theorem with remainder, for $t \geq T^*$

$$(3.2) \quad (2n-1)! R(t) = (2n-1)! y(t) + \int_{T^*}^t (t-s)^{2n-1} y_\tau(s) f(s, y_\tau(s)) ds,$$

where

$$R(t) = \sum_{k=0}^{2n-1} \frac{1}{k!} D^k y(T^*) (t - T^*)^k .$$

Since $y_+(s)$, and hence $-D^{2n}y(s)$, is positive for $s > T^*$, condition (iii) together with $(t - T^*) > (t - s) > 0$ imply that

$$\begin{aligned} &(2n - 1)! R(t) \\ &\leq (2n - 1)! y(t) + (t - T^*)^{2n-1} [D^{2n-1}y(T^*) - D^{2n-1}y(t)] . \end{aligned}$$

Dividing this by $(t - T^*)^{2n-1}$ and noting that

$$\lim_{t \rightarrow \infty} (2n - 1)! (t - T^*)^{1-2n} R(t) = D^{2n-1}y(T^*) ,$$

it follows upon passage to the limit that

$$\begin{aligned} &D^{2n-1}y(T^*) \\ &\leq \varliminf_{t \rightarrow \infty} (2n - 1)! (t - T^*)^{1-2n} y(t) + D^{2n-1}y(T^*) - \lim_{t \rightarrow \infty} D^{2n-1}y(t) . \end{aligned}$$

Hence,

$$(3.3) \quad \lim_{t \rightarrow \infty} D^{2n-1}y(t) \leq \varliminf_{t \rightarrow \infty} (2n - 1)! (t - T^*)^{1-2n} y(t) .$$

To prove the reverse inequality let σ be chosen such that $T^* < \sigma < t$. By restricting s to lie in the interval $[T^*, \sigma]$, we have $(t - s)^{2n-1} \geq (t - \sigma)^{2n-1}$ and

$$\begin{aligned} (2n - 1)! R(t) &\geq (2n - 1)! y(t) + (t - \sigma)^{2n-1} \int_{T^*}^{\sigma} y_+(s) f(s, y_+(s)) ds \\ &= (2n - 1)! y(t) + (t - \sigma)^{2n-1} [D^{2n-1}y(T^*) - D^{2n-1}y(\sigma)] . \end{aligned}$$

Multiplying this by $(t - \sigma)^{1-2n}$, keeping σ fixed and letting $t \rightarrow \infty$ through a sequence of points for which $(t - \sigma)^{1-2n} y(t)$ tends to its limit superior, we obtain

$$\begin{aligned} &D^{2n-1}y(T^*) \\ &\geq \overline{\lim}_{t \rightarrow \infty} (2n - 1)! (t - T^*)^{1-2n} y(t) + D^{2n-1}y(T^*) - D^{2n-1}y(\sigma) \end{aligned}$$

from which it follows that

$$\overline{\lim}_{t \rightarrow \infty} (2n - 1)! (t - T^*)^{1-2n} y(t) \leq D^{2n-1}y(\sigma) .$$

Since σ is arbitrary and $\lim_{t \rightarrow \infty} D^{2n-1}y(t)$ exists, it follows that

$$\overline{\lim}_{t \rightarrow \infty} (2n - 1)! (t - T^*)^{1-2n} y(t) \leq \lim_{t \rightarrow \infty} D^{2n-1}y(t) .$$

Combining this with (3.3) yields the desired result.

THEOREM 3.2. *Equation (3.1) has a positive solution $y(t)$ satisfying*

$$(3.4) \quad \lim_{t \rightarrow \infty} t^{1-2n} y(t) = k, \quad 0 < k < \infty,$$

if, and only if, for some $C > 0$

$$(3.5) \quad \int_{T_0}^{\infty} t^{2n-1} f(t, Ct^{2n-1}) dt < \infty.$$

Proof. Suppose (3.5) holds. A lower limit T_0 can be chosen sufficiently large so that

$$(3.5') \quad \int_{T_0}^{\infty} t^{2n-1} f(t, Ct^{2n-1}) dt < (2n-1)! - 1/2.$$

If $y(t) = y(t, T_0)$ is the solution of (3.1) defined by $D^k y(T_0) = 0$, $k = 0, \dots, 2n-2$, $D^{2n-1} y(T_0) = (2n-1)! C$ and $y(t) = 0$ for $T_0 - T \leq t \leq T_1$, then $y(t, T_0)$ is positive on some open interval whose left endpoint is T_0 . Let $t = T_1$ be the first zero of $y(t, T_0)$ in (T_0, ∞) . By Taylor's theorem we have

$$(3.6) \quad \begin{aligned} (2n-1)! C(t - T_0)^{2n-1} &= (2n-1)! y(t, T_0) \\ &+ \int_{T_1}^t (t-s)^{2n-1} y_{\tau}(s) f(s, y_{\tau}(s)) ds. \end{aligned}$$

Since $y(s) > 0$ for $T_0 - T \leq s \leq T_1$, $y_{\tau}(s) > 0$ for $T_0 - T \leq s - \tau(s) \leq T_1$, i.e., for $s \geq T_0 - T + \tau(s)$. Hence $y_{\tau}(s) > 0$ for $s \geq T_0$. A similar argument shows that $y_{\tau}(s) > 0$ for $s \leq T_1$. Thus

$$(3.7) \quad y(t) = y(t, T_0) \leq C(t - T_0)^{2n-1}, \quad T_0 \leq t \leq T_1.$$

Moreover, letting $t = T_1$ in (3.6) we have

$$(3.8) \quad \begin{aligned} (2n-1)! C(T_1 - T_0)^{2n-1} &= \int_{T_0}^{T_1} (T_1 - s)^{2n-1} y_{\tau}(s) f(s, y_{\tau}(s)) ds \\ &\leq (T_1 - T_0)^{2n-1} \int_{T_0}^{T_1} y_{\tau}(s) f(s, y_{\tau}(s)) ds. \end{aligned}$$

By condition (iii) and (3.7)

$$y_{\tau}(s) f(s, y_{\tau}(s)) \leq C s^{2n-1} f(s, C s^{2n-1}),$$

so that (3.8) yields

$$(2n-1)! \leq \int_{T_0}^{T_1} s^{2n-1} f(s, C s^{2n-1}) ds,$$

which contradicts (3.5') and demonstrates the existence of a positive solution satisfying (3.4).

To prove necessity let $y(t)$ be a positive solution of (3.1) satisfying (3.4). It follows from Lemma 3.1 that

$$(3.9) \quad \int_{T_1}^{\infty} y_{\tau}(s)f(s, y_{\tau}(s))ds = D^{2n-1}y(T_1) - (2n - 1)! k .$$

(3.4) shows that for any $\varepsilon > 0$ there is a $T_* \geq T_1$ such that $y(t) \geq (k - \varepsilon)t^{2n-1}$ for all $t \geq T_*$ so that $(k - \varepsilon)(t - T)^{2n-1} \leq y_{\tau}(t)$ for $t > T^* = T_* + T$. Also, it follows from (ii) that

$$f(s, y_{\tau}(s)) \geq f[s, (k - \varepsilon)(s - T)^{2n-1}] .$$

Since (3.9) is valid with T_1 replaced by T^* ,

$$\begin{aligned} & D^{2n-1}y(T^*) - (2n - 1)! k \\ & \geq (k - \varepsilon) \int_{T^*}^{\infty} (s - T)^{2n-1} f[s, (k - \varepsilon)(s - T)^{2n-1}] ds . \end{aligned}$$

For so $s \geq 2T$,

$$\begin{aligned} & \int_{\infty}^{\infty} s^{2n-1} f(s, Cs^{2n-1}) ds < 2^{2n-1} \\ & \quad \times \int_{\infty}^{\infty} (s - T)^{2n-1} f[s, 2^{2n-1}C(s - T)^{2n-1}] ds \leq N_1 , \end{aligned}$$

where $N_1 = 2^{2n-1}(k - \varepsilon)^{-1}[D^{2n-1}y(T^*) - (2n - 1)!k]$ and $C = (k - \varepsilon)/2^{2n-1}$. The conclusion follows.

For the linear equation (2.5) with $\lambda = 0$, $n = 2$, $r(t) \equiv 1$, and $\tau(t) \equiv 0$, this result reduces to that of Leighton and Nehari [6]. As in the discussion following Corollary 2.8, we observe that when $\lambda = 0$ and $r(t) \equiv 1$, (2.5) is not a special case of (3.1). However, Theorem 3.2 remains valid and the proof given above may be easily modified to yield the result. The details of this are omitted for brevity. For the nonlinear equation (3.1) with $\tau(t) \equiv 0$ and $n = 2$, see also Wong [10]. The case of (3.1) with $\tau(t) \neq 0$ and $n = 2$ is treated by Terry and Wong [8].

4. In this section we shall apply the direct method of Lyapunov to obtain nonoscillation criteria for (1.1). For convenience, we first introduce some notation which will be used in the section. Let $R_a = [a, \infty)$, $a \geq 0$, $R^* = (0, \infty)$, $R_* = (-\infty, 0)$ and $R^1 = R = (-\infty, \infty)$. We shall abbreviate the cartesian products of these intervals as follows:

$$\begin{aligned} R^{p*} &= R^* \times R^* \times \dots \times R^*, \text{ } p \text{ times} \\ R_{p*} &= R_* \times R_* \times \dots \times R_*, \text{ } p \text{ times} \\ R_*^* &= R_* \times R^*; R^*_* = R^* \times R_* . \end{aligned}$$

Other products may be defined in terms of these, e.g.,

$$R_a^{p*} = R_a \times R^{p*} .$$

To begin with we shall consider an arbitrary $2n$ th order equation of the form

$$(4.1) \quad D^n[r(t)D^n y(t)] + F(t, \sigma(t), \sigma_\tau(t)) = 0,$$

where

$$\sigma(t) = (y_0(t), \dots, y_{2n-1}(t)) \text{ and } \sigma_\tau(t) = \sigma(t - \tau(t)).$$

A real-valued function $V(t, \sigma)$ will be called a Lyapunov function if $V(t, \sigma)$ is continuous in its domain and locally Lipschitzian in σ . Following Yoshizawa [11, 12], we define the trajectory derivative $\dot{V}_{(1)}$ of V along solutions $\sigma(t)$ of (4.1) by

$$(4.2) \quad \dot{V}_{(1)}(t, \sigma) = \overline{\lim}_{h \downarrow 0} \frac{1}{h} [V(t+h, \sigma(t+h)) - V(t, \sigma(t))].$$

The first result of this section is an extension of Yoshizawa's theorem to (4.1) and the proof is based on his.

THEOREM 4.1. *Let V be a real-valued continuous function defined on $R_a^{(2j+2)*} \times (R_*^*)^{n-j-1}$ for some $a > 0$ such that*

(i) *V tends to infinity uniformly for $\sigma \in R^{(2j+2)*} \times (R_*^*)^{n-j-1}$ as t tends to infinity; and*

(ii) *for each solution $y(t)$ of (4.1) such that $(t, \sigma(t)) \in R_b^{(2j+2)*} \times (R_*^*)^{n-j-1}$ for some $b \geq a$, $\dot{V}_{(1)}(t, \sigma(t)) \leq 0$.*

Then (4.1) has no solution of type B_j .

Proof. Let $y(t)$ be a solution of type B_j . There is a positive a for which $\sigma(t) \in R_a^{(2j+2)*} \times (R_*^*)^{n-j-1}$ if $t \geq a$. By (ii) for t sufficiently large, i.e., for $t \geq b \geq a$,

$$V(t, \sigma(t)) \leq V(b, \sigma(b)).$$

On the other hand, (i) implies that there is a $c \geq b$ for which

$$V(t, \sigma(t)) > V(b, \sigma(b))$$

if $t > c$, which is a contradiction.

As in other applications of the direct method of Lyapunov, the key to applying this result is the construction of suitable Lyapunov functions V having the requisite properties. In this case of (1.1) Theorem 2.5 may be regarded as a special case of this result, for if $y(t)$ is a B_j -solution there are constants $C > 0$ and $K > 0$ such that for sufficiently large t , $y_\tau(t) \geq NKt^{2j}y_{2j}(t) \geq Ct^{2j}$. It follows that by taking a suitable T^* the function

$$V(t, \sigma(t)) = y_{2n-1}(t)[y_{2j}(t)]^{-1} + NK \int_{T^*}^t s^{2j} f(s, Cs^{2j}) ds$$

will be a Lyapunov function satisfying the conditions of Theorem 4.1 provided (2.4) holds.

Another application of Theorem 4.1 to equation (4.1) is given by the following result.

THEOREM 4.2. *Suppose there exist functions $p(t)$ and $\rho(u)$ of class $C(R_a)$ and $C^1(R)$ respectively such that*

(i) $\rho'(u) \geq 0; u\rho(u) > 0, u \neq 0;$

(ii) $F(t, \sigma(t)) \geq p(t)\rho(y), y \geq 0;$

(iii) $\int_a^\infty p(t)dt = +\infty.$

Then (4.1) has no solutions of type B_j .

Proof. Suppose $y(t)$ is a solution of (4.1) of type B_j . Then $(t, \sigma(t)) \in R_b^{(2j+2)*} \times (R_*^*)^{n-j-1}$ for some $b \geq a$. Let $V(t, \sigma)$ be the function defined by

$$V(t, \sigma(t)) = y_{2n-1}(t)[\rho(y(t))]^{-1} + \int_b^t p(s)ds.$$

In view of hypotheses (i), (ii), and (iii), V will clearly satisfy condition (i) of Theorem 4.1. Moreover, since both $y'(t)$ and $y_{2n-1}(t)$ are positive for large t , a simple calculation with the help of (i) and (ii) shows that

$$\begin{aligned} \dot{V}_{(1)} &= \frac{Dy_{2n-1}(t)}{\rho[y(t)]} + p(t) - \frac{\rho'[y(t)]}{\rho^2[y(t)]}y_{2n-1}(t)y'(t) \\ &\leq p(t) - \frac{F(t, \sigma(t))}{\rho[y(t)]} \leq 0. \end{aligned}$$

Theorem 4.1 is thus applicable and we conclude that (4.1) cannot have any solution of type B_j .

Acknowledgment. This paper is part of a thesis submitted to the Graduate School of Michigan State University in partial fulfillment of the requirements for the degree of Doctor of Philosophy. The author wishes to thank Dr. Pui-Kei Wong for his guidance during the writing of the thesis.

REFERENCES

1. J. S. Bradley, *Oscillation theorems for a second order delay equation*, J. Differential Equations, **8** (1970), 397-403.
2. F. Burkowski, *Oscillation theorems for a second order nonlinear functional differential equation*, J. Math. Anal. Appl., **33** (1971), 258-262.
3. H. E. Gollwitzer, *On nonlinear oscillations for a second order delay equation*, J. Math. Anal. Appl., **26** (1969), 385-389.

4. I. T. Kiguradze, *Oscillation properties of solutions of certain ordinary differential equations*, Dokl. Akad. Nauk SSSR, **144** (1962), 33-36; translated in Soviet Math. Dokl., **3** (1962), 649-652.
5. G. Ladas, *Oscillation and asymptotic behavior of solutions of differential equations with retarded arguments*, J. Differential Equations, **10** (1971), 281-290.
6. W. Leighton and Z. Nehari, *On the oscillation of solutions of self-adjoint linear differential equations of the fourth order*, Trans. Amer. Math. Soc., **89** (1958), 325-377.
7. V. A. Staikos and A. G. Petsoulas, *Some oscillation criteria for second order nonlinear delay differential equations*, J. Math. Anal. Appl., **30** (1970), 695-701.
8. R. D. Terry and P. K. Wong, *Oscillatory properties of a fourth order delay differential equation*, Funkcialaj Ekvacioj, **15** (1972), 209-221.
9. J. S. W. Wong, *Second order oscillation with retarded argumenes*, Proceedings of Conference on Ordinary Differential Equations, June 14-23, (1971), edited by L. Weiss, Academic Press, New York, 1972, 581-596.
10. P. K. Wong, *On a class of nonlinear fourth order differential equations*, Ann. Mat. Pura Appl., **81** (1969), 331-346.
11. T. Yoshizawa, *Oscillatory property of solutions of second order differential equations*, Tohoku Math. J., **22** (170), 619-634.
12. ———, *Oscillatory property for second order differential equations*, Proceedings of Conference on Ordinary Differential Equations, June 14-23, (1971), edited by L. Weiss, Academic Press, New York, 1972, 315-327.

Received December 18, 1972 and in revised form July 23, 1973.

GEORGIA INSTITUTE OF TECHNOLOGY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RICHARD ARENS (Managing Editor)
University of California
Los Angeles, California 90024

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. A. BEAUMONT
University of Washington
Seattle, Washington 98105

D. GILBARG AND J. MILGRAM
Stanford University
Stanford, California 94305

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NAVAL WEAPONS CENTER

David R. Adams, <i>On the exceptional sets for spaces of potentials</i>	1
Philip Bacon, <i>Axioms for the Čech cohomology of paracompacta</i>	7
Selwyn Ross Caradus, <i>Perturbation theory for generalized Fredholm operators</i>	11
Kuang-Ho Chen, <i>Phragmén-Lindelöf type theorems for a system of nonhomogeneous equations</i>	17
Frederick Knowles Dashiell, Jr., <i>Isomorphism problems for the Baire classes</i>	29
M. G. Deshpande and V. K. Deshpande, <i>Rings whose proper homomorphic images are right subdirectly irreducible</i>	45
Mary Rodriguez Embry, <i>Self adjoint strictly cyclic operator algebras</i>	53
Paul Erdős, <i>On the distribution of numbers of the form $\sigma(n)/n$ and on some related questions</i>	59
Richard Joseph Fleming and James E. Jamison, <i>Hermitian and adjoint abelian operators on certain Banach spaces</i>	67
Stanley P. Gudder and L. Haskins, <i>The center of a poset</i>	85
Richard Howard Herman, <i>Automorphism groups of operator algebras</i>	91
Worthen N. Hunsacker and Somashekhar Amrith Nainpally, <i>Local compactness of families of continuous point-compact relations</i>	101
Donald Gordon James, <i>On the normal subgroups of integral orthogonal groups</i>	107
Eugene Carlyle Johnsen and Thomas Frederick Storer, <i>Combinatorial structures in loops. II. Commutative inverse property cyclic neofields of prime-power order</i>	115
Ka-Sing Lau, <i>Extreme operators on Choquet simplexes</i>	129
Philip A. Leonard and Kenneth S. Williams, <i>The septic character of 2, 3, 5 and 7</i>	143
Dennis McGavran and Jingyal Pak, <i>On the Nielsen number of a fiber map</i>	149
Stuart Edward Mills, <i>Normed Köthe spaces as intermediate spaces of L_1 and L_∞</i>	157
Philip Olin, <i>Free products and elementary equivalence</i>	175
Louis Jackson Ratliff, Jr., <i>Locally quasi-unmixed Noetherian rings and ideals of the principal class</i>	185
Seiya Sasao, <i>Homotopy types of spherical fibre spaces over spheres</i>	207
Helga Schirmer, <i>Fixed point sets of polyhedra</i>	221
Kevin James Sharpe, <i>Compatible topologies and continuous irreducible representations</i>	227
Frank Siwiec, <i>On defining a space by a weak base</i>	233
James McLean Sloss, <i>Global reflection for a class of simple closed curves</i>	247
M. V. Subba Rao, <i>On two congruences for primality</i>	261
Raymond D. Terry, <i>Oscillatory properties of a delay differential equation of even order</i>	269
Joseph Dinneen Ward, <i>Chebyshev centers in spaces of continuous functions</i>	283
Robert Breckenridge Warfield, Jr., <i>The uniqueness of elongations of Abelian groups</i>	289
V. M. Warfield, <i>Existence and adjoint theorems for linear stochastic differential equations</i>	305