



Invited Review

Osteoarthritis of the hip and knee in former male professional soccer players

Stefano Petrillo[†], Rocco Papalia[†], Nicola Maffulli^{‡,§,*}, Piero Volpi[¶], and Vincenzo Denaro[†]

[†]Department of Orthopaedic and Trauma Surgery, Campus Bio-Medico University, Via Alvaro del Portillo, 200, 00128 Trigoria, Rome, Italy, [‡]Department of Musculoskeletal Disorders, University of Salerno School of Medicine and Surgery, Via Giovanni Paolo II, 132 - 84084 Fisciano, Salerno, Italy, [§]Centre for Sports and Exercise Medicine, Barts and The London School of Medicine and Dentistry, Mile End Hospital, 275 Bancroft Road, London E1 4DG, England, and [¶]Knee Surgery and Sports Traumatology Unit, Humanitas Research Hospital, Via Manzoni 56, 20089 Rozzano, Milan, Italy

*Correspondence address. Department of Musculoskeletal Disorders, University of Salerno School of Medicine and Surgery, Via Giovanni Paolo II, 132 - 84084 Fisciano, Salerno, Italy. E-mail: n.maffulli@qmul.ac.uk

Editorial Decision 28 December 2017; Accepted 1 January 2018

Abstract

Background: Professional soccer (PS) players are at great risk of osteoarthritis (OA) of the knee and hip.

Sources of data: Following the PRISMA guidelines, the key words 'osteoarthritis' and 'soccer' or 'football' were matched with 'players' or 'former' or 'retired' and with 'hip' or 'knee' on December 24, 2017 in the following databases: PubMed, Cochrane, Google scholar, Embase and Ovid. Only comparative studies reporting the prevalence rate of OA of both hip and knee joint in former PS athletes (fPSa) and age and sex matched controls were considered.

Areas of agreement: In fPSa, the prevalence rate of OA of both hip and knee is significantly higher compared to age and sex matched controls.

Areas of controversy: The pathological pathways responsible for the development of OA of the hip and knee in PS athletes (PSa) are still not clearly understood.

Growing points: The prevalence rate of clinical OA of the hip was 8.6% in fPSa and 5.6% in controls (odd ratio (OR) = 1.5; 95% CI: 1.06–2.31). The radiographic rate of OA was 21.2% in fPSa and 9.8% in controls (OR = 2.4;

95% CI: 1.66–3.69). A total of 14.6 and 53.7% of fPSa presented clinical and radiographic signs of OA of the knee, respectively, vs 12.9% (OR = 1.16; 95% CI: 0.86–1.55) and 31.9% (OR = 2.47; 95% CI: 2.03–3.00) of controls. Sonographic evidence of OA of the knee was found in 52% of fPSa and 33% of controls (OR = 2.2; 95% CI: 1.24–3.89).

Areas timely for developing research: Preventive training programmes should be developed to reduce the number of fPSa presenting early OA.

Key words: osteoarthritis, hip, knee, soccer, football, athletes

Introduction

Osteoarthritis (OA) of the hip and knee is the fourth and eighth most prevalent pathology in female and male patients respectively, impacting negatively on work capacity and quality of life.¹ Following the increase of mean age and life expectancy of the general population, OA is expected to become the most frequent orthopaedic pathology in the elderly.^{2,3} Moreover, patients with OA are at high risk to suffer from other medical conditions,^{4–6} producing elevated costs for the National Health Systems.¹

Factors such as ageing, female sex, obesity, smoking habit, rheumatological or haematological diseases, lower limb malalignment, trauma and sports activities are associated with OA of the lower limb.^{4,7} Furthermore, in the last few decades, athletic activities, and in particular professional soccer (PS), have been recognized as a risk factor of OA of the lower limb.^{8–13} Indeed, PS players are at greater risk for knee OA respect to athletes involved in other sports.¹⁴

Worldwide, ~200 000 PS athletes (PSa) are active every year.^{15,16} PSa are exposed to a high risk to sustain lower limb joints trauma both during training and matches.^{11,17–19} The most common type of injuries are muscle lesions,^{20–23} followed by contusion and joints sprains involving the hip, knee and ankle.^{17,24,25} The incidence of injuries per 1000 training hours in PSa was estimated to be 10–35.5, significantly higher than other sports.²⁶ Furthermore, several studies have demonstrated that the joints exposed to weight bearing and rotational motions

such as the hip and knee are at high risk to develop early OA.^{18,27–30} PS activity seems to play a crucial role in both aetiology and epidemiology of OA of the hip and knee, because of the synergy of overusing physical activity and high frequency of joint trauma.

Previous systematic reviews have highlighted that fPSa exhibit a high prevalence rate of OA of the ankle,³¹ knee,^{14,31} spine and hip.³² However, none of these specifically evaluated the prevalence rate of OA of both hip and knee in male fPSa and age-sex matched controls.

This systematic review ascertained whether the prevalence of both hip and knee OA in male fPSa is higher compared to age and sex matched controls.

Materials and methods

The present systematic review was conducted in accordance with the PRISMA guidelines. The literature search strategy is summarized in Figure 1 (PRISMA flow diagram).

The literature search was performed on December 24, 2017 in the following databases: PubMed, Cochrane, Google scholar, Embase and Ovid. The key words matched were: ‘osteoarthritis’ and ‘soccer’ or ‘football’ with ‘players’ or ‘former’ or ‘retired’ and with ‘hip’ or ‘knee’. All articles in English, Italian, Spanish and Portuguese language published in peer review journals from 1980 to 2017 were considered. Inclusion and exclusion criteria are summarized in Table 1.

The abstract of each article was screened. Articles without an abstract were excluded. If the




Fig. 1 PRISMA flow diagram.

Table 1 Inclusion and exclusion criteria of the study

Inclusion criteria	
Database	Medline, Google Scholar, Embase, Ovid, Pubmed
Type of study	Observational with age and sex matched control group
Language	English
Year of publication	1980–2017
Number of patients enroled	≥20
Sex of patients enroled	male
Duration of professional career	≥5
Diagnosis of OA	Radiographic, sonographic, clinical examination, clinical questionnaires
Exclusion criteria	
Type of study	Literature reviews, letters to editor, observational without control group
No of patients enroled	<20
Sex of patients enroled	Female
Duration of professional career	<5 years

abstract did not allow to assess the compliance with the defined inclusion and exclusion criteria, the full-text article was retrieved. Moreover, to identify

other relevant articles eligible for the study, cross-reference search of the selected studies was performed. Two investigators (S.P. and R.P.) extracted

independently and in a blind fashion the following information from the included articles: (i) year of publication of the study; (ii) demographic details of the patients enrolled in the studies (number of fPSa and controls, sex, age); (iii) role in the soccer field of fPSa; (iv) body mass index (BMI) of both fPSa and controls; (v) duration of professional career of fPSa; (vi) age at the time of diagnosis of hip and/or knee OA; (vii) prevalence rate of hip and/or knee OA in both fPSa and controls; (viii) method of diagnosis of hip and/or knee OA; and (ix) method of classification of radiographic results.

Data presentation

The continuous variables evaluated (age; duration of PS career; BMI) were expressed as mean value \pm standard deviation (range). The prevalence rate of OA of both hip and knee joint in fPSa and controls was expressed as percentage (%). Odd ratio (OR) was reported as mean with 95% confidence interval.

Results

A total of 17 studies were identified as eligible for our systematic review. However, seven studies did not comply with our inclusion and exclusion criteria (six studies^{8,9,33–36} without a control group; one study³⁷ in German language). Therefore, 10 articles^{38–47} published from 1980 to 2017 were included.

All the included studies were observational, and included fPSa and an age and sex matched control group (Level III).

Demographic results

Overall, there were 3997 male individuals, of which 1406 (35.2%) were fPSa and 2591 (64.8%) were controls. The mean age was reported in 1406 (100%) fPSa and 2398 (92.5%) controls, and it was 54.7 ± 8.8 years (range: 44–70 years) and 56.3 ± 10.3 years (range: 43.7–70), respectively. The demographics

characteristics of the individuals enrolled in the studies are shown in Table 2.

Assessment of hip and knee OA

The prevalence rate of OA of the hip was estimated in five studies^{39,40,42,44,45} involving 2470 (61.8%) individuals [fPSa: 688 (27.8%); controls: 1782 (72.2%)]. The prevalence rate of OA of the knee was estimated in eight studies^{38–40,43,45–48} involving 3580 (89.5%) individuals [fPSa: 1267 (35.4%); controls: 2313 (64.6%)].

A total of eight studies^{38,39,41–45,47} involving 2161 (54.1%) individuals [fPSa: 935 (43.3%); controls: 1226 (56.7%)] estimated the prevalence rate of OA of the hip and/or knee with radiographs. Of these, one study³⁹ involving 303 individuals [fPSa: 121 (39.9%); controls: 182 (60.1%)] used both radiographs and clinical examination, while another study⁴⁰ involving 1636 (43.6%) individuals [fPSa: 371 (22.7%); controls: 1265 (77.3%)] used a self-reported clinical questionnaire. In one study⁴⁶ involving 200 individuals [fPSa: 100 (10.7%); controls: 100 (4.7%)] the prevalence rate of OA of the knee was estimated with knee sonography.

The OA of the hip and/or knee was graded with the Kellgren–Lawrence⁴⁹ classification in three studies,^{38,39,48} while the Croft,⁴² Ahlback⁴³ and Danielsson⁴⁴ classification was used in one study each. In another study,⁴⁵ the radiographic diagnosis of OA of the hip and knee was made analysing the presence of joint space narrowing, subchondral sclerosis and cysts. Another study⁵⁰ used both the Nottingham Line Drawing Atlas (NLDA)⁵⁰ and Kellgren–Lawrence classification to grade knee OA, while in only one study⁴⁶ the presence and extent of OA of the knee was assessed with sonography according to published guidelines.^{51–53}

Prevalence rate of hip and knee OA

The prevalence rate of OA of both hip and knee was higher in fPSa than controls (Table 3). The prevalence rate of clinical and radiographic OA of the hip and knee, and of sonographic OA of the

Table 2 Demographic details of the patients

	Reported in N (%)		Result	
	fPSa	Controls	fPSa	Controls
Patients enrolled	1406 (100%)	2591 (100%)	1406	2591
Mean age ^{38–42,45–47}	1406 (100%)	2398 (92.5%)	54.7 ± 8.8	56.3 ± 10.3
Role in the soccer field ^{38,41}	67 (8%)	n.a.		
Goalkeepers			11	n.a.
Defenders			21	n.a.
Midfielders			19	n.a.
Strikers			16	n.a.
Duration of PS career ^{38,39,41,42,45,46}	323 (38.6%)	n.a.	15.8 ± 4.6	n.a.
BMI ^{38,39,47}				
During PS career	148 (17.7%)	n.a.	22.7 ± 0.5	n.a.
During investigation	1130 (28.2%)	2118 (52.6%)	26 ± 0.7	26.7 ± 1.4

fPSa, former professional soccer athletes; PS, professional soccer; BMI, body mass index.

Table 3 Overall prevalence rate of clinical and radiographic OA of the hip and knee, and of sonographic OA of the knee in fPSa and controls

	Evaluated in N patients		Results		OR (95% CI)
	fPSa	Controls	fPSa	Controls	
Hip OA ^{39,40,42,44,45}					
Clinical ^{39,42}	492	1447	8.6%	5.6%	1.5 (1.06–2.31)
Radiographic ^{39,40,42,44,45}	317	513	21.2%	9.8%	2.4 (1.66–3.69)
Knee OA ^{38–40,43,45,46,48}					
Clinical ^{39,41}	492	1447	14.6%	12.9%	1.16 (0.86–1.55)
Radiographic ^{38–41,43,45,47}	796	948	53.7%	31.9%	2.47 (2.03–3.00)
Sonographic ⁴⁶	100	100	52%	33%	2.2 (1.24–3.89)

OA, osteoarthritis; fPSa, former professional soccer athletes; OR, odds ratio; 95% CI, confidence interval.

knee in both fPSa and controls is reported in Table 3, with associated OR and 95% CI.

Discussion

The present systematic review assessed whether the prevalence rate of OA of both hip and knee in male fPSa was higher than that of age and sex matched individuals. fPSa present a higher prevalence rate of OA of both hip and knee compared to controls at a mean age of 55 years. The prevalence rate of clinical OA of the hip in fPSa was 8.6% (vs 5.6% of controls), while it was 21.2% for radiographic

features of OA of the hip (vs 9.8% of controls). A total of 14.6 and 53.7% of fPSa had a clinical and radiographic diagnosis of OA of the knee respectively (vs 12.9 and 31.9% of controls). Moreover, the prevalence rate of sonographic OA of the knee was 52% in fPSa and 33% in controls. These results about prevalence rate of OA of the hip and knee are significantly higher than those reported in non-PSa.⁵⁴

The results of our study are similar to those reported in different systematic reviews on the topic. Lohkamp *et al.*³² demonstrated that the prevalence of hip OA and hip replacements is

significantly higher in former PS players compared to a control group. Similar findings were reported by Driban *et al.*¹⁴ and Kuijt *et al.*³¹ regarding OA of the knee.

The prevalence rate of OA of the hip in the general population ranges from 0.7 to 4.4%, and it increases to 3.2–4% in patients older than 60 years.⁵⁴ The prevalence rate of OA of the knee ranges from 6.1 to 21%, and it increases to 10.1% in individuals older than 70 years. Furthermore, in the United Kingdom, clinical OA of the knee in non-professional athletes aged 45 years was 12.5%, while radiographic OA of the knee in individuals older than 50 years was 18%.⁵⁴

We included a total of 10 comparative studies in the present systematic review. The instruments used to evaluate the prevalence rate of OA strongly influence the results. Radiographic diagnosis was associated with a higher prevalence rate of OA than clinical diagnosis. Plain radiography of the hip and knee were the most common diagnostic method used in the studies included in our systematic review. However, no agreement was found within these studies in the classification of radiographic results. The Kellgren–Lawrence⁴⁹ classification was used in three studies,^{38,39,48} while the Croft,⁴² Ahlback⁴³ and Danielsson⁴⁴ classification was used in one study each, while in another study⁴⁷ the OA of the knee was scored with both Kellgren–Lawrence⁴⁹ classification and NLDA.⁵⁰ These findings suggest a huge heterogeneity of outcomes measurement, producing a high variability of results. Moreover, radiographs had an intra- and inter-rater variability of 0.85–0.91 and 0.91–0.93, respectively, to diagnose OA.³² Furthermore, the diagnosis of OA through radiographs, especially regarding the knee, is influenced by the type of views used.³²

Professional athletic activity stressing lower limb joints is associated with a high prevalence rate of lower limb OA.^{18,55,56} Iosifidis *et al.*³⁹ reported a statistically significant higher prevalence rate of radiographic OA in elite athletes compared with non-sports controls (36.6% in athletes vs 23.9% in controls). However, they reported that former professional athletes without history of trauma have

similar prevalence rate of lower limb OA of an age-matched control group of non-professional athletes.³⁹ Tveit *et al.*⁴⁰ confirmed that OA of the hip and knee is common in former PSa, while OA of the knee in impact athletes was associated with previous knee injuries.

When considering soccer, several studies have explained a relationship between overuse, repetitive loading, trauma and the development of OA of the lower limb.⁵⁷ The risk of injury is higher during matches because of contacts and increased competitiveness,¹⁷ and PSa have an injury risk 1000 times greater compared with industrial workers.^{9,33}

OA of the hip develops earlier in relatively young PSa, while OA of the knee is strongly related with history of trauma and previous surgery.^{58–64} Modifications of the hip anatomy begin in young PSa because of overuse and high repetitive torsional loadings.⁶⁵ Moreover, an increased prevalence of femoro-acetabular impingement was found in PSa.⁶⁶ All these conditions determine abnormal hip biomechanics leading to cartilage degeneration and early onset of OA.^{10,67–69} On the other hand, PSa are at a high risk to sustain anterior cruciate ligament (ACL) or menisci lesions during their career.^{70,71} In these patients, surgery is often performed to restore knee function and allow resumption of sports participation.^{72–77} However, ACL reconstruction or meniscectomy was related to the development of early OA of the knee in 50 and 92% of the patients, respectively.^{2,70,71,78–82} Moreover, isolated ACL or meniscal lesions produce a 10-fold increased risk to develop OA compared to uninjured individuals.^{70,71,82} Furthermore, genu varus was associated with the development of OA of the knee in young soccer players.⁸³

The major strength of our systematic review is that it was conducted following PRISMA guidelines. Moreover, to assess whether the prevalence rate of OA of both hips and knees was higher in fPSa compared to non-PS individuals, only comparative studies with an age and sex matched control group were included. Another strength is that both literature search and analysis of results was performed by two independent investigators in a blind fashion. Furthermore, we have included only male fPSa, and

the amount of patients included in our study is large enough to guarantee a large enough sample for the scope of our study.

However, we are aware that the main limitation of our study is represented by its design. Another important limitation is that our results were not stratified according to history of trauma, previous surgery of the hip or knee, BMI, smoking habit, rheumatologic or hematologic diseases, which are well-known risk factors for the development of OA. Unfortunately, risk factors were rarely reported in the primary studies. Nevertheless, our results are similar to those reported in other studies about the topic, suggesting a higher prevalence of OA of both hip and knee in fPSa compared to controls and compared to the general population.

Conclusions

In fPSa, the prevalence rate of OA of both hip and knee is significantly higher compared to age and sex matched controls.

Based on the evidence available in the current peer reviewed published literature, OA of the hip is associated with overuse and repetitive loadings, resulting in morphological changes of hip anatomy which begin in youth age. Furthermore, earlier trauma, ligaments or meniscal injuries requiring surgery demonstrate evidence of a statistically significant association with OA of the knee.

The pathological pathways involved in the development of OA of the hip and knee in PSa are not clearly understood yet. These may be useful to develop effective prevention programmes to reduce the number of fPSa developing early OA.

Conflict of interest statement

The authors have no potential conflicts of interest.

References

1. Murray CJ, Lopez AD, Jamison DT. The global burden of disease in 1990: summary results, sensitivity analysis and future directions. *Bull World Health Organ* 1994; 72:495–509.
2. Maffulli N, Osti L. ACL stability, function, and arthritis: what have we been missing? *Orthopedics* 2013;36: 90–2.
3. Piccirilli E, Oliva F, Murè MA, et al. Viscosupplementation with intra-articular hyaluronic acid for hip disorders. A systematic review and meta-analysis. *Muscles Ligaments Tendons J* 2016;6:293–9.
4. Kadam UT, Croft PR. Clinical comorbidity in osteoarthritis: associations with physical function in older patients in family practice. *J Rheumatol* 2007;34: 1899–904.
5. Keller K, Engelhardt M. Strength and muscle mass loss with aging process. Age and strength loss. *Muscles Ligaments Tendons J* 2013;3:346–50.
6. Rosemann T, Laux G, Szecsenyi J. Osteoarthritis: quality of life, comorbidities, medication and health service utilization assessed in a large sample of primary care patients. *J Orthop Surg Res* 2007;2:12.
7. Kadam UT, Croft PR. Group NSGC. Clinical multimorbidity and physical function in older adults: a record and health status linkage study in general practice. *Fam Pract* 2007;24:412–9.
8. Drawer S, Fuller CW. Propensity for osteoarthritis and lower limb joint pain in retired professional soccer players. *Br J Sports Med* 2001;35:402–8.
9. Turner AP, Barlow JH, Heathcote-Elliott C. Long term health impact of playing professional football in the United Kingdom. *Br J Sports Med* 2000;34: 332–6.
10. Comba F, Yacuzzi C, Ali PJ, et al. Joint preservation after hip arthroscopy in patients with FAI. Prospective analysis with a minimum follow-up of seven years. *Muscles Ligaments Tendons J* 2016;6:317–23.
11. Pangrazio O, Forriol F. Epidemiology of soccer players traumatic injuries during the 2015 America Cup. *Muscles Ligaments Tendons J* 2016;6:124–30.
12. Maffulli N. Intensive training in young athletes. The orthopaedic surgeon's viewpoint. *Sports Med* 1990;9: 229–43.
13. Maffulli N, King JB. Effects of physical activity on some components of the skeletal system. *Sports Med* 1992; 13:393–407.
14. Driban JB, Hootman JM, Sitler MR, et al. Is participation in certain sports associated with knee osteoarthritis? A systematic review. *J Athl Train* 2017;52: 497–506.
15. Engström B, Johansson C, Törnkvist H. Soccer injuries among elite female players. *Am J Sports Med* 1991;19: 372–5.
16. Junge A, Dvorak J. Soccer injuries: a review on incidence and prevention. *Sports Med* 2004;34:929–38.

17. Volpi P, Pozzoni R, Galli M. The major traumas in youth football. *Knee Surg Sports Traumatol Arthrosc* 2003;11:399–402.
18. Maffulli N, Longo UG, Gougoulas N, et al. Sport injuries: a review of outcomes. *Br Med Bull* 2011;97:47–80.
19. Pangrazio O, Forriol F. Epidemiology of injuries sustained by players during the 16th Under-17 South American Soccer Championship. *Rev Esp Cir Ortop Traumatol* 2016;60:192–9.
20. Chan O, Del Buono A, Best TM, et al. Acute muscle strain injuries: a proposed new classification system. *Knee Surg Sports Traumatol Arthrosc* 2012;20:2356–62.
21. De Carli A, Volpi P, Pelosini I, et al. New therapeutic approaches for management of sport-induced muscle strains. *Adv Ther* 2009;26:1072–83.
22. Melegati G, Tornese D, Gevi M, et al. Reducing muscle injuries and reinjuries in one Italian professional male soccer team. *Muscles Ligaments Tendons J* 2013;3:324–30.
23. Volpi P, Melegati G, Tornese D, et al. Muscle strains in soccer: a five-year survey of an Italian major league team. *Knee Surg Sports Traumatol Arthrosc* 2004;12:482–5.
24. Osti L, Del Buono A, Maffulli N. Arthroscopic debridement of the ankle for mild to moderate osteoarthritis: a midterm follow-up study in former professional soccer players. *J Orthop Surg Res* 2016;11:37.
25. Rolf CG, Barclay C, Riyami M, et al. The importance of early arthroscopy in athletes with painful cartilage lesions of the ankle: a prospective study of 61 consecutive cases. *J Orthop Surg Res* 2006;1:4.
26. Keller CS, Noyes FR, Buncher CR. The medical aspects of soccer injury epidemiology. *Am J Sports Med* 1988;16:S105–12.
27. Volpi P. Why so many injuries in professional football players? *J Sports Med Phys Fitness* 2013;53:101–2.
28. Angioi M, Maffulli GD, McCormack M, et al. Early signs of osteoarthritis in professional ballet dancers: a preliminary study. *Clin J Sport Med* 2014;24:435–7.
29. Maffulli N, Longo UG, Spiezia F, et al. Aetiology and prevention of injuries in elite young athletes. *Med Sport Sci* 2011;56:187–200.
30. Oryan A, Alidadi S, Moshiri A, et al. Bone regenerative medicine: classic options, novel strategies, and future directions. *J Orthop Surg Res* 2014;9:18.
31. Kuijt MT, Inklaar H, Gouttebarger V, et al. Knee and ankle osteoarthritis in former elite soccer players: a systematic review of the recent literature. *J Sci Med Sport* 2012;15:480–7.
32. Lohkamp M, Kromer TO, Schmitt H. Osteoarthritis and joint replacements of the lower limb and spine in ex-professional soccer players: a systematic review. *Scand J Med Sci Sports* 2017;27:1038–49.
33. Krajnc Z, Vogrin M, Recnik G, et al. Increased risk of knee injuries and osteoarthritis in the non-dominant leg of former professional football players. *Wien Klin Wochenschr* 2010;122:40–3.
34. Kujala UM, Kettunen J, Paananen H, et al. Knee osteoarthritis in former runners, soccer players, weight lifters, and shooters. *Arthritis Rheum* 1995;38:539–46.
35. Oztürk A, Ozkan Y, Ozdemir RM, et al. Radiographic changes in the lumbar spine in former professional football players: a comparative and matched controlled study. *Eur Spine J* 2008;17:136–41.
36. Chantraine A. Knee joint in soccer players: osteoarthritis and axis deviation. *Med Sci Sports Exerc* 1985;17:434–9.
37. Brouwer PJ, Geesink RG, Prompers LA, et al. [Symptoms and lesions of knees and ankles in former 1956 professional soccer players]. *Ned Tijdschr Geneeskde* 1981;125:694–7.
38. Arliani GG, Astur DC, Yamada RK, et al. Early osteoarthritis and reduced quality of life after retirement in former professional soccer players. *Clinics (Sao Paulo)* 2014;69:589–94.
39. Iosifidis MI, Tsarouhas A, Fylaktou A. Lower limb clinical and radiographic osteoarthritis in former elite male athletes. *Knee Surg Sports Traumatol Arthrosc* 2015;23:2528–35.
40. Tveit M, Rosengren BE, Nilsson J, et al. Former male elite athletes have a higher prevalence of osteoarthritis and arthroplasty in the hip and knee than expected. *Am J Sports Med* 2012;40:527–33.
41. Elleuch MH, Guermazi M, Mezghanni M, et al. [Knee osteoarthritis in 50 former top-level soccer players: a comparative study]. *Ann Readapt Med Phys* 2008;51:169–73.
42. Shepard GJ, Banks AJ, Ryan WG. Ex-professional association footballers have an increased prevalence of osteoarthritis of the hip compared with age matched controls despite not having sustained notable hip injuries. *Br J Sports Med* 2003;37:80–1.
43. Roos H, Lindberg H, Gärdsell P, et al. The prevalence of gonarthrosis and its relation to meniscectomy in former soccer players. *Am J Sports Med* 1994;22:219–22.
44. Lindberg H, Roos H, Gärdsell P. Prevalence of coxarthrosis in former soccer players. 286 players compared with matched controls. *Acta Orthop Scand* 1993;64:165–7.

45. Klünder KB, Rud B, Hansen J. Osteoarthritis of the hip and knee joint in retired football players. *Acta Orthop Scand* 1980;51:925–7.
46. Paxinos O, Karavasilis A, Delimpasis G, et al. Prevalence of knee osteoarthritis in 100 athletically active veteran soccer players compared with a matched group of 100 military personnel. *Am J Sports Med* 2016;44:1447–54.
47. Fernandes GS, Parekh SM, Moses J, et al. Prevalence of knee pain, radiographic osteoarthritis and arthroplasty in retired professional footballers compared with men in the general population: a cross-sectional study. *Br J Sports Med* 2017;0:1–7.
48. Elleuch MH, Guermazi M, Mezghanni M, et al. Knee osteoarthritis in 50 former top-level soccer players: a comparative study. *Ann Readapt Med Phys* 2008;51:174–8.
49. Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthritis. *Ann Rheum Dis* 1957;16:494–502.
50. Nagaosa Y, Mateus M, Hassan B, et al. Development of a logically devised line drawing atlas for grading of knee osteoarthritis. *Ann Rheum Dis* 2000;59:587–95.
51. Iagnocco A, Meenagh G, Riente L, et al. Ultrasound imaging for the rheumatologist XXIX. Sonographic assessment of the knee in patients with osteoarthritis. *Clin Exp Rheumatol* 2010;28:643–6.
52. Mermerci BB, Garip Y, Uysal RS, et al. Clinic and ultrasound findings related to pain in patients with knee osteoarthritis. *Clin Rheumatol* 2011;30:1055–62.
53. Möller I, Bong D, Naredo E, et al. Ultrasound in the study and monitoring of osteoarthritis. *Osteoarthritis Cartilage* 2008;16:S4–7.
54. Cooper C, Snow S, McAlindon TE, et al. Risk factors for the incidence and progression of radiographic knee osteoarthritis. *Arthritis Rheum* 2000;43:995–1000.
55. Chawla A, Twycross-Lewis R, Maffulli N. Microfracture produces inferior outcomes to other cartilage repair techniques in chondral injuries in the paediatric knee. *Br Med Bull* 2015;116:93–103.
56. Gougoulas N, Khanna A, Maffulli N. Sports activities after lower limb osteotomy. *Br Med Bull* 2009;91:111–21.
57. Volpi P, Taioli E. The health profile of professional soccer players: future opportunities for injury prevention. *J Strength Cond Res* 2012;26:3473–9.
58. Longo UG, Franceschetti E, Maffulli N, et al. Hip arthroscopy: state of the art. *Br Med Bull* 2010;96:131–57.
59. Papalia R, Del Buono A, Franceschi F, et al. Femoroacetabular impingement syndrome management: arthroscopy or open surgery? *Int Orthop* 2012;36:903–14.
60. Vendittoli PA, Young DA, Stitson DJ, et al. Acetabular rim lesions: arthroscopic assessment and clinical relevance. *Int Orthop* 2012;36:2235–41.
61. Osti L, Papalia R, Del Buono A, et al. Good results five years after surgical management of anterior cruciate ligament tears, and meniscal and cartilage injuries. *Knee Surg Sports Traumatol Arthrosc* 2010;18:1385–90.
62. Osti L, Papalia R, Del Buono A, et al. Surgery for ACL deficiency in patients over 50. *Knee Surg Sports Traumatol Arthrosc* 2011;19:412–7.
63. Ramjug S, Ghosh S, Walley G, et al. Isolated anterior cruciate ligament deficiency, knee scores and function. *Acta Orthop Belg* 2008;74:643–51.
64. Rathbone S, Maffulli N, Cartmell SH. Most british surgeons would consider using a tissue-engineered anterior cruciate ligament: a questionnaire study. *Stem Cells Int* 2012;2012:303724.
65. Agricola R, Bessems JH, Ginai AZ, et al. The development of Cam-type deformity in adolescent and young male soccer players. *Am J Sports Med* 2012;40:1099–106.
66. Agricola R, Heijboer MP, Ginai AZ, et al. A cam deformity is gradually acquired during skeletal maturation in adolescent and young male soccer players: a prospective study with minimum 2-year follow-up. *Am J Sports Med* 2014;42:798–806.
67. Mardones R, Via AG, Rivera A, et al. Arthroscopic treatment of femoroacetabular impingement in patients older than 60 years. *Muscles Ligaments Tendons J* 2016;6:397–401.
68. Fukushima K, Uchiyama K, Takahira N, et al. Prevalence of radiographic findings of femoroacetabular impingement in the Japanese population. *J Orthop Surg Res* 2014;9:25.
69. Fukushima K, Takahira N, Imai S, et al. Prevalence of radiological findings related to femoroacetabular impingement in professional baseball players in Japan. *J Orthop Sci* 2016;21:821–5.
70. Smith MV, Nepple JJ, Wright RW, et al. Knee osteoarthritis is associated with previous meniscus and anterior cruciate ligament surgery among Elite College American Football Athletes. *Sports Health* 2017;9:247–51.
71. van Meer BL, Oei EH, Meuffels DE, et al. Degenerative changes in the knee 2 years after anterior cruciate ligament rupture and related risk factors: a prospective observational follow-up study. *Am J Sports Med* 2016;44:1524–33.
72. Andia I, Maffulli N. Biological therapies in regenerative sports medicine. *Sports Med* 2017;47:807–28.

73. Capuano L, Hardy P, Longo UG, et al. No difference in clinical results between femoral transfixation and bio-interference screw fixation in hamstring tendon ACL reconstruction. A preliminary study. *Knee* 2008;15: 174–9.
74. Johnson DH, Maffulli N, King JB, et al. Anterior cruciate ligament reconstruction: a cynical view from the British Isles on the indications for surgery. *Arthroscopy* 2003;19:203–9.
75. Maffulli N, King JB. Anterior cruciate ligament injury. *Br J Sports Med* 1998;32:266.
76. Tudisco C, Bisicchia S, Cosentino A, et al. Knee stability, athletic performance and sport-specific tasks in non-professional soccer players after ACL reconstruction: comparing trans-tibial and antero-medial portal techniques. *Muscles Ligaments Tendons J* 2015;5:175–80.
77. Malliaropoulos N, Kakoura L, Tsitas K, et al. Active knee range of motion assessment in elite track and field athletes: normative values. *Muscles Ligaments Tendons J* 2015;5:203–7.
78. Forriol F, Longo UG, Hernández-Vaquero D, et al. The effects of previous meniscus and anterior cruciate ligament injuries in patients with total knee arthroplasty. *Ortop Traumatol Rehabil* 2010;12:50–7.
79. Maffulli N, Binfield PM, King JB. Articular cartilage lesions in the symptomatic anterior cruciate ligament-deficient knee. *Arthroscopy* 2003;19:685–90.
80. Maffulli N. A European ACL register. *Knee Surg Sports Traumatol Arthrosc* 2007;15:685.
81. Maffulli N, Longo UG, Gougoulas N, et al. Long-term health outcomes of youth sports injuries. *Br J Sports Med* 2010;44:21–5.
82. van der Hart CP, van den Bekerom MP, Patt TW. The occurrence of osteoarthritis at a minimum of ten years after reconstruction of the anterior cruciate ligament. *J Orthop Surg Res* 2008;3:24.
83. Krajnc Z, Rupreht M, Drobnič M. Quantitative evaluation of growth plates around the knees of adolescent soccer players by diffusion-weighted magnetic resonance imaging. *Biomed Res Int* 2015;2015:482017.