

# Outcome of COVID-19 patients with haematological malignancies after the introduction of vaccination and monoclonal antibodies: results from the HM-COV 2.0 study

**Alessandra Oliva** (✉ [alessandra.oliva@uniroma1.it](mailto:alessandra.oliva@uniroma1.it))

Department of Public Health and Infectious Diseases, Sapienza University of Rome

**Francesco Cogliati Dezza**

Department of Public Health and Infectious Diseases, Sapienza University of Rome

**Flavia Petrucci**

Department of Public Health and Infectious Diseases, Sapienza University of Rome

**Francesco Eugenio Romani**

Department of Public Health and Infectious Diseases, Sapienza University of Rome

**Matteo Morviducci**

Department of Public Health and Infectious Diseases, Sapienza University of Rome

**Flavio Marco Mirabelli**

Department of Public Health and Infectious Diseases, Sapienza University of Rome

**Francesca Cancelli**

Department of Public Health and Infectious Diseases, Sapienza University of Rome

**Emanuele Valeriani**

Department of Public Health and Infectious Diseases, Sapienza University of Rome

**Giulia Marcelli**

Unit of Emergency Radiology, Department of Radiological, Oncological and Pathological Sciences, Sapienza University of Rome

**Francesco Pugliese**

Department of Anesthesia and Critical Care Medicine, Sapienza University of Rome, Policlinico Umberto I

**Ombretta Turriziani**

Microbiology and Virology Laboratory, Department of Molecular Medicine, Sapienza University of Rome

**Paolo Ricci**

Unit of Emergency Radiology, Department of Radiological, Oncological and Pathological Sciences, Sapienza University of Rome

**Mario Venditti**

Department of Public Health and Infectious Diseases, Sapienza University of Rome

**Paolo Palange**

Department of Public Health and Infectious Diseases, Sapienza University of Rome

**Claudio Maria Mastroianni**

Department of Public Health and Infectious Diseases, Sapienza University of Rome

---

## Research Article

**Keywords:** COVID-19, haematological malignancies, outcome, vaccines, monoclonal antibodies

**Posted Date:** January 24th, 2023

**DOI:** <https://doi.org/10.21203/rs.3.rs-2492556/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Purpose.** Patients with hematological malignancies (HM) and SARS-CoV-2 infection present a higher risk of severe COVID-19 and mortality. The aim of the study was to investigate whether vaccination and monoclonal antibodies (mAbs) have modified the outcomes of HM patients with COVID-19.

**Methods.** Single-center retrospective study in HM patients hospitalized due to SARS-CoV-2 infection from March 2020 to April 2022. Patients were divided into PRE-V-mAb group (patients hospitalized before the introduction of vaccination and mAbs) and POST-V-mAb group (patients hospitalized after the use of vaccine and mAbs).

**Results.** A total of 126 patients were included (65 PRE-V-mAb and 61 POST-V-mAb). POST-V-mAb patients showed a significantly lower risk of Intensive Care Unit (ICU) admission (8.2% vs 27.7%,  $p=0.005$ ), shorter viral shedding [17 (IQR 10-28) vs 24 days (IQR 15-50),  $p=0.011$ ] and shorter hospitalization length [13 (IQR 7-23) vs 20 (IQR 14-41) days,  $p=0.0003$ ] compared to the PRE-V-mAb group. Nevertheless, both in-hospital and 30-day mortality rates did not significantly differ between the two groups (29.5% POST-V-mAb vs 36.9% PRE-V-mAb and 21.3% POST-V-mAb vs 29.2% PRE-V-mAb, respectively). At the multivariable analysis an active malignancy ( $p=0.042$ ), a critical COVID-19 at admission ( $p=0.025$ ) and the need for high-level of oxygen support at respiratory worsening [either HFNC/CPAP ( $p=0.022$ ) or Mechanical Ventilation ( $p=0.011$ )] were independently associated with in-hospital mortality. In the subgroup of POST-V-mAb patients, receiving therapy with mAbs was a protective factor ( $p=0.033$ ).

**Conclusion.** Despite the new therapeutic and preventive strategies available, HM patients with COVID-19 disease represent an extremely vulnerable group with still high mortality rates.

## Introduction

SARS-CoV-2 is the etiological agent of COVID-19 and since December 2019 it has rapidly spread around the world causing the most significant global pandemic in the last century [1]. Nowadays, the international knowledge about COVID-19 continues to evolve, and recently, data suggest that severe and critical disease can occur in up to 15% and 5% of patients, respectively [2].

However, immunosuppression could lead to more severe disease and mortality [3], with patients suffering from haematological malignancies (HM) presenting the highest risk of adverse outcomes compared to the general population [4–7].

Indeed, HM patients have an increased chance to be admitted to Intensive Care Units (ICUs), a prolonged viral shedding with longer hospitalization time, and a higher mortality rates than subjects without HM [4, 6, 8–12], especially in the presence of acute myeloid leukemia and myelodysplastic syndromes [4, 13].

The vulnerability of this population is explained by the severe immunosuppression resulting from both chemo-immunotherapy and the underlying disease [3]; in addition, the long-lasting persistence of the

virus may contribute to worse outcome and selection of new variants.

Since the first pandemic wave, the scenario has radically changed due to the widespread of SARS-CoV-2 vaccination and the availability of early therapies including antiviral drugs and monoclonal antibodies (mAbs) [14]. However, data has shown a low rate of seroconversion in patients with HM fully vaccinated with two doses of anti-SARS-CoV-2 mRNA vaccine, especially if previously treated with anti-CD20 antibodies [15]. Moreover, a third vaccine does not induce seroconversion in patients who have not responded before even if it cannot be ruled out the presence of possible protective cellular T-cell response [16].

Monoclonal antibodies (mAb) such as sotrovimab, casirivimab–imdevimab and bamlanivimab-etesevimab or the recently introduced long acting one have provided valuable options for the treatment of COVID-19 disease; this may be especially true for patients with immunocompromised conditions [17, 18], who may also benefit from convalescent plasma [19, 20].

However, whether these new available strategies have changed the outcomes in HM patients with SARS-CoV-2 is still under investigation.

Based on these considerations, we aimed to evaluate the clinical characteristics and outcomes (in-hospital mortality, ICU admission and duration of viral shedding) in hospitalized patients with HM and COVID-19 after the introduction of vaccination and mAb. Furthermore, we evaluated the risk factors for prolonged viral shedding in this population.

## Materials And Methods

From March 2020 to April 2022, a retrospective, single-center study was performed in a cohort of patients with HM and COVID-19 hospitalized at an Academic Hospital in Rome. Inclusion criteria were: i) diagnosis of COVID-19 by means of molecular/antigen tests, ii) hospitalization, and iii) age > 18 years. Patients with haematological diseases other than malignancy and age < 18 years were excluded from the study. The cohort of patients was further divided in two groups: the PRE-V-mAb group (patients hospitalized before the introduction of vaccination, mAbs and oral antiviral) and the POST-V-mAb group (patients hospitalized after the use of vaccine, mAbs and oral antiviral).

At our Institution, vaccination to patients with HM started in January 2021 whereas mAb and oral antivirals were available from April 2021 and January 2022, respectively. Therefore, the PRE-V-mAb group included patients enrolled from March 2020 to February 2021 and the POST-V-mAb group from May 2021 to April 2022 (Fig. 1).

Nasopharyngeal swab samples were collected and SARS-CoV-2 RNA was detected using real-time RT-PCR assay (RealStar SARS-CoV2 RT-PCR, Altona Diagnostics). For each patient, laboratory and clinical data at hospital admission and during hospitalization were collected and recorded anonymously in an electronic database. Therapeutic regimens including mAbs, antiviral therapy and the use of steroids were based on

the international and local guidelines available at that time [14, 21] and on clinical judgement, also following consultation with haematologists, when appropriate. Apart from guidelines indication, the use of monoclonal antibodies including casirivimab/imdevimab, bamlanivimab/etesevimab or sotrovimab was also based on hospital pharmacy availability and dominant circulating variants of concern at the time. In this regard, the first Italian Omicron case was detected on 27 November 2021 and the variant became dominant in Italy (99% of cases) at the end of January 2022. So, at the time patients in the POST-V-mAb group were infected, the most widely circulating variants in Italy were firstly Delta and, for a lesser extent, Omicron. [22]

## Definitions

Severity of infection was defined according to the WHO classification available at the time of enrolment [23]. Use of corticosteroids within the previous 30 days included therapy with prednisone or its equivalent at a dose > 0.5 mg/kg/day for at least 1 month. Prior infection and antibiotic therapy were defined as a diagnosis of infection and/or the receipt of antibiotics in the 30 days prior to hospital admission, respectively. Status of haematological malignancy was defined as new diagnosis, remission, refractory/relapsing disease or yet to define, according to the guidelines of European Society for Medical Oncology [24]. Active malignancy was defined as patients with new diagnosis or refractory/relapsing disease [25]. Prior active treatment included the receipt of chemotherapy or immunotherapy, or both, in the previous 90 days. Immunotherapy included the receipt of monoclonal antibodies (rituximab, daratumumab, obinutuzumab) and tyrosine kinase inhibitors (imatinib, ibrutinib, ruxolitinib, venetoclax). Worsening of respiratory conditions was based on the change of PaO<sub>2</sub>/FiO<sub>2</sub> and was defined as: i) need of supplementary oxygen therapy or ii) need of increasing oxygen therapy supplementation in a patient with SARS-CoV2 infection for reasons directly related to the infection, as it was already reported [8].

Time of viral shedding was defined as the number of days from the first viral detection by RT-PCR on nasopharyngeal specimen until the first negative result.

The study was approved by the local Ethics Committee (ID Prot. 109/2020).

## Radiology

Two multidetector CT scanners (Siemens Somatom go.Now 32 and Somatom Sensation 64; Siemens Healthineers) were used. COVID-19 pneumonia was confirmed by means of the following CT parameters: ground glass opacity, crazy-paving pattern and pulmonary consolidation. A semi-quantitative CT severity score to identify the percentage of lung parenchyma involved by the infective process was therefore calculated and then three main groups were obtained: minimum involvement (0–30%), medium involvement (31–60%) and high involvement (> 60%) [25].

## Statistical analyses

Continuous data were expressed as median (Interquartile Range, IQR) and categorical data were summarized as number of observations (n) and percentages (%). Univariable analysis was used to

identify risk factors and predictors for all cause in-hospital mortality. Baseline predictors possibly associated with the outcome at univariable comparison and variables considered clinically significant were considered for multivariable Cox regression analysis. Survival was analysed by Kaplan–Meier curves and statistical significance of the differences between the groups was assessed using the log-rank test. A logistic multivariable model was constructed to evaluate the predictors of prolonged viral shedding (PVS), which has been considered > 20 days according to the median value of viral shedding in our study population (20.5 days). All statistical analyses were performed with STATA/IC software (StataCorp) version 17.

## Results

### *General characteristics*

Overall, 126 patients were included in the study, of which 65 PRE-V-mAb group and 61 POST-V-mAb group, respectively. General characteristics of the population are shown in Table 1. Gender distribution as well as patients' comorbidities, symptoms, and disease severity at the admission did not differ significantly between the two groups, while patients in the POST-V-mAb were older [71 (58-80) vs 64 (50-74) years,  $p=0.012$ ]. Notably, a similar percentage of patients in both groups had pneumonia at hospital admission (84.6% vs 78.6%,  $p=0.389$ ) and developed a respiratory worsening during hospitalization (58.5 vs 44.3%,  $p=0.111$ ).

Non-Hodgkin Lymphoma was the most frequent underlying HM in both groups (40% vs 44.3%,  $p=0.628$ ) followed by Chronic Lymphocytic Leukemia (10.8% vs 14.7%,  $p=0.598$ ) and Multiple Myeloma (15.4% vs 6.6%,  $p=0.115$ ). No difference was observed between the two groups when considering active malignancy rates (41.5% vs 50.8% in PRE- and POST-V-mAb groups, respectively,  $p=0.223$ ), in detail, patients in the PRE-V-mAb group were less likely to have a relapsing or refractory disease (18.5% vs 34.4%,  $p=0.042$ ) than patients in POST-V-mAb group and more likely to have a complete or partial remission of underlying HM (53.8% vs 34.4%,  $p=0.028$ ). A similar proportion of patients in both groups received prior active treatment (55.4% vs 52.5%,  $p=0.742$ ) (Supplementary Table 1).

The use of corticosteroids (91.0% vs 63.9%,  $p=0.001$ ) and antibiotics (83.0% vs 70.5%,  $p=0.04$ ) were significantly higher in the PRE-V-mAb group, while no difference was observed in treatment with remdesivir (49.2% vs 50.8%,  $p=0.861$ ), enoxaparin (75.4% vs 57.3%,  $p=0.283$ ), tocilizumab (9.2% vs 4.9%,  $p=0.348$ ) and baricitinib (1.6% vs 1.6%,  $p=0.924$ ) (Table 1).

In the POST-V-mAb group, 37 (63.8%) patients received mAb, 31 (50.8%) remdesivir, 23 (62.2%) received remdesivir plus mAbs. The majority of patients (55, 90.5%) had been previously vaccinated against SARS-CoV2, 22 of whom (40.0%) have received the last vaccine dose more than 120 days before hospital admission. The median number of vaccine doses was 3 (2-3).

### *Clinical outcomes and risk of in-hospital mortality*

Patients in the PRE-V-mAb group showed a significantly higher risk of ICU admission (27.7% vs 8.2%,  $p=0.005$ ) and longer hospital length (20 vs 13 days,  $p=0.0003$ ) than patients in POST-V-mAb group. Conversely, both in-hospital and 30-day mortality (36.9% vs 29.5% and 29.2% vs 21.3% in PRE- and POST-V-mAb groups,  $p=0.378$ ,  $p=0.307$ , respectively) rates did not significantly differ between the two groups, even though they tend to be lower in the POST-V-mAb group.

At multivariable analysis, an active malignancy (Hazard ratio - HR 2.20, 95% confidence interval - CI - 1.02-4.72,  $p=0.042$ ), a critical COVID-19 at admission (HR 2.99, 95% CI, 1.15-7.80,  $p=0.025$ ) and the need for high level of oxygen support at respiratory worsening [either high flow nasal cannula/Continuous Positive Airway Pressure, HFNC/CPAP, (HR 5.96, 95% CI, 1.29-27.52,  $p=0.022$ ) or Mechanical Ventilation (HR 7.63, 95% CI, 1.59-36.62,  $p=0.011$ )] were independently associated with an increased risk of in-hospital mortality in the overall population (Table 2).

When just the POST-V-mAb group was included in the model, mAbs administration (HR 0.03, 95% CI, 0.002-0.76,  $p=0.033$ ) was independently associated with a reduced risk of in-hospital mortality whereas the severity of infection at admission (HR 5.75, 95% CI 1.01-32.74,  $p=0.048$ ) confirmed its association with a worse outcome (Supplementary Table 2).

### *Duration and risk of prolonged viral shedding (PVS)*

Overall, median duration of viral shedding was 20.5 days (IQR, 18.7-22.2) and was higher in PRE-V-mAb than in POST-V-mAb group [24 (IQR, 15-50) vs 17 (IQR, 10-28) days,  $p=0.011$ ]. Baseline characteristics of patients stratified according to viral shedding (i.e., < 20 vs > 20 days) are reported in Table 3.

According to the logistic regression model shown in Table 4, active treatment for the HM in the previous 90 days (OR 3.55, 95% CI, 1.35-9.32,  $p=0.010$ ), respiratory worsening after more than 10 days (OR, 2.92, 95% CI, 1.05-8.05,  $p=0.038$ ), use of corticosteroids (OR, 7.59, 95% CI, 1.73-33.24,  $p=0.007$ ) and its duration for >7 days (OR, 4.92, 95% CI, 1.59-15.22,  $p=0.006$ ) were risk factors independently associated with a prolonged viral shedding.

## **Discussion**

In the present study, we observed that hospitalized patients with HM and COVID-19 continued to present high mortality rates despite the introduction of vaccination and mAbs. However, we could show a significant reduction in terms of ICU admission, hospitalization length and duration of viral shedding compared with the first pandemic waves. Notably, our results highlight how both the severity of COVID-19 at admission and at respiratory worsening and the presence of an active malignancy at the time of SARS-CoV2 diagnosis represent the major drivers of mortality.

In the literature, the mortality rate reported on hospitalized patients widely varied, ranging from 12–23%, but still significantly lower than the previous periods [6, 26, 27]. More recently, results from the international platform EPICoVIDHEA [28] reported mortality rates of 11.5% and 16% in a vaccinated

hospitalized population affected by HM in 2021 and patients hospitalized due to Omicron infection, respectively [26, 27]. On the other hand, preliminary data from a Danish study focusing only on Omicron infected Chronic Lymphocytic Leukemia patients and a Japanese case series observed a 23% and 22% 30-day mortality rate, respectively [29, 30].

In our study, the mortality rates were higher, although a slight, but not significant, reduction was observed after vaccination and mAb. However, given the rapid pandemic evolution and the succeeding of new variants, which may overlap during the same period of time and differ in circulation amongst countries, consistent and homogeneous data on outcomes in patients with HM and the effects of vaccination, mAbs and oral antivirals are still lacking. Furthermore, recent studies focused on both non-hospitalized and hospitalized patients or only on a particular type of HM or a specific variant, circumstances that may by definition influence the outcome. Consequently, it's very difficult to make a direct and crude comparison among different studies.

We found that the independent predictors of mortality were the severity of infection at admission and at respiratory worsening and the presence of an active malignancy, prompting that COVID-19 and HM might have a co-attributable role in the mortality of hospitalized patients with HM despite advance in SARS-CoV-2 prevention and treatment.

These findings indeed highlight how the COVID-19 still represent *itself* a major challenge in HM patients, since, although reduced after the implementation of vaccination and mAb, a not negligible rate of subjects experienced a respiratory worsening needing oxygen support exclusively as a consequence of infection's progression. At the same time, an active malignancy at the time of SARS-CoV2 diagnosis may concur to a worse outcome [31]. Indeed, in our cohort of patients more than half of subjects had an active malignancy. When looking in depth, there was a statistically significant higher percentage of patients with relapsing or refractory disease in the POST-V-mAb group, which may have contributed to the observed high mortality in this setting.

Furthermore, patients in the POST-V-mAb group were admitted at a time when first the Delta and only later the Omicron variants were predominant in Italy. Even though Omicron variant has a higher transmission rate but less severity than previous ones, the presence of Delta variant could also explain our higher fatality rate compared with last published studies strictly under conditions of Omicron dominance.

In the subgroup of patients in the POST-V-mAb, we could observe that receiving mAb was a protective factors for mortality, suggesting the advantage of early therapies in this high-risk group of patients and in line with a recent published paper by the EPICOVIDHEA group [32].

In line with the literature [18, 33–36], we found a reduction on ICU admission rates. These findings may be explained with the lower need of high-level oxygen support (ie, CPAP/HFNC and MV) observed in the POST-V-mAb group.



Another interesting finding was the analysis of the duration of viral shedding in this high-risk population. First, we confirmed that HM patients exhibit a prolonged shedding of viral RNA from the upper respiratory tract [8], which has a crucial impact on the decision whether to start or continue the treatment of the HM, especially in patients with active malignancy. Nevertheless, the observed reduction of viral shedding observed in the POST-V-mAb group has important consequences: on one hand, it increases the opportunity for the patients to receive the treatment of underlying disease; on the other hand, it may reduce the possibility of selecting viral variants, which may be less susceptible to patients' immunity and to drugs' activity [37, 38]. Second, we assessed the predictors of PVS and found that not only corticosteroid use (either for the treatment of the underlying disease or COVID-19) but also its duration for more than 7 days were independently associated with PVS [39, 40].

This study undoubtedly presents several limitations. Firstly, its retrospective and single centre nature; furthermore, the small study population did not allow us to stratify our patients by the different types of HM. Secondly, we were unable to identify the SARS-CoV-2 variant of concern in all infected patients; therefore, the variant was indirectly defined based on the dominant one in Italy in the same period. However, the aim of the study was not to investigate the effect of a specific variant on the outcome, which, in our opinion, deserves targeted studies. Finally, it's quite complicated to speculate whether the treatment of the underlying HM would have affected mortality in those who were not treated. In addition, there was a lack of data on the efficacy of oral antivirals and the long-acting mAbs, which need further targeted studies in the haematological population.

In conclusion, we showed that vaccination and mAbs reduce ICU admission, hospitalization length, and viral shedding in COVID-19 patients with HM, who, however, still remain a high-vulnerable population with high mortality rates. Strict adherence to non-pharmacological interventions and vaccinations remains mandatory for this frail population in the time where SARS-CoV-2 is continuously circulating in the community.

### **Financial support**

This research was supported by EU funding within the NextGeneration EU-MUR PNRR Extended Partnership initiative on Emerging Infectious Diseases (Project no. PE00000007, INF-ACT).

### **Ethical Approval**

**Statement:** The study was conducted in accordance with the Declaration

of Helsinki and approved by the Ethics Committee of Policlinico Umberto I (ID Prot. 109/2020).

## **Declarations**

**Financial support:** This research was supported by EU funding within the NextGeneration EU-MUR PNRR Extended Partnership initiative on Emerging Infectious Diseases (Project no. PE00000007, INF-ACT).

**Ethical Approval Statement:** The study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of Policlinico Umberto I (ID Prot. 109/2020).

**Potential conflict of interest:** The authors declare no potential conflict of interest. Oliva A. No conflict, Cogliati Dezza F. No conflict, Petrucci F. No conflict, Romani FE. No conflict, Morviducci M. No conflict, Mirabelli FM. No conflict, Cancelli F. No conflict, Valeriani E. No conflict, Marcelli G. No conflict, Pugliese F. No conflict, Ricci P. No conflict, Venditti M. No conflict, Palange P. No conflict, Mastroianni CM. No conflict.

### **Author contribution**

Conceptualization of the study: AO, FCD, data collection: FP, FER, MM, MFM, EV, FC, radiology analyses: GM, PR, virological analyses: OT, manuscript draft writing: AO, FCD, FP, supervision: FP, MV, PP, CMM.

**Acknowledgements:** Authors thank all the physicians devoted to the cure of patients with haematological malignancies and COVID-19 belonging to the Emergency Department and Infectious Diseases, Pneumology and Intensive Care Units of Azienda Policlinico Umberto I, Sapienza University of Rome. Authors thank all the Hematologists who contributed to the cure and assistance of these patients. Authors also thank all the nurse staff for their assistance to COVID-19 patients.

## **References**

1. World Health Organization. Coronavirus disease (COVID-19) – World Health Organization [Internet]. 2022 [cited 2022 Oct 21]. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>
2. Long B, Carius BM, Chavez S, et al. Clinical update on COVID-19 for the emergency clinician: Presentation and evaluation. *Am J Emerg Med.* 2022,54:46–57.
3. El-Sharkawi D, Iyengar S. Haematological cancers and the risk of severe COVID-19: Exploration and critical evaluation of the evidence to date. *Br J Haematol.* 2020,190:336–45.
4. Pagano L, Salmanton-García J, Marchesi F, et al. COVID-19 infection in adult patients with hematological malignancies: a European Hematology Association Survey (EPICOVIDEHA). *J Hematol Oncol.* 2021,14:168.
5. Goldman JD, Robinson PC, Uldrick TS, Ljungman P. COVID-19 in immunocompromised populations: implications for prognosis and repurposing of immunotherapies. *J Immunother Cancer. BMJ Specialist Journals,* 2021,9:e002630.
6. Vijenthira A, Gong IY, Fox TA, et al. Outcomes of patients with hematologic malignancies and COVID-19: a systematic review and meta-analysis of 3377 patients. *Blood.* 2020,136:2881–92.
7. Cogliati Dezza F, Oliva A, Mauro V, et al. Real-life use of remdesivir-containing regimens in COVID-19: a retrospective case-control study. *Infez Med.* 2022,30:211–22.

8. Oliva A, Curtolo A, Volpicelli L, et al. Clinical course of Coronavirus Disease-19 in patients with haematological malignancies is characterized by a longer time to respiratory deterioration compared to non-haematological ones: results from a case-control study. *Infection*. 2022,
9. Passamonti F, Cattaneo C, Arcaini L, et al. Clinical characteristics and risk factors associated with COVID-19 severity in patients with haematological malignancies in Italy: a retrospective, multicentre, cohort study. *Lancet Haematol*. 2020,7:e737–45.
10. He W, Chen L, Chen L, et al. COVID-19 in persons with haematological cancers. *Leukemia*. 2020,34:1637–45.
11. Sanchez-Pina JM, Rodríguez Rodríguez M, Castro Quismondo N, et al. Clinical course and risk factors for mortality from COVID-19 in patients with haematological malignancies. *European Journal of Haematology*. 2020,105:597–607.
12. Carrara E, Razzaboni E, Azzini AM, et al. Predictors of clinical evolution of SARS-CoV-2 infection in hematological patients: A systematic review and meta-analysis. *Hematological Oncology [Internet]*. 2022 [cited 2022 Oct 21],n/a. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/hon.3084>
13. García-Suárez J, de la Cruz J, Cedillo Á, et al. Impact of hematologic malignancy and type of cancer therapy on COVID-19 severity and mortality: lessons from a large population-based registry study. *J Hematol Oncol*. 2020,13:133.
14. Bartoletti M, Azap O, Barac A, et al. ESCMID COVID-19 living guidelines: drug treatment and clinical management. *Clin Microbiol Infect*. 2022,28:222–38.
15. Re D, Barrière J, Chamorey E, Delforge M, et al. Low rate of seroconversion after mRNA anti-SARS-CoV-2 vaccination in patients with hematological malignancies. *Leukemia & Lymphoma*. Taylor & Francis, 2021,62:3308–10.
16. Re D, Seitz-Polski B, Brglez V, et al. Humoral and cellular responses after a third dose of SARS-CoV-2 BNT162b2 vaccine in patients with lymphoid malignancies. *Nat Commun*. 2022,13:864.
17. Owen C, Robinson S, Christofides A, Sehn LH. A Canadian Perspective: Monoclonal Antibodies for Pre- and Post-Exposure Protection from COVID-19 in Vulnerable Patients with Hematological Malignancies. *Current Oncology*. Multidisciplinary Digital Publishing Institute, 2022,29:3940–9.
18. Boeckel GR, Hölscher SD, Bürger C, et al. Comprehensive Treatment of Hematological Patients with SARS-CoV-2 Infection Including Anti-SARS-CoV-2 Monoclonal Antibodies: A Single-Center Experience Case Series. *Curr Oncol*. 2022,29:2312–25.
19. Oliva A, Cancelli F, Brogi A, et al. Convalescent plasma for haematological patients with SARS-CoV-2 pneumonia and severe depletion of B-cell lymphocytes following anti-CD20 therapy: a single-centre experience and review of the literature. *New Microbiol*. 2022,45:62–72.
20. Thompson MA, Henderson JP, Shah PK, et al. Association of Convalescent Plasma Therapy With Survival in Patients With Hematologic Cancers and COVID-19. *JAMA Oncol*. 2021,7:1167–75.
21. Focà E, Ripamonti D, Rizzi M, et al. SEZIONE REGIONE LOMBARDIA. 2020,8.

22. EpiCentro. Monitoraggio delle varianti del virus SARS-CoV-2 di interesse in sanità pubblica in Italia [Internet]. 2022 [cited 2022 Oct 21]. Available from: <https://www.epicentro.iss.it/coronavirus/sars-cov-2-monitoraggio-varianti-indagini-rapide>
23. Agarwal A, Rochwerg B, Lamontagne F, et al. A living WHO guideline on drugs for covid-19. *BMJ. British Medical Journal Publishing Group*, 2020,370:m3379.
24. ESMO. ESMO Clinical Practice Guidelines: Haematological Malignancies [Internet]. 2022 [cited 2022 Oct 21]. Available from: <https://www.esmo.org/guidelines/guidelines-by-topic/haematological-malignancies>
25. Francone M, Iafrate F, Masci GM, et al. Chest CT score in COVID-19 patients: correlation with disease severity and short-term prognosis. *Eur Radiol*. 2020,30:6808–17.
26. Pagano L, Salmanton-García J, Marchesi F, et al. COVID-19 in vaccinated adult patients with hematological malignancies: preliminary results from EPICOVIDEHA. *Blood*. 2022,139:1588–92.
27. Blennow O, Salmanton-García J, Nowak P, et al. Outcome of infection with omicron SARS-CoV-2 variant in patients with hematological malignancies: An EPICOVIDEHA survey report. *Am J Hematol*. 2022,97:E312–7.
28. Salmanton-García J, Busca A, Cornely OA, et al. EPICOVIDEHA: A Ready to Use Platform for Epidemiological Studies in Hematological Patients With COVID-19. *Hemasphere*. 2021,5:e612.
29. Niemann CU, da Cunha-Bang C, Helleberg M, Ostrowski SR, Brieghel C. Patients with CLL have a lower risk of death from COVID-19 in the Omicron era. *Blood*. 2022,140:445–50.
30. Taenaka R, Obara T, Kohno K, Aoki K, Ogawa R. Infections with the SARS-CoV-2 Omicron variant show a similar outcome as infections with the previous variants in patients with hematologic malignancies. *Ann Hematol*. 2022,101:1877–8.
31. Lee M, Quinn R, Pradhan K, et al. Impact of COVID-19 on case fatality rate of patients with cancer during the Omicron wave. *Cancer Cell*. 2022,40:343–5.
32. Pagano L, Salmanton-García J, Marchesi F, et al. Breakthrough COVID-19 in vaccinated patients with hematologic malignancies: results from EPICOVIDEHA survey. *Blood* [Internet]. 2022 [cited 2022 Oct 4], Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9492383/>
33. Salvini M, Damonte C, Mortara L, et al. Immunogenicity and clinical efficacy of anti-SARS-CoV-2 vaccination in patients with hematological malignancies: Results of a prospective cohort study of 365 patients. *American Journal of Hematology*. 2022,97:E321–4.
34. Marasco V, Piciocchi A, Candoni A, et al. Neutralizing monoclonal antibodies in haematological patients paucisymptomatic for COVID-19: The GIMEMA EMATO-0321 study. *Br J Haematol*. 2022,
35. Bronstein Y, Gat R, Levi S, et al. COVID-19 in patients with lymphoproliferative diseases during the Omicron variant surge. *Cancer Cell*. 2022,40:578–80.
36. Bradwell S, Hone L, Thorneycroft K, et al. 2022 update on the clinical outcome of coronavirus disease 2019 in haemato-oncology patients. *Leuk Res*. 2022,119:106908.

37. Siqueira JD, Goes LR, Alves BM, et al. SARS-CoV-2 genomic analyses in cancer patients reveal elevated intrahost genetic diversity. *Virus Evol.* 2021,7:veab013.
38. Baang JH, Smith C, Mirabelli C, et al. Prolonged Severe Acute Respiratory Syndrome Coronavirus 2 Replication in an Immunocompromised Patient. *J Infect Dis.* 2020,jiaa666.
39. Li J, Liao X, Zhou Y, et al. Association between glucocorticoids treatment and viral clearance delay in patients with COVID-19: a systematic review and meta-analysis. *BMC Infectious Diseases.* 2021,21:1063.
40. Cogliati Dezza F, Oliva A, Cancelli F, et al. Determinants of prolonged viral RNA shedding in hospitalized patients with SARS-CoV-2 infection. *Diagn Microbiol Infect Dis.* 2021,100:115347.

## Tables

**Table 1.** General features of study population.

	Total (n=126)	PRE-V-mAb (N=65)	POST-V-mAb (N=61)	<i>p</i> value
<b>Demographics</b>				
Gender (Female), n (%)	65 (51.5)	32 (49.2)	33 (50.7)	0.597
Age, median (IQR)	66 (52-77)	64 (50-74)	71 (58-80)	<b>0.012</b>
Steroids in the last 30 days before admission, n (%)	43 (34.4)	25 (38.5)	18 (29.5)	0.320
Infection in the last 30 days before admission, n (%)	23 (18.3)	13 (20)	10 (16.4)	0.600
Antibiotic therapy in the last 30 days (excluded prophylaxis), n (%)	39 (31.0)	21 (32.3)	18 (29.5)	0.734
Antibiotic prophylaxis therapy, n. (%)	42 (33.3)	23 (35.4)	19 (31.1)	0.614
Days of hospitalization, median (IQR)	16 (9-38)	20 (14-44)	13 (7-23)	<b>0.0003</b>
Days from symptoms onset to hospitalization, median (IQR)	3 (1-7)	3 (1-7)	3 (1-7)	0.769
<b>Comorbidities, n (%)</b>				
Hypertension	52 (41.3)	22 (33.8)	30 (49.2)	0.081
Coronary artery disease	28 (22.2)	14 (21.5)	14 (22.9)	0.849
Diabetes	17 (13.5)	8 (12.3)	9 (14.8)	0.688
Chronic Obstructive Pulmonary Disease	8 (6.3)	4 (6.1)	4 (6.56)	0.926
Chronic Renal Failure	9 (7.1)	6 (9.2)	3 (4.9)	0.348
<b>Disease severity at the admission, n (%)</b>				
Moderate	80 (63.5)	41 (63.08)	39 (63.93)	0.920
Severe	31 (24.6)	18 (27.7)	13 (21.3)	0.406
Critical	8 (3.4)	3 (4.6)	5 (8.2)	0.410
Pneumonia	103 (81.8)	55 (84.6)	48 (78.7)	0.389
<b>Symptoms at the admission, n (%)</b>				
Dyspnea	51 (40.4)	26 (40)	25 (41)	0.910
Cough	51 (40.4)	24 (36.9)	27 (44.3)	0.402
Asthenia	45 (35.7)	28 (43)	17 (27.9)	0.075
Anosmia/ageusia	6 (4.8)	4 (6.1)	2 (3.3)	0.449

Fever	83 (65.9)	46 (70.8)	37 (60.7)	0.231
Conjunctivitis	6 (4.8)	2 (3)	4 (6.6)	0.365
Asymptomatic	10 (7.9)	4 (6.1)	6 (9.8)	0.445
<b>Respiratory features at the admission, median (IQR)</b>				
SpO <sub>2</sub>	97 (95-98)	97 (95-98)	97 (95-98)	0.658
PO <sub>2</sub> /FiO <sub>2</sub>	367 (290-448)	362 (290-443)	400 (290-457)	0.375
FiO <sub>2</sub>	21 (21-21)	21 (21-21)	21 (21-21)	0.125
<b>Laboratory findings at the admission, median (IQR)</b>				
Haemoglobin, g/dl	11.7 (9.7-13.6)	11.4 (9.4-13.6)	12 (10.2-13.55)	0.948
White Blood Cells, x 10 <sup>6</sup> /L	6380 (3790-10660)	5630 (3140-9390)	7330 (4110-12605)	0.175
Neutrophils, x 10 <sup>6</sup> /L	4130 (2210-6820)	3650 (1950-7040)	4375 (2445-6765)	0.384
Lymphocytes, x 10 <sup>6</sup> /L	900 (480-1630)	820 (410-1140)	1055 (505-2890)	<b>0.031</b>
Monocytes, x 10 <sup>6</sup> /L	310 (160-490)	260 (120-410)	375 (175-570)	0.05
Platelets, x 10 <sup>9</sup> /L	155 (105-207)	160 (113-202)	152 (103.5-215)	0.847
Thrombocytopenia (<150 x 10 <sup>9</sup> /L), n (%)	55 (43.6)	26 (40)	29 (48.3)	0.348
Neutropenia (<500 x 10 <sup>9</sup> /L), n (%)	12 (9.5)	9 (13.9)	3 (5)	0.093
Creatinin, mg/dl	0.9 (0.7-1.3)	0.9 (0.7-1.2)	0.95 (0.7-1.3)	0.653
Albumin, g/dl	36 (33-40)	36 (32-39)	37 (33-41)	0.323
D-dimer, µg/L	864 (478-1691.5)	831.5 (385-1551)	896 (623-1961)	0.163
C-Reactive Protein, mg/dL	4.5 (1.8-9.6)	3.46 (1.45-7.5)	4.8 (1.85-10.23)	0.209
Ferritin, µg/L	602.5 (210-1741)	935.5 (288-1743)	512 (140-1513)	0.186

LDH, U/L	296 (229-398)	290 (241-392)	311 (201-433)	0.971
Procalcitonin, ng/ml	0.24 (0.07-0.61)	0.24 (0.11-0.48)	0.24 (0.06-1.38)	0.999
<b>Features at the respiratory worsening*</b>				
Worsening during hospitalization, n (%)	65 (51.6)	38 (58.46)	27 (44.26)	0.111
FiO <sub>2</sub> median (IQR)	35 (21-60)	35 (21-50)	30.5 (21-100)	0.508
PO <sub>2</sub> /FiO <sub>2</sub> , median (IQR)	242 (146-350)	248 (163-374)	204 (89-324)	0.204
Days from SARS-CoV2 diagnosis to worsening, median (IQR)	8 (3-14)	7.5 (2-16)	6 (1-12)	0.412
Days from admission to worsening, median (IQR)	7 (2-17)	10 (3-17)	4 (1-12)	0.088
Late worsening (>10 days), n (%)	88 (69.8)	47 (72.3)	41 (67.2)	0.533
Type of worse ventilation, n (%)				<b>0.046</b>
No oxygen	34 (27.0)	12 (18.5)	22 (36.1)	
Oxygen support by Venturi Mask	46 (35.5)	24 (36.9)	22 (36.1)	
HFNC/CPAP	32 (25.4)	18 (27.7)	14 (22.9)	
Mechanical ventilation	14 (11.1)	11 (16.2)	3 (4.9)	
<b>Radiological findings at admission</b>				
Lung involvement, %, median (IQR)	18.8 (5-40)	15 (5-35)	20 (5-50)	0.216
Classes of lung involvement, n (%)		48 (73.85)	39 (63.93)	0.484
Low (0-30%)	87 (69.9)	11 (16.92)	14 (22.95)	
Medium (31-60%)	25 (19.0)	6 (9.23)	8 (13.11)	
High (>60%)	14 (11.1)			
<b>Therapy</b>				
Overall remdesivir therapy, n (%)	68 (53.9)	32 (49.2)	36 (59)	0.273
3 days remdesivir, n (%)	5 (3.9)	NA	5 (8.2)	<b>0.026</b>
5 days remdesivir, n (%)	63 (50)	32 (49.2)	31 (50.8)	0.900
Monoclonal antibodies, n (%)	37 (29.4)	NA	37 (63.8)	-
Remdesivir plus monoclonal antibodies, n (%)	23 (18.2)	NA	23 (62.2)	-



Oral antivirals, n (%)	1 (0.7)	NA	1 (1.64)	-
Tocilizumab, n (%)	9 (7.1)	6 (9.23)	3 (4.92)	0.348
Enoxaparin, n (%)	84 (66.6)	49 (75.4)	35 (57.3)	0.283
Corticosteroids, n (%)	98 (77.8)	59 (91)	39 (63.9)	<b>0.001</b>
Days of corticosteroids, median (IQR)	7 (2-17)	10.5 (8-20)	8 (6-10)	<b>0.002</b>
Baricitinib, n (%)	2 (1.6)	1 (1.6)	1 (1.64)	0.924
Convalescent plasma, n (%)	8 (6.3)	8 (12)	0 (0)	<b>0.004</b>
Antibiotics, n (%)	97 (77.0)	54 (83)	43 (70.5)	<b>0.04</b>
Vaccination, n (%)	55 (43.6)	NA	55 (90.2)	-
Number of vaccine doses, n (%)	3 (2-3)	NA	3 (2-3)	-
Last vaccine dose >120 days, n (%)	22 (17.5)	NA	22 (40)	-
<b>Outcomes</b>				
Intensive Care Unit admission, n (%)	23 (18.3)	18 (27.7)	5 (8.2)	<b>0.005</b>
In-hospital mortality, n (%)	42 (33.3)	24 (36.9)	18 (29.51)	0.378
30-d mortality, n (%)	32 (25.4)	19 (29.2)	13 (21.3)	0.307
Days of viral shedding, median (IQR)	41 (32.5)	24 (15-50)	17 (10-28)	<b>0.011</b>
Secondary infections, n (%)	48 (38.1)	25 (38.5)	23 (37.7)	0.930
MDRO colonization, n (%)	14 (11.1)	9 (13.8)	5 (8.1)	0.399
Opportunistic infections, n (%)	14 (11.1)	8 (12.3)	6 (9.84)	0.659

\*: respiratory worsening was defined as as: i) the need of supplementary oxygen therapy or ii) the need of increasing oxygen therapy supplementation in a patient with SARS-CoV2 infection for reasons directly related to the infection. A careful evaluation of causes of supplementary oxygen therapy for reasons other than SARS-CoV2 infection (i.e. cardiac failure, bacterial superinfections) was performed. In the case of doubt, a panel discussion was performed. MDRO: Multidrug resistant organisms. HFNC/CPAP: High Flow nasal cannula, Continuous Positive Airway Pressure.

**Table 2.** Analysis of risk factors for in-hospital mortality in patients with haematological malignancies.

Cox regression model	HRs (CIs 95%)	<i>p</i> -value
Sex (female)	0.67 (0.31-1.44)	0.307
Age>65y	2.02 (0.86-4.72)	0.103
Severity of infection (critical vs non-critical)	2.99 (1.15-7.80)	<b>0.025</b>
Type of haematological malignancy (compared to ALL)		
AML	2.21 (0.23-21.00)	0.490
CLL		0.989
MM	0.98 (0.10-9.83)	0.992
NHL	1.01 (0.10-9.60)	0.959
HL		0.834
Myelodysplastic syndrome	0.94 (0.11-7.72)	0.658
Myelofibrosis	1.37 (0.06-27.08)	0.835
	0.51 (0.28-9.47)	
	1.37 (0.07-27.32)	
Active malignancy°	2.20 (1.02-4.72)	<b>0.042</b>
Group of radiological involvement, % (compared to minimum, 0-30%)		
Medium (31-60%)		
High (>60%)	0.85 (0.35-2.06)	0.734
	1.29 (0.48-3.45)	0.611
Type of worse oxygen support at respiratory worsening* (compared to no oxygen support)		
Oxygen	1.11 (0.21-5.83)	0.898
HFNC/CPAP		<b>0.022</b>
MV	5.96 (1.29-27.52)	<b>0.011</b>

° Active malignancy includes patients with new diagnosis or relapsing/refractory disease, \*: respiratory worsening was defined as: i) the need of supplementary oxygen therapy or ii) the need of increasing oxygen therapy supplementation in a patient with SARS-CoV2 infection for reasons directly related to the infection. A careful evaluation of causes of supplementary oxygen therapy for reasons other than SARS-CoV2 infection (i.e. cardiac failure, bacterial superinfections) was performed. In the case of doubt, a panel discussion was performed. ALL: Acute lymphocytic leukemia, AML: Acute myeloid leukemia, CLL: Chronic lymphocytic leukemia, MM: multiple myeloma, NHL: Non-Hodgkin lymphoma, HL: Hodgkin lymphoma, HFNC: High Flow Nasal Cannula, CPAP: Continuous Positive Airway Pressure, MV: mechanical ventilation.

**Table 3.** Features of study population according to short (≤20 days) or prolonged (>21 days) viral shedding.

	<b>Short VS (N=62)</b>	<b>Long VS (N=64)</b>	<b>p value</b>
<b>Demographics</b>			
Gender (Female), n (%)	24 (38.7)	27 (42.2)	0.691
Age, median (IQR)	68.5 (57-79)	68.5 (54-76)	0.705
Steroids in the last 30 days before admission, n (%)	19 (31.2)	24 (37.5)	0.455
Infection in the last 30 days before admission, n (%)	9 (14.5)	14 (21.9)	0.285
Antibiotic therapy in the last 30 days (excluded prophylaxis), n (%)	17 (27.4)	22 (34.4)	0.398
Days of hospitalization, median (IQR)	10 (7-16)	35.5 (16.5-50)	<b>0.0001</b>
Days from symptoms onset to hospitalization, median (IQR)	2 (1-5)	3 (1-7)	0.206
Active treatment (previous 90-d)	27 (43.6)	41 (64.1)	<b>0.021</b>
Active treatment: chemotherapy only	18 (29.0)	22 (34.38)	0.520
Active treatment: immunotherapy only	4 (6.45)	17 (26.56)	<b>0.002</b>
Active treatment: chemotherapy plus immunotherapy	6 (9.7)	9 (14.1)	0.346
Chemotherapy (previous 30-d)	18 (29.5)	17 (25.6)	0.714
<b>Comorbidities, n (%)</b>			
Hypertension	30 (48.4)	22 (34.4)	0.110
Coronary artery disease	16 (25.8)	12 (18.8)	0.341
Diabetes	7 (11.3)	9 (15.6)	0.476
Chronic Obstructive Pulmonary Disease	4 (6.5)	4 (6.3)	0.963
Chronic Renal Failure	6 (9.7)	3 (4.7)	0.277
<b>Disease severity at the admission, n (%)</b>			
Moderate	43 (69.4)	37 (57.8)	0.179
Severe	14 (22.6)	17 (26.6)	0.604
Critical	4 (6.5)	4 (6.3)	0.963

Pneumonia	51 (82.3)	52 (81.3)	0.884
<b>Features at the respiratory worsening*</b>			
Worsening during hospitalization, n (%)	29 (46.8)	36 (56.3)	0.287
Days from SARS-CoV2 diagnosis to worsening, median (IQR)	3.5 (1-9)	13 (4-30)	<b>0.0005</b>
Days from admission to worsening, median (IQR)	3 (1-7)	14.5 (3-24.5)	<b>0.0003</b>
Late worsening (>10 days), n (%)	38 (61.3)	50 (78.1)	<b>0.040</b>
<b>Therapy</b>			
Overall remdesivir therapy, n (%)	40 (64.5)	27 (43.6)	<b>0.019</b>
Duration of remdesivir therapy, days (median, IQR)	5 (5-5)	5 (5-5)	0.421
Monoclonal antibodies, n (%)	21 (34.4)	16 (25.8)	0.297
Remdesivir plus monoclonal antibodies, n (%)	18 (45.0)	5 (17.2)	<b>0.016</b>
Oral antivirals, n (%)	0 (0)	1 (1.5)	-
Tocilizumab, n (%)	3 (4.8)	6 (9.4)	0.323
Corticosteroids, n (%)	42 (72.4)	56 (93.3)	<b>0.002</b>
Days of corticosteroids, median (IQR)	8 (6-11)	10 (8-22)	<b>0.001</b>
Duration of corticosteroids >7 days	44 (72.1)	55 (85.9)	0.057
Baricitinib, n (%)	0 (0)	2 (3.2)	0.154
Convalescent plasma, n (%)	0 (0)	8 (12.7)	<b>0.004</b>
Antibiotics, n (%)	45 (72.6)	52 (83.9)	0.128
Vaccination, n (%)	33 (53.2)	22 (34.4)	0.033
Last vaccine dose >120 days, n (%)	18 (36.0)	26 (46.4)	0.277
<b>Outcomes</b>			
Intensive Care Unit admission, n (%)	11 (17.7)	12 (18.8)	0.884
In-hospital mortality, n (%)	22 (35.5)	20 (31.3)	0.614
Secondary infections, n (%)	20 (32.3)	28 (43.7)	0.184

\*: respiratory worsening was defined as as: i) the need of supplementary oxygen therapy or ii) the need of increasing oxygen therapy supplementation in a patient with SARS-CoV2 infection for reasons directly related to the infection. A careful evaluation of causes of supplementary oxygen therapy for reasons


other than SARS-CoV2 infection (i.e. cardiac failure, bacterial superinfections) was performed. In the case of doubt, a panel discussion was performed. VS: viral shedding.

**Table 4.** Analysis of risk factors for prolonged (>20 days) viral shedding.

<b>Logistic regression model<sup>°</sup></b>	<b>ORs (CIs 95%)</b>	<b><i>p</i>-value</b>
Active treatment (previous 90-d)	3.55 (1.35-9.32)	<b>0.010</b>
SARS-CoV-2 vaccination	0.94 (0.26-3.33)	0.926
Corticosteroids	7.59 (1.73-33.24)	<b>0.007</b>
Respiratory worsening* >10 days	2.92 (1.05-8.05)	<b>0.038</b>
Duration of corticosteroid therapy >7 days	4.92 (1.59-15.22)	<b>0.006</b>
Type of therapy for SARS CoV2 infection (compared to no therapy)		
RDV only	0.30 (0.09-0.955)	<b>0.042</b>
Monoclonal antibodies only	2.09 (0.31-13.81)	0.444
RDV+Monoclonal antibodies	0.16 (0.02-1.07)	0.060

<sup>°</sup>: model adjusted for age and sex. \*: respiratory worsening was defined as: i) the need of supplementary oxygen therapy or ii) the need of increasing oxygen therapy supplementation in a patient with SARS-CoV2 infection for reasons directly related to the infection. A careful evaluation of causes of supplementary oxygen therapy for reasons other than SARS-CoV2 infection (i.e. cardiac failure, bacterial superinfections) was performed. In the case of doubt, a panel discussion was performed. RDV: remdesivir.

## Figures



*Legend:* HM, haematological malignancies; other haematological diseases: haematological diseases other than malignancy (as MGUS or thrombocytopenia)

**Figure 1**

Flow chart of the study population. HM: haematological malignancies

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarydata.docx](#)