

# Outer membrane composition of a lipopolysaccharide-deficient *Neisseria meningitidis* mutant

Liana Steeghs<sup>1,2,3</sup>, Hans de Cock<sup>2</sup>,  
Evert Evers<sup>4</sup>, Bert Zomer<sup>4</sup>, Jan Tommassen<sup>2</sup>  
and Peter van der Ley<sup>1</sup>

<sup>1</sup>Laboratory of Vaccine Research and <sup>4</sup>Laboratory of Organic-Analytical Chemistry, National Institute of Public Health and the Environment, RIVM, Antonie van Leeuwenhoeklaan 9, PO Box 1, 3720 BA Bilthoven and <sup>2</sup>Department of Molecular Microbiology and Institute of Biomembranes, Utrecht University, Padualaan 8, 3584 CH Utrecht, The Netherlands

<sup>3</sup>Corresponding author at: Laboratory of Vaccine Research, National Institute of Public Health and the Environment, RIVM, Antonie van Leeuwenhoeklaan 9, PO Box 1, 3720 BA Bilthoven, The Netherlands  
e-mail: liana.steeghs@rivm.nl

**In the pathogen *Neisseria meningitidis*, a completely lipopolysaccharide (LPS)-deficient but viable mutant can be obtained by insertional inactivation of the *lpxA* gene, encoding UDP-GlcNAc acyltransferase required for the first step of lipid A biosynthesis. To study how outer membrane structure and biogenesis are affected by the absence of this normally major component, inner and outer membranes were separated and their composition analysed. The expression and assembly of integral outer membrane proteins appeared largely unaffected. However, the expression of iron limitation-inducible, cell surface-exposed lipoproteins was greatly reduced. Major changes were seen in the phospholipid composition, with a shift towards phosphatidylethanolamine and phosphatidylglycerol species containing mostly shorter chain, saturated fatty acids, one of which was unique to the LPS-deficient outer membrane. The presence of the capsular polysaccharide turned out to be essential for viability without LPS, as demonstrated by using a strain in which LPS biosynthesis could be switched on or off through a *tac* promoter-controlled *lpxA* gene. Taken together, these results can help to explain why meningococci have the unique ability to survive without LPS.**

**Keywords:** capsular polysaccharide/lipopolysaccharide/*Neisseria meningitidis*/outer membrane proteins/  
phospholipids

## Introduction

The outer membrane of Gram-negative enteric bacteria is an asymmetrical bilayer composed of phospholipids and lipopolysaccharide (LPS) in the inner and outer leaflet, respectively (Kamio and Nikaido, 1976). LPS consists of three moieties: the hydrophobic lipid A, which anchors it to the outer membrane; a generally well-conserved surface-exposed hydrophilic oligosaccharide, designated the core; and a hypervariable polysaccharide, the O-antigen. The O-antigen is lacking in many mucosal surface-colonizing bacteria, such as *Neisseria*

*meningitidis*. Moreover, the asymmetry of the outer membrane of the Neisseriaceae is less pronounced than in the enteric bacteria since phospholipids are also present in the outer leaflet of the outer membrane (Lysko and Morse, 1981). The core oligosaccharide of LPS is negatively charged, resulting in a strong affinity for divalent cations (Schindler and Osborn, 1979). The resulting network of divalent cations and LPS forms a strong permeability barrier in which outer membrane proteins (OMPs) are inserted to facilitate the influx of nutrients.

Among the major OMPs are the porins, which form pores to allow for the non-specific uptake of small compounds by passive diffusion. The basic structure of these porins is a trimer, in which each monomer forms a  $\beta$ -barrel (Weiss *et al.*, 1991; Cowan *et al.*, 1992). The porins of *N.meningitidis* are designated PorA and PorB. Additional OMPs are required for the uptake of specific nutrients. For example, growth of *N.meningitidis* under iron limitation results in the expression of receptors for iron-binding proteins of the host, such as lactoferrin and transferrin. These receptors generally consist of an integral OMP with sequence similarity to the siderophore receptors of *Escherichia coli*, and a cell surface-exposed lipoprotein (Schryvers and Morris, 1988; Legrain *et al.*, 1993; Pettersson *et al.*, 1994; Pettersson *et al.*, 1998). The outer membrane of *N.meningitidis* is anchored to the peptidoglycan in the periplasm by the RmpM protein, a homologue of the *E.coli* OmpA (Klugman *et al.*, 1989). However, in contrast to OmpA, which is anchored in the outer membrane via an N-terminal  $\beta$ -barrel domain (Pautsch and Schulz, 1998), RmpM lacks such a  $\beta$ -barrel domain but is firmly associated with the porins (Jansen *et al.*, 2000) and the iron limitation-inducible OMPs (Prinz and Tommassen, 2000).

The cell surface of many Gram-negative bacteria is covered by a capsular polysaccharide (CPS). In serogroup B *N.meningitidis* strains, such as our model strain H44/76, this capsule is composed of  $\alpha$ -2,8-linked polysialic acid, which is anchored in the outer membrane via a phospholipid at the reducing end of the polysaccharide chain (Gotschlich *et al.*, 1981; Frosch and Muller, 1993). The genes involved in the biosynthesis, regulation and transport of CPS are clustered on the chromosome (Frosch *et al.*, 1989).

Various *in vivo* and *in vitro* studies have shown that in *E.coli*, LPS is required for the correct assembly of OMPs (Koplow and Goldfine, 1974; Ried *et al.*, 1990; Sen and Nikaido, 1991; Laird *et al.*, 1994; de Cock and Tommassen, 1996; de Cock *et al.*, 1999a). Disruption of the genes involved in the early steps of lipid A biosynthesis appears impossible and only conditionally lethal mutants have been described. Therefore, it was suggested that this part of LPS is essential for bacterial

growth (Raetz, 1990). However, we have recently isolated an LPS-deficient, but viable mutant of *N.meningitidis* by insertional inactivation of *lpxA*, encoding the enzyme required for the first step in lipid A biosynthesis (Steeghs *et al.*, 1998). This mutant still produced an outer membrane, which could be clearly discerned by electron microscopy. Preliminary data did not show major deviations in the OMP profile of this LPS-deficient mutant compared with that of the wild-type strain.

In this study, we have characterized the outer membrane composition of the LPS-deficient mutant in detail. Our results demonstrate that in *N.meningitidis*, LPS is not essential for correct assembly of integral OMPs, such as the porins, although it is required for biogenesis of cell surface-exposed lipoproteins. Furthermore, the phospholipid composition appears to be altered in the absence of LPS. Interestingly, CPS appears to be required for viability of the LPS-deficient *N.meningitidis* mutant.


## Results

### Separation of inner and outer membranes of the LPS-deficient mutant

To analyse the cell envelope composition of the LPS-deficient *N.meningitidis* strain, inner and outer membranes of strain H44/76 and its *lpxA* mutant derivative were separated by isopycnic sucrose gradient centrifugation. Since it is generally assumed that the higher buoyant density of the outer membranes, as compared with the inner membranes, is due to the presence of LPS, we were not sure whether separation of the membranes of the LPS-deficient mutant was possible by this method. For this analysis, 1 ml gradient fractions were collected and analysed for the presence of lactate dehydrogenase (LDH) activity as a marker for the inner membrane (Osborn *et al.*, 1972), and for the presence of the porins PorA and PorB as outer membrane markers. LDH activity peaked in fraction 3, followed by a gradual decline in fractions 4–6, with no significant difference between the membranes of the two strains (Figure 1A). Thus, the inner membranes of both the wild-type and the mutant strain localized in the low-density region, i.e. fractions 1–6, of the gradient. For both strains, the majority of PorA and PorB was found in the high-density fractions 7–12 (Figure 1B), which contained hardly any LDH activity, while some PorA and PorB was present in the low-density fractions 4–6, which have intermediate LDH activity (Figure 1). Apparently, the absence of LPS did not influence the separation of inner and outer membranes by isopycnic sucrose gradient centrifugation. For further analysis, gradient fractions were pooled in pure inner membrane fraction P1 (fractions 1–3), inner membrane fraction P2 (fractions 4–6), which is slightly contaminated with outer membranes, outer membrane fraction P3 (fractions 7–8), which is slightly contaminated with inner membranes, and pure outer membrane fraction P4 (fractions 9–12), as indicated in Figure 1A.


### Assembly of outer membrane proteins in the LPS-deficient mutant

LPS has been implicated in the biogenesis of Gram-negative bacterial porins, and therefore defects in the assembly of PorA and PorB in the LPS-deficient strain



**Fig. 1.** Analysis of fractions 1–12, collected after isopycnic sucrose gradient centrifugation, to separate the membranes of wild-type strain H44/76 and its LPS-deficient mutant. (A) LDH per milligram of protein and sucrose concentration of the fractions. P1–4 indicate the fractions that were pooled for detailed analysis. (B) SDS-PAGE analysis of the fractions. Positions of molecular weight standard proteins are indicated on the left in kDa; positions of the porins PorA and PorB are indicated on the right.


were to be expected. The oligomeric state of PorA in outer membrane fractions P3 and P4 was analysed by SDS-PAGE under denaturing and non-denaturing conditions followed by western blotting. Analysis of outer membranes of the LPS-deficient strain under denaturing conditions revealed, besides the full-length protein of the expected apparent molecular weight, a partial degradation product of PorA with a slightly higher electrophoretic mobility (Figure 2A, lanes a). This degradation product was detected with monoclonal antibody (mAb) MN5C11G (Figure 2A) and MN23G2.38 (results not shown), which recognize epitopes in the putative cell surface-exposed loops 4 and 3 of PorA, respectively. No reaction was found with mAb MN14C11G, which recognizes loop 1, indicating that proteolysis had occurred at the N-terminal end of the protein (results not shown). In spite of this partial degradation, trimerization of PorA appeared unaffected in




**Fig. 2.** Western immunoblot of outer membrane fractions P3 and P4 of the wild-type strain H44/76 and of its LPS-deficient mutant, incubated with anti-PorA mAb MN5C11G (A) and anti-RmpM mAb MN2D6D (B). The samples were incubated at 100°C (lanes a) or room temperature (lanes b) in the presence of 2% SDS before SDS-PAGE. The positions of the molecular weight markers (kDa), the PorA-containing oligomeric complexes, and the PorA and the RmpM monomer are indicated.

the LPS-deficient mutant. Both the PorA and its degradation product migrated as trimers with the same electrophoretic mobility as the trimers in the wild-type strain, when the samples were not boiled before electrophoresis (Figure 2A, lanes b). The PorA trimers from the H44/76 membranes have been shown to remain associated with the RmpM protein during non-denaturing SDS-PAGE (Jansen *et al.*, 2000). Western blot analysis with the RmpM-specific mAb MN2D6D confirmed the presence of RmpM in the trimeric porin complexes of both the wild-type and the mutant strain (Figure 2B). Thus, porin assembly is not affected in the LPS-deficient strain. In addition, no effect was found on expression and assembly of other integral OMPs, such as the opacity proteins and the outer membrane phospholipase A (results not shown). Expression of the secretin PilQ could also be clearly demonstrated in the LPS-deficient mutant, although apparently at a slightly lower level as compared with the wild-type strain (Figure 3). However, no difference in pili expression was found for the LPS-deficient mutant when compared with the wild type as measured by a dot-blot assay with the pilin-specific mAb SM1 (M.Christoudoulides, unpublished observation).

We also studied the expression of the cell surface-exposed lipoproteins LbpB and TbpB, which are parts of the lactoferrin and transferrin receptors, respectively. To express these receptors, strain H44/76 and the LPS-deficient mutant were grown under iron limitation, and the membranes were isolated by isopycnic sucrose gradient centrifugation. Western blot analysis did not reveal any significant differences between the two strains with respect to the expression level of the integral membrane component LbpA of the lactoferrin receptor (Figure 4A). In contrast, the amounts of the lipoproteins LbpB and



**Fig. 3.** Western immunoblot of membrane fractions P1–P4 of the wild-type strain H44/76 and its LPS-deficient mutant, incubated with anti-PilQ monoclonal antibody.




**Fig. 4.** Western immunoblot of membrane fractions P1–P4 of the wild-type strain H44/76 and its LPS-deficient mutant, incubated with anti-LbpA antiserum, anti-LbpB antiserum and anti-TbpB antiserum (A) and anti-CtrA mAb 2619 (B).

TbpB were drastically reduced in the outer membranes of the mutant (Figure 4A). The LbpB and TbpB proteins did not accumulate in the inner membrane fractions P1 and P2, in the soluble fraction or in the extracellular medium (results not shown) and are probably degraded upon mislocalization. In contrast, no difference was found in the expression of other outer membrane-associated lipoproteins, such as CtrA (Figure 4B), presumed to be part of the CPS transport complex (Frosch *et al.*, 1992). Similarly, expression of the lipoprotein H.8 associated with the periplasmic side of the outer membrane appeared unaffected (R.Moxon and J.Richards, unpublished observation). In conclusion, these results show that the biogenesis of cell surface-exposed lipoproteins is clearly disturbed in the LPS-deficient mutant, whereas the biogenesis of integral OMPs, RmpM and other lipoproteins is not affected.

#### Phospholipid composition of inner and outer membranes in the LPS-deficient mutant

To study the effect of LPS deficiency on the phospholipid composition of the cell envelope, phospholipids were isolated from crude membrane fractions and analysed by thin-layer chromatography (TLC). The majority of the phospholipids found in both H44/76 and the LPS-deficient mutant were phosphatidylethanolamine (PE) and, to a lesser extent, phosphatidylglycerol (PG). Only trace amounts of cardiolipin (CL) and no detectable amounts of lysophospholipids were present in both strains (results not shown). In both strains, the distribution of these phospholipid classes in inner and outer membranes was similar. However, analysis of the fatty acyl chains of PE in



**Fig. 5.** Distribution of the PE and PG species in membrane fractions P1–P4 of the wild-type strain H44/76 and its LPS-deficient mutant. The estimated error in the measurements of the fatty acid contents of the membrane phospholipids is 10%. Example of abbreviations of phospholipids: PE(C32:0) indicates a phosphatidyl-ethanolamine in which the total number of carbons for both fatty acyl residues is 32.

the isolated membrane fractions P1–P4 showed a remarkable difference between the two strains (Figure 5). In both inner membranes (fractions P1 and P2) and outer membranes (fractions P3 and P4) of the LPS-deficient strain, an increase in the relative amounts of the saturated PE species PE(C30:0) [PE(C30:0) indicates a saturated phosphatidyl-ethanolamine in which the total number of carbons for both fatty acyl residues is 30] and PE(C32:0), and a concomitant decrease in the unsaturated PE species PE(C32:1), PE(C32:2), PE(C34:1) and PE(C34:2), was apparent. The reduced amount of unsaturated PE was most striking for PE(C32:1), representing 50% of the PE species in H44/76 outer membranes (fractions P3 and P4), but only 30% in the outer membrane of the mutant strain. Remarkably, PE(C32:0), which was barely detectable in

outer membranes of the wild-type strain, represented 30% of the PE species in those of the mutant.


A similar analysis of the PG content revealed that the fatty acyl chain length in both strains was generally shorter than in the major PE species: PG(C28) versus PE(C32) (Figure 5). As for PE, an increase in saturated fatty acids was apparent in PG in the outer membranes of the LPS-deficient mutant, but this was not the case in the inner membrane. PG(C28:0), representing 50 and 35% of the PG species in inner and outer membranes of the wild-type strain, respectively, was reduced to 30% and increased to 50%, respectively, in those of the mutant. In conclusion, these data show that the LPS-deficient mutant preferentially incorporates saturated PE and PG species with shorter fatty acyl chains in its cell envelope, and of these phospholipids, PE(C32:0) is uniquely present in the LPS-deficient outer membrane.

### **CPS is essential for viability of the LPS-deficient mutant**


The expression of the CPS in the LPS-deficient mutant was comparable to CPS expression in wild-type H44/76, as measured by whole-cell enzyme-linked immunosorbent assay (ELISA) with a CPS-specific antibody. Capsule-deficient mutants of the wild-type strain H44/76 could readily be obtained by complete deletion of the *cps* cluster or by insertional inactivation of the *siaD* gene, encoding the polysialyl transferase involved in multimerization of the polysialic B CPS as described previously (Edwards *et al.*, 1994). However, we were not able to isolate a CPS-deficient mutant of the *lpxA* mutant strain in either way. This raised the possibility that the CPS is essential for viability of the LPS-deficient mutant. To test this hypothesis, we constructed a derivative of strain H44/76, designated HA3003, which carries an isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG)-inducible *lpxA* gene on the chromosome, allowing for depletion of LPS (Figure 6).

To test for *lacI<sup>q</sup>-tac* control of *lpxA* in strain HA3003, LPS was extracted from lysates of whole cells, which were grown in Mueller–Hinton broth, and the LPS was analysed by Tricine–SDS–PAGE (Figure 7). Surprisingly, no difference in LPS expression was observed between HA3003 grown with or without IPTG. Possibly, the casein hydrolysate present in Mueller–Hinton broth contained enough lactose to derepress the *tac* promoter. Therefore, other growth media, i.e. GC agar and Meningococcal Medium, neither of which contains casein, were tested as well. As expected, LPS was barely, if at all, detected when HA3003 was grown in either of these media without IPTG, but its biosynthesis was restored to the wild-type level upon induction with IPTG (Figure 7). Furthermore, the growth properties of HA3003 were compared with those of the wild-type strain and the *lpxA* mutant in Meningococcal Medium with and without IPTG (Figure 8). The reduced growth rate previously reported for the *lpxA* mutant (Steeghs *et al.*, 1998) was again observed (results not shown) and was also seen, even to a stronger degree, for HA3003 in the absence of IPTG. However, the growth rate of HA3003 was largely restored upon addition of IPTG (Figure 8), confirming that a tight regulation of *lpxA* was achieved in this strain.


Subsequently, a CPS-deficient mutant of strain HA3003 was constructed in the presence of IPTG, by



**Fig. 6.** Schematic representation of the chromosomal arrangement of the *lpxD-fabZ-lpxA* locus of wild-type strain H44/76 and of the *lacI<sup>Q</sup>-tac*-regulated *lpxA* strain HA3003. The transcriptional terminator is indicated by a black triangle.



**Fig. 7.** Silver-stained Tricine-SDS-PAGE gel showing the LPS pattern of proteinase K-treated whole-cell lysates of HA3003 grown in the absence (a) or presence (b) of 1 mM IPTG on Mueller-Hinton broth (MHB), Gonococcal Agar (GC) or Meningococcal Medium (Men Med).




**Fig. 8.** Growth curves of the *lacI<sup>Q</sup>-tac*-regulated *lpxA* strain HA3003 (circles), its CPS-deficient derivative HA2104 (triangles) and wild-type strain H44/76 (squares) in Meningococcal Medium in the absence (grey lines) or presence (black lines) of 1 mM IPTG.

transformation and allelic replacement with plasmid pMF32.35::T5 (Frosch *et al.*, 1989) carrying a *Tn1725* insertion in the *siaD* gene. Absence of the capsule in one of the transformants was confirmed by ELISA with a CPS-specific antibody and this strain was designated HA2104. This strain was then used to study the influence of CPS deficiency in an LPS-deficient background. Whereas a *siaD* mutant derivative of the wild-type strain H44/76 grew like its parental strain in the presence or absence of IPTG (results not shown), HA2104 did not grow at all when IPTG was absent, whereas growth was restored in the presence of IPTG (Figure 8). Thus, it can be concluded that the capsule is essential for the viability of an LPS-deficient *N.meningitidis* mutant.

#### Outer membrane permeability barrier

In order to determine the effect of LPS deficiency on the barrier function of the outer membrane, the susceptibility



**Fig. 9.** Antibiotic susceptibility of wild-type strain H44/76 and its LPS-deficient mutant. The diameter of the inhibition zone around filter paper discs containing each of four different antibiotics is shown. The data shown represent one of three experiments.

of the LPS-deficient mutant to four hydrophobic antibiotics was determined (Figure 9). Larger growth inhibition zones around filter discs containing these antibiotics were observed for the LPS-deficient mutant than for the wild-type strain, indicating that the barrier function of the outer membrane is compromised.

#### Discussion

In contrast to most Gram-negative bacteria, *N.meningitidis* is viable without LPS (Steeghs *et al.*, 1998). Since LPS determines the asymmetry of the outer membrane and many important functions in membrane biogenesis have been assigned to this molecule, one might expect that the composition of the LPS-deficient mutant outer membrane is drastically changed and/or that the role of LPS is taken over by other components. Therefore, we studied the outer membrane composition of the LPS-deficient mutant in detail. Inner and outer membranes were separated by isopycnic sucrose gradient centrifugation. Although it is generally assumed that the presence of LPS determines the higher buoyant density of the outer membrane as compared with the inner membrane (Osborn *et al.*, 1972), the buoyant density of the LPS-deficient outer membranes appeared unchanged. Apparently, other outer membrane components contribute to the higher buoyant density of the *N.meningitidis* outer membrane. In this respect, the CPS, which is anchored in the outer membrane via a phospholipid tail (Gotschlich *et al.*, 1981; Frosch and Muller, 1993) and peptidoglycan, which is presumably associated to the outer membrane via the RmpM protein (Klugman *et al.*, 1989), are the most likely candidates.

Analysis of the phospholipid composition of the separated membranes revealed a remarkable shift in both inner and outer membranes of the LPS-deficient mutant towards phospholipids with shorter and saturated fatty acyl chains. This effect was most pronounced for the major phospholipid PE. The preference for the shorter fatty acids in the phospholipids in the outer membrane of the *lpxA* mutant can be rationalized by the presence of generally shorter fatty acids in LPS than in phospholipids. For the hydrophobic matching with the integral outer membrane proteins, those lipidic components replacing LPS should not affect the overall thickness of the outer membrane. Furthermore, the fatty acyl chains of LPS are generally saturated. Thus, the increased saturation of the fatty acyl

chains of the phospholipids might compensate for the membrane fluidity changes invoked by the LPS deficiency. The observed increase in susceptibility to hydrophobic antibiotics is consistent with such a replacement of the missing LPS by phospholipids.

The failure to disrupt CPS synthesis suggested an essential role for the CPS in the LPS-deficient mutant strain. To verify this, strain HA3003 carrying the *lpxA* gene under the control of an IPTG-inducible promoter was constructed. When this strain was grown in the absence of IPTG, LPS was barely detectable and growth was even more impaired than that of the *lpxA* knock-out strain. A possible explanation for this observation is that the insertion of a transcriptional terminator immediately downstream of *fabZ* has affected its expression and/or regulation either at the transcriptional or translational level. Consequently, expression of FabZ, an essential enzyme involved in fatty acid biosynthesis (Mohan *et al.*, 1994), might be reduced, with a negative effect on the biosynthesis of phospholipids. Furthermore, we can not exclude the existence of secondary compensating mutations in the *lpxA* knock-out strain resulting in a higher growth rate of this strain compared with strain HA3003. Upon induction with IPTG, LPS synthesis was restored to the wild-type level in HA3003, but there was still a slight growth defect apparent. HA2104, a *siaD* mutant of HA3003, was not viable in the absence of IPTG, directly demonstrating that CPS is essential for viability in an LPS-deficient *N.meningitidis* mutant. This would explain our unsuccessful attempts to construct an LPS-deficient mutant of the closely related bacterium *Neisseria gonorrhoeae*, which lacks a CPS (our unpublished observation). Interestingly, it has been reported that glycosphingolipids that are present in the outer membrane of the naturally LPS-deficient Gram-negative rod *Sphingomonas paucimobilis* exhibit a high degree of similarity to LPS in their contribution to outer membrane function (Wiese *et al.*, 1996). Like the CPS of H44/76, these glycosphingolipids contain two fatty acyl chains covalently linked to a sugar moiety, in contrast to the six or seven fatty acyl chains normally present in LPS. In this respect, CPS seems structurally and functionally analogous to sphingolipids. Furthermore, CPS could possibly maintain the typical asymmetry of the outer membrane, by which it can be distinguished from the inner membrane, thus allowing for insertion of outer membrane proteins in the correct membrane. In this way, the role of LPS in the biogenesis of porins, which has been described previously in *E.coli* and *Salmonella typhimurium* (Koplow and Goldfine, 1974; Ried *et al.*, 1990; Sen and Nikaido, 1991; Laird *et al.*, 1994; de Cock and Tommassen, 1996; de Cock *et al.*, 1999a), might be taken over by CPS. In deep-rough LPS mutants of *E.coli* and *S.typhimurium*, a reduction in the amounts of OMPs has been observed (Koplow and Goldfine, 1974). In contrast, such a reduction was not observed in a comparable heptose-deficient *rfaC* mutant of *N.meningitidis* (P.van der Ley, unpublished data), already indicating that OMP assembly proceeds differently in this organism. Moreover, our data also show that the lipid A moiety of LPS is not required for assembly of integral outer membrane proteins, such as the porins. Although a minor degradation product of PorA was detected, the expression and assembly of the major OMPs appeared

unaffected in the *N.meningitidis* LPS-deficient mutant. This degradation product was also observed in the *rfaC* mutant of *N.meningitidis* (P.van der Ley, unpublished observation), pointing to a proteolytic cleavage site within PorA that becomes accessible in the absence of the complete oligosaccharide core. If the meningococcal porins do indeed need a glycolipid for their assembly, the CPS would be the obvious candidate.

The only group of OMPs whose expression was greatly reduced in the LPS-deficient mutant encompasses the cell surface-exposed lipoproteins, such as LbpB and TbpB. These lipoproteins did not accumulate in the inner membrane and were not detected in the soluble fraction or extracellular medium. These proteins are probably degraded when they are not properly localized. In *E.coli*, the LolA-E system is required for the sorting of lipoproteins from the inner membrane to the outer membrane (Yokota *et al.*, 1999; Yakushi *et al.*, 2000). Homologues of the Lol system are present in the *N.meningitidis* chromosome (Tettelin *et al.*, 2000), indicating that a similar route for the translocation of outer membrane lipoproteins exists in *N.meningitidis*. Since normal expression of the lipoproteins CtrA and H.8, both of which are associated with the outer membrane but are not fully cell surface exposed, was found in the LPS-deficient mutant, it can be concluded that lipoprotein transporting to the outer membrane by the Lol system is not affected. However, no component of the Lol system involved in the translocation of the surface-exposed lipoproteins from the periplasmic side of the outer membrane to the cell surface has yet been identified, a step which is not known to occur in *E.coli*. LPS could be involved in this step by binding these lipoproteins on their way to the outer leaflet of the outer membrane. Another possibility is that these lipoproteins do not correctly insert and/or are easily released and degraded, due to the altered biophysical properties of the outer membrane.

OMPs are considered important candidates for the development of a vaccine against *N.meningitidis*. However, the presence of LPS, also known as endotoxin, is a drawback in the use of outer membrane vesicles for immunization. With the availability of an LPS-deficient strain, this problem may be circumvented. However, we have previously shown that the immunogenicity of outer membrane complexes of an LPS-deficient *N.meningitidis* mutant is greatly reduced (Steeghs *et al.*, 1999). Furthermore, the requirement for the presence of the polysialic acid CPS, which is homologous to host antigens, constitutes another drawback in the use of this LPS-deficient *N.meningitidis* mutant for a whole-cell or vesicle vaccine. However, by using the newly constructed strain HA3003 with an IPTG-inducible *lpxA* gene, it might be feasible to find an optimal balance between endotoxin content and immunogenicity even when the CPS is not present.

*Neisseria meningitidis* is the only Gram-negative bacterium described so far in which the synthesis of LPS can be disrupted completely. The viability of the mutant is apparently related to the fact that *N.meningitidis* carries a CPS, which is anchored in the outer membrane via a phospholipid. The structure of the *E.coli* K1 capsule is similar to the *N.meningitidis* B capsule (Kasper *et al.*, 1973) and it would be interesting to investigate whether an

LPS-deficient *E.coli* K1 mutant is also viable. On the other hand, in contrast to *E.coli*, *N.meningitidis* and also *N.gonorrhoeae* contain phospholipids in the outer leaflet of the outer membrane (Lysko and Morse, 1981), indicating that the typical asymmetry of the outer membrane normally found in Gram-negative bacteria is less pronounced in the Neisseriaceae. Consequently, Neisseriaceae are apparently more flexible than other bacterial species in accepting lipidic components other than LPS in the outer leaflet of the outer membrane. Nevertheless, Sprong *et al.* (2001) recently demonstrated that the LPS content of *N.meningitidis* strain H44/76 is similar to that previously reported for *E.coli*. It can thus be ruled out that LPS normally represents only a very minor component of the outer membrane of *N.meningitidis*, and would therefore be more easily replaced by phospholipids than in *E.coli*. However, as compared with *E.coli*, greater heterogeneity of the membrane phospholipids has been found in *N.meningitidis* and *N.gonorrhoeae* (Rahman *et al.*, 2000). In particular, the pool of neisserial phospholipids having short-chain fatty acyl substituents, i.e. C12:0, C14:0 and C14:1, is much greater than previously reported for *E.coli* (Wilkinson, 1988). This diversity of meningococcal phospholipid acyl chains presumably results from the different substrate specificities of its multiple lysophosphatidic acid (LPA) acyltransferases. At least two LPA acyltransferases were demonstrated in *N.meningitidis* (Shih *et al.*, 1998), in contrast to only a single one in *E.coli* (Coleman, 1992). Regulation of these LPA acyltransferases in *N.meningitidis* might help to compensate for defects occurring after the loss of LPS. Taken together, the ability of *N.meningitidis* to synthesize such a heterogeneous phospholipid pool and the existence of the CPS, being a glycolipid itself, apparently permit this bacterium to survive without LPS.

## Materials and methods

### Bacterial strains and growth conditions

*Escherichia coli* NM522 (Promega), used for the propagation of plasmids, was grown in LB medium containing ampicillin or kanamycin (100 µg/ml) at 37°C. *Neisseria meningitidis* strains were grown overnight at 37°C on GC medium base (Difco Laboratories) supplemented with IsoVitalX (Becton Dickinson) in a humid atmosphere containing 5% CO<sub>2</sub>. For the isolation of inner and outer membranes, bacteria were grown in Meningococcal Medium (van der Ley *et al.*, 1993), supplemented, if appropriate, with 39 µM ethylene-diamine-dihydroxyphenyl-acetic acid (Sigma) to impose iron restriction. Bacteria were grown until late log phase and inactivated by incubation for 2 h with 200 µg/ml tetracycline in 2% ethanol (final concentration) at 37°C under aeration. Transformation of *N.meningitidis* was carried out as described previously (van der Ley and Poolman, 1992) with selection on kanamycin (100 µg/ml) or chloramphenicol (10 µg/ml). To induce the *tac* promoter, strains were grown in Mueller-Hinton broth or Meningococcal Medium in the presence of 1 mM IPTG (Roche).

### Construction of plasmids

Plasmids were constructed by using standard recombinant DNA techniques (Sambrook *et al.*, 1989). Plasmid DNA was isolated using the Wizard kit (Promega). Plasmid pLA19 is a pUC18 derivative carrying a 1.9 kb insert, containing nucleotides 656–2472 of the *lpxD-fabZ-lpxA* locus of strain H44/76 (Steeghs *et al.*, 1997). This plasmid was the basis for the construction of plasmid pLSK10, containing *lpxA* under an IPTG-inducible promoter. First, a 64mer linker named Termlink1 (5'-CGC-GTGGTTTGAGTGAAGTTAAGGGATGCATAAACTGCATCCCTT-AACTTGTTTTTTCGCGAA-3'), which possesses *MluI* overhangs and contains the last 12 bp (including stop codon) of *fabZ*, a transcriptional terminator and a unique *NruI* site, was cloned into the *MluI* site located

12 bp upstream of the stop codon of *fabZ* (Figure 6), resulting in plasmid pLS7. In this way, the complete *fabZ* open reading frame was maintained, while a transcriptional terminator was introduced 3 bp downstream of the *fabZ* stop codon. The orientation of Termlink1 in pLS7 was verified by sequence analysis with an Applied Biosystems automatic sequencer 310 on double-stranded plasmid DNA templates with a cycle sequencing protocol and primers hybridizing with *fabZ* and *lpxA*. Subsequently, the *lacI<sup>t</sup>-tac* control region released from plasmid pJF118EH (Fürste *et al.*, 1986) by a *NruI*-*SmaI* digest was inserted into the unique *NruI* site in pLS7 upstream of *lpxA*, resulting in plasmid pLS8. The orientation of the *lacI<sup>t</sup>-tac* control region was checked by restriction analysis with *NruI*, which had to be maintained after this cloning procedure. pLS8 was then used to insert a kanamycin-resistance gene, released from plasmid pUC4K (Vieira and Messing, 1982) by *HindIII* digestion, in the opposite transcriptional orientation of the *tac* promoter to create pLSK10. This plasmid was then used to transform *N.meningitidis* H44/76 (van der Ley and Poolman, 1992) to create strain HA3003. Plasmid pMF32.35:T5, containing a Tn1725 insertion element in the *siaD* gene, was a generous gift of Dr M.Frosch (Universität Würzburg, Germany).

### Membrane separation

Inner membranes and outer membranes were isolated and separated by sucrose density gradient centrifugation as described (Masson and Holbein, 1983). Briefly, bacteria were harvested by centrifugation in a Sorvall SS34 rotor at 6000 r.p.m. at 4°C, and pellets were washed three times with phosphate-buffered saline (PBS). Pellets were resuspended in 50 mM Tris-HCl pH 8.0, and 50 µg/ml RNase (Roche) and DNase (Roche) were added. Bacteria were passed twice through a French pressure cell at 15 000 p.s.i. Unbroken cells were removed by centrifugation in a Sorvall SS34 rotor at 6000 r.p.m. at 4°C, and the supernatant was loaded on top of a discontinuous sucrose gradient consisting of a 6 ml 55% (w/w) sucrose cushion and a 9 ml 15% (w/w) sucrose top layer, both in 3 mM EDTA pH 8.0. After centrifugation for 2 h at 50 000 r.p.m. in a Beckman Ti60 rotor at 4°C, the crude membrane fraction was collected from the top of the cushion with a J-shaped Pasteur pipette. The sucrose concentration of this crude membrane fraction was lowered to 30% (w/w) sucrose with 3 mM EDTA pH 8.0, followed by separation on a second discontinuous sucrose gradient, consisting of 2.9 ml layers of 45, 40 and 35% (w/w) sucrose on top of a 1.9 ml 50% (w/w) sucrose cushion. The gradient was centrifuged for 36 h at 33 200 r.p.m. in an SW41 Beckman rotor at 4°C, and fractionated in 12 1-ml fractions from top to bottom. Sucrose concentrations in the fractions were determined by using a refractometer. Subsequently, the sucrose concentration of either the individual fractions or pooled fractions as depicted in Figure 1 was lowered to <10% (w/w) sucrose with 3 mM EDTA pH 8.0. Subsequently, membranes were collected by centrifugation for 2 h at 50 000 r.p.m. in a Beckman Ti60 rotor at 4°C and resuspended in buffer L (50 mM triethanolamine acetate pH 7.5, 250 mM sucrose, 1 mM dithiothreitol). The protein content of the membranes was determined using a bicinchoninic acid protein assay reagent (Pierce Chemical Co.) with bovine serum albumin as a standard. Membranes were stored at -20°C.

### Lactate dehydrogenase activity assay

Lactate dehydrogenase (LDH) activity was measured as described (Osborn *et al.*, 1972).

### SDS-PAGE and western blotting

Prior to electrophoresis, fractions were incubated in sample buffer (Lugtenberg *et al.*, 1975) containing 2% SDS (final concentration). Incubations were performed for 10 min at room temperature or 100°C. SDS-polyacrylamide gels were prepared according to Lugtenberg *et al.* (1975). The gels were run at 20 mA in a temperature-controlled room at 4°C to prevent denaturation of various folded forms of the proteins as a result of heating of the gels during electrophoresis. Western blotting was performed as described (de Cock *et al.*, 1999b). Prior to the blotting procedure, gels were heated under steam (Brok *et al.*, 1995) to denature folded proteins. The PorA-specific mAbs used were MN5C11G (P1.16) or MN14C11.6 (P1.7) (Scholten *et al.*, 1993). The RmpM-specific mAb used was MN2D6D (Poolman *et al.*, 1985). The PilQ-specific antiserum was raised in mice against purified PilQ. The LbpA-specific mouse antiserum was directed against loop 2 of LbpA (Prinz *et al.*, 1999). The LbpB-specific antiserum was raised in rabbits (A.Pettersson, P.Voet, P.Nadin, J.T.Poolman and J.Tommassen, in preparation). The TbpB-specific rabbit antiserum was a generous gift of Dr Anthony Schryvers (University of Calgary, Canada).

### Phospholipid analysis

Total lipids were isolated from membrane fractions as described (Bligh and Dyer, 1959). Phospholipids were separated on TLC plates (HPTLC silicagel 60, Merck) with chloroform/methanol/acetic acid in a ratio of 65:25:10. Lipids were visualized by spraying with 50% sulfuric acid followed by charring. The fatty acyl content of PE and PG species was determined by using fast atom bombardment neutral loss tandem mass spectrometry (Cole and Enke, 1991). The specific loss of 141 u and 172 u was used to assign specific ions as those derived from PE and PG lipids, respectively.

### LPS analysis

Tricine-SDS-PAGE was performed in 4% stacking and 16% separating gels as described (Lesse et al., 1990). Proteinase K-treated, boiled bacterial cells were used as samples (Apicella et al., 1994). The gels were run for 17 h at a constant current of 20 mA and silver stained (Tsai and Frasch, 1982).

### Whole-cell ELISA

Expression of CPS and LPS was tested by whole-cell ELISA (Abdillahi and Poolman, 1987). The group B CPS- and LPS-specific mAbs used were 101C11 (Hurpin et al., 1992) and Mn4A8B2 (Scholten et al., 1994), respectively.

### Determination of antibiotic sensitivity

Meningococcal strains were grown overnight on GC agar plates. Three colonies of each strain were resuspended in 200 µl of Meningococcal Medium and 175 µl of this solution were spread on fresh GC plates. Filter paper discs (Oxoid) containing rifampicin (5 µg), bacitracin (10 U), tetracycline (10 µg) or novobiocin (30 µg) were placed on the agar surface. After overnight incubation at 37°C, the halo of growth inhibition around each disc was measured.

### Acknowledgements

We gratefully acknowledge Dr M.Frosch (Universität Würzburg, Germany) for providing plasmid pMF32.35::T5 and anti-CtrA mAb 2619, and Dr Anthony Schryvers (University of Calgary, Canada) for providing the TbpB-specific rabbit antiserum.

### References

Abdillahi, H. and Poolman, J. (1987) Whole cell ELISA for typing *Neisseria meningitidis* with monoclonal antibodies. *FEMS Microbiol. Lett.*, **48**, 367–371.

Apicella, M.A., Griffiss, J.M. and Schneider, H. (1994) Isolation and characterization of lipopolysaccharides, lipooligosaccharides, and lipid A. *Methods Enzymol.*, **235**, 242–252.

Bligh, E.G. and Dyer, W.J. (1959) A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.*, **37**, 911–917.

Brok, R.G., Dekker, N., Gerrits, N., Verheij, H.M. and Tommassen, J. (1995) A conserved histidine residue of *Escherichia coli* outer-membrane phospholipase A is important for activity. *Eur. J. Biochem.*, **234**, 934–938.

Cole, M.J. and Enke, C.G. (1991) Direct determination of phospholipid structures in microorganisms by fast atom bombardment triple quadrupole mass spectrometry. *Anal. Chem.*, **63**, 1032–1038.

Coleman, J. (1992) Characterization of the *Escherichia coli* gene for 1-acyl-sn-glycerol-3-phosphate acyltransferase (*plsC*). *Mol. Gen. Genet.*, **232**, 295–303.

Cowan, S.W., Schirmer, T., Rummel, G., Steiert, M., Ghosh, R., Pauptit, R.A., Jansonius, J.N. and Rosenbusch, J.P. (1992) Crystal structures explain functional properties of two *E.coli* porins. *Nature*, **358**, 727–733.

de Cock, H. and Tommassen, J. (1996) Lipopolysaccharides and divalent cations are involved in the formation of an assembly-competent intermediate of outer-membrane protein PhoE of *E.coli*. *EMBO J.*, **15**, 5567–5573.

de Cock, H., Brandenburg, K., Wiese, A., Holst, O. and Seydel, U. (1999a) Non-lamellar structure and negative charges of lipopolysaccharides required for efficient folding of outer membrane protein PhoE of *Escherichia coli*. *J. Biol. Chem.*, **274**, 5114–5119.

de Cock, H., Schafer, U., Potgeter, M., Demel, R., Muller, M. and Tommassen, J. (1999b) Affinity of the periplasmic chaperone Skp of *Escherichia coli* for phospholipids, lipopolysaccharides and

non-native outer membrane proteins. Role of Skp in the biogenesis of outer membrane protein. *Eur. J. Biochem.*, **259**, 96–103.

Edwards, U., Muller, A., Hammerschmidt, S., Gerardy-Schahn, R. and Frosch, M. (1994) Molecular analysis of the biosynthesis pathway of the  $\alpha$ -2,8 polysialic acid capsule by *Neisseria meningitidis* serogroup B. *Mol. Microbiol.*, **14**, 141–149.

Frosch, M. and Muller, A. (1993) Phospholipid substitution of capsular polysaccharides and mechanisms of capsule formation in *Neisseria meningitidis*. *Mol. Microbiol.*, **8**, 483–493.

Frosch, M., Weisgerber, C. and Meyer, T.F. (1989) Molecular characterization and expression in *Escherichia coli* of the gene complex encoding the polysaccharide capsule of *Neisseria meningitidis* group B. *Proc. Natl Acad. Sci. USA*, **86**, 1669–1673.

Frosch, M., Muller, D., Bousset, K. and Muller, A. (1992) Conserved outer membrane protein of *Neisseria meningitidis* involved in capsule expression. *Infect. Immun.*, **60**, 798–803.

Fürste, J.P., Pansegrau, W., Frank, R., Blocker, H., Scholz, P., Bagdasarian, M. and Lanka, E. (1986) Molecular cloning of the plasmid RP4 primase region in a multi-host-range *tacP* expression vector. *Gene*, **48**, 119–131.

Gotschlich, E.C., Fraser, B.A., Nishimura, O., Robbins, J.B. and Liu, T.Y. (1981) Lipid on capsular polysaccharides of gram-negative bacteria. *J. Biol. Chem.*, **256**, 8915–8921.

Hurpin, C.M., Carosella, E.D. and Cazenave, P.A. (1992) Bactericidal activity of two IgG2a murine monoclonal antibodies with distinct fine specificities for group B *Neisseria meningitidis* capsular polysaccharide. *Hybridoma*, **11**, 677–687.

Jansen, C., Wiese, A., Reubsæet, L., Dekker, N., de Cock, H., Seydel, U. and Tommassen, J. (2000) Biochemical and biophysical characterization of *in vitro* folded outer membrane porin PorA of *Neisseria meningitidis*. *Biochim. Biophys. Acta*, **1464**, 284–298.

Kamio, Y. and Nikaido, H. (1976) Outer membrane of *Salmonella typhimurium*: accessibility of phospholipid headgroups to phospholipase C and cyanogen bromide activated dextran in the external medium. *Biochemistry*, **15**, 2561–2570.

Kasper, D.L., Winkelhake, J.L., Zollinger, W.D., Brandt, B.L. and Arstenstein, M.S. (1973) Immunochemical similarity between polysaccharide antigens of *Escherichia coli* 07: K1(L):NM and group B *Neisseria meningitidis*. *J. Immunol.*, **110**, 262–268.

Klugman, K.P., Gotschlich, E.C. and Blake, M.S. (1989) Sequence of the structural gene (*rmpM*) for the class 4 outer membrane protein of *Neisseria meningitidis*, homology of the protein to gonococcal protein III and *Escherichia coli* OmpA, and construction of meningococcal strains that lack class 4 protein. *Infect. Immun.*, **57**, 2066–2071.

Koplow, J. and Goldfine, H. (1974) Alterations in the outer membrane of the cell envelope of heptose-deficient mutants of *Escherichia coli*. *J. Bacteriol.*, **117**, 527–543.

Laird, M.W., Kloser, A.W. and Misra, R. (1994) Assembly of LamB and OmpF in deep rough lipopolysaccharide mutants of *Escherichia coli* K-12. *J. Bacteriol.*, **176**, 2259–2264.

Legrain, M., Mazarin, V., Irwin, S.W., Bouchon, B., Quentin-Millet, M.J., Jacobs, E. and Schryvers, A.B. (1993) Cloning and characterization of *Neisseria meningitidis* genes encoding the transferrin-binding proteins Tbp1 and Tbp2. *Gene*, **130**, 73–80.

Lesse, A.J., Campagnari, A.A., Bittner, W.E. and Apicella, M.A. (1990) Increased resolution of lipopolysaccharides and lipooligosaccharides utilizing tricaine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *J. Immunol. Methods*, **126**, 109–117.

Lugtenberg, B., Meijers, J., Peters, R., van der Hoek, P. and van Alphen, L. (1975) Electrophoretic resolution of the 'major outer membrane protein' of *Escherichia coli* K12 into four bands. *FEBS Lett.*, **58**, 254–258.

Lysko, P.G. and Morse, S.A. (1981) *Neisseria gonorrhoeae* cell envelope: permeability to hydrophobic molecules. *J. Bacteriol.*, **145**, 946–952.

Masson, L. and Holbein, B.E. (1983) Physiology of sialic acid capsular polysaccharide synthesis in serogroup B *Neisseria meningitidis*. *J. Bacteriol.*, **154**, 728–736.

Mohan, S., Kelly, T.M., Eveland, S.S., Raetz, C.R. and Anderson, M.S. (1994) An *Escherichia coli* gene (*fabZ*) encoding (3R)-hydroxymyristoyl acyl carrier protein dehydrase. Relation to *fabA* and suppression of mutations in lipid A biosynthesis. *J. Biol. Chem.*, **269**, 32896–32903.

Osborn, M.J., Gander, J.E., Parisi, E. and Carson, J. (1972) Mechanism of assembly of the outer membrane of *Salmonella typhimurium*. Isolation and characterization of cytoplasmic and outer membrane. *J. Biol. Chem.*, **247**, 3962–3972.



- Pautsch,A. and Schulz,G.E. (1998) Structure of the outer membrane protein A transmembrane domain. *Nature Struct. Biol.*, **5**, 1013–1017.
- Petersson,A., Maas,A. and Tommassen,J. (1994) Identification of the *iroA* gene product of *Neisseria meningitidis* as a lactoferrin receptor. *J. Bacteriol.*, **176**, 1764–1766.
- Petersson,A., Prinz,T., Umar,A., van der Biezen,J. and Tommassen,J. (1998) Molecular characterization of LbpB, the second lactoferrin-binding protein of *Neisseria meningitidis*. *Mol. Microbiol.*, **27**, 599–610.
- Poolman,J.T., Wientjes,F.B., Hopman,C.T.P. and Zanen,H.C. (1985) Influence of the length of lipopolysaccharide molecules on the surface exposure of class 1 and 2 outer membrane proteins of *Neisseria meningitidis* strain 2996 (B:2b:P1.2). In Schoolnik,G.K. (ed.), *The Pathogenic Neisseria*. American Society for Microbiology, Washington, DC, pp. 562–570.
- Prinz,T. and Tommassen,J. (2000) Association of iron-regulated outer membrane proteins of *Neisseria meningitidis* with the RmpM (class 4) protein. *FEMS Microbiol. Lett.*, **183**, 49–53.
- Prinz,T., Meyer,M., Petersson,A. and Tommassen,J. (1999) Structural characterization of the lactoferrin receptor from *Neisseria meningitidis*. *J. Bacteriol.*, **181**, 4417–4419.
- Raetz,C.R. (1990) Biochemistry of endotoxins. *Annu. Rev. Biochem.*, **59**, 129–170.
- Rahman,M.M., Kolli,V.S., Kahler,C.M., Shih,G., Stephens,D.S. and Carlson,R.W. (2000) The membrane phospholipids of *Neisseria meningitidis* and *Neisseria gonorrhoeae* as characterized by fast atom bombardment mass spectrometry. *Microbiology*, **146**, 1901–1911.
- Ried,G., Hindennach,I. and Henning,U. (1990) Role of lipopolysaccharide in assembly of *Escherichia coli* outer membrane proteins OmpA, OmpC, and OmpF. *J. Bacteriol.*, **172**, 6048–6053.
- Sambrook,J., Fritsch,E.F. and Maniatis,T. (1989) *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Schindler,M. and Osborn,M.J. (1979) Interaction of divalent cations and polymyxin B with lipopolysaccharide. *Biochemistry*, **18**, 4425–4430.
- Scholten,R.J., Bijlmer,H.A., Poolman,J.T., Kuipers,B., Caugant,D.A., van Alphen,L., Dankert,J. and Valkenburg,H.A. (1993) Meningococcal disease in The Netherlands, 1958–1990: a steady increase in the incidence since 1982 partially caused by new serotypes and subtypes of *Neisseria meningitidis*. *Clin. Infect. Dis.*, **16**, 237–246.
- Scholten,R.J., Kuipers,B., Valkenburg,H.A., Dankert,J., Zollinger,W.D. and Poolman,J.T. (1994) Lipo-oligosaccharide immunotyping of *Neisseria meningitidis* by a whole-cell ELISA with monoclonal antibodies. *J. Med. Microbiol.*, **41**, 236–243.
- Schryvers,A.B. and Morris,L.J. (1988) Identification and characterization of the transferrin receptor from *Neisseria meningitidis*. *Mol. Microbiol.*, **2**, 281–288.
- Sen,K. and Nikaïdo,H. (1991) Lipopolysaccharide structure required for *in vitro* trimerization of *Escherichia coli* OmpF porin. *J. Bacteriol.*, **173**, 926–928.
- Shih,G.C., Kahler,C.M., Swartley,J.S., Rahman,M.M., Coleman,J., Carlson,R.W. and Stephens,D.S. (1998) Multiple lysophosphatidic acid acyltransferases in *Neisseria meningitidis*. *Mol. Microbiol.*, **32**, 874–880.
- Sprong,T., Stikkelbroeck,N., van der Ley,P., Steeghs,L., van Alphen,L., Klein,N., Netea,M.G., van der Meer,J.W.M. and van Deuren,M. (2001) The relative contribution of lipopolysaccharide and non-LPS components of *Neisseria meningitidis* to the proinflammatory cytokine response. *J. Leukoc. Biol.*, **70**, 283–288.
- Steeghs,L., Jennings,M.P., Poolman,J.T. and van der Ley,P. (1997) Isolation and characterization of the *Neisseria meningitidis* *lpxD-fabZ-lpxA* gene cluster involved in lipid A biosynthesis. *Gene*, **190**, 263–270.
- Steeghs,L., den Hartog,R., den Boer,A., Zomer,B., Roholl,P. and van der Ley,P. (1998) Meningitis bacterium is viable without endotoxin. *Nature*, **392**, 449–450.
- Steeghs,L., Kuipers,B., Hamstra,H.J., Kersten,G., van Alphen,L. and van der Ley,P. (1999) Immunogenicity of outer membrane proteins in a lipopolysaccharide-deficient mutant of *Neisseria meningitidis*: influence of adjuvants on the immune response. *Infect. Immun.*, **67**, 4988–4993.
- Tettelin,H. *et al.* (2000) Complete genome sequence of *Neisseria meningitidis* serogroup B strain MC58. *Science*, **287**, 1809–1815.
- Tsai,C.M. and Frasch,C.E. (1982) A sensitive silver stain for detecting lipopolysaccharides in polyacrylamide gels. *Anal. Biochem.*, **119**, 115–119.
- van der Ley,P. and Poolman,J.T. (1992) Construction of a multivalent meningococcal vaccine strain based on the class 1 outer membrane protein. *Infect. Immun.*, **60**, 3156–3161.
- van der Ley,P., van der Biezen,J., Hohenstein,P., Peeters,C. and Poolman,J.T. (1993) Use of transformation to construct antigenic hybrids of the class 1 outer membrane protein in *Neisseria meningitidis*. *Infect. Immun.*, **61**, 4217–4224.
- Vieira,J. and Messing,J. (1982) The pUC plasmids, an M13mp7-derived system for insertion mutagenesis and sequencing with synthetic universal primers. *Gene*, **19**, 259–268.
- Weiss,M.S., Abele,U., Weckesser,J., Welte,W., Schiltz,E. and Schulz,G.E. (1991) Molecular architecture and electrostatic properties of a bacterial porin. *Science*, **254**, 1627–1630.
- Wiese,A., Reiners,J.O., Brandenburg,K., Kawahara,K., Zahringer,U. and Seydel,U. (1996) Planar asymmetric lipid bilayers of glycosphingolipid or lipopolysaccharide on one side and phospholipids on the other: membrane potential, porin function, and complement activation. *Biophys. J.*, **70**, 321–329.
- Wilkinson,S.G. (1988) Gram-negative bacteria. In Ratledge,C. and Wilkinson,S.G. (eds), *Microbial Lipids*. Academic Press, London, UK, pp. 299–488.
- Yakushi,T., Masuda,K., Narita,S., Matsuyama,S. and Tokuda,H. (2000) A new ABC transporter mediating the detachment of lipid-modified proteins from membranes. *Nature Cell Biol.*, **2**, 212–218.
- Yokota,N., Kuroda,T., Matsuyama,S. and Tokuda,H. (1999) Characterization of the LolA–LolB system as the general lipoprotein localization mechanism of *Escherichia coli*. *J. Biol. Chem.*, **274**, 30995–30999.

Received February 19, 2001; revised October 22, 2001;  
accepted November 1, 2001