

# Ovarian and pituitary function in dogs after hysterectomy

B. Hoffmann<sup>1</sup>, R. Höveler<sup>1</sup>, S. H. Hasan<sup>2</sup> and K. Failing<sup>3</sup>

<sup>1</sup>*Ambulatorische und Geburtshilfliche Veterinärklinik Justus-Liebig-Universität Gießen, 6300 Gießen, Germany;* <sup>2</sup>*Schering AG, Müllerstraße 170–178, 1000 Berlin, Germany;* and <sup>3</sup>*Institut für Veterinär-Physiologie, Arbeitsgruppe Biomathematik, Justus-Liebig-Universität Gießen, 6300 Gießen, Germany*

**Summary.** In studies of five hysterectomized and five control dogs, hysterectomy shortened the anoestrous interval ( $96.6 \pm 28.0$  versus  $149.4 \pm 50.9$  days,  $P < 0.05$ ). No differences in hormone concentrations (progesterone, oestradiol, prolactin and growth hormone) were observed between the control and hysterectomized dogs except for a brief fall in progesterone concentrations over 8 days immediately after surgery, between days 35 and 40 after onset of pro-oestrous bleeding; only these animals developed symptoms of overt pseudopregnancy. It is concluded that, in dogs, luteal regression occurs independently of a uterine luteolysin, but that the uterus may play a role in control of duration of anoestrus. Pseudopregnancy seems to be initiated by a fall in progesterone concentrations rather than by other hormonal changes.

**Keywords:** dog; hysterectomy; ovary; growth hormone; prolactin

## Introduction

Domestic dogs are monoestrous and predominantly aseasonal breeders, with several months of anoestrus between active reproductive phases (Stabenfeldt & Shille, 1977; Siegel, 1982). In contrast to other domestic animal species, luteal function is almost identical in pregnant and nonpregnant females, except that pregnant animals attain baseline concentrations of progesterone earlier owing to the immediate prepartum decline of this hormone (Concannon *et al.*, 1978). The development of lactatio falsa (overt pseudopregnancy) in many nonpregnant bitches during the latter half of dioestrus might be related to this extended luteal lifespan (Shille & Stabenfeldt, 1980; Concannon, 1987; Gerres *et al.*, 1988). Endocrine control of gonadal, in particular luteal, function in dogs is poorly understood. The decline of plasma progesterone concentrations after hypophysectomy during the early or midluteal phase (Okkens *et al.*, 1986) demonstrates the necessity of pituitary support. Both luteinizing hormone (LH) and prolactin must be considered luteotrophic. During the later luteal phase or late pregnancy, inhibition of prolactin by treatment with the dopamine agonist CB-154 significantly reduced progesterone concentrations (by >90%) or induced interruption of pregnancy (Conley & Evans, 1984; Concannon *et al.*, 1987). Injection of anti-LH-antiserum significantly reduced progesterone concentrations of plasma (by 76–93%) in pregnant bitches (Concannon *et al.*, 1987), but similar treatments during the early luteal phase induced only a temporary and less distinct decline of progesterone concentrations (Conley & Evans, 1984; Concannon *et al.*, 1987), showing that the luteotrophic activity of LH and prolactin is related to the reproductive stage. The observation that prostaglandin  $F_{2\alpha}$  of endometrial origin induces luteal regression in most domestic animals (Anderson *et al.*, 1969) led to studies in dogs. Both Hadley (1975) and Baker *et al.* (1980) observed a shortened luteal lifespan after hysterectomy and a luteotrophic factor of uterine origin was postulated by Hadley (1975). Chaffaux & Thibier (1978) observed a decline in progesterone concentrations after hysterectomy in normal pregnant dogs, but not in dogs with pyometra or other pathological uterine contents and concluded that there was a

luteotrophic factor of fetoplacental origin. Olson *et al.* (1984), however, observed no changes in luteal function after hysterectomy of pregnant and nonpregnant dogs, whereas Okkens *et al.* (1985) reported a reduction of progesterone concentrations without a shortened luteal lifespan. Jöchle *et al.* (1973) could not interrupt pregnancy after treatments with prostaglandin  $F_{2\alpha}$ , but observed only a temporary decrease in progesterone concentrations. Repeated treatments, however, induced abortions in some dogs at late stages of pregnancy; in the nonaborting animals, progesterone concentrations were reduced only temporarily (Concannon & Hansel, 1977). These data are conflicting and allow only limited conclusions on whether and to what extent prostaglandin  $F_{2\alpha}$  of uterine origin acts as a luteolytic agent in dogs. In the present experiments, we attempted to define maintenance and resumption of gonadal function in intact and hysterectomized dogs by observing the animals over a long period and by monitoring progesterone and oestradiol concentrations. After determining prolactin in selected samples, it was hoped to gain related information on the availability of this gonadotrophic factor. Since pseudopregnancy is an inherent part of canine reproduction, possibly also resulting from interactions between luteal function and growth hormone (El Etreby, 1979), growth hormone was also determined in a series of samples.

## Materials and Methods

### Animals and experimental design

The experiments were performed over 2 years. Healthy bitches with proven fertility (3–6 years old; bodyweight 27–35 kg) were removed successively from a pack of English Foxhounds and housed in groups of up to four in a kennel, with about 100 m<sup>2</sup> outdoor space. A standard diet was provided with water *ad libitum*. Sexual behaviour, external genitalia and mammary gland development were examined once or twice a day. In addition to the developmental stage tested by inspection and palpation, the secretory activity of the mammary glands was recorded (colour and type of secreted fluid, spontaneous release or in response to applied pressure). Vaginal cytology and introduction of a male dog served for confirmatory purposes. Five control dogs were monitored through two complete reproductive periods (cycles I and II) with the exception of one animal, which had to be removed from the experiment during cycle II. In four other animals, hysterectomy was performed in cycle I between days 35 and 40 (day  $37.7 \pm 2.2$ ) after onset of pro-oestrous bleeding and, in another dog, added later to the experiment, at the end of the luteal phase (progesterone concentration  $16.3 \text{ nmol l}^{-1}$ ); in all five dogs, a complete cycle (cycle II) was observed after hysterectomy.

Onset of pro-oestrous bleeding was considered as day 0. Blood samples were taken at intervals of 2 days during pro-oestrus, oestrus, metoestrus, dioestrus and during a period of about 10 days before pro-oestrus (terminology according to Stabenfeldt & Shille, 1977); otherwise, sampling was at intervals of 4 days during anoestrus, which was defined as the period when progesterone concentrations were  $< 3.2 \text{ nmol l}^{-1}$  ( $= 1 \text{ ng ml}^{-1}$ ). Progesterone and oestradiol were determined in all samples collected; prolactin and growth hormone were assayed in six cycles of the control group and in three cycles during and after hysterectomy.


### Hormone assay

All hormones were determined by radioimmunoassay.

**Progesterone.** The method followed the procedure outlined by Hoffmann *et al.* (1973). Plasma (0.1 ml) was extracted with hexane, the antiserum used was directed against 4-pregnene-11 $\alpha$ -ol-3,20-dione-HS-bovine serum albumin (BSA) (final dilution 1:75 000,  $K_a = 3.42 \times 10^9 \text{ mol l}^{-1}$ ). Intra- and interassay coefficients of variation were 14 and 15%, respectively, the detection limit was at  $0.19 \text{ nmol l}^{-1}$ .

**Oestradiol.** Initially, for the determination of oestradiol, high-performance liquid chromatography (HPLC) was applied before radioimmunoassay. After solid-phase extraction of 0.5 ml plasma (Extrelut columns (Merck, Darmstadt, Germany), elution with 6.0 ml tertiary-butyl-methyl-ether) and evaporation to dryness, the samples were redissolved in 0.6 ml of 35% acetonitrile in water and an aliquot of 0.5 ml was injected into the HPLC system (reversed-phase HPLC; column: Lichrosorb-RP 18 (Merck, Darmstadt), isocratic (40% acetonitrile in water) and isothermal (40°C) conditions, flow  $0.4 \text{ ml min}^{-1}$ ). As established by runs with [<sup>3</sup>H]-standards, the fraction containing the oestradiol (retention 24 min) was collected in radioimmunoassay glass tubes and evaporated to dryness, and 0.1 ml BSA-phosphate buffer was added.

The radioimmunoassay was set up as a sequential assay (Strecker *et al.*, 1979). After the addition of 0.4 ml anti-serum dilution (antiserum Gi-E<sub>2</sub>-KI, obtained after immunization of rabbits with oestradiol-17 $\beta$ -6CMO-BSA; final dilution 1:20 000;  $K_a = 11.79 \times 10^9 \text{ mol l}^{-1}$ , crossreactions: oestrone, 1.30%; oestriol, 0.68%; all other tested non-phenolic steroids  $< 0.01\%$ ) to the samples and standards (0.74 fmol to 0.11 pmol per tube in 0.1 ml BSA-phosphate



**Fig. 1.** Mean values of progesterone concentration ( $\bar{x}_G \pm DF^{\pm 1}$ ) in peripheral plasma of (a) control dogs (number of cycles,  $n = 9$ ) and during (b) cycle I ( $n = 4$ ) and (c) cycle II ( $n = 5$ ) of hysterectomized bitches.

buffer), a first 'overnight' incubation (minimum 12 h) at 4°C was performed. [ $^3\text{H}$ ]oestradiol (0.132 pmol = 0.444 KBq per tube) was then added and the samples were again incubated for 45 min at 4°C (second incubation). After separation of free and bound steroid by addition of 0.2 ml of ice-cold charcoal suspension (0.5% charcoal, 0.05% dextrane) and centrifugation for 15 min at 2400  $g$  and 4°C, 0.6 ml of the supernatant was removed to measure the radioactivity of the bound fraction. Absolute binding of [ $^3\text{H}$ ]oestradiol was 15–20%. As determined from plasma samples, mean recovery of 14.7 ( $n = 8$ ), 29.4 ( $n = 8$ ), 58.9 ( $n = 6$ ) and 117.8 ( $n = 7$ ) pmol oestradiol  $\text{l}^{-1}$  was at 77%, the mean intra- and interassay coefficients of variations were 13 and 22%, respectively. The lower limit of sensitivity was at 12.6 pmol  $\text{l}^{-1}$ . During the course of the experiments, sample purification by HPLC was omitted, since virtually identical results were obtained with and without chromatography. Instead, before radioimmunoassay, 0.25 ml plasma was extracted twice with 3.0 ml toluene.

**Prolactin.** Prolactin was measured by the method of Gräf *et al.* (1977): 300  $\mu\text{l}$  of plasma in duplicate and 100  $\mu\text{l}$  of standard were incubated with 200  $\mu\text{l}$  tracer (9000 c.p.m.) for 24 h at room temperature. The antibody-bound labelled prolactin was precipitated by adding 200  $\mu\text{l}$  of anti-rabbit gamma globulin in sheep (Schering, Berlin) in a dilution of 1:60, which provided maximum separation and, after further incubation for 24 h, the resulting complex was then centrifuged at 3000  $g$  for 20 min at 4°C. The supernatant was discarded and the amount of precipitate was calculated. The minimum detectable concentration in the assay of these samples was 0.25 ng per tube. The interassay coefficient of variation was 22%.

**Growth hormone.** Growth hormone was measured essentially by the method of Cocola *et al.* (1976): 400  $\mu\text{l}$  of plasma in duplicate and 100  $\mu\text{l}$  of standard were incubated with 200  $\mu\text{l}$  of antibody of appropriate dilution and 200  $\mu\text{l}$  tracer (14 000 c.p.m.) for 48 h at room temperature. The antibody-bound labelled growth hormone was precipitated by adding 200  $\mu\text{l}$  of anti-monkey gamma globulin in goat (Paesel, Frankfurt) at a dilution of 1:10 which provided


**Table 1.** Duration of luteal phase\* and area under the curve (nmol day<sup>-1</sup> l<sup>-1</sup>) for peripheral progesterone for cycles I and II of control and hysterectomized dogs

Group	No. of dog	Cycle I		Cycle II	
		Luteal phase (days)	Area under curve	Luteal phase (days)	Area under curve
Control	101†	58	3165.4	—	—
	102	56	2956.4	59	2614.9
	103	60	2630.2	60	3104.6
	104	73	3418.2	75	5596.2
	105	102	3474.2	70	3323.7
	$\bar{x}_G$	68.0	3112.9	65.7	3505.5
Hysterectomized	DF	1.28	1.11	1.12	1.39
	301	68	2826.1	86	3887.9
	302‡	—	—	80	4282.5
	303	64	2031.7	70	2306.8
	304	68	3474.8	58	4686.0
	305	62	3132.3	64	4092.0
	$\bar{x}_G$	65.5	2800.6	70.9	3744.8
	DF	1.05	1.27	1.17	1.32

\*Period with progesterone > 3.2 nmol l<sup>-1</sup> (= 1.0 ng ml<sup>-1</sup>).


†Had to be removed during cycle II.

‡Surgery during late luteal phase.



**Fig. 2.** Mean values of progesterone concentration ( $\bar{x}_G \pm \text{DF}$ ) 10 days before and after hysterectomy during midcycle (days 35–40) (—■—,  $n = 4$ ) and in control dogs (—◆—,  $n = 9$ ) during the respective periods.

maximum separation and, after further incubation for 24 h, the resulting complex was centrifuged at 3000 *g* for 20 min at 4°C. The supernatant was discarded and the amount of precipitate was calculated. The minimum detectable concentration in the assay of these samples was 0.25 ng per tube. The interassay coefficient of variation was 18.5%.




**Fig. 3.** Mean values of oestradiol concentration ( $\bar{x}_G \cdot DF^{\pm 1}$ ) in peripheral plasma of (a) control dogs (number of cycles,  $n = 9$ ) and during (b) cycle I ( $n = 4$ ) and (c) cycle II ( $n = 5$ ) of hysterectomized bitches.

### Statistical evaluation

To cope with the asymmetrical distribution of hormone plasma values, which are often positively skewed, the geometric means ( $\bar{x}_G$ ) and dispersion factors (DF) were calculated (Sachs, 1984). Results were expressed as  $\bar{x}_G \cdot DF^{\pm 1}$ . Arithmetic means were calculated for the anoestrous intervals. A two-factorial analysis of variance with repeated measures on the factor time was used to test for time and group differences with respect to the experimental design. To describe availability of hormones over time as an indicator of hormone production, the area under the curve was calculated (in  $\text{nmol day}^{-1} \text{l}^{-1}$ ) (Dost, 1968). All statistical evaluations were done with the BMDP statistical software (Dixon, 1987).

### Results

No significant differences in the course of progesterone concentrations were evident between cycles (Fig. 1). Table 1 gives the individual data and the mean values for the duration of the luteal phase (progesterone  $> 3.2 \text{ nmol l}^{-1}$ ) and the respective area under the curve. However, there was a drop in progesterone concentrations over 8 days in the four animals hysterectomized between days 35 and 40 (Fig. 2); these dogs, but not the control animals, developed symptoms of overt



**Fig. 4.** Mean values of prolactin concentration ( $\bar{x}_G \pm DF^{\pm 1}$ ) in peripheral plasma of (a) control dogs (number of cycles,  $n = 6$ ) and during (b) cycle I ( $n = 3$ ) and (c) cycle II ( $n = 3$ ) of hysterectomized bitches.


pseudopregnancy, as was indicated by enlargement of mammary glands and secretion on pressure of the caudal mammary complexes.

No differences in concentrations of oestradiol, prolactin and growth hormone were found between control and hysterectomized animals or between cycles I and II in the hysterectomized dogs (Figs 3, 4 and 5).

Whereas progesterone, oestradiol and prolactin concentrations significantly changed over time ( $P < 0.0001$ ), no such effects could be established for growth hormone. The average anoestrous interval between cycle I and cycle II was significantly ( $P < 0.05$ ) longer in the control ( $149.4 \pm 50.9$  days) than in the hysterectomized dogs ( $96.6 \pm 28.0$  days) (Table 2). This was tested by means of the two-sample  $t$  test, not assuming equal variances.

## Discussion

There were no differences between the control group and the hysterectomized animals for any of the hormones determined (progesterone, oestradiol, prolactin and growth hormone). There was also no difference between the cycle when hysterectomy was performed and the next cycle. In



**Fig. 5.** Mean values of growth hormone concentration ( $\bar{x}_G \pm \text{DF}^{\pm 1}$ ) in peripheral plasma of control dogs (number of cycles,  $n = 6$ ) and during (b) cycle I ( $n = 3$ ) and (c) cycle II ( $n = 3$ ) of hysterectomized bitches.

**Table 2.** Onset and duration of anoestrus\* in control and hysterectomized dogs

Group	No. of dog	Anoestrus		$\bar{x} \pm \text{SD}$	Difference
		Onset	Length (days)		
Control	102	17 July	238	$149.4 \pm 50.9$	
	103	5 February	129		
	104	19 August	116		
	105	7 June	118		
	106	14 June	146		
Hysterectomized	301	27 July	79	$96.6 \pm 28.0$	$P < 0.05^\dagger$
	302	28 April	115		
	303	9 August	112		
	304	19 May	121		
	305	21 February	56		

\*Length between cycles with progesterone  $< 3.2 \text{ nmol l}^{-1}$  ( $= 1.0 \text{ ng ml}^{-1}$ ).

$^\dagger$ Compared with the control group.

agreement with previous observations (Olson *et al.*, 1984; Okkens *et al.*, 1985), these observations provide further evidence that in nonpregnant dogs luteal regression does not depend on a uterine luteolysin and that normal reproductive cycles follow hysterectomy. However, the uterus may have an influence on the length of the period of anoestrus, which was significantly shortened after hysterectomy (97 versus 149 days). Clearly, in view of the small number of dogs investigated, this observation needs further confirmation. Thus, high variability has been reported for the length of anoestrus in dogs (Christie & Bell, 1971; Concannon, 1987) and the season of the year may play a role, though in both groups of animals anoestrus commenced between February and August. Hysterectomy led to an immediate decrease of progesterone by about 50% compared with the control animals; after 8 days, progesterone concentrations in the control group and hysterectomized animals were the same. This temporary decrease of progesterone, which did not affect the general area under the curve established for the whole cycle, was associated with the appearance of symptoms of overt pseudopregnancy. Similar observations have been made by Okkens *et al.* (1985) and would support the conclusions of Gerres *et al.* (1988) (they could provoke onset of pseudopregnancy in dogs after treatment with the antigestagen RU 38486) that removal of progesterone is one of the triggers for overt pseudopregnancy. Since there were no differences between the cycles of the control animals and the second cycle of the hysterectomized animals, this temporary reduction in progesterone cannot be interpreted as a removal of a uterine luteotrophic factor (Hadley, 1975), but, rather, as a consequence of surgery, possibly by interfering with ovarian blood supply.

In contrast to progesterone, oestradiol and prolactin, no changes in relation to stage of the cycle could be established for growth hormone. Although only a few animals were investigated, this observation seems to indicate that growth hormone has little importance, if any, in the development of pseudopregnancy. Thus the increases in growth hormone concentrations and activity of growth hormone cells in the pituitary gland after treatment of dogs with progestagens (Gräf & El Etreby, 1979) must be interpreted carefully in respect of the mechanism underlying mammary gland development and onset of pseudopregnancy. In addition, the prolactin concentrations were not different between cycles; although the sampling intervals of 2 days allow only conclusions with respect to baseline concentrations, the lack of any shift does not point towards the role of this hormone in the development and initiation of pseudopregnancy.

The present data yield further evidence that in dogs luteal regression occurs independently of a uterine luteolysin. However, the uterus may play an active role in controlling the length of anoestrus. No evidence has been obtained that prolactin and growth hormone are associated with the initiation of overt pseudopregnancy, which seems to be a result of the decrease in progesterone concentrations during the second half of dioestrus.

With support of the German Research Foundation (DFG).

## References

- Anderson, L.L., Bland, K.P. & Melampy, R.M. (1969) Comparative aspects of uterine-luteal relationships. *Recent Progress in Hormone Research* **25**, 57–65.
- Baker, B.A., Archbald, L.F., Clooney, L.L., Lotz, K. & Godke, R.A. (1980) Luteal function in the hysterectomized bitch following treatment with prostaglandin F<sub>2a</sub> (PGF<sub>2a</sub>). *Theriogenology* **14**, 195–205.
- Chaffaux, S. & Thibier, M. (1978) Effets de l'hystérectomie sur la concentration de progesterone plasmatique chez la chienne. *Recueil de Médecine Vétérinaires* **154**, 933–938.
- Christie, D.W. & Bell, E.T. (1971) Some observations on the seasonal incidence and frequency of oestrus in breeding bitches in Britain. *Journal of Small Animal Practice* **12**, 159–167.
- Cocola, F., Udeschini, G., Secchi, C., Panerai, A.E., Neri, P. & Müller, E.E. (1976) A rapid radioimmunoassay method of growth hormone in dog plasma. *Proceedings of the Society for Experimental Biology and Medicine* **151**, 140–145.
- Concannon, P. (1987) The physiology of ovarian cycles, pregnancy and parturition in the domestic dog. In *Proceedings of the Annual Meeting of the Society for Theriogenology and The American College of Theriogenologists*, pp. 159–197. Society for Theriogenology, Hastings, NE, USA.



- Concannon, P. & Hansel, W. (1977) Prostaglandin  $F_{2\alpha}$ -induced luteolysis, hypothermia, and abortions in Beagle bitches. *Prostaglandins* **13**, 533–542.
- Concannon, P., Butler, W.R., Hansel, W., Knight, P.J. & Hamilton, J.M. (1978) Parturition and lactation in the bitch: serum progesterone, cortisol and prolactin. *Biology of Reproduction* **19**, 1113–1118.
- Concannon, P., Weinstein, P., Whaley, S. & Frank, D. (1987) Suppression of luteal function in dogs by luteinizing hormone antiserum and by bromocryptine. *Journal of Reproduction and Fertility* **81**, 175–180.
- Conley, A. & Evans, L. (1984) Bromergocryptine induced abortion in the bitch. *Proceedings of the 10th International Congress on Animal Reproduction and Artificial Insemination*, University College, Dublin, Ireland, Vol. 1, p. 504.
- Dixon, W.J. (1987) *BMDP Statistical Software Manual*. University of California Press, Berkeley, pp. 359–387.
- Dost, F.H. (1968) *Grundlagen der Pharmakologie*. Georg Thieme, Stuttgart, pp. 267–271.
- El Etreby, M.F. (1979) Effect of cyproterone acetate, levonorgestrel and progesterone on adrenal glands and reproductive organs in the Beagle bitch. *Cell and Tissue Research* **200**, 229–243.
- Gerres, S., Höveler, R., Evers, P. & Hoffmann, B. (1988) Investigations on the role of progesterone (P4) in the endocrine control of overt pseudopregnancy in the bitch, application of an antigestagen. *Proceedings of the 11th International Congress on Animal Reproduction and Artificial Insemination*, University of Illinois at Urbana, Champaign, USA, Vol. 2, p. 90.
- Gräf, K.J. & El Etreby, M.F. (1979) Endocrinology of reproduction in the female dog and its significance in mammary gland tumorigenesis *Acta Endocrinologica* **90**, Supplement 222, pp. 1–34.
- Gräf, K.J., Friedrich, E., Matthes, S. & Hasan, S.H. (1977) Homologous radioimmunoassay for canine prolactin and its application in various physiological states. *Journal of Endocrinology* **75**, 93–103.
- Hadley, J.C. (1975) The effects of serial uterine biopsies and hysterectomy on peripheral blood levels of total unconjugated oestrogen and progesterone in the bitch. *Journal of Reproduction and Fertility* **44**, 389–393.
- Hoffmann, B., Kyrein, H.J. & Ender, M.L. (1973) An efficient procedure for the determination of progesterone by radioimmunoassay applied to bovine peripheral plasma. *Hormone Research* **4**, 302–310.
- Jöchle, W., Tomlinson, R.V. & Andersen, A.C. (1973) Prostaglandin effects on plasma progesterone levels in the pregnant and cycling dog (Beagle). *Prostaglandins* **3**, 209–217.
- Okkens, A.C., Dieleman, S.J., Bevers, M.M. & Willemse, A.H. (1985) Evidence for the non-involvement of the uterus in the lifespan of the corpus luteum in the cyclic dog. *Veterinary Quarterly* **7**, 169–173.
- Okkens, A.C., Dieleman, S.J., Bevers, M.M., Lubberink, A.A.M.E. & Willemse, A.H. (1986) Influence of hypophysectomy on the lifespan of the corpus luteum in the cyclic dog. *Journal of Reproduction and Fertility* **77**, 187–192.
- Olson, P.N., Bowen, R.A., Behrendt, M.D., Olson, J.D. & Nett, T.M. (1984) Concentrations of progesterone and luteinizing hormone in the serum of diestrus bitches before and after hysterectomy. *American Journal of Veterinary Research* **45**, 149–153.
- Sachs, L. (1984) *Angewandte Statistik*. Springer Verlag, Berlin, pp. 70–89.
- Shille, V.M. & Stabenfeldt, G.H. (1980) Current concepts in reproduction of the dog and cat. *Advances in Veterinary Science and Comparative Medicine* **24**, 211–230.
- Siegel, E.T. (1982) *Endokrine Krankheiten des Hundes*. Paul Parey Verlag, Berlin, pp. 179–187.
- Stabenfeldt, G.H. & Shille, V.M. (1977) Reproduction in the dog and cat. In *Reproduction in Domestic Animals*, 3rd edn, pp. 499–527. Eds H. H. Cole & P. T. Cupps. Academic Press, New York.
- Strecker, H., Hachmann, H. & Seidel, L. (1979) Der Radioimmunotest (RIA), eine hochspezifische extrem empfindliche quantitative Analysenmethode. *Chemiker Zeitung* **103**, 53–68.

Received 12 December 1991