

Ovarian antral folliculogenesis during the human menstrual cycle: a review

Angela R. Baerwald^{1,*}, Gregg P. Adams², and Roger A. Pierson³

¹Department of Obstetrics, Gynecology and Reproductive Sciences, Royal University Hospital, University of Saskatchewan, Saskatoon, SK, Canada S7N 0W8 ²Department of Veterinary Biomedical Sciences, Western College of Veterinary Medicine, University of Saskatchewan, 52 Campus Drive, Saskatoon, SK, Canada S7N 5B4 ³Department of Obstetrics, Gynecology and Reproductive Sciences, Royal University Hospital, University of Saskatchewan, 103 Hospital Drive, Saskatoon, SK, Canada S7N 0W8

*Correspondence address. Tel: +1-306-966-8073; Fax: +1-306-966-8040; E-mail: angela.baerwald@usask.ca

Submitted on February 4, 2011; resubmitted on August 15, 2011; accepted on August 24, 2011

TABLE OF CONTENTS

- Introduction
- Methods
- Setting the stage: pre-antral and early antral follicle development
- Terminology used in describing folliculogenesis
- Antral follicle recruitment
- Follicle selection
- Pre-ovulatory follicle development
- Discussion

BACKGROUND: Ovarian follicles undergo dynamic morphologic and endocrinologic changes during the human menstrual cycle. The physiologic mechanisms underlying recruitment and selection of antral follicles in women are not fully elucidated.

METHODS: A comprehensive review of >200 studies was conducted using PubMed. The objective was to compare and contrast different perspectives on human antral folliculogenesis.

RESULTS: Antral folliculogenesis has been studied using histologic, endocrinologic and/or ultrasonographic techniques. Different theories of antral follicle recruitment include: (i) continuous recruitment throughout the menstrual cycle; (ii) recruitment of a 'cohort' of antral follicles once in the late-luteal phase or early-follicular phase of each cycle and (iii) recruitment of two or three 'cohorts' or 'waves' during each cycle. Generally, a single dominant follicle is selected in the mid-follicular phase of each cycle and this follicle ovulates at mid-cycle. However, a dominant follicle may also be selected during anovulatory waves that precede the ovulatory wave in some women.

CONCLUSIONS: There is increasing evidence to indicate that multiple waves of antral follicles develop during the human menstrual cycle. Ovarian follicular waves in women are comparable with those documented in several animal species; however, species-specific differences exist. Enhancing our understanding of the endocrine and paracrine mechanisms underlying antral follicular wave dynamics has clinical implications for understanding age-related changes in reproductive function, optimizing hormonal contraceptive and ovarian stimulation regimens and identifying non-invasive markers of the physiologic status of follicles which are predictive of oocyte competence and assisted reproduction outcomes.

Key words: antral / follicle / menstrual cycle / ovary

Introduction

The earliest descriptions of reproduction came from Hippocrates in the fifth century BC who attested that 'generation' occurred through the joining of the male ejaculate and female menstrual blood (Short, 1977; Cobb, 2006). Over 2 millennia later, William

Harvey proclaimed 'ex ovo omnia'—all things come from the egg (Cobb, 2006). In the late-1600s, a period that heralded the age of enlightenment, Regnier de Graaf recognized the ovary for what it is—the producer of eggs (Cobb, 2006). These early modern scientists, however, held the mistaken belief that the follicle itself was the egg. It was Karl Ernst von Baer, in 1827, who discovered that the mammalian

oocyte was enclosed within the ovarian follicle (Cobb, 2006). It was not until the twentieth century that the human oocyte was discovered and the first descriptions of human ovarian follicular development were made (Block, 1951; Short, 1977).

Much of the current knowledge on ovarian function in women has been extrapolated from studies performed in non-human primates, farm animals and rodents. However technologic advances over the past 60 years have provided non-invasive tools for evaluating ovarian function in women. Early studies to evaluate human folliculogenesis and menstrual cyclicality in the 1950–1970s were based on histologic and/or endocrinologic evaluation of ovarian function (Block, 1951, 1952; Baker, 1963; Bjersing, 1967; Mikhail, 1970; Ross et al., 1970; Baird and Fraser, 1974; Lintern-Moore et al., 1974; Ross, 1974; Baird et al., 1975; Sherman and Korenman, 1975; Gougeon, 1977, 1979; McNatty, 1978; McNatty et al., 1979, 1981a). Histologic assessment of the ovaries provides precise microscopic information about the structural and functional status of follicles at a given point in time. Serum endocrine profiling provides information about folliculogenesis at any given point or over multiple days during the cycle, albeit indirectly. Conceptual models for understanding ovarian function over the menstrual cycle were developed from data obtained using a combination of anatomical and endocrinologic techniques. However, it was not until the development of transabdominal ultrasonography in the late-1970s that the serial growth and regression of follicles could be visualized (Hackeloer and Robinson, 1978; Hackeloer et al., 1979; Hall et al., 1979; O'Herlihy, 1980; Queenan et al., 1980; Renaud et al., 1980; Kerin et al., 1981; Zegers-Hochschild et al., 1984; Lenz, 1985). High-resolution transvaginal ultrasonography, first used in the late-1980s, dramatically improved our ability to visualize ovarian structures *in situ*. Antral follicles as small as 2 mm can now be detected and the growth dynamics of individually identified follicles have been described (Andreotti et al., 1989; Pache et al., 1990; Pierson et al., 1990; Pierson and Chizen, 1991; Martinuk et al., 1992; Bakos et al., 1994; Gore et al., 1995). The resolving power of conventional ultrasonographic imaging is still much less than that of histology, and measurements of follicle size (i.e. diameter, volume) alone provide limited information about ovarian function. However, Doppler ultrasonographic measurements of blood flow (Jokubkiene et al., 2006) and computer-assisted image analyses (Singh et al., 1997, 1998; Rezaei et al., 2009) can provide information about some physiologic aspects of ovarian function. It is the synergistic use of histologic, endocrinologic and ultrasonographic approaches that has been invaluable for elucidating our understanding of antral follicular development during the menstrual cycle.

The objective of the present review is to provide a comprehensive discussion of the different perspectives on antral folliculogenesis during the human menstrual cycle. The patterns and mechanisms of follicular recruitment, selection and ovulation in women are deliberated. Knowledge gleaned from the study of antral folliculogenesis during the estrous cycles of other mammals are considered where human data are not available and for comparative purposes.

Methods

Literature search

A comprehensive review was conducted on ovarian antral follicle dynamics using PubMed. The search criteria used were: antral, antrum, atresia,

corpus, cycle, dynamics, endocrine, follicle, follicular, folliculogenesis, histologic, hormonal, hormone, luteal, luteum, menstrual, morphologic, ovary, ovarian, ovulation, paracrine, recruitment, regression, selection, ultrasonographic, ultrasound and/or wave. Searches were limited to the female gender and criteria were identified in the title and/or abstract of the publications. Literature available from 1950 to 2010 was included. Searches were conducted by the primary author and two trained research assistants. Additional journal articles were identified from the bibliographies of studies included as well as textbooks and hand searches of other source materials. No registration number for this review is available.

Study selection process

Approximately 4100 journal articles, case reports and reviews were initially identified. Articles written in English in the field of human medicine were selected first. A separate search was conducted using the same criteria indicated above to include results of research conducted in animals. Duplicated studies were removed. Articles written in a language other than English without available translations were excluded from our review. The titles were first reviewed for relevance to the study of ovarian follicular dynamics. The abstracts of ~610 English and non-English (translated) articles were reviewed. Research conducted to exclusively characterize antral follicle development during ovarian stimulation cycles or hormonal contraception cycles were excluded. Only review articles describing pre-antral follicular development were included. The full-text articles of ~320 studies were further examined. Of these, 246 were deemed relevant and included in our review.

Setting the stage: pre-antral and early antral follicle development

Early descriptions of ovarian follicular development in women were made using histologic evaluation of ovaries postmortem or following oophorectomy. From these studies, the entire duration of human folliculogenesis, from the primordial phase to the pre-ovulatory phase, has been estimated to span >175 days (Gougeon, 1986). Follicular development begins as early as the fourth month of fetal life (Baker, 1963). The primordial germ cells at this time have migrated from the yolk sac endoderm to the gonadal ridge, undergoing mitotic divisions. Once arriving at the gonadal ridge, the oogonia enter the first meiotic division and become primary oocytes. Somatic cells originating from the primitive gonad (i.e. surface epithelial cells, follicular granulosa and theca cells, interstitial cells, fibroblasts) surround the oogonia, forming rudimentary ovarian follicles (<0.1 mm in diameter) (Gougeon, 1979). Follicles containing oocytes arrested in the dictyate stage of meiosis I constitute the ovarian follicular reserve, which provides a woman with reproductive potential for her entire lifetime. The number of follicles occupying the follicular reserve is estimated to be ~7 million at 20 weeks gestation (Baker, 1963). Depletion of the ovarian follicular reserve begins during fetal life and continues throughout a woman's lifetime (Block, 1952; Baker, 1963; Gougeon et al., 1994; Hansen et al., 2008).

The mechanisms underlying the developmental progression of human primordial follicles from the resting to growing state are not fully understood, and are not the focus of the present review. Briefly, currently available evidence is broadly interpreted to mean that the destiny of each follicle is dependent on a delicate balance between the expression and action of factors promoting follicular cell proliferation, growth and differentiation and those promoting

apoptosis; some primordial follicles initiate growth while others remain quiescent for months or years [reviewed by: (Roy and Greenwald, 1996; Fortune, 2003; Aerts and Bols, 2010; Gougeon, *epub ahead of print*)]. Pre-antral follicles (~0.1–0.2 mm) develop independent of gonadotrophin support [reviewed in Gougeon (1979) and Craig *et al.*, (2007)]. At a diameter of ~0.2–0.4 mm, a fluid-filled cavity (i.e. antrum) begins to develop, and follicles become responsive to gonadotrophins (Gougeon, 1979; Craig *et al.*, 2007). Follicles may develop to the early antral stage throughout fetal life, infancy and childhood, but regress (Peters, 1979; Gougeon, 1996). At puberty, maturation of the hypothalamo-pituitary-gonadal axis results in pulsatile release of follicle-stimulating hormone (FSH) and luteinizing hormone (LH) from the pituitary which culminates in the cyclic development of antral follicles ≥ 2 mm, onset of ovulation and menstrual cyclicity [reviewed in Gougeon (2004) and Messinis (2006)].

Terminology used in describing folliculogenesis

Differing terminology used to describe antral follicular development among studies and among species has led to considerable confusion. For example, in the human literature, the term ‘recruitment’ has been used to describe three important, but distinctly different, physiologic events: (i) the initial transition of primordial follicles from the resting pool into the pre-antral growth phase, (ii) the cyclic recruitment of a cohort of antral follicles (2–5 mm) during the menstrual cycle following puberty and (iii) the preferential growth of the dominant ovulatory follicle. Follicles comprising a recruited cohort of 2–5 mm follicles have been referred to as ‘selectable’ follicles (Gougeon, 1996).

Similarly, the term ‘selection’ has been used to describe two different phenomena: (i) the recruitment of a cohort of 2–5-mm antral follicles and (ii) the preferential growth of a species-specific number of large antral follicles from the recruited cohort. The follicle that is selected from the recruited cohort has been referred to as the ‘dominant’ (Hodgen, 1982) or ‘privileged’ (Gougeon, 1986) follicle, while all other follicles of the cohort which undergo atresia have been termed ‘ordinary’ (Gougeon, 1986), ‘challenger’ (Gore *et al.*, 1997), ‘subdominant’ (Gore *et al.*, 1994), or ‘subordinate’ (Adams, 1999).

In the present review, the terms follicle ‘recruitment’ and ‘selection’ are used to represent two different physiologic events. Follicle ‘recruitment’ refers to the emergence of a group or cohort of medium-size (2–5 mm) antral follicles. ‘Selection’ refers to the preferential growth of the dominant follicle from the cohort of recruited antral follicles. ‘Subordinate’ follicles comprise all follicles of the recruited cohort, excluding the dominant follicle. The term ‘cohort’ is used interchangeably with ‘wave’. The term ‘recruitment’ is used interchangeably with ‘emergence’. The term ‘interovulatory interval’ (IOI) is used to define the time period between successive ovulations (Douglas and Ginther, 1975; Pierson and Ginther, 1987b; Knopf *et al.*, 1989; Baerwald *et al.*, 2003b). In this way, we have attempted to maintain consistency with terminologies used in the human and animal literature.

Antral follicle recruitment

Antral follicles 2–5 mm in diameter have been detected histologically and ultrasonographically throughout the human menstrual cycle (Gougeon and Lefevre, 1983; Chikazawa *et al.*, 1986; Pache *et al.*, 1990; Baerwald *et al.*, 2003a). The pattern of emergence of 2–5 mm follicles, however, is a matter of long-standing debate. Several studies have been conducted to characterize changes in the diameters and numbers of antral follicles during the menstrual cycle. Some investigators have suggested that antral follicles 2–5 mm develop continuously, while others have proposed that ‘cohorts’ or ‘waves’ of antral follicles develop in a cyclic manner during the menstrual cycle. Three separate theories of follicular recruitment have been proposed and are presented in detail below.

Continuous recruitment (Theory 1)

Early histological studies in rats (Mandle and Zuckerman, 1950), monkeys (Green and Zuckerman, 1951) and sheep (Turnbull *et al.*, 1977) led investigators to conclude that early antral follicle growth occurred continuously throughout the estrous/menstrual cycle. From these studies in animals, the theory of continuous follicular recruitment evolved for the human menstrual cycle. According to this theory, small antral follicles ≤ 4 –6 mm are recruited to grow continuously, at all stages of reproductive life, independent of gonadotrophins (McNatty, 1981a; Westergaard *et al.*, 1986; Baird, 1987). The follicle destined to ovulate is said to be selected, by chance, from the continuous supply of antral follicles by virtue of being at the right stage of maturity to respond to the rise in FSH (and subsequently LH) that occurs following luteal regression (Wiele *et al.*, 1970; Baird, 1983, 1987) (Fig. 1a).

Cyclic recruitment

Cyclic increases in the number of antral follicles, referred to as ‘recruitment of a follicular cohort’ or ‘follicle wave emergence’, have been observed at regular intervals during the menstrual cycle in women (Block, 1951; Gougeon and Lefevre, 1983; McNatty *et al.*, 1983; Baerwald *et al.*, 2003b) and non-human primates (Dizerega and Hodgen, 1981). There is inconsistency in the literature, however, regarding the number of recruitment episodes that occur during the human menstrual cycle.

Single recruitment episode (Theory 2)

Histologic, endocrinologic and early ultrasonographic investigations demonstrated that a cohort of 2–5 mm follicles is recruited from a continuous supply of antral follicles once during each menstrual cycle (Gougeon, 1979; O’Herlihy, 1980; Chikazawa *et al.*, 1986; Pache *et al.*, 1990) (Figs 1b, 2a and b). The single increase in the number of follicles 2–5 mm, detected after regression of the CL in the late-luteal phase or early follicular phase of the menstrual cycle, has been referred to as a ‘privileged’ phase of follicle development (Gougeon, 1979). It is thought that the recruited cohort represents a group of follicles at comparable, but not identical, stages of development (Fauser and Van Heusden, 1997). Follicles of the recruited cohort are believed to leave the pool of resting primordial follicles around the same period of time several months before (Fauser and Van Heusden, 1997).




Figure 1 Schematic illustrations of the three different theories of ovarian follicular recruitment during the menstrual/estrous cycle. The theory of continuous recruitment (a) suggests that small antral follicles grow and regress constantly throughout the interovulatory interval and the dominant ovulatory follicle is selected, by chance, from the pool following luteal regression. The theory of a single recruitment episode (b) suggests that an ovulatory follicle is selected from a single follicular cohort that emerges following luteal regression. The wave theory of follicle recruitment (c) suggests that two or more cohorts (i.e. waves) of antral follicles are recruited (i.e. emerge) during the ovarian cycle. The dominant follicle that develops in the final wave of the IOI ovulates while preceding waves are anovulatory; dominant follicles develop in a minority of anovulatory waves (dotted line). M, menses.

Following regression of the CL, luteal estradiol and inhibin fall resulting in a rise in circulating concentrations of FSH (Roseff et al., 1989; Le Nestour et al., 1993). The rise in FSH following luteolysis is

thought to be responsible for preventing atresia of a cohort of 2–5 mm antral follicles in each ovary (Koering, 1969; Hodgen, 1982; McNatty et al., 1983; Gougeon, 1984; Pache et al., 1990; Van Santbrink et al., 1995). The corollary to the notion of preventing atresia is the concept that recruitment is induced by rising FSH; it has been postulated that each follicle and/or woman has a threshold requirement for FSH below which follicle recruitment does not occur (Brown, 1978; Fauser and Van Heusden, 1997). Thus, recruitment of the follicular cohort (through stimulation of growth or prevention of atresia) rather than selection of the dominant follicle occurs in response to a transient elevation in circulating FSH.

Antral fluid from follicles of the recruited cohort contains low concentrations of estradiol but relatively high concentrations of androgens (Gougeon, 1996). Inhibin B produced by granulosa cells in follicles of the recruited cohort (Groome et al., 1996; Fraser et al., 1999; Laven and Fauser, 2004) acts in an endocrine manner to inhibit continued FSH secretion in the mid-follicular phase (Van Santbrink et al., 1995; Groome et al., 1996). By comparison, inhibin A concentrations are low during the follicular phase and maximal during the midluteal phase (Groome et al., 1996), suggesting that the CL is a source of inhibin A.

In the luteal phase of the menstrual cycle in women and non-human primates, estradiol and inhibin A production by the CL is thought to suppress FSH and thereby prevent the development of healthy follicles >4 mm (Savard et al., 1965; Mikhail, 1970; Gougeon and Lefevre, 1983; McNatty et al., 1983; Baird et al., 1984; Chikazawa et al., 1986; McLachlan et al., 1987; Sasano et al., 1989; Bassetti et al., 1990; Smith et al., 1990; Roberts et al., 1993; Groome et al., 1996; Sanders and Stouffer, 1997; Devoto et al., 2008). Most antral follicles observed histologically during the luteal phase in women were atretic, leading to the conclusion that there is a limited number of healthy antral follicles that can develop under the influence of the corpus luteum (McNatty et al., 1983). In addition, the development of healthy follicles in the luteal phase has been demonstrated when the CL was enucleated or when exogenous gonadotrophins were administered (Nilsson et al., 1982; Baird et al., 1984; Baird, 1987).

In contrast to primates, large antral follicles develop in some domestic animals throughout the luteal phase (i.e. sheep, cows) (Baird et al., 1975, 1984; Pierson and Ginther, 1987b; Sirois and Fortune, 1988). Estradiol and inhibin production by the CL was thought to be unique to primates (Savard et al., 1965; Mikhail, 1970; Baird et al., 1975; Bassetti et al., 1990). Thus, the development of luteal phase follicles and asymmetry in the lengths of the luteal and follicular phases observed in domestic animals (in particular, sheep) were attributed to the absence of estradiol production (and possibly inhibin) by the CL (Short, 1964; Kaltenbach et al., 1967; Baird et al., 1973, 1975). However, a more recent documentation of luteal aromatase activity in mares (Albrecht et al., 1997) has challenged the notion that luteal estradiol production is unique to primates.

Contrary to early histologic and ultrasonographic studies in which a single increase in the number of 2–5 mm follicles was detected in the late-luteal or early-follicular phase (reviewed above), the results of other studies indicated no clear change in the number of antral follicles across the menstrual cycle. In one study, the mean number of antral follicles (referred to as the antral follicle count—AFC) were not different in the early-follicular, late-follicular and luteal phases (Pache et al., 1990). In another study, the numbers of antral follicles 2–5 and




Figure 2 Data regarding the single recruitment episode theory of antral follicle development during the human menstrual cycle. Morphologic changes associated with the selection of a single dominant follicle from the recruited cohort in the mid-follicular phase, followed by ovulation at mid-cycle and limited luteal phase follicle development; data represent a single woman [a and b; Reproduced by permission from the American Society for Reproductive Medicine, Pache *et al.*, 1990 *Fertil Steril*, 54(4):638–42]. The number of antral follicles 2–5 and 2–10 mm in diameter at different phases of the menstrual cycle [c and d; Reproduced with permission from Oxford Journals, Disseldorp *et al.*, 2010, *Hum Reprod* 25(1): 221–7].

2–10 mm were not different between the early-, mid- and late-follicular, pre-ovulatory and early-, mid- and late-luteal phases of the cycle (van Disseldorp *et al.*, 2010) (Fig. 2c and d). Non-significant but observable increases in the mean number of antral follicles in the early-luteal phase of the cycle were attributed to inter-individual differences and were considered to be of limited biological importance (van Disseldorp *et al.*, 2010).

Anti-Mullerian hormone (AMH) is produced from the granulosa cells of primary, secondary, pre-antral and early antral follicles (≤ 4 mm). AMH inhibits the initiation of primordial follicle growth from the ovarian reserve (Weenen *et al.*, 2004; Nilsson *et al.*, 2007). Furthermore, in mice, AMH decreases the sensitivity of follicles to FSH thereby inhibiting FSH-induced antral follicle growth (Durlinger *et al.*, 2002). Although not fully understood, AMH may play a role in regulating recruitment of the antral follicular cohort (Visser and Themmen, 2005; Visser *et al.*, 2006).

Follicular waves (Theory 3)

In contrast to the notion of a single recruitment episode during the menstrual cycle in women, multiple cohorts or 'waves' of antral follicle recruitment have been described (Block, 1951; Hackeloer *et al.*, 1979; Dervain, 1980; Queenan *et al.*, 1980; Baerwald *et al.*, 2003a, b)

(Fig. 1c). A 'wave' of follicular development is defined as the synchronous growth of a group of antral follicles that occurs at regular intervals during the menstrual/estrous cycle; follicles in each wave are of similar, but not identical, diameters (Pierson and Ginther, 1987b; Ginther *et al.*, 1989a; Baerwald *et al.*, 2003a; Ginther *et al.*, 2004).

The earliest studies of human follicular dynamics involved histologic evaluations of excised ovaries (Block, 1951; McNatty, 1978, 1981a; McNatty *et al.*, 1983). Two waves of follicle growth > 1 mm were detected during the menstrual cycle (Block, 1951). The first wave occurred in the follicular phase and a second wave occurred in the luteal phase. The number of antral follicles > 4 mm increased in the late-follicular and late-luteal phases (McNatty, 1978). However, luteal phase follicles had fewer granulosa cells and produced less estradiol compared with those in the follicular phase (McNatty, 1978, 1981a). In another study, antral follicles 1–10 mm were detected in the luteal phase (McNatty *et al.*, 1983). Most follicles were atretic; however, an increase in the number of non-atretic antral follicles > 1 mm was reported in the early-luteal versus mid-luteal and late-follicular phases.

Waves of follicle development have been documented in healthy women using transabdominal ultrasonography (Hackeloer *et al.*, 1979; Dervain, 1980). Two waves were detected in women with regular 30–35 day cycles compared with one follicular wave in

women with 26–30 day cycles. Results of a more recent study using serial transvaginal ultrasonography and concurrent endocrine profiling further supported a wave theory of antral folliculogenesis in women (Baerwald et al., 2003b). Emergence of a wave of 4–14 follicles ≥ 4 –5 mm was detected either two or three times during the IOI in a large population of healthy women. Most women (68%) exhibited two waves of follicle recruitment during the IOI, while the remaining women (32%) exhibited three waves. Women with three follicular waves had a mean IOI of 29 days, which was significantly longer than the mean IOI of 27 days in women with two waves. In women with two follicular waves, an anovulatory wave emerged at the time of ovulation (i.e. early-luteal phase) followed by emergence of the ovulatory wave during the early-follicular phase. In women with three waves, an anovulatory wave emerged at the time of ovulation, a second anovulatory wave emerged during the mid- to late-luteal phase and a third wave (the ovulatory wave) emerged in the early- to mid-follicular phase (Baerwald et al., 2003b).

An elevation in circulating FSH appears to precede the recruitment of each follicular wave during the IOI in women (Baerwald et al., 2003a), consistent with previous reports in domestic animals (Adams et al., 1992b; Ginther et al., 1995; Adams, 1999; Gastal et al., 2000; Baerwald et al., 2003a). In cows, the wave-eliciting surge in circulating FSH spans a period of ~ 3 –4 days and is of similar magnitude among waves (Adams et al., 2008). In contrast to initial expectations, the height of the peak in circulating FSH in cows is inversely related to the number of follicles recruited into a wave (Singh et al., 2004; Burns et al., 2005). The inverse association between circulating FSH and number of follicles in a wave cannot be directly explained by alterations in the secretion of estradiol, inhibin, or insulin-like growth factor (IGF)-I (Burns et al., 2005), but is consistent with the age-related decrease in AFC and elevated serum FSH in women (Klein et al., 1996a; Gougeon, 1998), cattle (Malhi et al., 2005) and sheep (De Souza et al., 1998).

Inhibin B from the granulosa cells of the recruited cohort (Groome et al., 1996; Fraser et al., 1999; Laven and Fauser, 2004) inhibits continued FSH secretion in the mid-follicular phase (Van Santbrink et al., 1995; Groome et al., 1996). A second short-lived peak in inhibin B has been documented 2 days after the mid-cycle LH surge (Groome et al., 1996), providing evidence of a luteal phase wave of follicular growth in women. The precise roles of inhibin A, inhibin B and AMH in regulating the emergence of multiple follicular waves in women are not understood; investigations are currently underway (Baerwald et al., unpublished data).

Follicular waves have been recently detected in women during the perimenopausal transition (Hale et al., 2007; Baerwald et al., unpublished data) and in women undergoing ovarian stimulation therapy (Bentov et al., 2010). Further support for the concept of multiple follicular waves is provided by clinical reports of successful luteal phase oocyte retrieval and *in vitro* maturation as an optional procedure for urgent fertility preservation (Demirtas et al., 2008; Maman et al., 2011). The existence of healthy antral follicles in the luteal phase was attributed to the emergence of a new wave of follicle development in the luteal phase.

Observations of antral follicular waves in women are consistent with those previously documented in several animal species, including cattle (Pierson and Ginther, 1987b, 1988; Sirois and Fortune, 1988; Ginther et al., 1989a), mares (Pierson and Ginther, 1987a; Sirois et al., 1990;

Ginther, 1993), sheep and goats (Ginther et al., 1995; Evans et al., 2000), llamas and alpacas (Adams et al., 1990; Vaughan et al., 2004), musk oxen (Hoare et al., 1997), water buffalo (Taneja et al., 1996), as well as deer and wapiti (Asher et al., 1997; McCorkell et al., 2006). Evidence of follicular waves has also been reported in sub-human primates (Lujan et al., 2006; Bishop et al., 2009). The day of follicular wave emergence is recognized ultrasonographically when the largest follicle of the cohort first attains a diameter of 4–6 mm in women (Baerwald et al., 2003a; Ginther et al., 2004), compared with 4–5 mm in cows (Sirois and Fortune, 1988; Ginther et al., 1989b; Ginther, 2000), 4–5 mm in sheep (Evans et al., 2000) and 13–15 mm in mares (Sirois et al., 1990; Ginther et al., 2004).

Follicle selection

Follicle ‘selection’ is the process by which a single ‘dominant’ follicle is chosen from the recruited cohort or wave for preferential growth (Block, 1951; Hodgen, 1982; Chikazawa et al., 1986; Gougeon, 1986; Pache et al., 1990; Van Santbrink et al., 1995; Ginther et al., 2001a; Baerwald et al., 2003a). Follicle selection has generally been documented to occur once in the early- to mid- follicular phase of the menstrual cycle, leading to ovulation (Baird, 1990; Pache et al., 1990; Zeleznik, 2004) (Fig. 2a and b).

Follicle divergence

The dominant and largest subordinate follicles of the ovulatory wave undergo a common growth phase in women, consistent with observations in mares and cows (Ginther et al., 2001a). At the time of selection, the growth profile of the dominant follicle begins to ‘diverge’ as it continues to grow while the subordinate follicles undergo atresia (Adams et al., 1993; Ginther et al., 2001a). Divergence occurs when the dominant follicle reaches a diameter of ~ 10 mm on Day 6–9 of the follicular phase in women (Pache et al., 1990; Van Santbrink et al., 1995; Ginther et al., 2001a; Baerwald et al., 2003a). Morphologic changes in the growth dynamics of dominant and first subordinate follicles in women are similar to that described in other monovular species (cows and mares; Fig. 3) (Adams and Pierson, 1995; Ginther et al., 2001a).

Follicle dominance

Ablation of the dominant follicle in monkeys (Dizerega and Hodgen, 1981) and women (Nilsson et al., 1982; Araki et al., 1983) has been shown to cause a delay in the pre-ovulatory gonadotrophin surge and ovulation by ~ 2 weeks. This delay was interpreted to mean that the dominant follicle had already been selected at the time of cauterization and that no other follicle was competent to accommodate a timely ovulation. The delay reflected the period of time required for a new group of follicles to be recruited, out of which another dominant follicle was selected and ovulated. Similar follicle ablation studies in cattle have demonstrated that the dominant follicle suppresses the growth of subordinants of the same wave and suppressed emergence of the next follicular wave through an inhibitory effect on circulating FSH concentrations (Adams et al., 1992b, 1993). Collectively, these findings from human and animal studies support the concept that the dominant follicle exerts both morphologic and functional dominance once selection has occurred.




Figure 3 Comparative changes in the diameters of the dominant and first subordinate follicles during the process of follicle selection in women, mares/horses and heifers/cows (day 0 = follicle wave emergence). The dotted horizontal line indicates the diameter at which deviation in follicle growth rates between the dominant and first subordinate follicles occurs (Reproduced with permission from Elsevier, Ginther et al., 2001, *Biol Reprod*, 65:648–647).

Follicle-stimulating hormone

While an increase in circulating concentrations of FSH is requisite for recruitment of the follicular cohort or wave (Roseff et al., 1989; Adams et al., 1992a; Le Nestour et al., 1993; Van Santbrink et al., 1995; Baerwald et al., 2003a), the nature of the post-surge decline in FSH is a critical factor in selection of the dominant follicle (Adams et al., 1992a, 1993). The duration of the rise in FSH above a critical threshold determines the number of dominant follicles selected from the recruited cohort for preferential growth (Gibbons et al., 1997; Schipper et al., 1998). This concept has been termed the 'FSH Threshold/Window/Gate Concept' (Brown, 1978; Baird, 1987; Schipper et al., 1998) (Fig. 4). The duration that FSH is above the threshold is short for the development of a single dominant follicle. In contrast, widening the FSH window allows multiple follicles to be selected at the same time, as occurs in polyovular species and during ovarian stimulation therapy (Schipper et al., 1998).

There is evidence in women and domestic animals that the dominant follicle has an early size advantage over subordinate follicles (Ginther et al., 1997, 2004; Jaiswal et al., 2004). It has been postulated that the future dominant follicle may contain more granulosa cells and FSH receptors, making it more sensitive to FSH, compared with the subordinate follicles (Fauser and Van Heusden, 1997). Subordinate follicles are not able to thrive in an environment of declining FSH and therefore succumb to atresia (Yamoto et al., 1992a; Roberts et al., 1993; Van Santbrink et al., 1995; Magoffin and Jakimiuk, 1997; Schneyer et al., 2000; Baerwald et al., 2003a). Selection has been described as a phenomenon of avoiding atresia. That is, the selection process may be considered a hierarchical progression of follicle atresia over the period encompassing the rise and fall in FSH. The smallest follicles in the cohort are the least capable of competing for




Figure 4 Schematic representation of the FSH threshold (window) concept and follicle growth dynamics (recruitment, selection and dominance) during the follicular phase of the menstrual cycle. [Reproduced with permission from Elsevier, Fauser and Van Heusden, 1997, *Endocrine Reviews*, 18(1): 71–105; Originally adapted from Baird et al., 1987, *J Steroid Biochem*, 71(1): 15–23].

diminishing levels of FSH and therefore undergo atresia first, followed by regression of progressively larger follicles in the wave until ultimately only one (dominant) follicle survives (Adams et al., 1993; Jaiswal et al., 2004).

Estradiol

By Day 5–8 of the menstrual cycle, aromatase activity begins in granulosa cells of follicles larger than 6–8 mm, with the dominant follicle producing more estradiol-17 β than other follicles in the cohort (Mikhail, 1967; Lloyd et al., 1971; Baird and Fraser, 1975; McNatty et al., 1976; Hillier et al., 1981; Chikazawa et al., 1986). The follicular fluid of dominant follicles contains more estrogen and less androgen, while atretic subordinate follicles exhibit a greater androgen/estrogen ratio (Westergaard et al., 1986; Mango et al., 1988; Gougeon, 1996; Van Dessel et al., 1996). LH-induced production of androgens in thecal cells provides the substrate for estradiol production in granulosa cells (Ryan, 1979; Baird, 1987; Adashi, 1994). Estradiol-17 β produced from the dominant follicle provides negative feedback on pituitary FSH secretion which contributes, in part, to the mid-follicular phase decrease in circulating FSH and inhibition of subordinate follicle growth (Ginther et al., 2000a). LH receptors form on granulosa cells of the dominant follicle following estradiol secretion (Zeleznik et al., 1974; Bodensteiner et al., 1996). The dominant follicle therefore becomes less dependent on FSH and more responsive to LH during the selection process (Yamoto et al., 1992b; Sullivan et al., 1999; Filicori et al., 2002). Greater production of estradiol and a shift from FSH to LH sensitivity in the dominant follicle in women is consistent with observations in cows and mares (Bodensteiner et al., 1996; Gastal et al., 1999, 2000; Ginther et al., 2000a, 2001b, c).

Transforming growth factor- β (TGF- β) superfamily

Bi-directional communication between the oocyte and cumulus cells regulates both folliculogenesis and oogenesis (Eppig, 2001; Senbon et al., 2003). Oocyte and follicle-derived factors have stage-specific roles in regulating antral follicular development and oocyte competence. Members of the transforming growth factor- β (TGF- β) superfamily, including inhibin, activin, follistatin, TGF- β , bone morphogenic protein (BMP), growth differentiation factor (GDF) and AMH, are among the many growth factors which operate via paracrine and/or autocrine mechanisms to regulate follicular development and oocyte maturation.

All follicles of the cohort produce inhibin B, which contributes to the decrease in FSH that occurs prior to selection (Gibbons et al., 1997; Fraser et al., 1999; Ginther et al., 2000b). However, the role of activin in dominant follicle selection in women is not well elucidated. Activin B is produced by the anterior pituitary and stimulates FSH production from the gonadotropes by autocrine/paracrine mechanisms [reviewed in Bilezikjian et al. (2004)]. Activins are also produced by the granulosa cells and have been shown to stimulate granulosa cell proliferation in large and small follicles in rats (Li et al., 1995; Miro and Hillier, 1996). Activin A was shown to cause a time- and dose-dependent inhibitory effect on LH-induced production of progesterone and oxytocin production in bovine granulosa cells *in vitro*, indicating that activin A inhibits spontaneous luteinization in mature antral follicles (Findlay, 1993). Granulosa cell activin A in rats (Hsueh et al., 1987) and humans (Hillier et al., 1991a) was also shown to attenuate LH-dependent androgen production *in vitro* in thecal cells of small to medium size antral follicles, with this effect reversed by follistatin/activin-binding protein. Conversely, granulosa cell inhibin A of selected antral follicles in rodents (Hsueh et al., 1987), domestic animals (Wrathall and Knight, 1995) and humans (Hillier et al., 1991b) increased LH-induced thecal androgen production, which was then thought to serve as the substrate for dominant follicle estradiol production. In contrast to inhibin A, follicular fluid inhibin B peaks at a follicle diameter of 9–10 mm in women (Andersen et al., 2010). Thus, it has been proposed that an orderly transition from an inhibin B/activin follicular environment to an inhibin A/follistatin environment is critical for dominant follicle development in women (Roberts et al., 1993; Groome et al., 1996; Schneyer et al., 2000). Inhibin and activin are involved in regulating follicle selection in domestic animals; however, species-specific differences have been detected in the isoforms involved [reviewed in Ginther et al. (2001a) and Mihm and Evans (2008)]. Differences and similarities between the secretion of gonadotrophins, inhibin and estradiol during follicle selection in women, compared and contrasted to that in domestic farm animals, are illustrated in Fig. 5.

Intrafollicular AMH gradually decreases during antral follicular growth until 8–10 mm (i.e. approximate time of selection), when AMH profoundly decreases and remains low thereafter (Andersen et al., 2010). The decrease in intrafollicular AMH during selection of the ovulatory follicle occurs in association with a rise in granulosa cell aromatase expression (Nielsen et al., 2010); higher AMH concentrations have been detected in cumulus versus granulosa cells (Grondahl et al., 2011). It is plausible that the decrease in AMH around the time of selection may release inhibition of aromatase

expression; however, the precise roles of AMH in dominant follicle selection are not known.

Research conducted primarily in rodents suggests that granulosa-cell activin and BMP-6 act via autocrine/paracrine mechanisms while oocyte-derived GDF-9, BMP-15, BMP-6 act via paracrine routes to promote granulosa cell proliferation and modulate FSH-dependent follicle function [reviewed in Knight and Glister (2006)]. It is proposed that differential exposure to these signaling molecules may be one of the ways in which the dominant follicle is sensitized to FSH and thereby selected for preferential growth.

Insulin-like growth factor (IGF) system

The IGF system plays an important role in the intraovarian regulation of antral follicular development. Circulating levels of IGF-I and IGF-II have not been shown to differ during the human menstrual cycle (Jesionowska et al., 1990). However, changes in follicular fluid IGF concentrations have been reported (Eden et al., 1988). Both IGF-I and II have been shown *in vitro* to have effects on ovarian follicular function; however, IGF-II appears to be the primary IGF in primate ovaries. IGF-I and II mRNA have been detected in thecal cells from small antral follicles, but only IGF II mRNA has been detected in the granulosa cells of dominant follicles (El-Roeiy et al., 1993).

IGF-II increases in the dominant follicle at the time of selection. Simultaneously, its bioavailability is increased via proteolysis of insulin-like growth factor-binding protein-4 (IGFBP-4) by IGFBP-4 protease (Poretsky et al., 1999; Giudice, 2001). IGF II (and also IGF-I) stimulates aromatase activity, estradiol and progesterone production in human granulosa cells and promotes androgen production in the thecal cells of the growing dominant follicle. In subordinate follicles, IGF is sequestered by IGFBP-4, granulosa and thecal cell steroidogenesis is inhibited, and atresia ensues (Iwashita et al., 1996; Mason et al., 1998; Poretsky et al., 1999; Hourvitz et al., 2000). The role of IGF-I and II in regulating antral follicle development in women is similar to that described in rodent and domestic animal species (Poretsky et al., 1999; Fortune et al., 2004; Beg and Ginther, 2006). There is further evidence in rodents to suggest that follicular activin, GDF-9, AMH and BMPs may regulate dominant follicle selection by modulating granulosa cell IGF-dependent signaling pathways [reviewed in Knight and Glister (2006)].

Although current data on the role of paracrine and autocrine factors in regulating physiologic selection of the dominant follicle show general similarities among species, inconsistencies have been reported. Continued research is required to elucidate the precise roles of intraovarian factors in modulating antral folliculogenesis.

Luteal influences on follicle selection

Numerous studies have been conducted to determine whether the presence of the CL influences selection of the dominant follicle. Seven out of eight dominant follicles were reported to develop contralateral to the CL from the previous ovulation in women (Chikazawa et al., 1986). Dominant follicles which developed contralateral to the previous ovulation have been shown to exhibit higher estradiol/androstenedione ratios than dominant follicles that developed ipsilateral to the previous CL (Fukuda et al., 1996). Furthermore, pregnancy rates in natural cycles following insemination and IVF were found to be higher when the ovulatory follicle developed contralateral to the CL




Figure 5 Temporal interrelationships between FSH, estradiol and inhibin during the process of dominant follicle selection in mares, women and cows. DF, dominant follicle; SF, subordinate follicle [Reproduced with permission from Wiley Blackwell Publishing, Mihm and Evans, 2008, *Reprod Dom Anim*, 43(Suppl 2): 48–56].


from the previous ovulation (Fukuda *et al.*, 1996). Evaluations of ovulation in both infertile and fertile women, however, support the notion that follicle selection and subsequent ovulation occurs randomly between the right and left ovaries (Baird, 1987; Check *et al.*, 1991; Ecochard and Gougeon, 2000; Ojha *et al.*, 2000; Baerwald *et al.*, 2003a), similar to findings in domestic animal species (Adams *et al.*, 2008).

Selection of a dominant follicle has been generally described to occur once in the early- to mid-follicular phase of the menstrual cycle (Gougeon and Lefevre, 1983; Baird, 1990). However, selection has been observed more than once during natural menstrual cycles in some healthy women (Baerwald *et al.*, 2003a). Major and minor waves of follicle growth were characterized in healthy women of reproductive age (Baerwald *et al.*, 2003a; Ginther *et al.*, 2004). Major waves were those in which a dominant follicle was selected, and minor waves were those in which dominance was not manifest. Most women developed a major ovulatory wave in the follicular phase and 1 or 2 minor anovulatory waves in the preceding luteal phase. However, 1 or 2 dominant follicles developed and subsequently regressed prior to selection of the ovulatory follicle in nearly a quarter of natural menstrual cycles (Baerwald *et al.*, 2003a). Reports of major anovulatory waves developing spontaneously prior to the ovulatory follicular wave in some, but not all, women are similar to follicular dynamics in mares (Ginther *et al.*, 2004), despite differences in luteal phase lengths between these species.

The role of the CL in regulating follicular wave dynamics has been studied in women and domestic farm animals. No differences in the size or lifespan of the CL, progesterone secretion or luteal phase estradiol secretion were detected in women with two versus three waves or in women with major versus minor waves preceding the ovulatory wave (Baerwald *et al.*, 2005). However, the presence of the CL appeared to influence dominant follicle selection in women

with three waves. When the second of three waves emerged in the mid-luteal phase, selection of a dominant follicle did not occur (i.e. a minor anovulatory wave developed) (Baerwald *et al.*, 2003a). By comparison, when the second wave emerged in the late-luteal or early-follicular phase, a dominant follicle was selected (i.e. a major anovulatory wave developed) (Baerwald *et al.*, 2003a). In contrast to women, one or two major anovulatory waves develop during the luteal phase of all estrous cycles in cows; minor waves have not been observed (Pierson and Ginther, 1987b; Ginther *et al.*, 1989a, 2004; Baerwald *et al.*, 2003a). Luteal regression and progesterone withdrawal occurs later in cows with three versus two follicular waves, at which time the viable dominant follicle present goes on to ovulate [reviewed in Adams *et al.* (2008)].

The functional status of dominant follicles that develop during anovulatory waves preceding the ovulatory wave in women is not fully understood. Estradiol levels have been shown to increase in association with the emergence of anovulatory follicle waves during the luteal and early-follicular phases of the cycle (Baerwald *et al.*, 2003a); however, the source of the estradiol (i.e. follicle versus corpus luteum) is not known. In a preliminary study, no differences in circulating estradiol concentrations were observed in the luteal phase of women with anovulatory major waves versus minor waves (Ginther *et al.*, 2005); however, data from a larger sample of women are required to either confirm or refute the preliminary inference. Ultrasonographic image attributes of dominant follicles have been found to differ between anovulatory and ovulatory waves (Rezaei *et al.*, 2009). Taken together, these data suggest that dominant follicles of anovulatory waves may exhibit different physiologic characteristics than dominant follicles of ovulatory waves in women. The morphologic and endocrinologic changes occurring in association with the development of follicle waves during the human menstrual cycle are illustrated in Fig. 6.



Pre-ovulatory follicle development

The dominant follicle of the final follicular phase wave continues to develop after it is selected and reaches pre-ovulatory status at a diameter of 16–29 mm in the late-follicular phase (Renaud et al., 1980; Kerin et al., 1981; Lenz, 1985; Gougeon, 1986; Pache et al., 1990; Bakos et al., 1994; Baerwald et al., 2003b). The ovulatory follicle grows at a rate of 1–4 mm/day, with reports of increases, decreases or no change in growth rate in the few days leading up to ovulation (Hackeloer et al., 1979; Renaud et al., 1980; Gougeon and Lefevre, 1983; Bomsel-Helmreich, 1985; Rossavik and Gibbons, 1985; Doody et al., 1987; Pache et al., 1990; Bakos et al., 1994; Baerwald et al., 2009). The dominant ovulatory follicle

grows slightly faster after selection compared with before (Baerwald et al., 2009).

Preferential growth of the dominant follicle in the mid-late-follicular phase is associated with increased aromatase activity and a rapid elevation of circulating and follicular fluid estradiol-17 β (Bomsel-Helmreich et al., 1979; Hackeloer et al., 1979; Kerin et al., 1981; McNatty, 1981a; Chikazawa et al., 1986; Bakos et al., 1994; Van Dessel et al., 1996; Baerwald et al., 2003a). Greater gonadotrophin responsiveness in the dominant follicle, compared with subordinate follicles, is responsible for mediating dominant follicle granulosa cell estradiol production, LH receptor expression and continued pre-ovulatory growth (McNatty, 1981b; Baird, 1983; Yamoto et al., 1992b; Van Santbrink et al., 1995; Van Dessel et al., 1996; Sullivan et al., 1999; Ginther et al., 2000a; Filicori, 2002). The dominant follicle




Figure 6 Continued.

is responsible for over 90% of the estrogen production in the pre-ovulatory period (Baird and Fraser, 1974).

Both intraovarian and endocrine factors contribute to pre-ovulatory follicle growth. The increase in granulosa cell aromatase mRNA expression in the dominant pre-ovulatory follicle is inversely related to follicular fluid AMH concentrations (Nielsen *et al.*, 2010). Increasing levels of inhibin A stimulate thecal cell androgen production, which acts as the substrate for continued estradiol secretion (Hillier *et al.*, 1991a; Roberts *et al.*, 1993; Schneyer *et al.*, 2000). IGF-II mRNA is abundantly expressed by granulosa cells of pre-ovulatory follicles (Geisthovel *et al.*, 1989) and stimulates aromatase activity (Hernandez *et al.*, 1992). Oocyte-derived GDF-9 and BMP-15 appear to be required for follicle development to the ovulatory stage (Juengel *et al.*, 2002), and BMP-15 may be involved in cumulus expansion (Gueripel *et al.*, 2006).

Estradiol production from the dominant follicle peaks the day before the LH surge (Viele *et al.*, 1970; Bomsel-Helmreich *et al.*, 1979) providing positive feedback at the hypothalamus and pituitary to stimulate the surge of LH necessary for inducing ovulation. The highly vascularized pre-ovulatory follicle, which has acquired LH

receptors, is now able to respond to the mid-cycle rise in LH (McNatty, 1978; Bomsel-Helmreich *et al.*, 1979; Tsang *et al.*, 1979). Ovulation occurs, on average, within 24 h of the LH peak (Kerin *et al.*, 1981). Serum progesterone concentrations begin to rise after the pre-ovulatory estradiol peak but before the LH surge, and indicate the onset of follicular luteinization (McNatty, 1978; Bomsel-Helmreich *et al.*, 1979; Brailly *et al.*, 1981; Chikazawa *et al.*, 1986; Westergaard *et al.*, 1986). Interactions between LH and EGF in the granulosa cell cytoplasm stimulate expression of extracellular-regulated protein kinases 1 and 2, which are believed to mediate a decrease in granulosa cell proliferation and estrogen synthesis, an increase in cumulus expansion and progesterone secretion and ultimately ovulation (Duggavathi and Murphy, 2009).

Discussion

Research conducted over the past 60 years using histologic, endocrinologic and ultrasonographic techniques has profoundly increased our understanding of antral ovarian folliculogenesis in women.

Different theories have emerged regarding the patterns in which antral follicles develop during the human menstrual cycle. However, comprehensive review of the literature shows similar trends among studies and among species. The following sections highlight common themes among research and suggest areas where continued investigations are required.

Animal models for the study of antral follicular dynamics in women

The study of human ovarian function is limited by ethical and practical considerations. Therefore, non-human primate, rodent and domestic farm animal models have been used to make inferences about follicular dynamics in women (Mandle and Zuckerman, 1950; McNatty, 1981b; Hodgen, 1982; Adams and Pierson, 1995; Bartlewski et al., 1999; Menezo and Herubel, 2002; Ginther et al., 2004; Ginther et al., 2005; Baerwald, 2008). While the use of animal models has increased our understanding of antral follicular dynamics in women, it is important to recognize similarities and differences in physiologic function as well as differences in research methodologies employed between species. In women, the presence of endometrial shedding and symmetric luteal and follicular phases are different from that observed during the estrous cycles of domestic farm animals. Despite these differences, general similarities in antral follicular dynamics exist. It is important to note that a continuous pattern of antral follicle development was originally proposed in domestic livestock species, similar to that shown in Fig. 1a. However, the use of frequent serial high-resolution ultrasonography and simultaneous endocrine profiling in these animal species has resulted in a broad understanding of follicular wave dynamics (Fig. 1c). Follicular waves have now been described in every species (including humans) in which this approach has been used. The number of follicular waves that develop appears to be species-specific and is positively correlated with the length of the menstrual/estrous cycle. Furthermore, despite a proportionately longer luteal phase in cows and mares, the growth phase of the dominant ovulatory follicle (defined as the time period from wave emergence to ovulation) is comparable in the human, cow and mare (Adams, 1999; Baerwald et al., 2003a; Ginther et al., 2004).

Critical studies of the characteristics and control of ovarian follicular and luteal dynamics in domestic farm animals have involved frequent (i.e. daily or multiple times a day) blood sampling and ultrasonography (Pierson and Ginther, 1987b; Sirois and Fortune, 1988; Ginther et al., 1989b; Sirois et al., 1990; Adams et al., 1992a; Ginther et al., 1998, 2000a, 2001c; Adams et al., 2008). Studies of this nature in women are difficult, and often unethical, to conduct. Differences in antral folliculogenesis between humans and animals appear to be more in detail rather than in essence, and may reflect differences in intrinsic physiology and/or differences in our ability to detect changes in ovarian function. Future studies should be conducted to compare endocrinologic and morphologic aspects of antral follicular wave dynamics at focused periods of the menstrual and estrous cycles in women and domestic farm species. This approach will continue to provide valuable insight for hypothesis testing and experimental design.

The relatively large diameters of antral follicles in cows and mares, compared with monkeys, sheep and rodents provide greater feasibility for characterizing antral follicular dynamics ultrasonographically. Thus, at present, cows and mares may be the most appropriate models for

the ultrasonographic study of antral follicular dynamics in women (Adams and Pierson, 1995; Ginther et al., 2004)

Continuous versus wave patterns of follicular recruitment

Discrepancies among studies about the continuous versus cyclic nature of antral follicular recruitment in women may, in part, be attributed to methodological challenges associated with serially monitoring follicles <6 mm. Serial ultrasonographic evaluation of antral follicle growth 2–5 mm is technically difficult due to the large quantity of these follicles present in the ovaries throughout the cycle. Follicle waves were most readily detected when follicles ≥ 5 mm were evaluated; waves were not as obvious when follicles <5 mm were analyzed (Baerwald et al., 2003a). Quantifying the growth of 2–3 mm follicles is near the detection limit of current ultrasound equipment, and there is a great deal of error in detecting these small follicles in women. It is interesting that 1–3 mm follicles have been shown to develop in a wave-like manner in association with surges in plasma concentrations of FSH waves in cattle (Jaiswal et al., 2004). Similar ultrasonographic studies in women have not been conducted. As non-invasive imaging technology improves, future research should be directed at carefully characterizing the dynamics of antral follicles <5 mm during the menstrual cycle. Further characterization of the endocrine and paracrine mechanisms underlying follicular waves in women is also required. Elucidation of a third subtle, but significant, FSH peak and fluctuations in luteal phase estradiol in women with three follicle waves is of particular interest (Baerwald et al., 2003a).

Comprehensive comparisons of data that describe continuous, single and multiple waves of antral follicular recruitment show similar trends. We believe that inconsistencies in descriptions of antral follicular growth are attributable to differences in study designs employed, statistical methods used and interpretations of data. For example, it is plausible that the data from one woman, interpreted as a single recruitment episode (Fig. 2a and b), may actually represent two follicle waves, a major ovulatory wave in the follicular phase followed by a minor anovulatory wave in the luteal phase. No differences in AFC were detected during the cycle in this study; however, AFC was compared at only the early-follicular, late-follicular and luteal phases of the cycle. The grouping of data into three categories versus daily data collection for the entire cycle could have masked changes in the number of antral follicles that occurred. Similarly, when AFC data were grouped into seven different phases of the cycle (Fig. 2c and d), no differences in AFC (2–5 or 2–10 mm) were detected. A non-significant increase in AFC in the early-luteal phase was observed, but was attributed to inter-individual differences and not considered to be biologically relevant. In contrast, significant changes in AFC during the cycle have been reported in other studies (Block, 1951; McNatty et al., 1983; Baerwald et al., 2003b). It therefore appears that evaluation of daily or more frequent changes in AFC, follicle size and/or hormone concentrations throughout the cycle may more readily enable the detection of follicle waves while data grouped into phases of the cycle may not.

Further characterization of the daily changes in AFC during the menstrual cycle has important implications for determining the most appropriate time to consistently measure AFC as a predictor of response to ovarian stimulation (Verhagen et al., 2008). Knowledge

of multiple follicular waves during the menstrual cycle provides a rationale for the notion that ovarian stimulation therapy can be initiated at different times during the cycle. The plausibility of initiating ovarian stimulation therapy during the luteal phase in women is supported by reports that multiple dominant follicles were induced to grow following FSH administration in the early-luteal phase of monkeys (Dizerega and Hodgen, 1980). Rationale for luteal phase ovarian stimulation protocols in women is further provided by knowledge of the routine early-luteal phase initiation of ovarian stimulation in domestic farm animals (Adams *et al.*, 1994b). Synchronizing ovarian stimulation therapy with wave emergence has been shown to optimize follicular responses in both cattle (Nasser *et al.*, 1993) and in women with a history of a poor response to treatment (Baerwald *et al.*, 2010).

Follicle selection in anovulatory waves

It has been generally accepted that follicle selection occurs only once during the menstrual cycle (Gougeon and Lefevre, 1983; Baird, 1990; Pache *et al.*, 1990). However, more recent research suggests that selection may occur more than once in approximately one-quarter of apparently healthy women (Baerwald *et al.*, 2003a). These findings are similar to domestic farm animals; in particular, mares (Ginther, 1993; Ginther *et al.*, 2004). It is plausible that follicular dynamics in women of reproductive age may be more variable than previously thought. The functional status and ovulatory potential of follicles comprising anovulatory follicular waves requires further evaluation, considering earlier evidence that the majority of luteal phase follicles in women were atretic (McNatty *et al.*, 1983). In animals, antral follicle dynamics are influenced by environmental factors such as energy balance, temperature and exposure to endocrine disrupting chemicals (Frisch, 1985; Mackey *et al.*, 2000; Armstrong *et al.*, 2003; De *et al.*, 2008; Kwintkiewicz and Giudice, 2009). However, the effects of environmental factors on human antral folliculogenesis are not well understood.

Role of the corpus luteum in regulating antral follicular development

Observations of multiple follicle waves during the menstrual cycle and the development of more than one dominant follicle in some women have challenged the notion of luteal inhibition of antral follicular development in primates. Therefore, future research should be conducted to confirm species-specific differences in luteal regulation of antral follicular dynamics. Evidence of aromatase activity in the mare CL (Albrecht *et al.*, 1997) provides a rationale for research to confirm whether the CL of other domestic animals produces estradiol and to determine its potential roles in regulating antral folliculogenesis. It is important to clarify whether the estradiol increase that occurs in association with anovulatory waves in women is of follicular or luteal origin. Furthermore, the precise roles of luteal inhibin in regulating antral follicular wave dynamics in women are unknown.

Repeatability in the follicular wave pattern

Knowledge about the repeatability of follicular waves in women is fundamental for designing future studies to elucidate the mechanisms underlying follicular wave dynamics and to predict the number of waves that a woman may develop in a given cycle. The consistency, or lack thereof, in the number of waves that develop per cycle has

clinical implications for optimizing strategies which manipulate ovarian follicular development, such as hormonal contraceptive and ovarian stimulation regimens. Studies in cattle have revealed that the number of follicle waves per estrous cycle is fairly consistent within individuals (Singh *et al.*, 2004; Jaiswal *et al.*, 2009). Similar research in women is ongoing (Baerwald *et al.*, unpublished data).

The pre-pubertal and pubertal period

Human follicular development occurs to the early antral stage during infancy and childhood (Peters *et al.*, 1978). The first few menstrual cycles during puberty may be anovulatory due to incomplete maturation of the hypothalamo-pituitary-axis and the inability to adequately produce LH. In comparison, repeated anovulatory waves of antral follicle development have been documented in pre-pubertal calves, as early as 2 weeks of age (Adams *et al.*, 1994a; Evans *et al.*, 1994). Follicular dynamics in pre-pubertal and pubertal girls remain poorly understood due to ethical and technical constraints on conducting reproductive research in this population. Our understanding of pubertal disorders would be greatly enhanced by the elucidation of ovarian function during the prepubertal and pubertal phases of development.

Reproductive senescence

The age-related decrease in ovarian reserve correlates with a decrease in the number of antral follicles 2–10 mm in diameter (Scheffer *et al.*, 1999). A decrease in the AFC in perimenopausal women results in a reduction in circulating inhibin B, AMH and IGF-I and a correspondent increase in FSH and cycle irregularity; estradiol levels remain the same or are elevated (Lee *et al.*, 1988; Ahmed Ebbyary *et al.*, 1994; Klein *et al.*, 1996a, b, c; Scheffer *et al.*, 1999; Klein *et al.*, 2000; Erdem *et al.*, 2002; Broekmans *et al.*, 2004; Van Rooij *et al.*, 2005; Visser *et al.*, 2006). Continued depletion of the AFC results in a continued decline in inhibin B, increasing FSH and LH, decreasing estradiol and progesterone and loss of menstrual cyclicity culminating in menopause at ~51 years of age [reviewed in Broekmans *et al.* (2009)].

In women, increasing age has been associated with higher mitotic activity of granulosa cells in small antral follicles (Gougeon, 1998), earlier selection of the dominant follicle, a shorter follicular phase, shorter menstrual cycle, anovulation and an increased incidence of dizygotic twinning (Lenton *et al.*, 1984; Klein *et al.*, 1996a, 2002; Beemsterboer *et al.*, 2006; Brodin *et al.*, 2007). Earlier selection of the dominant ovulatory follicle in aging women has been attributed to a faster growth rate of the dominant follicle (Gougeon, 2005) or earlier emergence of the follicular cohort in the luteal phase of the preceding cycle (Van Zonneveld *et al.*, 2003). Similar age-related changes in ovarian function have been reported in mares (Carnevale, 2008) and cows (Malhi *et al.*, 2005, 2006, 2007).

Age-related changes in antral follicle dynamics are not fully understood. Research to characterize age-related changes in follicular wave dynamics may explain the earlier selection of the dominant follicle and shorter follicular phases (described above), as well as abnormal ovulation, estradiol secretion and cyclicity observed in women of advanced reproductive age (Santoro *et al.*, 2003; Hale *et al.*, 2009). It is plausible that women with more follicular waves during their cycles may undergo menopause earlier; however, research to test this hypothesis has not yet been conducted.

Summary

Theories of continuous versus cyclic antral follicle recruitment during the human menstrual cycle have been proposed. There is increasing evidence to suggest that multiple cohorts (also referred to as 'waves') of antral follicles are recruited (i.e. emerge) during the menstrual cycle. Observations of ovarian follicular waves in women are comparable with those documented in several animal species; however, some species-specific differences appear to exist. Enhancing our understanding of the physiologic, biochemical and molecular mechanisms underlying antral follicular wave dynamics is important for understanding age-related changes in reproductive function and will have clinical implications for the development of safer, more efficacious and patient-friendly hormonal contraceptive and ovarian stimulation regimens. Elucidation of endocrine and paracrine regulation of follicular wave dynamics in women is also important for understanding follicle–oocyte interactions and identifying non-invasive markers of the physiologic status of follicles, which are predictive of oocyte competence and assisted reproduction outcomes.

Authors' roles

All of the authors have made substantial contributions to the present review article in the following manner: (i) conception and design, acquisition of data or analysis and interpretation of data, (ii) drafting or revising the article and (iii) final approval of the version to be published.

Acknowledgements

The authors would like to thank Dr Sharon Mortimer for editorial assistance and Lindsey Servinski for graphic design used in the preparation of this manuscript.

Funding

This work is supported through grants from the following agencies/institutions: University of Saskatchewan, Canadian Institutes of Health Research, Natural Sciences and Engineering Council of Canada, Canadian Foundation for Women's Health and the Saskatchewan Health Research Foundation.

Conflict of interest

The authors have no conflicts of interest in publishing this article.

References

- Adams GP. Comparative patterns of follicle development and selection in ruminants. *J Reprod Fertil Suppl* 1999;**54**:17–32.
- Adams GP, Pierson RA. Bovine model for study of ovarian follicular dynamics in humans. *Therio* 1995;**43**:113–120.
- Adams GP, Sumar J, Ginther OJ. Effects of lactational and reproductive status on ovarian follicular waves in llamas. *J Reprod Fertil* 1990;**90**:535–545.
- Adams GP, Matteri RL, Kastelic JP, Ko JC, Ginther OJ. Association between surges of follicle-stimulating hormone and the emergence of follicular waves in heifers. *J Reprod Fertil* 1992a;**94**:177–188.
- Adams GP, Matteri RL, Kastelic JP, Ko JCH, Ginther OJ. Association between surges of follicle-stimulating hormone and the emergence of follicular waves in heifers. *J Reprod Fertil* 1992b;**94**:177–188.
- Adams GP, Kot K, Smith CA, Ginther OJ. Selection of a dominant follicle and suppression of follicular growth in heifers. *Anim Reprod Sci* 1993;**30**:259–271.
- Adams GP, Evans AC, Rawlings NC. Follicular waves and circulating gonadotrophins in 8-month-old prepubertal heifers. *J Reprod Fertil* 1994a;**100**:27–33.
- Adams GP, Nasser LF, Bo GA, Garcia A, Del Campo MR, Mapletoft RJ. Superovulatory response of ovarian follicles of wave 1 versus wave 2 in heifers. *Therio* 1994b;**42**:1103–1113.
- Adams GP, Jaiswal R, Singh J, Malhi P. Progress in understanding ovarian follicular dynamics in cattle. *Therio* 2008;**69**:72–80.
- Adashi EY. Endocrinology of the ovary. *Hum Reprod* 1994;**9**:815–827.
- Aerts JM, Bols PE. Ovarian follicular dynamics: a review with emphasis on the bovine species. Part I: folliculogenesis and pre-antral follicle development. *Reprod Dom Anim* 2010;**45**:171–179.
- Ahmed Ebbyary NA, Lenton EA, Cooke ID. Hypothalamic-pituitary ageing: progressive increase in FSH and LH concentrations throughout the reproductive life in regularly menstruating women. *Clin Endocrinol* 1994;**41**:199–206.
- Albrecht BA, Macleod JN, Daels PF. Differential transcription of steroidogenic enzymes in the equine primary corpus luteum during diestrus and early pregnancy. *Biol Reprod* 1997;**56**:821–829.
- Andersen CY, Schmidt KT, Kristensen SG, Rosendahl M, Byskov AG, Ernst E. Concentrations of AMH and Inhibin-B in relation to follicular diameter in normal human small antral follicles. *Hum Reprod* 2010;**25**:1282–1287.
- Andreotti RF, Thompson GH, Janowitz W, Shapiro AG, Zusmer NR. Endovaginal and transabdominal sonography of ovarian follicles. *J Ultrasound Med* 1989;**8**:555–560.
- Araki S, Chikazawa K, Akabori A, Ijima K, Tamada T. Hormonal profile after removal of the dominant follicle and corpus luteum in women. *Endocrinol Japon* 1983;**30**:55–70.
- Armstrong DG, Gong JG, Webb R. Interactions between nutrition and ovarian activity in cattle: physiological, cellular and molecular mechanisms. *Reproduction Suppl* 2003;**61**:403–414.
- Asher GW, Scott IC, O'Neill KT, Smith JF, Inskip EK, Townsend EC. Ultrasonographic monitoring of antral follicle development in red deer (*Cervus elaphus*). *J Reprod Fertil* 1997;**111**:91–99.
- Baerwald A. Human antral folliculogenesis: what we have learned from the bovine and equine models. *Anim Reprod* 2008;**6**:20–29.
- Baerwald A, Adams G, Pierson R. Characteristics of ovarian follicular wave dynamics in women. *Biol Reprod* 2003a;**69**:1023–1031.
- Baerwald A, Adams G, Pierson R. A new model for ovarian follicular development during the human menstrual cycle. *Fertil Steril* 2003b;**80**:116–122.
- Baerwald AR, Adams GP, Pierson RA. Form and function of the corpus luteum during the human menstrual cycle. *Ultrasound Obstet Gynecol* 2005;**25**:498–507.
- Baerwald AR, Walker RA, Pierson RA. Growth rates of ovarian follicles during natural menstrual cycles, oral contraception cycles, and ovarian stimulation cycles. *Fertil Steril* 2009;**91**:440–449.
- Baerwald A, Anderson P, Mcvee J, Case A, Yuzpe A, Fluker M. Synchronization of ovarian stimulation with follicle wave emergence in patients with a history of a poor response to treatment. In: *Proceedings of the 56th Annual Meeting of the Canadian Fertility and Andrology Society*. Vancouver, BC, 2010. Abstract: TC04.
- Baird DT. Factors regulating the growth of the preovulatory follicle in the sheep and human. *J Reprod Fertil* 1983;**69**:343–352.
- Baird D. A model for follicular selection and ovulation: lessons from superovulation. *J Ster Biochem* 1987;**27**:15–23.
- Baird D. The selection of the follicle of the month. In Evers J, Heineman M (eds). *From Ovulation to Implantation Proceedings of the VII Regnier de Graaf Symposium*. Maastricht, the Netherlands: Excerpta Medica, 1990.
- Baird D, Fraser I. Blood production and ovarian secretion rates of estradiol and estrone in women throughout the menstrual cycle. *J Clin Endocrinol Metab* 1974;**38**:1009–1017.
- Baird D, Fraser IS. Concentration of oestrone and oestradiol in follicular fluid and ovarian venous blood of women. *Clin Endocrinol* 1975;**4**:259–266.
- Baird D, McCracken J, Goding J. Studies in steroid synthesis and secretion with the autotransplanted sheep ovary and adrenal. In: Pieerpoint C (ed). *The Endocrinology of Pregnancy and Parturition*. Cardiff: Alpha Omega Alpha Publishing Co., 1973.

- Baird DT, Baker TG, McNatty KP, Neal P. Relationship between the secretion of the corpus luteum and the length of the follicular phase of the ovarian cycle. *J Reprod Fertil* 1975;**45**:611–619.
- Baird DT, Backstrom T, Mcneilly AS, Smith SK, Wathen AG. Effect of enucleation of the corpus luteum at different stages of the luteal phase of the human menstrual cycle on subsequent follicular development. *J Reprod Fertil* 1984;**70**:615–624.
- Baker T. A quantitative and cytological study of germ cells in human ovaries. *Proc R Soc Lond [Biol]* 1963;**158**:417–433.
- Bakos O, Lundkvist O, Wide L, Bergh T. Ultrasonographical and hormonal description of the normal ovulatory menstrual cycle. *Acta Obstet Gynecol Scand* 1994;**73**:790–796.
- Bartlewski PM, Beard AP, Cook SJ, Chandolia RK, Honaramooz A, Rawlings NC. Ovarian antral follicular dynamics and their relationships with endocrine variables throughout the oestrous cycle in breeds of sheep differing in prolificacy. *J Reprod Fertil* 1999;**115**:111–124.
- Basseti SG, Winters SJ, Keeping HS, Zeleznik AJ. Serum immunoreactive inhibin levels before and after luteectomy in the cynomolgus monkey (*Macaca fascicularis*). *J Clin Endocrinol Metab* 1990;**70**:590–594.
- Beemsterboer SN, Homburg R, Gorter NA, Schats R, Hompes PG, Lambalk CB. The paradox of declining fertility but increasing twinning rates with advancing maternal age. *Hum Reprod* 2006;**21**:1531–1532.
- Beg MA, Ginther OJ. Follicle selection in cattle and horses: role of intrafollicular factors. *Reproduction* 2006;**132**:365–377.
- Bentov Y, Efsandiari N, Gokturk A, Burstein E, Fainaru O, Casper RF. An ongoing pregnancy from two waves of follicles developing during a long follicular phase of the same cycle. *Fertil Steril* 2010;**94**:350 e8–11.
- Bilezikjian LM, Blount AL, Leal AM, Donaldson CJ, Fischer WH, Vale WW. Autocrine/paracrine regulation of pituitary function by activin, inhibin and follistatin. *Mol Cell Endocrinol* 2004;**225**:29–36.
- Bishop CV, Sparman ML, Stanley JE, Bahar A, Zelinski MB, Stouffer RL. Evaluation of antral follicle growth in the macaque ovary during the menstrual cycle and controlled ovarian stimulation by high-resolution ultrasonography. *Amer J Primatol* 2009;**71**:384–392.
- Bjersing L. On the morphology and endocrine function of granulosa cells of ovarian follicles and corpora lutea. *Acta Endocrinol* 1967;**125**:5.
- Block E. Quantitative morphological investigations of the follicular system in women: variations in the different phases of the sexual cycle. *Acta Endocrinol* 1951;**8**:33–54.
- Block E. Quantitative morphological investigations of the follicular system in women: variations at different ages. *Acta Anat* 1952;**14**:108–123.
- Bodensteiner KJ, Wiltbank MC, Bergfelt DR, Ginther OJ. Alterations in follicular estradiol and gonadotropin receptors during development of bovine antral follicles. *Therio* 1996;**45**:499–512.
- Bomsel-Helmreich O. Ultrasound and the preovulatory human follicle. *Oxford Rev Reprod Biol* 1985;**7**:1–72.
- Bomsel-Helmreich O, Gougeon A, Thebault A, Satarelli D, Milgrom E, Frydman R, Papiernik E. Healthy and atretic human follicles in the preovulatory phase: differences in evolution of follicular morphology and steroid content of follicular fluid. *J Clin Endocrinol Metab* 1979;**48**:686–694.
- Brailly S, Gougeon A, Milgrom E, Bomsel-Helmreich O, Papiernik E. Androgens and progestins in the human ovarian follicle: differences in the evolution of preovulatory, healthy nonovulatory, and atretic follicles. *J Clin Endocrinol Metab* 1981;**53**:128–134.
- Brodin T, Bergh T, Berglund L, Hadziosmanovic N, Holte J. Menstrual cycle length is an age-independent marker of female fertility: results from 6271 treatment cycles of *in vitro* fertilization. *Fertil Steril* 2007;**90**:1656–1661.
- Broekmans FJ, Faddy MJ, Scheffer G, Te Velde ER. Antral follicle counts are related to age at natural fertility loss and age at menopause. *Menopause* 2004;**11**:607–14.
- Broekmans FJ, Soules MR, Fauser BC. Ovarian aging: mechanisms and clinical consequences. *Endocr Rev* 2009;**30**:465–93.
- Brown JB. Pituitary control of ovarian function: concepts derived from gonadotrophin therapy Austr NZ. *J Obstet Gynaecol* 1978;**18**:47–54.
- Burns DS, Jimenez-Krassel F, Ireland JL, Knight PG, Ireland JJ. Numbers of antral follicles during follicular waves in cattle: evidence for high variation among animals, very high repeatability in individuals, and an inverse association with serum follicle-stimulating hormone concentrations. *Biol Reprod* 2005;**73**:54–62.
- Carnevale EM. The mare model for follicular maturation and reproductive aging in the woman. *Therio* 2008;**69**:23–30.
- Check J, Dietterich C, Houck MA. Ipsilateral versus contralateral ovary selection of dominant follicle in succeeding cycle. *Obstet Gynecol* 1991;**77**:247–249.
- Chikazawa K, Araki S, Tamada T. Morphological and endocrinological studies on follicular development during the human menstrual cycle. *J Clin Endocrinol Metab* 1986;**62**:305–313.
- Cobb M. *Generation: The Seventeenth Century Scientists Who Unraveled the Secrets of Sex, Life and Growth*. New York: Bloomsbury Publishing, 2006.
- Craig J, Orisaka M, Wang H, Orisaka S, Thompson W, Zhu C, Kotsuji F, Tsang BK. Gonadotropin and intra-ovarian signals regulating follicle development and atresia: the delicate balance between life and death. *Front Biosci* 2007;**12**:3628–3639.
- De ST-JJR, De FaPM, De Sa WF, De MFA, Viana JH, Camargo LS, Ramos AA, Folhadella IM, Polisseni J, De Freitas C et al. Effect of maternal heat-stress on follicular growth and oocyte competence in *Bos indicus* cattle. *Therio* 2008;**69**:155–166.
- De Souza CJ, Campbell BK, Baird DT. Incipient ovarian failure associated with raised levels of follicle stimulating hormone and reduced levels of inhibin A in older sheep. *Hum Reprod* 1998;**13**:3016–3022.
- Demirtas E, Elizur SE, Holzer H, Gidoni Y, Son WY, Chian RC, Tan SL. Immature oocyte retrieval in the luteal phase to preserve fertility in cancer patients. *Reprod Biomed Online* 2008;**17**:520–523.
- Dervain I. *Etude echographique de la croissance du follicle ovarien normal et detection de l'ovulation*. These de medicine: Universite Louis Pasteur, Strasbourg, 1980.
- Devoto L, Fuentes A, Kohen P, Cespedes P, Palomino A, Pommer R, Munoz A, Strauss JF III. The human corpus luteum: life cycle and function in natural cycles. *Fertil Steril* 2008;**92**:1067–1079.
- Dizerega GS, Hodgen GD. Cessation of folliculogenesis during the primate luteal phase. *J Clin Endocrinol Metab* 1980;**51**:158–160.
- Dizerega G, Hodgen G. Folliculogenesis in the primate ovarian cycle. *Endocr Rev* 1981;**2**:27–49.
- Doody MC, Gibbons WE, Zamah NM. Linear regression analysis of ultrasound follicular growth series: statistical relationship of growth rate and calculated date of growth onset to total growth period. *Fertil Steril* 1987;**47**:436–440.
- Douglas RH, Ginther OJ. Route of prostaglandin F2 alpha injection and luteolysis in mares (38519). *Proc Soc Exper Biol Med Soc Exp Bio Med* 1975;**148**:263–269.
- Duggavathi R, Murphy BD. Development. Ovulation signals. *Science* 2009;**324**:890–891.
- Durlinger AL, Visser JA, Themmen AP. Regulation of ovarian function: the role of anti-Mullerian hormone. *Reproduction* 2002;**124**:601–609.
- Ecochard R, Gougeon A. Side of ovulation and cycle characteristics in normally fertile women. *Hum Reprod* 2000;**15**:752–755.
- Eden JA, Jones J, Carter GD, Alagband-Zadeh J. A comparison of follicular fluid levels of insulin-like growth factor-I in normal dominant and cohort follicles, polycystic and multicystic ovaries. *Clin Endocrinol* 1988;**29**:327–336.
- El-Roeiy A, Chen X, Roberts VJ, Leroith D, Roberts CT Jr, Yen SS. Expression of insulin-like growth factor-I (IGF-I) and IGF-II and the IGF-I, IGF-II, and insulin receptor genes and localization of the gene products in the human ovary. *J Clin Endocrinol Metab* 1993;**77**:1411–1418.
- Eppig J. Oocyte control of ovarian follicular development and function in mammals. *Reproduction* 2001;**122**:829–838.
- Erdem A, Erdem M, Biberoglu K, Hayit O, Arslan M, Gursoy R. Age-related changes in ovarian volume, antral follicle counts and basal FSH in women with normal reproductive health. *J Reprod Med* 2002;**47**:835–839.
- Evans AC, Adams GP, Rawlings NC. Follicular and hormonal development in prepubertal heifers from 2 to 36 weeks of age. *J Reprod Fertil* 1994;**102**:463–470.
- Evans AC, Duffy P, Hynes N, Boland MP. Waves of follicle development during the estrous cycle in sheep. *Therio* 2000;**53**:699–715.
- Fausser B, Van Heusden A. Manipulation of human ovarian function: physiological concepts and clinical consequences. *Endocr Rev* 1997;**18**:71–106.
- Filicori M. The potential value of mid-follicular phase LH. *Hum Reprod* 2002;**17**:517–520.
- Filicori M, Cognigni GE, Tabarelli C, Pocognoli P, Taraborrelli S, Spettoli D, Ciampaglia W. Stimulation and growth of antral ovarian follicles by selective LH activity administration in women. *J Clin Endocrinol Metab* 2002;**87**:1156–1161.
- Findlay JK. An update on the roles of inhibin, activin, and follistatin as local regulators of folliculogenesis. *Biol Reprod* 1993;**48**:15–23.
- Fortune JE. The early stages of follicular development: activation of primordial follicles and growth of pre-antral follicles. *Anim Reprod Sci* 2003;**78**:135–163.

- Fortune JE, Rivera GM, Yang MY. Follicular development: the role of the follicular microenvironment in selection of the dominant follicle. *Anim Reprod Sci* 2004; **82-83**:109–126.
- Fraser HM, Groome NP, Mcneilly AS. Follicle-stimulating hormone-inhibin B interactions during the follicular phase of the primate menstrual cycle revealed by gonadotropin-releasing hormone antagonist and antiestrogen treatment. *J Clin Endocrinol Metab* 1999; **84**:1365–1369.
- Frisch RE. Fatness, menarche, and female fertility. *Perspect Biol Med* 1985; **28**:611–633.
- Fukuda M, Fukuda K, Anderson CY, Byskov AG. Contralateral selection of dominant follicle favours pre-embryo development. *Hum Reprod* 1996; **11**:1958–1962.
- Gastal EL, Gastal MO, Wiltbank MC, Ginther OJ. Follicle deviation and intrafollicular and systemic estradiol concentrations in mares. *Biol Reprod* 1999; **61**:31–39.
- Gastal EL, Gastal MO, Nogueira GP, Bergfelt DR, Ginther OJ. Temporal interrelationships among luteolysis, FSH and LH concentrations and follicle deviation in mares. *Therio* 2000; **53**:925–940.
- Geisthovel F, Moretti-Rojas I, Asch RH, Rojas FJ. Expression of insulin-like growth factor-II (IGF-II) messenger ribonucleic acid (mRNA), but not IGF-I mRNA, in human preovulatory granulosa cells. *Hum Reprod* 1989; **4**:899–902.
- Gibbons JR, Wiltbank MC, Ginther OJ. Functional interrelationships between follicles greater than 4 mm and the follicle-stimulating hormone surge in heifers. *Biol Reprod* 1997; **57**:1066–1073.
- Ginther OJ. Major and minor follicular waves during the equine estrous cycle. *J Equine Veter Sci* 1993; **13**:18–25.
- Ginther OJ. Selection of the dominant follicle in cattle and horses. *Anim Reprod Sci* 2000; **60-61**:61–79.
- Ginther OJ, Kastelic JP, Knopf L. Composition and characteristics of follicular waves during the bovine estrous cycle. *Anim Reprod Sci* 1989a; **20**:187–200.
- Ginther OJ, Knopf L, Kastelic JP. Temporal associations among ovarian events in cattle during oestrous cycles with two and three follicular waves. *J Reprod Fertil* 1989b; **87**:223–230.
- Ginther OJ, Kot K, Wiltbank MC. Associations between emergence of follicular waves and fluctuations in FSH concentrations during the estrous cycle in ewes. *Therio* 1995; **43**:689–703.
- Ginther OJ, Kot K, Kulick LJ, Wiltbank MC. Emergence and deviation of follicles during the development of follicular waves in cattle. *Therio* 1997; **48**:75–87.
- Ginther OJ, Bergfelt DR, Kulick LJ, Kot K. Pulsatility of systemic FSH and LH concentrations during follicular-wave development in cattle. *Therio* 1998; **50**:507–519.
- Ginther OJ, Bergfelt DR, Kulick LJ, Kot K. Selection of the dominant follicle in cattle: role of estradiol. *Biol Reprod* 2000a; **63**:383–389.
- Ginther OJ, Bergfelt DR, Kulick LJ, Kot K. Selection of the dominant follicle in cattle: role of two-way functional coupling between follicle-stimulating hormone and the follicles. *Biol Reprod* 2000b; **62**:920–927.
- Ginther OJ, Beg MA, Bergfelt DR, Danadeu FX, Kot K. Follicle selection in monovular species. *Biol Reprod* 2001a; **65**:638–647.
- Ginther OJ, Bergfelt DR, Beg MA, Kot K. Effect of LH on circulating oestradiol and follicular fluid factor concentrations during follicle deviation in cattle. *Reproduction* 2001b; **122**:103–110.
- Ginther OJ, Bergfelt DR, Beg MA, Kot K. Follicle selection in cattle: role of luteinizing hormone. *Biol Reprod* 2001c; **64**:197–205.
- Ginther OJ, Gastal EL, Gastal MO, Bergfelt DR, Baerwald AR, Pierson RA. Comparative study of the dynamics of follicular waves in mares and women. *Biol Reprod* 2004; **71**:1195–1201.
- Ginther OJ, Beg MA, Gastal EL, Gastal MO, Baerwald AR, Pierson RA. Systemic concentrations of hormones during the development of follicular waves in mares and women: a comparative study. *Reproduction* 2005; **130**:379–388.
- Giudice LC. Insulin-like growth factor family in Graafian follicle development and function. *J Soc Gynecol Investig* 2001; **8**:S26–S29.
- Gore M, Nayudu P, Vlasisavljevic V, Thomas N. Dominance is not what it seems: individual identification of ovarian follicles during the follicular phase. In: *Proceedings of the 10th Annual Meeting of the ESHRE*. Brussels, 1994, p.43.
- Gore MA, Nayudu PL, Vlasisavljevic V, Thomas N. Ovarian follicle dynamics *in vivo*: dominant follicle position and the subdominant follicle population dynamics. *Biol Reprod* 1995; **52**:81.
- Gore M, Nahudu P, Vlasisavljevic V. Attaining dominance *in vivo*: distinguishing dominant from challenger follicles in humans. *Hum Reprod* 1997; **12**:2741–2747.
- Gougeon A. Steroid 3 β -ol-dehydrogenase activity in the largest healthy and atretic follicles in the human ovary during the menstrual cycle. *Ann Biol Anim Bioch Biophys* 1977; **17**:1095–1099.
- Gougeon A. Qualitative changes in medium and large antral follicles in the human ovary during the menstrual cycle. *Annals Biol Anim Bioch Biophys* 1979; **19**:1464–1468.
- Gougeon A. Influence of cyclic variations in gonadotrophin and steroid hormones on follicular growth in the human ovary. In: De Brux J, Gautray J (eds). *Clinical Pathology of the Endocrine Ovary*. Lancaster, UK: MTP Press, 1984.
- Gougeon A. Dynamics of follicular growth in the human: a model from preliminary results. *Hum Reprod* 1986; **1**:81–87.
- Gougeon A. Regulation of ovarian follicular development in primates: facts and hypotheses. *Endocr Rev* 1996; **17**:121–155.
- Gougeon A. Ovarian follicular growth in humans: ovarian ageing and population of growing follicles. *Maturitas* 1998; **30**:137–142.
- Gougeon A. Dynamics of human follicular growth: morphologic, dynamic, and functional aspects. In: Leung P, Adashi E (eds). *The Ovary*. 2nd edn. Amsterdam: Elsevier Academic Press, 2004.
- Gougeon A. The biological aspects of risks of infertility due to age: the female side. *Revue d'epidemiologie et de sante publique* 2005; **53** Spec No 2:2537–2545.
- Gougeon A. Human ovarian follicular development: from activation of resting follicles to preovulatory maturation. *Ann d'endocrinol* 2010; **71**:132–143.
- Gougeon A, Lefevre B. Evolution of the diameters of the largest healthy and atretic follicles during the human menstrual cycle. *J Reprod Fertil* 1983; **69**:497–502.
- Gougeon A, Echochard R, Thalbard JC. Age-related changes of the population of human ovarian follicles: increase in the disappearance rate of non-growing and early-growing follicles in aging women. *Biol Reprod* 1994; **50**:653–663.
- Green SH, Zuckerman S. The number of oocytes in the mature rhesus monkey (*Macaca mulatta*). *J Endocrinol* 1951; **7**:194–202.
- Grondahl ML, Nielsen ME, Dal Canto MB, Fadini R, Rasmussen IA, Westergaard LG, Kristensen SG, Yding Andersen C. Anti-Mullerian hormone remains highly expressed in human cumulus cells during the final stages of folliculogenesis. *Reprod Biomed Online* 2011; **22**:389–398.
- Groome NP, Illingworth PJ, O'Brien M, Pai R, Rodger FE, Mather JP, Mcneilly AS. Measurement of dimeric inhibin B throughout the human menstrual cycle. *J Clin Endocrinol Metab* 1996; **81**:1401–1405.
- Gueripel X, Brun V, Gougeon A. Oocyte bone morphogenetic protein 15, but not growth differentiation factor 9, is increased during gonadotropin-induced follicular development in the immature mouse and is associated with cumulus oophorus expansion. *Biol Reprod* 2006; **75**:836–843.
- Hackeloer B, Robinson H. Ultraschalldarstellung des wachsenden follikels und corpus luteum im normalen physiologischen zyklus [author's translation: ultrasound examination of the growing ovarian follicle and of the corpus luteum during the normal physiologic menstrual cycle]. *Geburtshilfe Frauenheilkd* 1978; **38**:163–168.
- Hackeloer BJ, Fleming R, Robinson HP, Adam AH, Coutts JRT. Correlation of ultrasonic and endocrinologic assessment of human follicular development. *Amer J Obstet Gynecol* 1979; **135**:122–128.
- Hale GE, Zhao X, Hughes CL, Burger HG, Robertson DM, Fraser IS. Endocrine features of menstrual cycles in middle and late reproductive age and the menopausal transition classified according to the staging of reproductive aging workshop (STRAW) staging system. *J Clin Endocrinol Metab* 2007; **92**:3060–3067.
- Hale GE, Hughes CL, Burger HG, Robertson DM, Fraser IS. Atypical estradiol secretion and ovulation patterns caused by luteal out-of-phase (LOOP) events underlying irregular ovulatory menstrual cycles in the menopausal transition. *Menopause* 2009; **16**:50–59.
- Hall DA, Hann LE, Ferrucci JT Jr, Black EB, Braitman BS, Crowley WF, Nikrul N, Kelley JA. Sonographic morphology of the normal menstrual cycle. *Radiology* 1979; **133**:185–188.
- Hansen KR, Knowlton NS, Thyer AC, Charleston JS, Soules MR, Klein NA. A new model of reproductive aging: the decline in ovarian non-growing follicle number from birth to menopause. *Hum Reprod* 2008; **23**:699–708.
- Hernandez ER, Hurwitz A, Vera A, Pellicer A, Adashi EY, Leroith D, Roberts CT Jr. Expression of the genes encoding the insulin-like growth factors and their receptors in the human ovary. *J Clin Endocrinol Metab* 1992; **74**:419–425.
- Hillier SG, Reichert LE Jr, Van Hall EV. Control of preovulatory follicular estrogen biosynthesis in the human ovary. *J Clin Endocrinol Metab* 1981; **52**:847–856.

- Hillier S, Yong E, Illingworth P, Baird D, Schwall R, Mason A. Effect of recombinant activin on androgen synthesis in cultured human thecal cells. *J Clin Endocrinol Metab* 1991a;**72**:1206–1211.
- Hillier SG, Yong EL, Illingworth PJ, Baird DT, Schwall RH, Mason AJ. Effect of recombinant inhibin on androgen synthesis in cultured human thecal cells. *Mol Cell Endocrinol* 1991b;**75**:R1–R6.
- Hoare EK, Parker SE, Flood PF, Adams GP. Ultrasonic imaging of reproductive events in muskoxen. *Rangifer* 1997;**17**:116–123.
- Hodgen G. The dominant ovarian follicle. *Fertil Steril* 1982;**38**:281–300.
- Hourvitz A, Widger AE, Filho FL, Chang RJ, Adashi EY, Erickson GF. Pregnancy-associated plasma protein-A gene expression in human ovaries is restricted to healthy follicles and corpora lutea. *J Clin Endocrinol Metab* 2000;**85**:4916–4920.
- Hsueh AJ, Dahl KD, Vaughan J, Tucker E, Rivier J, Bardin CW, Vale W. Heterodimers and homodimers of inhibin subunits have different paracrine action in the modulation of luteinizing hormone-stimulated androgen biosynthesis. *Proc Natl Acad Sci* 1987;**84**:5082–5086.
- Iwashita M, Kudo Y, Yoshimura Y, Adachi T, Katayama E, Takeda Y. Physiological role of insulin-like-growth-factor-binding protein-4 in human folliculogenesis. *Hormone Res* 1996;**46**:31–36.
- Jaiswal RS, Singh J, Adams GP. Developmental pattern of small antral follicles in the bovine ovary. *Biol Reprod* 2004;**71**:1244–1251.
- Jaiswal R, Singh J, Marshall L, Adams G. Repeatability of 2- and 3- wave patterns during the interovulatory interval in cattle. *Therio* 2009;**72**:81–90.
- Jesionowska H, Hemmings R, Guyda HJ, Posner BI. Determination of insulin and insulin-like growth factors in the ovarian circulation. *Fertil Steril* 1990;**53**:88–91.
- Jokubkiene L, Sladkevicius P, Rovas L, Valentini L. Assessment of changes in volume and vascularity of the ovaries during the normal menstrual cycle using three-dimensional power Doppler ultrasound. *Hum Reprod* 2006;**21**:2661–2668.
- Juengel JL, Hudson NL, Heath DA, Smith P, Reader KL, Lawrence SB, O'Connell AR, Laitinen MP, Cranfield M, Groome NP et al. Growth differentiation factor 9 and bone morphogenetic protein 15 are essential for ovarian follicular development in sheep. *Biol Reprod* 2002;**67**:1777–1789.
- Kaltenbach CC, Cook B, Niswender GD, Nalbandov AV. Effect of pituitary hormones on progesterone synthesis by ovine luteal tissue *in vitro*. *Endocrinology* 1967;**81**:1407–1409.
- Kerin J, Edmonds D, Warnes G, Cox L, Seamark R, Mathews C, Young G, Baird D. Morphological and functional relations of Graafian follicle growth to ovulation in women using ultrasonic, laparoscopic and biochemical measurements. *Br J Obstet Gynaecol* 1981;**88**:81–90.
- Klein NA, Battaglia DE, Fujimoto VY, Davis GS, Bremner WJ, Soules MR. Reproductive aging: accelerated ovarian follicular development associated with a monotropic follicle-stimulating hormone rise in normal older women. *J Clin Endocrinol Metab* 1996a;**81**:1038–1045.
- Klein NA, Battaglia DE, Miller PB, Soules MR. Circulating levels of growth hormone, insulin-like growth factor-I and growth hormone binding protein in normal women of advanced reproductive age. *Clin Endocrinol* 1996b;**44**:285–292.
- Klein NA, Illingworth PJ, Groome NP, Mcneilly AS, Battaglia DE, Soules MR. Decreased inhibin B secretion is associated with the monotropic FSH rise in older, ovulatory women: a study of serum and follicular fluid levels of dimeric inhibin A and B in spontaneous menstrual cycles. *J Clin Endocrinol Metab* 1996c;**81**:2742–2745.
- Klein NA, Battaglia DE, Woodruff TK, Padmanabhan V, Giudice LC, Bremner WJ, Soules MR. Ovarian follicular concentrations of activin, follistatin, inhibin, insulin-like growth factor I (IGF-I), IGF-II, and IGF-binding protein-2 (IGFBP-2), IGFBP-3, and vascular endothelial growth factor in spontaneous menstrual cycles of normal women of advanced reproductive age. *J Clin Endocrinol Metab* 2000;**85**:4520–4525.
- Klein NA, Harper AJ, Houmar BS, Sluss PM, Soules MR. Is the short follicular phase in older women secondary to advanced or accelerated dominant follicle development? *J Clin Endocrinol Metab* 2002;**87**:5746–5750.
- Knight PG, Glister C. TGF-Beta superfamily members and ovarian follicle development. *Reproduction* 2006;**132**:191–206.
- Knopf L, Kastelic JP, Schallenger E, Ginther OJ. Ovarian follicular dynamics in heifers: test of two-wave hypothesis by ultrasonically monitoring individual follicles. *Domest Anim Endocrinol* 1989;**6**:111–119.
- Koering MJ. Cyclic changes in ovarian morphology during the menstrual cycle in *Macaca mulatta* Amer. *J Anat* 1969;**126**:73–101.
- Kwintkiewicz J, Giudice LC. The interplay of insulin-like growth factors, gonadotropins, and endocrine disruptors in ovarian follicular development and function. *Sem Reprod Med* 2009;**27**:43–51.
- Laven JS, Fauser BC. Inhibins and adult ovarian function. *Mol Cell Endocrinol* 2004;**225**:37–44.
- Le Nestour E, Marraoui J, Lahlou N, Roger M, De Ziegler D, Bouchard P. Role of estradiol in the rise in follicle-stimulating hormone levels during the luteal-follicular transition. *J Clin Endocrinol Metab* 1993;**77**:439–442.
- Lee J, Lenotn EA, Sexton L, Cooke IE. The effect of age on the cyclical patterns of plasma LH, FSH, oestradiol and progesterone in women with regular menstrual cycles. *Hum Reprod* 1988;**3**:851–855.
- Lenton EA, Landgren BM, Sexton L, Harper R. Normal variation in the length of the follicular phase of the menstrual cycle: effect of chronological age. *Br J Obstet Gynaecol* 1984;**91**:681–684.
- Lenz S. Ultrasonic study of follicular maturation, ovulation and development of corpus luteum during normal menstrual cycles. *Acta Obstet Gynecol Scand* 1985;**64**:15–19.
- Li R, Phillips DM, Mather JP. Activin promotes ovarian follicle development *in vitro*. *Endocrinology* 1995;**136**:849–856.
- Lintern-Moore S, Peters H, Moore G, Faber M. Follicular development in the infant human ovary. *J Reprod Fertil* 1974;**9**:53–64.
- Lloyd CW, Lobotsky J, Weisz J, Baird DT, McCracken JA, Pupkin M, Zanartu J, Puga J. Concentration of unconjugated estrogens, androgens and gestagens in ovarian and peripheral venous plasma of women: the normal menstrual cycle. *J Clin Endocrinol Metab* 1971;**32**:155–166.
- Lujan M, Dierschke D, Pierson R, Adams G. Ultrasonographic characterization of follicle and luteal dynamics during the menstrual cycle in rhesus monkeys. In: *Proceedings of the Annual Meeting of the Canadian Fertility and Andrology Society*. Ottawa, Ontario, 2006.
- Mackey DR, Wylie AR, Sreenan JM, Roche JF, Diskin MG. The effect of acute nutritional change on follicle wave turnover, gonadotropin, and steroid concentration in beef heifers. *J Anim Sci* 2000;**78**:429–442.
- Magoffin DA, Jakimiuk AJ. Inhibin A, inhibin B and activin A in the follicular fluid of regularly cycling women. *Hum Reprod* 1997;**12**:1714–1719.
- Malhi PS, Adams GP, Singh J. Bovine model for the study of reproductive aging in women: follicular, luteal, and endocrine characteristics. *Biol Reprod* 2005;**73**:45–53.
- Malhi PS, Adams GP, Pierson RA, Singh J. Bovine model of reproductive aging: response to ovarian synchronization and superstimulation. *Therio* 2006;**66**:1257–1266.
- Malhi PS, Adams GP, Mapletoft RJ, Singh J. Oocyte developmental competence in a bovine model of reproductive aging. *Reproduction* 2007;**134**:233–239.
- Maman E, Meirou D, Brengauz M, Raanani H, Dor J, Hourvitz A. Luteal phase oocyte retrieval and *in vitro* maturation is an optional procedure for urgent fertility preservation. *Fertil Steril* 2011;**95**:64–67.
- Mandle A, Zuckerman S. Numbers of normal and atretic oocytes in unilaterally spayed rats. *J Endocrinol* 1950;**7**:194–202.
- Mango D, Scirpa P, Spina MA, Battaglia F, Tartaglia E, Manna P, Ricci S, Iacona T. Ultrasonic and endocrinologic relationships in spontaneous and induced follicular phase. *J Endocrinol Invest* 1988;**11**:7–13.
- Martinuk S, Chizen D, Pierson R. Ultrasonographic morphology of the human preovulatory follicle wall prior to ovulation. *Clinical Amer* 1992;**5**:339–352.
- Mason HD, Cwyfan-Hughes S, Holly JM, Franks S. Potent inhibition of human ovarian steroidogenesis by insulin-like growth factor binding protein-4 (IGFBP-4). *J Clin Endocrinol Metab* 1998;**83**:284–287.
- Mccorkell R, Woodbury M, Adams GP. Ovarian follicular and luteal dynamics in wapi during the estrous cycle. *Therio* 2006;**65**:540–556.
- Mclachlan RI, Robertson DM, Healy DL, Burger HG, De Kretser DM. Circulating immunoreactive inhibin levels during the normal human menstrual cycle. *J Clin Endocrinol Metab* 1987;**65**:954–961.
- McNatty KP. Cyclic changes in antral fluid hormone concentrations in humans. *Clinics Endocrinol Metab* 1978;**7**:577–600.
- McNatty KP. Hormonal correlates of follicular development in the human ovary. *Austral J Biol Sci* 1981a;**34**:249–468.
- McNatty KP. The Regnier de Graaf Lecture: Ovarian follicular development from the onset of luteal regression in humans and sheep. In Rolland R, Van Hall EV, Hillier SG, McNatty KP, Shoemaker J (eds). *Proceedings of the 14th Regnier de Graaf Symposium: 'Follicular Maturation and Ovulation'*, Mijmegen, 1981b:1–18.

- McNatty KP, Baird DT, Bolton A, Chambers P, Corker CS, Mclean H. Concentration of oestrogens and androgens in human ovarian venous plasma and follicular fluid throughout the menstrual cycle. *J Endocrinol* 1976; **71**:77–85.
- McNatty KP, Smith DM, Makris A, Osathanondh R, Ryan KJ. The microenvironment of the human antral follicle: interrelationships among the steroid levels in antral fluid, the population of granulosa cells, and the status of the oocyte *in vivo* and *in vitro*. *J Clin Endocrinol Metab* 1979; **49**:851–860.
- McNatty KP, Hillier SG, Boogaard AMVD, Trimbos-Kemper TC, Reichert LK, Hall EVV. Follicular development during the luteal phase of the human menstrual cycle. *J Clin Endocrinol Metab* 1983; **56**:1022–1031.
- Menezo YJ, Herubel F. Mouse and bovine models for human IVF. *Reprod Biomed Online* 2002; **4**:170–175.
- Messinis IE. From menarche to regular menstruation: endocrinological background. *Ann NY Acad Sci* 2006; **1092**:49–56.
- Mihm M, Evans AC. Mechanisms for dominant follicle selection in monovulatory species: a comparison of morphological, endocrine and intraovarian events in cows, mares and women. *Reprod Dom Anim* 2008; **43**(Suppl. 2):48–56.
- Mikhail G. Sex steroids in blood. *Clinical Obstet Gynecol* 1967; **10**:29–39.
- Mikhail G. Hormone secretion by the human ovaries. *Gynecol Invest* 1970; **1**:5–20.
- Miro F, Hillier SG. Modulation of granulosa cell deoxyribonucleic acid synthesis and differentiation by activin. *Endocrinology* 1996; **137**:464–468.
- Nasser LF, Adams GP, Bo GA, Mapletoft RJ. Ovarian superstimulatory response relative to follicular wave emergence in heifers. *Therio* 1993; **40**:713–724.
- Nielsen E, Rasmussen I, Fukuda M, Westergaard L, Andersen C. Concentrations of anti-Müllerian hormone in fluid from small human antral follicles show a negative correlation with CYP19 mRNA expression in the corresponding granulosa cells. *Molec Hum Reprod* 2010; **16**:637–643.
- Nilsson L, Wikland M, Hamberger L. Recruitment of an ovulatory follicle in the human following follicle-ectomy and luteectomy. *Fertil Steril* 1982; **37**:30–34.
- Nilsson E, Rogers N, Skinner MK. Actions of anti-Müllerian hormone on the ovarian transcriptome to inhibit primordial to primary follicle transition. *Reproduction* 2007; **134**:209–221.
- O'herlihy C. Monitoring ovarian follicular development with real-time ultrasound. *Br J Obstet Gynaecol* 1980; **87**:613–618.
- Ojha K, Nargund G, Sladkevicius P, Scaramuzzi RJ. Pulsed Doppler ultrasonography to assess follicular growth and the pattern of emergence of the dominant follicle and to determine ovarian follicular and stromal blood flow parameters in relation to follicular size. *J Reprod Fertil Abstract* 2000; **25**:45.
- Pache T, Wladimiroff J, Dejong F, Hop W, Fauser B. Growth patterns of nondominant ovarian follicles during the normal menstrual cycle. *Fertil Steril* 1990; **54**:638–642.
- Peters H. The human ovary in childhood and early maturity. *Eur J Obstet Gynecol Reprod Biol* 1979; **9**:137–144.
- Peters H, Byskov AG, Grinstead J. Follicular growth in fetal and prepubertal ovaries of humans and other primates. In: Ross GT, Lipsett MB (eds). *Clinics in Endocrinology and Metabolism*. London: W B Saunders Co. Ltd., 1978.
- Pierson R, Ginther O. Follicular population dynamics during the estrous cycle of the mare. *Anim Reprod Sci* 1987a; **14**:219–231.
- Pierson RA, Ginther OJ. Follicular populations during the estrous cycle in heifers I. Influence of day. *Anim Reprod Sci* 1987b; **14**:165–176.
- Pierson R, Ginther O. Ultrasonic imaging of the ovaries and uterus in cattle. *Therio* 1988; **29**:21–37.
- Pierson RA, Chizen DR. Transvaginal diagnostic ultrasonography in evaluation and management of infertility. *JOGC* 1991; **13**:37–49.
- Pierson R, Martinuk S, Chizen D, Simpson C. Ultrasonographic visualization of human ovulation. In: Evers J, Heineman M (eds). *From Ovulation to Implantation. Proceedings of the 7th Reignier de Graaf Symposium*. Amsterdam: Elsevier Science Publishers (Biomedical Division), 1990.
- Poretzky L, Cataldo NA, Rosenwaks Z, Giudice LC. The insulin-related ovarian regulatory system in health and disease. *Endocr Rev* 1999; **20**:535–582.
- Queenan JT, O'brien GD, Bains LM, Simposn J, Collins WP, Campbell S. Ultrasound scanning of ovaries to detect ovulation in women. *Fertil Steril* 1980; **34**:99–105.
- Renaud RL, Macler J, Dervain I, Ehret MC, Aron C, Plas-Roser S, Spira A, Pollack H. Echographic study of follicular maturation and ovulation during the normal menstrual cycle. *Fertil Steril* 1980; **33**:272–276.
- Rezaei E, Baerwald A, Pierson R. Ultrasonographic image analysis of ovarian follicles during the human menstrual cycle. In: *Proceedings of the 16th International Student Congress of Medical Sciences*. June 2–5, The Netherlands: , 2009.
- Roberts VJ, Barth S, El-Roeiy A, Yen SS. Expression of inhibin/activin subunits and follistatin messenger ribonucleic acids and proteins in ovarian follicles and the corpus luteum during the human menstrual cycle. *J Clin Endocrinol Metab* 1993; **77**:1402–1410.
- Roseff SJ, Bangah ML, Kettel LM, Vale W, Rivier J, Burger HG, Yen SS. Dynamic changes in circulating inhibin levels during the luteal-follicular transition of the human menstrual cycle. *J Clin Endocrinol Metab* 1989; **69**:1033–1039.
- Ross GT. Gonadotropins and preantral follicular maturation in women. *Fertil Steril* 1974; **25**:522–543.
- Ross GT, Cargille CM, Lipsett MB, Rayford PL, Marshall JR, Strott CA, Rodbard D. Pituitary and gonadal hormones in women during spontaneous and induced ovulatory cycles. *Rec Prog Horm Res* 1970; **26**:1–62.
- Rosavik IK, Gibbons WE. Variability of ovarian follicular growth in natural menstrual cycles. *Fertil Steril* 1985; **44**:195–199.
- Roy S, Greenwald G. Follicular development through preantral stages: signalling via growth factors. *J Reprod Fertil* 1996; **50**:83–94.
- Ryan KJ. Granulosa-thecal cell interaction in ovarian steroidogenesis. *J Ster Biochem* 1979; **11**:799–800.
- Sanders SL, Stouffer RL. Localization of steroidogenic enzymes in macaque luteal tissue during the menstrual cycle and simulated early pregnancy: immunohistochemical evidence supporting the two-cell model for estrogen production in the primate corpus luteum. *Biol Reprod* 1997; **56**:1077–1087.
- Santoro N, Isaac B, Neal-Perry G, Adel T, Weingart L, Nussbaum A, Thakur S, Jinnai H, Khosla N, Barad D. Impaired folliculogenesis and ovulation in older reproductive aged women. *J Clin Endocrinol Metab* 2003; **88**:5502–5509.
- Sasano H, Okamoto M, Mason JI, Simpson ER, Mendelson CR, Sasano N, Silverberg SG. Immunolocalization of aromatase, 17 alpha-hydroxylase and side-chain-cleavage cytochromes p-450 in the human ovary. *J Reprod Fertil* 1989; **85**:163–169.
- Savard K, Marsh JM, Rice BF. Gonadotropins and ovarian steroidogenesis. *Rec Prog Horm Res* 1965; **21**:285–365.
- Scheffer GJ, Broekmans FJ, Dorland M, Habbema JD, Looman CW, Te Velde ER. Antral follicle counts by transvaginal ultrasonography are related to age in women with proven natural fertility. *Fertil Steril* 1999; **72**:845–851.
- Schipper I, Hop S, Fauser B. The follicle-stimulating hormone (FSH) threshold/window concept examined by different interventions with exogenous FSH during the follicular phase of the normal menstrual cycle: duration, rather than magnitude, of FSH increase affects follicle development. *J Clin Endocrinol Metab* 1998; **83**:1292–1298.
- Schneyer AL, Fujiwara T, Fox J, Welt CK, Adams J, Messerlian GM, Taylor AE. Dynamic changes in the intrafollicular inhibin/activin/follistatin axis during human follicular development: relationship to circulating hormone concentrations. *J Clin Endocrinol Metab* 2000; **85**:3319–3330.
- Senbon S, Hirao Y, Miyano T. Interactions between the oocyte and surrounding somatic cells in follicular development: lessons from *in vitro* culture. *J Reproduc Dev* 2003; **49**:259–269.
- Sherman BM, Korenman SG. Hormonal characteristics of the human menstrual cycle throughout reproductive life. *J Clin Invest* 1975; **55**:699–706.
- Short RV. Ovarian steroid synthesis and secretion *in vivo*. *Rec Prog Horm Res* 1964; **20**:303–340.
- Short R. The discovery of the ovaries. In: Zuckerman S, Weir B (eds). *The Ovary*. 2nd ed. New York: Academic Press, 1977.
- Singh J, Pierson R, Adams G. Ultrasound image attributes of the bovine corpus luteum: structural and functional correlates. *J Reprod Fertil* 1997; **109**:35–44.
- Singh J, Pierson RA, Adams GP. Ultrasound image attributes of bovine ovarian follicles and endocrine and functional correlates. *J Reprod Fertil* 1998; **112**:19–29.
- Singh J, Dominguez M, Jaiswal R, Adams GP. A simple ultrasound test to predict the superstimulatory response in cattle. *Therio* 2004; **62**:227–243.
- Sirois J, Fortune JE. Ovarian follicular dynamics during the estrous cycle in heifers monitored by real-time ultrasonography. *Biol Reprod* 1988; **39**:308–317.
- Sirois J, Ball B, Fortune J. Pattern of growth and regression of ovarian follicles during the oestrus cycle and after hemiovariectomy in mares. *Equine Vet J Suppl* 1990; **8**:43–48.
- Smith KB, Lunn SF, Fraser HM. Inhibin secretion during the ovulatory cycle and pregnancy in the common marmoset monkey. *J Endocrinol* 1990; **126**:489–495.

- Sullivan MW, Stewart-Akers A, Krasnow JS, Berga SL, Zeleznik AJ. Ovarian responses in women to recombinant follicle-stimulating hormone and luteinizing hormone (LH): a role for LH in the final stages of follicular maturation. *J Clin Endocrinol Metab* 1999;**84**:228–232.
- Taneja M, Ali A, Singh G. Ovarian follicular dynamics in water buffalo. *Therio* 1996;**46**:121–130.
- Tsang BK, Moon YS, Simpson CW, Armstrong DT. Androgen biosynthesis in human ovarian follicles: cellular source, gonadotropic control, and adenosine 3',5'-monophosphate mediation. *J Clin Endocrinol Metab* 1979;**48**:153–158.
- Turnbull KE, Braden AW, Mattner PE. The pattern of follicular growth and atresia in the ovine ovary. *Austr J Biol Sci* 1977;**30**:229–241.
- Van Dessel HJHMT, Schipper I, Pache TD, Geldorp HV, Jong FHD, Fauser BCJM. Normal human follicle development: an evaluation of correlations with oestradiol, androstendione and progesterone levels in individual follicles. *Clin Endocrinol* 1996;**44**:191–198.
- Van Disseldorp J, Lambalk CB, Kwee J, Looman CW, Eijkemans MJ, Fauser BC, Broekmans FJ. Comparison of inter- and intra-cycle variability of anti-Mullerian hormone and antral follicle counts. *Hum Reprod* 2010;**25**:221–227.
- Van Rooij IA, Broekmans FJ, Scheffer GJ, Looman CW, Habbema JD, De Jong FH, Fauser BJ, Themmen AP, Te Velde ER. Serum anti-Mullerian hormone levels best reflect the reproductive decline with age in normal women with proven fertility: a longitudinal study. *Fertil Steril* 2005;**83**:979–987.
- Van Santbrink E, Hop W, Dessel TV, Jong FD, Fauser B. Incremental follicle-stimulating hormone and dominant follicle development during the normal menstrual cycle. *Fertil Steril* 1995;**64**:37–43.
- Van Zonneveld P, Scheffer GJ, Broekmans FJ, Blankenstein MA, De Jong FH, Looman CW, Habbema JD, Te Velde ER. Do cycle disturbances explain the age-related decline of female fertility? Cycle characteristics of women aged over 40 years compared with a reference population of young women. *Hum Reprod* 2003;**18**:495–501.
- Vaughan JL, Macmillan KL, D'occhio MJ. Ovarian follicular wave characteristics in alpacas. *Anim Reprod Sci* 2004;**80**:353–361.
- Verhagen TE, Hendriks DJ, Bancsi LF, Mol BW, Broekmans FJ. The accuracy of multivariate models predicting ovarian reserve and pregnancy after *in vitro* fertilization: a meta-analysis. *Hum Reprod Update* 2008;**14**:95–100.
- Visser JA, Themmen AP. Anti-Mullerian hormone and folliculogenesis. *Mol Cell Endocrinol* 2005;**234**:81–86.
- Visser JA, De Jong FH, Laven JS, Themmen AP. Anti-Mullerian hormone: a new marker for ovarian function. *Reproduction* 2006;**131**:1–9.
- Weenen C, Laven JS, Von Bergh AR, Cranfield M, Groome NP, Visser JA, Kramer P, Fauser BC, Themmen AP. Anti-Mullerian hormone expression pattern in the human ovary: potential implications for initial and cyclic follicle recruitment. *Mol Hum Reprod* 2004;**10**:77–83.
- Westergaard L, Christensen IJ, McNatty KP. Steroid levels in ovarian follicular fluid related to follicle size and health status during the normal menstrual cycle in women. *Hum Reprod* 1986;**1**:227–232.
- Wiele RLV, Bogumil J, Dyrenfurth I, Ferin M, Jewelewicz R, Warren M, Ritzallah T, Mikhail G. Mechanisms regulating the menstrual cycle in women. *Rec Prog Horm Res* 1970;**26**:63–103.
- Wrathall JH, Knight PG. Effects of inhibin-related peptides and oestradiol on androstenedione and progesterone secretion by bovine theca cells *in vitro*. *J Endocrinol* 1995;**145**:491–500.
- Yamamoto M, Minami S, Nakano R, Kobayashi M. Immunohistochemical localization of inhibin/activin subunits in human ovarian follicles during the menstrual cycle. *J Clin Endocrinol Metab* 1992a;**74**:989–993.
- Yamamoto M, Shima K, Nakano R. Gonadotropin receptors in human ovarian follicles and corpora lutea throughout the menstrual cycle. *Horm Res* 1992b;**37**:5–11.
- Zegers-Hochschild F, Gomez Lira C, Parada M, Altieri Lorenzini E. A comparative study of the follicular growth profile in conception and nonconception cycles. *Fertil Steril* 1984;**41**:244–247.
- Zeleznik AJ. The physiology of follicle selection. *Reprod Biol Endocrinol* 2004;**2**:31.
- Zeleznik AJ, Midgley AR Jr, Reichert LE Jr. Granulosa cell maturation in the rat: increased binding of human chorionic gonadotropin following treatment with follicle-stimulating hormone *in vivo*. *Endocrinology* 1974;**95**:818–825.