

Overdominant Epistatic Loci Are the Primary Genetic Basis of Inbreeding Depression and Heterosis in Rice. I. Biomass and Grain Yield

Zhi-Kang Li,^{*,†} L. J. Luo,[†] H. W. Mei,[†] D. L. Wang,[§] Q. Y. Shu,[§] R. Tabien,^{*} D. B. Zhong,[‡]
C. S. Ying,[‡] J. W. Stansel,^{*} G. S. Khush[†] and A. H. Paterson^{*,**}

[†]Plant Breeding, Genetics, and Biochemistry Division, International Rice Research Institute, Metro Manila, The Philippines, ^{*}Department of Soil and Crop Sciences, Texas A&M University, College Station, Texas 77843, [‡]China National Rice Research Institute, 310006 Hangzhou, China, [§]Department of Agronomy, Zhejiang Agricultural University, 310029 Hangzhou, China and ^{**}Department of Crop and Soil Sciences, University of Georgia, Athens, Georgia 30602

Manuscript received July 1, 1999

Accepted for publication May 18, 2001

ABSTRACT

To understand the genetic basis of inbreeding depression and heterosis in rice, main-effect and epistatic QTL associated with inbreeding depression and heterosis for grain yield and biomass in five related rice mapping populations were investigated using a complete RFLP linkage map of 182 markers, replicated phenotyping experiments, and the mixed model approach. The mapping populations included 254 F₁₀ recombinant inbred lines derived from a cross between Lemont (japonica) and Teqing (indica) and two BC and two testcross hybrid populations derived from crosses between the RILs and their parents plus two testers (Zhong 413 and IR64). For both BY and GY, there was significant inbreeding depression detected in the RI population and a high level of heterosis in each of the BC and testcross hybrid populations. The mean performance of the BC or testcross hybrids was largely determined by their heterosis measurements. The hybrid breakdown (part of inbreeding depression) values of individual RILs were negatively associated with the heterosis measurements of their BC or testcross hybrids, indicating the partial genetic overlap of genes causing hybrid breakdown and heterosis in rice. A large number of epistatic QTL pairs and a few main-effect QTL were identified, which were responsible for >65% of the phenotypic variation of BY and GY in each of the populations with the former explaining a much greater portion of the variation. Two conclusions concerning the loci associated with inbreeding depression and heterosis in rice were reached from our results. First, most QTL associated with inbreeding depression and heterosis in rice appeared to be involved in epistasis. Second, most (~90%) QTL contributing to heterosis appeared to be overdominant. These observations tend to implicate epistasis and overdominance, rather than dominance, as the major genetic basis of heterosis in rice. The implications of our results in rice evolution and improvement are discussed.

INBREEDING depression and heterosis are related phenomena of fundamental importance to evolutionary biology and applied genetics. Inbreeding depression refers to reduced fitness of progenies resulting from inbreeding (STEBBINS 1958; WRIGHT 1977). In contrast, heterosis, or hybrid vigor, is defined as the superiority of an F₁ hybrid over its parents (STUBER 1994). Both heterosis and inbreeding depression are widely observed in both animal and plant kingdoms. In evolution, inbreeding depression may contribute to formation of reproductive barriers between species and populations, while heterosis may be an important force in maintenance of genetic variation in populations (CROW 1986). In applied genetics, exploitation of heterosis has played a major role in the genetic improvement of many crop plants and animals (FALCONER 1981; STUBER 1994). Heterosis and inbreeding depression are considered

two aspects of the same phenomenon (FALCONER 1981; MATHER and JINKS 1982). Heterosis is clearly related to heterozygosity, but it has long been debated how heterozygosity results in heterosis. Two predominant theories were proposed as the genetic basis of heterosis. The overdominance hypothesis (SHULL 1908; EAST 1936) states that heterozygosity at single loci confers properties that are superior to either homozygote. In contrast, the dominance hypothesis (BRUCE 1910; KEEBLE and PELLEW 1910; JONES 1917) proposes that dominant factors from either parent mask deleterious recessive mutations from the other parent in the heterozygous F₁. In both cases, inbreeding depression is due to segregation and expression of deleterious recessive mutations in inbred progenies (ALLARD 1960; SIMMONDS 1979). A third, less widely embraced hypothesis suggests that heterosis may arise from epistasis between alleles at different loci (STUBER 1994; GOODNIGHT 1999).

Historically, heterosis and inbreeding depression are related to fitness and are influenced by many genes

Corresponding author: Zhi-Kang Li, Plant Breeding, Genetics, and Biochemistry Division, International Rice Research Institute, DAPO Box 7777, Metro Manila, The Philippines. E-mail: z.li@cgiar.org

as well as by environments (MATHER and JINKS 1982). Recent quantitative trait loci (QTL) mapping studies sought to gain insights into the genetic basis of heterosis and inbreeding depression in crop plants. Using restriction fragment length polymorphism (RFLP) markers, STUBER *et al.* (1992) mapped quantitative trait loci (QTL) contributing to grain yield and its components in a backcross (BC) population derived from crosses between the F_3 progeny from a cross (B73 \times Mo17) and their parental lines. Main-effect QTL with overdominance or pseudooverdominance effects were found to be largely responsible for heterosis in grain yield and its components in maize. XIAO *et al.* (1994) investigated the genetic basis of heterosis in two rice BCF_1 populations between 198 F_3 recombinant inbred lines and their parents. Ten QTL influencing grain yield components detected in both BC populations were completely or partially dominant. Recombinant inbred lines (RILs) having phenotypic values superior to the F_1 hybrid between the parental lines were found for all traits evaluated. These results led the authors to the conclusion that dominance complementation is the major genetic basis of heterosis in rice.

Results from other studies suggested that epistasis may be an important genetic basis of heterosis. LI *et al.* (1997a,b) reported that hybrid breakdown (part of inbreeding depression) in the F_4 progeny from an interspecific rice cross may be largely due to disharmonious interactions between alleles at many epistatic loci. YU *et al.* (1997) reported overdominance at several main-effect QTL and pronounced additive epistasis affecting grain yield and its components in F_3 progeny from the most widely grown hybrid cultivar in China, Minhui 63.

The goal of the experiments described here was to study inbreeding depression and heterosis in a diverse sampling of germplasm using an improved statistical methodology to shed light on the relative importance of main-effect QTL and digenic epistatic loci associated with inbreeding depression and heterosis in biomass and grain yield of rice.

MATERIALS AND METHODS

Plant materials: Five related mapping populations were used in this study. These included 254 F_{10} RILs derived by single seed descent from a cross between Lemont (*japonica*) and Teqing (*indica*), two BC_1F_1 populations, and two testcross populations. Two BC_1F_1 populations included 172 Lemont (LT) BCF_1 hybrids (the RILs \times Lemont) and 177 Teqing (TQ) BCF_1 hybrids (the RILs \times Teqing). Two testcross populations were derived from testcrossing the RILs with two testers, Zhong 413 (a widely compatible restorer line developed in China) and IR64 (an *indica* cultivar developed in IRRI), which included 192 Z413 F_1 hybrids (the RILs \times Zhong 413) and 187 IR64 F_1 hybrids (the RILs \times IR64). In addition, the parents (Lemont and Teqing), the F_1 (Lemont \times Teqing), and a hybrid cultivar, Shanyou63 (the most widely grown commercial hybrid cultivar in China), were used as checks in the phenotyping experiments.

Phenotyping experiments: The materials were evaluated in two separate experiments at two locations, Zhejiang Agricultural University (ZAU) and China National Rice Research Institute (CNRRI) in 1996. In the ZAU experiment, the RILs, parents, F_1 plants, the two BC_1F_1 populations (LT BCF_1 s and TQ BCF_1 s), and the check hybrid were planted in the seedling nursery on May 25, 1996. The 25-day-old seedlings were transplanted into three-row plots each consisting of a single row of the female RIL and the two BC_1F_1 hybrids (the RIL \times Lemont and Teqing). The plots were arranged in a randomized complete block design with two replications. Each row within a plot consisted of 15 plants with a spacing of 20 cm between the plants within each row and 35 cm between rows. Four check plots consisting of Lemont, Teqing, F_1 , and Shanyou63 were randomly arranged in each replication. In the CNRRI experiment, the same three-row plots, each consisting of a single row of a RIL, and two rows of testcross hybrids (the RIL \times Z413 and IR64) were used. In addition, the six check plots consisting of Lemont, Teqing, F_1 , Z413, IR64, and Shanyou63 were also included in each replication. The field arrangement in CNRRI was the same as the ZAU experiment except that three replications were used.

At the maturity stage, three representative plants from the middle of each row plot were sampled and dried in an oven. Each sampled plant was evaluated for grain yield per plant (GY), biomass per plant (BY), and other grain yield components. Data for BY and GY were converted to tons per hectare (t/ha). Both the original data and \log_e -transformed data for BY and GY were used in the data analyses.

Genotyping and construction of the RFLP linkage map: Genomic DNA of the RILs, parental lines, and testers were extracted from freshly harvested leaves of 25-day-old seedlings grown in the greenhouse at Texas A&M University, College Station, Texas. RFLP mapping was conducted using published procedures (LI *et al.* 1995) and 179 well-distributed RFLP markers from Cornell University and the Japanese Rice Genome Research Program. The RILs were also evaluated for two morphological markers, *C* (apiculus color) and *gl-1* (glabrous leaf), in the field. An additional marker, the reactions of the RILs, parents, and testers to phenol (*Ph*), was evaluated by soaking 10 grains of each test material in 1% phenol solution for 24 hr. Black grain color indicated a positive reaction while unchanged (yellow) grain color indicated a negative reaction. Mapmaker version 3.0 (LINCOLN and LANDER 1992) was used to construct a complete linkage map with 182 markers covering 12 rice chromosomes. Linkage between markers was determined by the group command with a LOD threshold >6.0 and a recombination fraction of 0.25.

Data analyses and QTL mapping: Data of the RI, BC, and testcross populations were analyzed separately. SAS PROC GLM (SAS INSTITUTE 1996) was used to test the differences among the RILs and the BC/testcross hybrids. Equations for calculating values of hybrid breakdown (HB, a component of inbreeding depression) of individual RILs and the midparental heterosis for BY and GY of individual BC/testcross hybrids are listed in Table 1. In addition, two other relative heterosis measurements were calculated as follows: the better parental heterosis $H_{BP} = 100 \times (F_1 - BP)/BP$ and the competitive heterosis $H_C = 100 \times (F_1 - \text{Shanyou63})/\text{Shanyou63}$, where BP and Shanyou63 were the better parent and check hybrid. For mapping main-effect and epistatic QTL, data from each of the mapping populations were analyzed separately. Hybrid breakdown values of individual RILs for BY and GY were used as input data to identify QTL associated with hybrid breakdown. The midparental heterosis H_{MP} values and the mean values of individual BC and testcross F_1 hybrids for BY and GY were used to identify QTL contributing to heterosis.

A mixed linear model for simultaneous mapping with back-

TABLE 1
Equations for calculating hybrid breakdown of the recombinant inbred lines
and the heterosis of the BCF₁ and testcross F₁ populations

Population ^a	N	Equation for measurement ^b
RILs	254	HB = RIL - MP, where MP = (Lemont + Teqing)/2
LTBCF ₁	172	H _{MP} = F ₁ - MP, where MP = (RIL + Lemont)/2
TQBCF ₁	177	H _{MP} = F ₁ - MP, where MP = (RIL + Teqing)/2
Z413BCF ₁	192	H _{MP} = F ₁ - MP, where MP = (RIL + Z413)/2
IR64BCF ₁	187	H _{MP} = F ₁ - MP, where MP = (RIL + IR64)/2

RIL, recombinant inbred line; H_{MP}, heterosis.

^aLTBCF₁ and TQBCF₁s are two BCF₁ populations obtained by crossing the RILs with the parents, Lemont (LT) and Teqing (TQ); while Z413F₁s and IR64F₁s are two testcross F₁ populations obtained by crossing the RILs with the testers, Zhong 413 (Z413) and IR64.

^bHB and H_{MP} are estimates of hybrid breakdown and midparental heterosis. F₁'s are mean trait values of individual BC or testcross hybrids while RIL is the corresponding female RIL parent for each of the BC or testcross hybrids.

ground genetic variation control was used for interval mapping of both main-effect and digenic epistatic QTL for GY and BY segregating in the RI, BC, and testcross populations (WANG *et al.* 1999). The model can be expressed as

$$y_k = \mu + a_i x_{A_{ik}} + a_j x_{B_{jk}} + aa_{ij} x_{AA_{ijk}} + \sum_f u_{M_{fk}} e_{M_j} + \sum_l u_{MM_{lk}} e_{MM_l} + \epsilon_k,$$

where y_k is the phenotypic value of a quantitative trait measured on the k th individual ($k = 1, 2, \dots, n$); μ is the population mean; a_i and a_j are the main effects (fixed) of the two putative QTL (Q_i and Q_j), respectively; aa_{ij} is the epistatic effect (fixed) between Q_i and Q_j ; $x_{A_{ik}}$, $x_{B_{jk}}$, and $x_{AA_{ijk}}$ are coefficients of QTL effects derived according to the observed genotypes of the markers (M_{i-} , M_{i+} and M_{j-} , M_{j+}) and the test positions ($r_{M_i-Q_i}$ and $r_{M_j-Q_j}$); $e_{M_j} \approx N(0, \sigma_{M_j}^2)$ is the random effect of marker f with indicator coefficient $u_{M_{fk}}$ (1 for $M_i M_j$ and -1 for $m_i m_j$); $e_{MM_l} \approx N(0, \sigma_{MM}^2)$ is the random effect of the l th marker interaction (between marker K_i and marker L_l) with indicator coefficient $u_{MM_{lk}}$ (1 for $M_K M_K M_L M_L$ or $m_K m_K m_L m_L$ and -1 for $M_K M_K m_L m_L$ or $m_K m_K M_L M_L$). $\epsilon_k \approx N(0, \sigma_\epsilon^2)$ is the random residual effect. The inclusion of e_{M_j} and e_{MM_l} in the model is intended to absorb additive and epistatic effects of background QTL (additional segregating QTL other than the loci searched) for controlling the noise caused by the background QTL (WANG *et al.* 1999).

A new computer software, QTLMAPPER version 1.0, was developed on the basis of the above model (WANG *et al.* 1999), which allows simultaneous interval mapping of both main effect and digenic epistatic QTL in a RI, doubled haploid (DH), or BC population (with two genotypes at each marker locus). QTL mapping was carried out in three steps using the computer software. First, significant markers were identified across the genome using stepwise regression analyses based on single marker genotypes for putative main-effect QTL and based on all possible pairwise marker pairs for epistatic QTL with a threshold of $P \leq 0.005$. Then, all putative main-effect and epistatic QTL were identified using composite interval mapping in genomic regions centered at the markers (covering two marker intervals in each QTL region) identified in the first step with all QTL fixed in the model to control the background genetic variation. In this way, each of the QTL included in the model were significant at a threshold of $P \leq 0.002$ and $R^2 > 5\%$. This threshold was shown to have a very low probability of false positives (WANG *et al.* 1999). Finally, genetic parameters (effects and test statistics) associated with significant main-effect and epistatic QTL were estimated at the positions of respective LOD peaks in individual putative

QTL regions (each putative QTL region covered two marker intervals) using the model and the restricted maximum-likelihood estimation method (PATTERSON and THOMPSON 1971, 1974; WANG *et al.* 1999).

The genetic expectations of the parameters estimated in the above model differ according to the nature of the mapping population and the input data. For the RI population, the main effects a_i and a_j are the additive effects of the two putative QTL (Q_i and Q_j), and aa_{ij} is the additive epistatic effect between Q_i and Q_j (WANG *et al.* 1999). For the BCF₁ populations, however, a_i and a_j are the combined effects of both additive and dominance gene actions ($\frac{1}{2}d - \frac{1}{2}a$) when estimated from the F₁ mean values and the QTL dominance effects ($\frac{1}{2}d$) when estimated from the midparental heterosis (H_{MP}) values. Similarly, the estimated epistatic effect using H_{MP} measurements is the dominance \times dominance (dd_{ij}) effect between epistatic QTL, while those from the mean F₁ values contained both additive and nonadditive epistatic components (MATHER and JINKS 1982). The assumptions underlying the estimation of the epistatic effect are $aa_{ii} = aa_{jj} = -aa_{ij} = -aa_{ji}$ for the RI population and $aa_{ij} = dd_{ij} = -ad_{ij} = -da_{ij}$ for the BCF₁ populations, where aa_{ij} , dd_{ij} , ad_{ij} , and da_{ij} are additive \times additive, dominance \times dominance, additive \times dominance, and dominance \times additive digenic epistatic effects between Q_i and Q_j .

RESULTS

RFLP linkage map construction: The complete linkage map of 182 markers (Figure 2) spanned 1918.7 cM and covered 12 rice chromosomes with an average interval of 11.3 cM between markers. There was a single gap of 54.8 cM on chromosome 9. The linear orders agreed largely with those obtained from the F₂ population of the same cross (LI *et al.* 1995). A total of 46 (25.8%) markers showed segregation distortion, largely clustered in terminal regions of chromosomes 6–11. On average, Lemont alleles accounted for $47.4 \pm 7.7\%$ of the genome, ranging from 16.1 to 62.3%.

Inbreeding depression in the RILs and heterosis in the BC and testcross hybrids: The paternal parent of the RILs, Teqing (*indica*), had significantly higher BY and GY than the maternal parent, Lemont (*japonica*),

TABLE 2
Summary statistics on inbreeding depression of the Lemont/Teqing RILs
and H_{MP} of their backcross/testcross F_1 populations

	Biomass (t/ha)			Grain yield (t/ha)		
	Mean	SD	Range	Mean	SD	Range
The ZAU experiment						
Lemont (LT)	2.74	0.64	—	1.46	0.43	—
Teqing (TQ)	5.46	1.08	—	3.47	0.64	—
F_1 (LT × TQ)	9.25	1.32	—	5.42	0.71	—
H_{MP}	5.15			2.96		
CK (SY63)	8.82	1.17	—	4.62	0.86	—
LTBCF ₁	5.96	2.00	~2.53–16.69	3.22	1.27	~0.87–9.89
(LTBCF ₁) H_{MP}	2.36	1.90	~-2.04–12.19	1.23	1.21	~-0.96–7.02
TQBCF ₁	7.80	2.20	~2.24–15.20	4.40	1.31	~1.34–7.92
(TQBCF ₁) H_{MP}	4.24	2.24	~-0.76–10.15	1.46	1.37	~-0.50–6.02
RILs	3.01	1.61	~1.13–9.63	1.47	0.78	~0.34–4.33
HB (RIL – MP) ^a	-1.13	1.61	~-3.00–5.52	-0.99	0.78	~-2.13–1.86
The CNRRI experiment						
Lemont (LT)	3.86	1.19		1.76	0.50	
Teqing (TQ)	15.63	1.89		8.24	0.80	
F_1 (LT × TQ)	13.93	1.00		8.09	1.76	
H_{MP}	4.19			3.09		
CK (SY63)	17.44	2.96		8.37	1.30	
Z413	14.07	0.34		8.87	0.44	
IR64	8.61	0.50		4.04	0.36	
Z413F ₁	13.09	3.99	~4.37–29.12	6.91	2.23	~2.24–13.05
(Z413F ₁) H_{MP}	2.79	4.14	~-9.38–19.63	1.13	2.34	~-4.22–7.87
IR64F ₁	14.76	3.11	~4.02–26.35	7.50	1.79	~1.79–13.73
(IR64F ₁) H_{MP}	7.81	3.26	~-1.49–18.13	4.27	1.84	~-0.69–9.60
RILs	6.09	2.27	~2.13–13.83	2.63	1.14	~0.46–7.55
HB (RIL – MP) ^a	-3.66	2.27	~-7.60–4.09	-2.37	1.14	~-4.54–2.54

^a HB = RIL – MP, where HB is hybrid breakdown and MP = (Lemont + Teqing)/2.

in both experiments (Table 2). The BY and GY values of the F_1 plants in the ZAU experiment were 9.25 and 5.42 t/ha, significantly higher than both parents, but 13.93 and 8.09 t/ha in the CNRRI experiment, similar to the better parent, Teqing. The midparental heterosis of the F_1 plants was 4.65 t/ha (101.1%) for BY and 2.96 t/ha (120.0%) for GY in ZAU, and 4.2 t/ha (42.9%) for BY and 3.09 t/ha (61.8%) in CNRRI, respectively.

Inbreeding depression of the RILs: Significant reductions for both BY and GY were observed, as a result of hybrid breakdown, in the RI population in both ZAU and CNRRI experiments (Table 2). In the ZAU experiment, hybrid breakdown values were -1.13 (-35.9%) and -0.99 (-40.1%) t/ha for BY and GY, respectively. None of the RILs had higher BY and GY than the F_1 plants, but two RILs had significantly higher BY and GY than the better parent, Teqing. Compared to the ZAU experiment, all the materials in the CNRRI experiment had much higher BY and GY and the RILs showed a greater degree of inbreeding depression. Hybrid breakdown values of the RILs were normally distributed (Figure 1) with mean values of -3.66 (-37.5%) and -2.37 (-47.4%) t/ha for BY and GY, respectively. None of the RILs had higher BY or GY than Teqing.

Heterosis in the BC and testcross hybrid populations: Highly significant heterosis for both BY and GY were observed in the BC and testcross hybrid populations, and heterosis values of the BCF₁ hybrids were distributed normally (Table 2 and Figure 1). On average, the IR64F₁ population showed the highest level of heterosis, the LTBCF₁ population the second, the TQBCF₁ the third, and the Z413F₁ the lowest. Within each of the populations, individual F_1 hybrids varied considerably in their mean values and heterosis values (Figure 1). Most BC or testcross hybrids showed highly significant positive heterosis. However, hybrids showing significant negative heterosis for BY and GY were observed but were much less frequent in all four F_1 populations.

In the ZAU experiment, the mean BY and GY values of the LTBCF₁ population were 5.96 and 3.22 t/ha. The heterosis values of individual hybrids were normally distributed with the mean of 2.36 t/ha (107.5%) for BY and 1.23 t/ha (120.1%) for GY. In particular, the top 10 hybrids had mean BY and GY values of 10.49 and 6.25 t/ha, giving a mean heterosis of 6.51 t/ha (265.1%) for BY and 4.55 t/ha (326.7%) for GY, respectively. The mean better parental heterosis and the competitive heterosis of the top 10 hybrids were 249.4 and 18.9%




FIGURE 1.—Frequency distribution of hybrid breakdown (HB = RIL - F₁) of the Lemont/Teqing recombinant inbred lines and midparental heterosis for biomass and grain yield per plant of their backcross/testcross F₁ populations. MP, mean midparental values.

for BY and 324.7 and 35.4% for GY, respectively. The TQBCF₁ population had a mean value of 7.80 and 4.40 t/ha for BY and GY. The heterosis values of individual BCF₁s were normally distributed with a mean of 4.24 t/ha (84.3%), ranging from -0.76 to 10.15 t/ha for BY and 1.46 t/ha (78.1%) ranging from -0.50 to 6.02 t/ha for GY. The top 10 hybrids had mean BY and GY of 12.07 and 7.17 t/ha, with mean heterosis of 7.89 t/ha (184.3%) and 4.70 t/ha (191.0%), respectively. The better parental heterosis and competitive heterosis of the top 10 hybrids were 119.5 and 36.9% for BY and 106.6 and 55.4% for GY, respectively.

In the CNRRI experiment, the Z413F₁ population had very high F₁ mean value for both BY (13.09 t/ha) and GY (6.91 t/ha) but the lowest heterosis for BY (2.79 t/ha, or 29.8%) and GY (1.13 t/ha, or 20.3%). The top 10 hybrids in the Z413F₁ population had mean BY and GY of 21.58 and 11.47 t/ha and mean heterosis of 12.1 t/ha (112.8%) and 6.12 t/ha (199.4%), respectively.

The better parental heterosis and competitive heterosis of the top 10 hybrids were 52.4 and 22.9% for BY and 29.2 and 37.0% for GY, respectively. The IR64F₁ population had the highest mean value for both BY (14.76 t/ha) and GY (7.50 t/ha) and the highest heterosis for BY (7.81 t/ha, or 100.8%) and GY (4.27 t/ha, or 124.8%). IR64 performed relatively poorly (with mean BY and GY of 8.61 and 4.04 t/ha). The top 10 hybrids of this population had mean BY and GY of 21.58 and 11.47 t/ha, with the mean heterosis of 12.1 t/ha (176.4%) and 6.12 t/ha (230.2%), respectively. The better parental heterosis and competitive heterosis of the top 10 hybrids were 135.8 and 16.5% for BY and 172.5 and 31.6% for GY, respectively.

The relationships between hybrid breakdown, heterosis, and F₁ performance: The mean performance of individual BC and testcross hybrids for BY and GY was largely determined by the levels of heterosis of individual hybrids instead of the mean performance of their

maternal RILs (Table 3). The correlations between the F₁ mean values and heterosis in the LTBCF₁, TQBCF₁, Z413F₁, and IR64F₁ populations were 0.76, 0.82, 0.85, and 0.81 for BY and 0.79, 0.81, 0.86, and, 0.83 for GY, respectively. There was no correlation between the F₁ mean values and the mean performances of their maternal RILs (Table 3).

In both ZAU and CNRRI experiments, hybrid breakdown and heterosis for GY and BY were distributed on both sides of the midparental value (at the zero point) with little overlapping (Figure 1). The hybrid breakdown values of individual RILs were negatively correlated with their heterosis values across all four F₁ populations. The correlation was highly significant ($P < 0.0001$) but moderate in magnitude ($r = -0.39, -0.59, -0.50, \text{ and } -0.52$ for BY and $-0.47, -0.63, -0.57, \text{ and } -0.49$ for GY; Table 3), suggesting that hybrid breakdown of the RILs and heterosis of their F₁ hybrids indeed shared a partially overlapping genetic basis. The mean performance of the paternal parents (Lemont, Teqing, Z413, and IR64) was also negatively associated with the midparental heterosis ($r = -0.84$ for BY and -0.97 for GY) but positively associated with the relative competitive heterosis of their F₁ hybrids ($r = 0.66$ for BY and 0.63 for GY).

Main-effect and epistatic QTL associated with hybrid breakdown in the RILs: The segregation of the RILs for BY and GY could be largely explained by many main-effect and epistatic QTL (Figure 2, Tables 4–6). In the ZAU experiment, two main-effect QTL affecting GY and BY were identified on chromosomes 2 and 11. However, a total of eight pairs of epistatic QTL were identified. Among these loci, two (on chromosomes 7 and 10) had significant additive effects (Table 6). The allele increasing GY and BY at all main-effect QTL but one was from Teqing. Five of the eight significant epistatic effects on GY and/or BY were positive and the remaining three were negative. In the CNRRI experiment, four main-effect QTL were mapped to chromosomes 3, 4, 6, and 9 (Table 5). Ten pairs of epistatic QTL affecting GY and/or BY were identified. Of these, four loci located on chromosomes 3, 5, 6, and 12 had significant main effects (Table 6). The Teqing allele at all main-effect QTL increased GY and BY. All significant epistatic effects but one were positive. The observation that most epistatic effects were positive indicated that hybrid breakdown for GY and BY in the RILs was largely due to the disharmonious interactions between alleles from different parents at these loci (Li *et al.* 1997a).

Main-effect and epistatic QTL associated with the F₁ performance and heterosis: Several main-effect QTL and many epistatic QTL were largely responsible for the phenotypic variation for BY and GY in the BCF₁ and testcross populations (Tables 4, 5, 7, Figure 2). In the LTBCF₁ population, a single main-effect QTL and seven pairs of epistatic loci affecting the F₁ performance and/or heterosis were identified (Tables 4 and 7). Of these

TABLE 3
Phenotypic correlation for BY and GY between the performance of the Lemont/Teqing RILs and H_{MP} in their BCF₁ and testcross F₁ populations

	Between RIL values and H _{MP}														
	Between RIL values and F ₁ values						Between H _{MP} and F ₁ values								
	Lemont		Teqing		Z413		IR64		Lemont		Teqing		Z413		IR64
GY	<i>r</i>	0.173	-0.060	0.004	-0.062	0.075	0.791	0.811	0.857	0.825	-0.466	-0.633	-0.567	-0.490	
	<i>R</i> ²	0.030	0.004	0.004	0.004	0.006	0.626	0.658	0.734	0.681	0.217	0.401	0.321	0.240	
BY	<i>r</i>	0.306	-0.017	0.030	0.067	0.067	0.760	0.819	0.851	0.814	-0.386	-0.587	-0.500	-0.519	
	<i>R</i> ²	0.094	0.000	0.001	0.004	0.004	0.578	0.671	0.724	0.663	0.149	0.345	0.250	0.269	

BY, biomass; GY, grain yield.

TABLE 4
Main-effect QTL affecting GY (in t/ha) and BY (in t/ha) of the Lemont/Teqing RILs and H_{MP} of their BCF₁ hybrids

Trait	Chromosome	Marker interval	RILs			BCF ₁ (Lemont)			H_{MP} (Lemont)			BCF ₁ (Teqing)			H_{MP} (Teqing)		
			LOD	<i>a</i>	<i>R</i> ²	LOD	<i>a</i> + <i>d</i>	<i>R</i> ²	LOD	<i>d</i>	<i>R</i> ²	LOD	<i>a</i> + <i>d</i>	<i>R</i> ²	LOD	<i>d</i>	<i>R</i> ²
BY	2	RG437-RZ476a	5.98	-0.53	15.8	4.28	0.72	10.2	3.30	0.69	9.6						
GY	2	RG437-RZ476a	5.23	-0.25	12.9	4.11	0.51	9.6	3.82	0.52	10.2						
BY	11	G44-RG1094b	2.45	-0.20	9.0												
GY	11	G44-RG1094b	2.28	-0.35	7.9												
BY	6	G294a-G1468b										7.48	1.04	19.5	1.08	22.3	
GY	6	G294a-G1468b										10.81	0.60	12.5	0.62	11.5	
BY	9	RG451-RZ404										2.46	0.74	6.2			
GY	9	RG451-RZ404										3.37	0.42	5.4			

^aIn the RI population, QTL effects were associated with the Lemont allele (due to replacement of the Teqing allele by the Lemont allele). In the BC populations, QTL effects for F₁ and H_{MP} were estimated by the heterozygote minus the homozygote. The genetic expectation of a QTL effect obtained is the additive gene effect (*a*) when estimated from the RI, the additive and dominance effects (*a* + *d*) from the F₁ mean values, and the dominance effect (*d*) from H_{MP} values.

epistatic QTL, eight loci (on chromosomes 1–3, 5, 7, and 8) had significant main effects on BY and/or GY (Table 7). The heterozygote at all main-effect QTL except one (near RG30 of chromosome 7) increased BY and/or GY. Four of the epistatic effects were positive, and the remaining three were negative.

In the TQBCF₁ population, two main-effect QTL (chromosomes 6 and 9) and eight pairs of epistatic loci affecting the F₁ mean and heterosis of BY and GY were identified and mapped to chromosomes 1, 4, 6–9, 11, and 12 (Tables 4 and 7, Figure 2). Six of these epistatic loci had significant main effects on the F₁ mean and/or heterosis of BY and GY. At all main-effect QTL except one (near RZ382 of chromosome 1), the heterozygote had greater BY and/or GY than the homozygous Teqing genotype. Five of the epistatic effects were positive, and the remaining three were negative.

For the Z413F₁ population, two main-effect QTL and nine pairs of digenic epistatic loci affecting the F₁ mean and/or heterosis were identified and mapped to nine of the rice chromosomes (1–8 and 12; Tables 5 and 7, Figure 2). Of the epistatic QTL, seven had significant main effects. The Lemont/Z413 (japonica/indica) heterozygote at all main-effect QTL except one (near RG653 on chromosome 6), had greater BY and/or GY than the Teqing/Z413 (indica/indica) heterozygote. Of the significant epistatic effects, five were positive and the other four were negative.

For the IR64F₁ population, two main-effect QTL and eight pairs of digenic epistatic loci affecting the F₁ mean and/or H_{MP} were identified and mapped to nine of the rice chromosomes (2–8, 11, and 12; Tables 5 and 7, Figure 2). Of these epistatic QTL, four had significant main effects on BY and/or GY. The Teqing/IR64 (indica/indica) heterozygote at all main-effect QTL had greater BY and/or GY than the Lemont/Z413 (japonica/indica) heterozygote. Of the significant epistatic effects, three were positive and the other five were negative.

CONCLUSIONS AND DISCUSSION

Decomposition of inbreeding depression in rice: Prevailing in outcrossing species, inbreeding depression has been intensively investigated. However, inbreeding depression in self-pollinated plant species like rice has received little attention. The 60–70% reduction of the Lemont/Teqing RI population from the F₁ was highly significant but underestimated the overall degree of inbreeding depression in the Lemont/Teqing cross. This was attributable to purging of deleterious alleles and/or less fit multilocus genotypes by natural selection during the development of the RILs (LI *et al.* 1995, 1997a,b). Genetically, the inbreeding depression values of individual RILs and the mean values of BC or testcross hybrids for BY and GY have two components. One is the deviation of the RILs from the midparental value,




FIGURE 2.—Genomic locations of main-effect QTL and epistatic loci affecting grain yield and biomass detected in the Lemont/Teqing recombinant inbred lines (RILs) and their backcross/testcross F_1 populations. LTBCF₁, TQBCF₁, Z413F₁, and IR64F₁ represent the two backcross and two testcross F_1 populations, respectively.




FIGURE 2.—Continued.

TABLE 5
Main-effect QTL affecting GY (in t/ha) and BY (in t/ha) of the Lemont/Teqing RILs and H_{AIP} of their testcross F_1 hybrids

Trait	Chromosome	Interval	RILs			F_1 (Z413)			H_{AIP} (Z413)			F_1 (IR64)			H_{AIP} (IR64)		
			LOD	Effect	R^2	LOD	Effect	R^2	LOD	Effect	R^2	LOD	Effect	R^2	LOD	Effect	R^2
BY	3	C515-RG348a	10.96	-0.94	14.8												
GY	3	C515-RG348	11.05	-0.48	14.3												
BY	3	G249-RG418	3.81	-0.58	5.6	2.92	1.19	9.4	4.04	1.29	11.2						
GY	3	G249-RG418	8.34	-0.46	12.2	3.95	0.57	10.8	4.07	0.65	12.7						
BY	6	HHU37-RZ682	4.79	-0.66	8.8												
GY	6	HHU37-RZ682	2.79	-0.30	5.3												
BY	9	RG451-RZ404	3.94	-0.54	5.8												
GY	9	RG451-RZ404	3.58	-0.34	7.2												
BY	12	RG574-G193				3.26	0.93	9.3	4.43	1.17	11.1						
GY	12	RG574-G193				4.25	0.63	11.0	4.10	0.65	11.6						
BY	11	RG16-RZ797b										2.45	-0.68	7.9	3.18	-0.73	8.8
GY	11	RG16-RZ797b										3.75	-0.43	11.1	4.86	-0.50	12.6
BY	12	RZ397-RZ257										2.17	-0.58	5.2	2.75	-0.65	6.2
GY	12	RZ397-RZ257										3.30	-0.37	6.3	3.01	-0.42	6.6

In the RI population, QTL effects were associated with the Lemont allele (the effects due to replacement of the Teqing allele by the Lemont allele). In the testcross populations, QTL effects for F_1 and H_{AIP} (midparental heterosis) were estimated by the Lemont/tester genotype minus the Teqing/tester genotype.

TABLE 6
Digenic epistatic QTL affecting hybrid breakdown of GY (in t/ha) and BY (in t/ha) in the Teqing/Lemont recombinant inbred population

Environment	Trait	Chromosome	Marker interval <i>i</i>	Chromosome	Marker interval <i>j</i>	LOD	a_i	a_j	aa_{ij}	R^2 (%)
ZAU	BY	2	RZ476b-RG83	7	RG678-G20	3.09			-0.37***	7.9
ZAU	GY					2.58			-0.19***	8.1
ZAU	BY	5	Y1049-R569a	10	RG1094f-C16	3.82			0.51***	13.9
ZAU	GY					4.37			0.30***	12.9
ZAU	BY	7	RG30-RG29	10	RG1094f-C16	3.94			-0.39***	8.8
ZAU	GY					4.61	0.16**		-0.24***	11.2
ZAU	BY	9	RZ698-G95	12	RG901-G402	3.61			-0.53***	14.8
ZAU	GY					4.24			-0.33***	14.1
ZAU	BY	1	RZ14-C944b	4	RZ740-G200b	2.78			0.37***	7.2
ZAU	BY	2	RZ446b-RZ446a	10	C16-RG561	3.24			0.37***	9.0
ZAU	GY	2	RZ599-RZ476b	3	RG100-RZ284	3.80			0.19***	8.5
ZAU	GY	6	RZ682-C236	11	L457b-G2132b	2.59	-0.13*		0.31***	6.9
CNRR1	BY	2	RG256-RZ260	12	RZ257-RZ797a	2.59			0.51***	6.2
CNRR1	GY					3.79			0.32***	8.5
CNRR1	BY	4	RZ740-G200b	11	RZ797b-RG1094d	3.03			0.44***	5.2
CNRR1	GY					3.95			0.30***	7.6
CNRR1	BY	5	<i>gl1</i> -Y1049	11	RZ536a-L457b	4.05	-0.52***		0.65***	8.5
CNRR1	GY					5.20	-0.30***		0.20***	5.2
CNRR1	BY	7	CDO405-CDO497	9	G103a-CDO395	3.30			0.47***	5.6
CNRR1	GY					3.52			0.22***	5.6
CNRR1	BY	3	RZ474-C746	6	C-G200a	2.86			0.47***	5.6
CNRR1	BY	3	RZ761-C249	12	L102-G1468a	4.76	-0.49***		0.36**	4.3
CNRR1	GY	2	RG139-C624x	3	RZ284-RZ403b	4.90			0.29***	7.4
CNRR1	GY	2	C624x-G45	9	RZ777-CDO82	3.73			-0.30***	8.0
CNRR1	GY	6	RZ667-C235a	12	RG901a-G402	3.47	-0.20**		0.26***	6.8
CNRR1	GY	11	RG16-RZ797b	12	G1106-RG901	3.99		-0.17*	0.24***	6.3

a_i and a_j were the main effects of the loci *i* and *j*, arising from the substitution of the Lemont allele by the Teqing allele, and aa_{ij} was the epistatic effect between loci *i* and *j*, as defined by MATHER and JINKS (1982). *, significance levels of $P < 0.05$. **, significance levels of $P < 0.001$. ***, significance levels of $P < 0.0001$.

TABLE 7
Digenic epistatic QTL affecting F_1 performance and H_{MP} of GY (in t/ha) and BY (t/ha) and BY (t/ha) and TQ BCF₁ and two testcross F_1 populations

Trait	Chromosome	Interval <i>i</i>	Chromosome	Interval <i>j</i>	F_1				H_{MP}					
					LOD	a_i	a_j	aa_{ij}	R^2 (%)	LOD	a_i	a_j	aa_{ij}	R^2 (%)
LT BY	1	R210-RZ382	2	RZ599-RZ476b	4.96	0.31*	0.68***	0.31*	5.1	2.52	0.30*	0.40**	0.39**	6.7
LT GY	1	C112-RG236	11	RZ53-RZ781	4.09	0.17*	0.39***	0.23*	6.4	3.15	0.38**	0.25*	0.30**	7.9
LT BY	2	RZ446b-RZ446a	7	G20-RG30	2.11	0.39*	-0.48**	-0.48**	7.5					
LT GY	2	RZ446b-RZ446a	7	G20-RG30	2.74	0.18*	-0.26**	-0.26**	7.5					
LT BY	3	C74a-RG450	5	CDSR49-RG346	5.10		0.54**	0.77***	12.0	4.13		0.59**	0.80***	13.4
LT GY	3	C74a-RG450	5	CDSR49-RG346	3.36		0.55***	0.44***	11.7	3.46		0.47**	0.44***	10.9
LT BY	5	RG556- <i>g1</i>	8	C424b-RZ143	2.43		0.38**	-0.55***	8.6	2.80		0.20*	-0.68***	11.8
LT GY	5	RG556- <i>g1</i>	8	C424b-RZ143	3.45			-0.41**	11.2	3.01	0.70**		-0.45***	11.6
LT BY	1	C944b-RG957	3	RZ474-C746	3.51	0.24*	0.24*	0.25**	7.1	2.34			-0.48***	12.3
LT GY	1	C944b-RG957	3	RZ474-C746	3.40		0.25*	0.34**	9.6	2.10		0.18*	0.34**	8.9
LT BY	6	C76-RZ516	8	G187-G56a	4.04	0.68**	-0.65***	-0.65***	8.5	3.97	0.70**		0.36**	9.4
TQ BY	1	CDO118-CDO455	1	RZ382-RG532a	5.28	0.39***	-0.44***	-0.44***	8.9	5.39	0.43***		-0.71***	10.9
TQ GY	1	RG236-RZ801	6	RZ762-C76	2.52		0.54**	0.54**	5.5	2.98			-0.44***	5.9
TQ BY	1	RG236-RZ801	6	RZ762-C76	5.09		0.40***	0.40***	7.4	5.01			0.59***	8.5
TQ GY	1	RZ288-C131	12	RG20q-RG91q	3.49		0.57***	0.57***	5.8	2.53			0.45***	6.6
TQ BY	1	RZ288-C131	12	RG20q-RG91q	5.72		0.57***	0.57***	8.1	3.64			0.54**	7.7
TQ GY	4	G177-RZ590b	9	CDO1081-RZ777	5.74		-0.48***	-0.48***	10.4	4.50	0.23*		0.51***	7.5
TQ BY	4	<i>Ph</i> -G379	11	RG1109-RZ537b	2.71	0.36*	0.56**	0.56**	6.8	2.11		0.23*	-0.40***	4.9
TQ GY	4	<i>Ph</i> -G379	11	RG1109-RZ537b	3.93	0.22*	0.28**	0.28**	5.8	3.21			0.53**	7.3
TQ BY	7	RG678-G20	8	RZ143-C825a	3.75	0.32*	-0.59***	-0.59***	6.9	3.42	0.40**		0.36**	5.4
TQ GY	7	RG678-G20	8	RZ143-C825a	5.69		-0.52***	-0.52***	12.5	4.41			-0.52**	6.8
TQ BY	1	RG472-RG447	12	RG91q-RG869	2.74		0.71**	0.71**	9.3	3.20			-0.48***	7.1
TQ GY	1	RG472-RG447	12	RG91q-RG869	3.29	0.51**	0.57**	0.57**	7.9				0.79***	11.4
TQ BY	8	G2140-RZ323a	9	CDO1081-RZ777	3.39		0.60*	0.94***	7.0	4.25		1.03***	0.93***	7.6
TQ GY	1	RZ14-C944b	6	G1468b-RG424	5.52		0.38**	0.68***	8.9	5.04		0.50**	0.67***	8.3
Z413 BY	2	RG139-C624x	8	CSU754-G104	3.77		1.21***	-0.62*	4.6	4.18		1.36***	-0.63*	5.1
Z413 GY	2	RG139-C624x	8	CSU754-G104	3.24		0.49**	-0.38*	5.0	3.62		0.59***	-0.41*	5.1
Z413 BY	3	CDO337-C944a	4	HHU39-RG143	2.72		-0.96***	-0.96***	7.1	3.48		0.51*	-1.16***	9.5
Z413 GY	3	CDO337-C944a	4	HHU39-RG143	2.99		-0.59***	-0.59***	6.5	4.10			-0.71***	8.8
Z413 BY	4	RZ69-RG190	6	C236-RG653	4.58	-1.62***	1.60***	-1.47***	10.9	4.19	-1.47***	1.55***	-1.47***	12.0
Z413 GY	4	RZ69-RG190	6	C236-RG653	3.22	-0.91**	0.67*	-0.72***	9.4	3.94	-0.96**	0.78*	-0.76***	9.4
Z413 BY	4	<i>Ph</i> -G379	8	G187-G56a	4.42		-1.16***	-1.16***	8.6	4.91		0.52*	-1.38***	11.3
Z413 GY	4	<i>Ph</i> -G379	8	G187-G56a	3.24		-0.67***	-0.67***	8.5	4.82			-0.80***	10.0
Z413 BY	1	C131-RG472	4	RG1094e-Y1065Lc	2.49		0.86**	0.86**	6.2					
Z413 GY	3	G249-RG418	5	RG556- <i>g1</i>	3.94	0.86***	0.86***	0.86***	6.4					

(continued)

TABLE 7
(Continued)

Trait	Chromosome	Interval <i>i</i>	Chromosome	Interval <i>j</i>	F_1					H_{MP}				
					LOD	a_i	a_j	aa_{ij}	R^2 (%)	LOD	a_i	a_j	aa_{ij}	R^2 (%)
Z413	7	C586-CDO405	12	RG901-G402	3.19			1.38***	10.2	3.52			1.64***	13.4
Z413	4	RG214- <i>Ph</i>	12	RG901-G402	3.98			0.81***	10.6	3.75			0.75***	9.3
IR64	2	RZ476a-RZ599	5	RG13-CDSR49	3.65	-0.45*		-0.84***	6.4	3.73			-0.96***	8.2
IR64	GY				3.93	-0.32**		-0.44***	5.3	3.88			-0.48***	6.0
IR64	2	RG520-RZ446b	8	C825a-CSU754	2.17			0.35**	4.9	3.13			0.56***	6.9
IR64	GY				2.66			0.85***	6.6	2.73			0.50***	6.4
IR64	3	RZ761-G249	8	C825a-CSU754	3.31			0.51***	7.0	3.41			0.77***	5.8
IR64	BY	G249-RG418	11	RG1109-RZ537b	3.66		-0.59**	-1.00***	9.0	3.13			0.48***	5.8
IR64	GY				3.83	-0.26*	-0.32*	-0.62***	9.1	3.65	-0.31*	-0.79**	-0.86**	6.5
IR64	4	<i>Ph</i> -G379	6	C-G200a	2.57			-0.75***	5.1	4.09			-0.61***	7.7
IR64	GY				3.15			-0.49***	6.5	3.94			-0.92***	7.4
IR64	6	C236-RG653	11	RG16-RZ797b	5.21			1.27***	14.6	5.03			-0.51***	6.6
IR64	BY				5.36			0.70***	13.3	5.15			1.19***	12.6
IR64	2	RG520-RZ446b	3	RZ284-RZ403b	3.06			-0.95***	10.9	3.25			0.65***	10.8
IR64	6	RZ762-C76	7	RC30-RG29	3.15		0.56**	-0.59***	8.6	3.22		0.66**	-1.19***	12.1
													-0.57**	7.2

a_i and a_j were the main effects of the loci i and j , arising from the substitution of the Lemont allele by the Teqing allele, and aa_{ij} was the epistatic effect between loci i and j , as defined by MATHER and JINKS (1982). *, significance levels of $P < 0.05$. **, significance levels of $P < 0.01$. ***, significance levels of $P < 0.0001$.

which is due to additive gene action, and causes hybrid breakdown (STEBBINS 1958; OKA 1988; LI *et al.* 1997a,b). Genes of this type are directly detectable in the RILs but confounded in the BC and testcross populations. The other is the deviation of the F_1 value from the midparental value or heterosis due to genes of non-additive action, which are segregating and contributing to heterosis in the BC and testcross populations but are not directly detectable in the RILs.

The decomposition of inbreeding depression into its additive (hybrid breakdown) and nonadditive (heterosis) components is very important in understanding the genetic basis of heterosis. The presence of hybrid breakdown in self-pollinated plant species such as rice has long been observed (STEBBINS 1958; OKA 1988; LI *et al.* 1997a,b) but not expressed mathematically as part of inbreeding depression and heterosis in quantitative genetic theory (FALCONER 1981). In our mapping populations, the genetic overlap between hybrid breakdown and heterosis was $\sim 30\%$ for GY and 25% for BY. This group of overlapping genes is of particular importance since they contributed positively to heterosis when in heterozygous status and negatively to the mean performance of the inbred RILs (resulting in hybrid breakdown) when in homozygous status. This also provided an explanation for the observation that the mean performance of the female RILs was not correlated with the mean performance of their BC/testcross hybrids. This is consistent with the observed heterosis in rice reported by Zeng, who found no correlation between F_1 mean and the midparental values for grain yield and biomass in 34 commercial rice hybrids (ZENG *et al.* 1979). In numerous classic quantitative genetic or breeding studies using diallel and/or test crosses, additive gene action was shown to be important to the mean performance of F_1 hybrids in rice and other crop species (*cf.* SIMMONDS 1979; VIRMANI 1994). This is not surprising since the materials used in most classical studies of test or diallel crosses had been more or less subjected to selection for improved performance (additive gene action). Thus, selection for improved performance of those inbred lines in breeding might have eliminated most hybrid breakdown genes or gene combinations observed in our base RI mapping population.

Genetic basis of heterosis in rice: Two unique features of this research are its experimental design and statistical methods used. Our crossing schemes and experiments using related RI, BC, and testcross populations were specifically designed to allow simultaneous mapping and characterization of loci contributing to inbreeding depression and heterosis. Data from the parents, RILs, BCF_1 hybrids, testcross hybrids, and testers in the same experiments provided direct measurements of hybrid breakdown and heterosis. In this way, both additive and nonadditive gene actions at the detected loci were more accurately resolved. For instance, the QTL main effects obtained using the heterosis values

of the BCF_1 populations were estimates of the QTL dominance effects ($\frac{1}{2}d$) while those obtained from the F_1 mean values contained both additive and dominance effects ($\frac{1}{2}d - \frac{1}{2}a$; MATHER and JINKS 1982). Similarly, for the epistatic loci, the estimated epistatic effects using heterosis measurements should be the dominance \times dominance effects of the epistatic QTL, while those from the mean F_1 values contained both additive and nonadditive epistatic components (MATHER and JINKS 1982). Use of two testcross populations offered additional advantages in understanding the genetic basis of heterosis since test crosses are the most common way to identify superior hybrids in animal and plant breeding programs. The main and epistatic effects of QTL obtained from the F_1 mean and heterosis values of the testcross populations also reflected the relative importance of different types of gene action in heterosis.

Second, the mixed model approach used in this study is an extension of the composite interval QTL mapping method (ZENG 1993, 1994) with inclusion of digenic epistasis and the appropriate background genetic variation control of all significant main-effect and epistatic QTL in the model (WANG *et al.* 1999). Computer simulation demonstrated that in a DH, RI, or BC population of 200 individuals, QTL with main and/or epistatic effects $>5\%$ in R^2 can be reliably detected and estimated (WANG *et al.* 1999). This was the basis of the threshold $P \leq 0.002$ and $R^2 \geq 5\%$ used in this study. Using this method, we were able to identify many main-effect and epistatic QTL responsible for $>70\%$ of the total phenotypic variation for BY and GY in each of the mapping populations. If we had used methods such as Map-Maker/QTL or regression, we would have reached similar results as XIAO *et al.* (1994) with one to three mapped main-effect QTL explaining $<30\%$ of the phenotypic variation in each of the mapping populations.

Epistasis is a common feature of most loci associated with inbreeding depression and heterosis: This conclusion was supported by the following three observations. First, in each of the mapping populations, the majority (43.7% for BY and 58.6% for GY in the RILs, 42.5 and 58.3% in $LTBCF_1$ s, 42.5 and 58.3% in $TQBCF_1$ s, 59.0 and 49.4% in $Z413F_1$ s, and 56.1 and 48.3% in $IR64F_1$ s) of the phenotypic variation in the F_1 mean and heterosis values was due to epistatic QTL, while a much smaller portion (28.5% for BY and 30.4% for GY in the RILs, 9.6 and 10.2% in $LTBCF_1$, 14.7 and 24.0% in $TQBCF_1$, 20.5 and 23.1% in $Z413F_1$, and 14.1 and 18.3% in $IR64F_1$) of the variation was due to main-effect QTL. Second, 25 (86%) of the 29 QTL with significant main-effects were involved in epistasis detected in one or more populations (Figure 2). Together, of at least 54 QTL identified in this study, only 4 were not involved in epistasis in any of the mapping populations (Figure 2). The pronounced epistasis detected in this study was not due to multiplicative gene action since all identified epistatic loci were detectable using both original and \log_e -transformed

data. In a similar experimental design, XIAO *et al.* (1994) identified a single main-effect QTL in each of the two rice BCF₁ populations, which had R^2 of $\sim 6\text{--}7\%$ for GY with the majority of the phenotypic variation unexplained. Apparently, their failure to detect epistasis was largely attributed to the unavailability of appropriate mapping methodology. With a similar experimental design, STUBER *et al.* (1992) reported 6 and 8 main-effect QTL responsible for $\sim 60\%$ of the phenotypic variation of GY in two maize BCF₁ populations, though epistasis was not adequately evaluated. These and other data suggest that main-effect QTL tend to explain a greater portion of phenotypic variation for GY in maize than in rice (STUBER *et al.* 1992; LIN *et al.* 1996; VELDBOOM and LEE 1996; LI *et al.* 1997a,b; YU *et al.* 1997). Epistasis for complex traits appears to be more pronounced in self-pollinated crop species than in outcrossing species. This is not surprising since coadapted gene complexes generated by epistasis between or among unlinked loci can be more easily maintained in the former than in the latter (ALLARD 1988).

Our results revealed several interesting properties of epistasis in rice. First, epistasis does not necessarily occur between main-effect QTL. For instance, of the 50 epistatic QTL pairs contributing to heterosis, 3 occurred between alleles at two main-effect QTL (type I), 20 between alleles at a main-effect QTL and a “background” locus (type II), and 27 between alleles at two complementary loci (type III). Of the 18 epistatic QTL pairs associated with hybrid breakdown, 6 were of type II and the rest (66.7%) were of type III. Second, all detected epistasis occurred between alleles at two unlinked QTL. These were consistent with the results on grain yield components observed in the F₄ progeny of the same cross (LI *et al.* 1997a). Third, the overall magnitude of the QTL epistatic effects detected in the present study was slightly greater than the mean main-effect QTL by 5.2% for BY and 16.3% for GY in the BC and testcross populations, but smaller by 19.1% for BY and 11.5% for GY in the RILs. It should be pointed out that the estimated epistatic effects obtained in this study almost certainly underestimated the true QTL epistatic effects. This is because the assumptions that $aa_{ii} = aa_{jj} = -aa_{ij} = -aa_{ji}$ for the RI population and $aa_{ij} = dd_{ij} = -ad_{ij} = -da_{ij}$ for the BCF₁ populations generally do not hold true (LI *et al.* 1997a). Our results that both positive and negative heterosis resulted from interactions between alleles at many epistatic loci suggest an explanation for the complexity of heterosis in rice and many other species.

Epistasis plays an important role in the evolution of rice: It is a long-debated issue tracked back to Wright’s shifting balance theory and Fisher’s large population size theory regarding the relative importance of epistasis as a genetic basis underlying evolutionary changes (*cf.* WADE and GOODNIGHT 1998). As discussed above, the pronounced epistasis for fitness traits in rice appeared to

be reflected in two aspects, the large portion of the total phenotypic variance in fitness (GY) and its components contributed by epistatic loci (LUO *et al.* 2001) and the predominant pattern by which most interactions occurred between alleles of complementary loci. In the former case, the contribution of epistasis to population variance was far greater than most theoreticians had assumed (GOODNIGHT 1995; WADE and GOODNIGHT 1998). In the latter case, our results lend strong support to WRIGHT’s (1969) statement that “Evolution depends on the fitting together of favorable complexes that can not be described in themselves as either favorable or unfavorable.”

Epistasis as the genetic basis for hybrid breakdown, outbreeding depression, and recombination load is demonstrated in almost all experimental studies involving interspecific crosses (*cf.* WADE and GOODNIGHT 1998). Similarly, breakdown of coadapted indica and japonica gene complexes by recombination appeared responsible for the significant level of hybrid breakdown observed in the RILs. This conclusion is supported by the observation that most epistatic effects, aa_{ij} , detected in the RI population were positive (>0). According to MATHER and JINKS (1982), this indicated that most recombinant type interactions between alleles from different parents at the epistatic QTL resulted in reduced BY and GY. In other words, hybrid breakdown of the RILs is due largely to incompatible interactions between *indica* (Teqing) and *japonica* (Lemont) alleles at unlinked epistatic QTL, as reported previously (LI *et al.* 1997a,b). It would be expected that main-effect QTL tend to increase the mean fitness of the RI population as a result of selection favoring the alleles for increased GY and/or BY.

The suggestion from the correlation analysis that a common group of genes contribute to both hybrid breakdown of the RILs and heterosis of the BC/testcross hybrids was supported by the close correspondence in genomic locations of detected QTL in the related populations (Figure 2). For instance, 7 of the 14 main-effect QTL and 10 of the 14 epistatic QTL associated with hybrid breakdown were also detected as main-effect and/or epistatic QTL affecting heterosis in at least one of the BC/testcross populations. We further noted a significant portion of QTL mapped in similar locations in different BC/testcross populations, even though no phenotypic correlation exists between the mean values of different hybrid populations (data not shown). For example, of the 29 main-effect QTL detected, 7 (24.1%) were detectable in more than one mapping population. When the epistatic loci were included, 25 (50%) of the 50 loci were detected in more than one mapping population.

As a predominantly selfing plant species, subdivision of populations into subspecies and different local ecotypes of rice was correlated with ecological variability and environmental heterogeneity. For instance, the

maximum diversity at both phenotypic (including isozymes) and molecular levels (RFLPs, randomly amplified polymorphic DNA and simple sequence repeats) of rice is along the Himalayas where the environments were most heterogeneous (our unpublished data; CHANG 1976; OKA 1988; LI and RUTGER 2000). In other words, predominantly selfing plant species present the most extreme form of the shifting balance process where there is more pronounced epistasis at multiple loci and subdivision of populations by inbreeding. It can be conceived that with strong epistasis and genotype-by-environment interactions for fitness traits, local adaptation can be readily achieved by rare multilocus genotypes arising from recombination of occasional outcrossing between subpopulations, leading to multiple fitness peaks in the diverse environments. In these cases, epistasis and genotype-by-environment interactions act as evolutionarily diverging forces (WADE and GOODNIGHT 1998), while recombination produces novel multilocus genotypes on which selection and inbreeding (or genetic drift) can operate. Thus, our results lend strong support to Wright's shifting balance theory and suggest that epistasis combined with genotype-by-environment interactions may have played a key role in the evolution of rice and other predominantly selfing plant species.

"Overdominance" is associated with most loci contributing to heterosis in rice: This conclusion comes from the following two results. First, 14 (58.3%) of the 24 main-effect QTL detected in the BC and testcross populations appeared to be overdominant as they were either only detectable using heterosis values or had a d/a ratio >2.0 . Two were dominant with d/a ratios of 1.43 and 1.11. The remaining 8 were additive as they were detected only by the F_1 mean values. Second, the dominance effects at all main-effect QTL detected in the two BCF₁ and Z413F₁ populations were positive (>0), resulting in increased GY and/or BY. Interestingly, at all 6 main-effect QTL detected in the IR64F₁ population, the indica heterozygotes (Teqing/IR64) had positive effects, resulting in increased GY and/or BY while all the japonica/indica heterozygotes (Lemont/IR64) had negative effects, resulting in reduced GY and/or BY. Similarly, 28 (87.5%) of the 32 detected epistatic QTL pairs appeared to be "overdominant" as the epistatic effects estimated from heterosis values were equal to or greater than those estimated from the F_1 mean values. In other words, most epistatic QTL contributing to heterosis showed only the dominance \times dominance gene action. There were only 4 pairs of additive epistatic loci that were detectable by only the F_1 mean values of the testcross hybrids.

These data strongly support the notion that heterosis for rice yield derives largely from epistatic interactions between loci that result in apparent overdominance. Pronounced overdominance at main-effect QTL for GY is reported also in maize and rice (STUBER *et al.* 1992; YU *et al.* 1997). Heterosis at single loci may result from

true overdominance at single loci or from pseudooverdominance generated by repulsion-phase linkage between partially dominant genes (CROW 1952). It is conceivable that overdominance at the main-effect QTL in maize is more likely true since outcrossing and selection do not favor a high frequency of repulsion linkage or a high level of epistasis between unlinked genes in maize populations. Our results on epistasis were certainly different from the situations described by SIMMONDS (1979), in which single-locus pseudooverdominance could arise from interactions between homozygous alleles at two loci. Heterosis can be generated by dominance \times dominance epistasis (GOODNIGHT 1999), but our results indicated that epistatic overdominance effects did not generally occur between loci having significant main effects (either additive or dominance effects). The overdominance at most main-effect and epistatic QTL observed in this study was unlikely due to repulsive linkage of completely or partially dominant genes; otherwise, one would have to explain why selection should favor such a high level of genetic load maintained by repulsion linkage in the rice genome.

Genetically, complete or partial dominance should be more likely for loci where there is a null allele (non-functional allele). However, studies on isozymes have indicated a high frequency of codominance and a very low frequency of null alleles at most isozyme loci in rice (LI and RUTGER 2000). Then, the long-debated issue on the genetic basis of heterosis would become the question of how codominance at the genic level in hybrids could lead to overdominance at the phenotypic level. Biochemical or physiological evidence and interpretation for the phenotypic overdominance resulting from the codominant heterotic genes/QTL should shed light on this important issue.

Implications for genetic improvement and marker-aided breeding for improved productivity in rice: Our results indicated that the genetic basis of hybrid breakdown and heterosis in rice is very complex, reflected by the large number of loci involved, their wide genomic distribution, and complex epistatic relationships. These results have important implications for genetic improvement of rice. Our observation that the top 10 hybrids in the BC and testcross populations out-yielded the best commercial hybrid cultivar, Shan you63, by 23.8% (BY) and 39.9% (GY) indicate that there is tremendous genetic variation and potential for heterosis in rice productivity. Thus, development of hybrid cultivars should be more efficient and promising than breeding for inbred varieties with regard to further increasing the productivity of rice through exploitation of intersubspecific heterosis for both increased biomass and its partitioning. To do so, however, backcross breeding should be more effective to introgress rare desirable alleles or allele combinations from distantly related donor parents and to overcome the genetic drag arising from incompatible epistasis. Marker-aided transfer of desirable QTL identi-

fied in this study to improve productivity is expected to be difficult because of epistasis, possible genotype-by-environment interactions, and few main-effect candidate QTL. Nevertheless, a main-effect QTL mapped between G249 and RG418 on chromosome 3 is of particular interest. This QTL was detected in three of the five mapping populations in this study and was mapped to the same genomic location in the F₄ progeny of the Lemont/Teqing cross and several other japonica/indica mapping populations (LIN *et al.* 1996; WU *et al.* 1996; LI *et al.* 1998). This QTL is associated with changes of the source size (leaf length and size) and sink capacity (LI *et al.* 1998). The large additive and dominance effects of this QTL justify its potential use in genetic improvement of both inbred and hybrid cultivars through marker-aided transfer in breeding programs.

We thank Dr. Clark and an anonymous reviewer for their valuable comments on the first draft of the manuscript. We are grateful to Drs. S. D. Tansley and S. McCouch of Cornell University and the Japanese Rice Genome Research Program for providing us with DNA probes. This research was supported by a grant from the Rockefeller Foundation (RF) to Z.-K. Li and A. H. Paterson, by a grant from the Chinese Ministry of Agriculture to C. S. Ying and L. J. Luo, and by a grant from the Texas High Education Board to Z.-K. Li and J. Stansel. L. J. Luo and H. W. Mei were also supported by the RF fellowships.

LITERATURE CITED

- ALLARD, R. W., 1960 *Principles of Plant Breeding*, pp. 213–223. John Wiley & Sons, New York.
- ALLARD, R. W., 1988 Genetic changes associated with the evolution of adaptedness in cultivated plants and their wild progenitors. *J. Hered.* **79**: 225–238.
- BRUCE, A. B., 1910 The Mendelian theory of heredity and the augmentation of vigor. *Science* **32**: 627–628.
- CHANG, T.-T., 1976 The origin, evolution, cultivation, and diversification of Asian and African rices. *Euphytica* **25**: 425–441.
- CROW, J. F., 1952 Dominance and overdominance, pp. 282–297 in *Heterosis*, edited by J. W. GOWEN. Iowa State College Press, Ames, IA.
- CROW, J. F., 1986 Basic concepts in population, quantitative, and evolutionary genetics. University of Wisconsin, Madison, WI/W. H. Freeman and Company, New York.
- EAST, E. M., 1936 Heterosis. *Genetics* **21**: 375–397.
- FALCONER, D. S., 1981 *Introduction to Quantitative Genetics*, Ed. 2. Longman, London/New York.
- JONES, D. F., 1917 Dominance of linked factors as a means of accounting for heterosis. *Proc. Natl. Acad. Sci. USA* **3**: 310–312.
- GOODNIGHT, C. J., 1995 Epistasis and the increase in additive genetic variance: implications for phase I of Wright's shifting balance process. *Evolution* **49**(3): 502–511.
- GOODNIGHT, C. J., 1999 Epistasis and heterosis, pp. 59–67 in *The Genetics and Exploitation of Heterosis in Crops*, edited by J. G. COORS and S. PANDEY. American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America, Madison, WI.
- KEEBLE, F., and C. PELLEW, 1910 The mode of inheritance of stature and of time of flowering in peas (*Pisum sativum*). *J. Genet.* **1**: 47–56.
- LI, Z., and J. N. RUTGER, 2000 Geographic distribution and multi-locus structure of isozyme variation in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* **101**: 379–387.
- LI, Z. K., S. R. M. PINSON, J. W. STANSEL and W. D. PARK, 1995 Identification of quantitative trait loci (QTL) for heading date and plant height in cultivated rice (*Oryza sativa* L.). *Theor. Appl. Genet.* **91**: 374–381.
- LI, Z., S. R. M. PINSON, A. H. PATERSON, W. D. PARK and J. W. STANSEL, 1997a Epistasis for three grain yield components in rice (*Oryza sativa* L.). *Genetics* **145**: 453–465.
- LI, Z., S. R. M. PINSON, A. H. PATERSON, W. D. PARK and J. W. STANSEL, 1997b Genetics of hybrid sterility and hybrid breakdown in an inter-subspecific rice (*Oryza sativa* L.) population. *Genetics* **145**: 1139–1148.
- LI, Z., S. R. M. PINSON, A. H. PATERSON and J. W. STANSEL, 1998 Genetic dissection of the source-sink relationship in rice. *Mol. Breed.* **4**: 419–426.
- LIN, H. X., H. R. QIAN, J. Y. ZHUANG, J. LU, S. H. MIN *et al.*, 1996 RFLP mapping of QTL for yield and related characters in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* **92**: 920–927.
- LINCOLN, S., M. DALY and E. LANDER, 1992 Constructing genetic maps with MAPMAKER/EXP 3.0, Ed. 3. Whitehead Technical Report.
- LUO, L. J., Z. K. LI, H. W. MEI, Q. Y. SHU, R. TABIEN *et al.*, 2001 Overdominant epistatic loci are the primary genetic basis of inbreeding depression and heterosis in rice. II. Grain yield components. *Genetics* **158**: 1755–1771.
- MATHER, K., and J. L. JINKS, 1982 *Biometrical Genetics*, Ed. 3. Chapman & Hall, London, New York.
- OKA, H. I., 1988 *Origin of Cultivated Rice*. Japan Scientific Society Press, Tokyo.
- PATTERSON, H. D., and R. THOMPSON, 1971 Recovery of inter-block information when block sizes are unequal. *Biometrika* **58**: 545–554.
- PATTERSON, H. D., and R. THOMPSON, 1974 Maximum likelihood estimation of components of variance. Proceedings of the 8th International Biometric Conference. Biometric Society, Washington, DC.
- SAS INSTITUTE, INC., 1996 *SAS Users Guide: Statistic*. SAS Institute, Cary, NC.
- SHULL, G. H., 1908 The composition of a field of maize. *Ann. Breed. Assoc.* **4**: 296–301.
- SIMMONDS, N. W., 1979 Principles of crop improvement. Longman Group Limited, London/New York.
- STEBBINS, G. L., 1958 The inviability weakness, and sterility of interspecific hybrids. *Adv. Genet.* **9**: 147–215.
- STUBER, C. W., 1994 Heterosis in plant breeding. *Plant Breed. Rev.* **12**: 227–251.
- STUBER, C. W., S. E. LINCOLN, D. W. WOLFF, T. HELENTJARIS and E. S. LANDER, 1992 Identification of genetic factors contributing to heterosis in a hybrid from two elite maize inbred lines using molecular markers. *Genetics* **132**: 832–838.
- VIRMANI, S., 1994 Heterosis and hybrid rice breeding. Springer Verlag, Berlin/Heidelberg/New York.
- WADE, M. J., and C. J. GOODNIGHT, 1998 The theories of Fisher and Wright in the context of metapopulations: when nature does many small experiments. *Evolution* **52**: 1537–1553.
- WANG, D. L., J. ZHU, Z. K. LI and A. H. PATERSON, 1999 Mapping QTL with epistatic effects and QTL × environment interactions by mixed model approaches. *Theor. Appl. Genet.* **99**: 1255–1264.
- WRIGHT, S., 1969 *Evolution and the Genetics of Populations*, Vol. 2. University of Chicago Press, Chicago.
- WRIGHT, S., 1977 *Evolution and the Genetics of Populations*, Vol. 4. University of Chicago Press, Chicago.
- WU, P., G. ZHANG and N. HUANG, 1996 Identification of QTL controlling quantitative characters in rice using RFLP markers. *Euphytica* **89**: 349–354.
- XIAO, J. H., J. LI, L. P. YUAN and S. D. TANKSLEY, 1994 Dominance is the major genetic basis of heterosis in rice as revealed by QTL analysis using molecular markers. *Genetics* **140**: 745–754.
- YU, S. B., J. X. LI, C. G. XU, Y. F. TAN, Y. J. GAO *et al.*, 1997 Importance of epistasis as the genetic basis of heterosis in an elite rice hybrid. *Proc. Natl. Acad. Sci. USA* **94**: 9226–9231.
- ZENG, S. X., Z. W. LU and X. Q. YANG, 1979 Relationships between F₁ heterosis and their parental performance in rice. *Acta Crop Sci. Sin.* **5**(3): 23–34.
- ZENG, Z.-B., 1993 Theoretical basis of separation of multiple linked gene effects on mapping quantitative trait loci. *Proc. Natl. Acad. Sci. USA* **90**: 10972–10976.
- ZENG, Z.-B., 1994 Precision mapping of quantitative trait loci. *Genetics* **136**: 1457–1468.