

 Open access • Journal Article • DOI:10.1038/77347

Overproduction of salicylic acid in plants by bacterial transgenes enhances pathogen resistance — [Source link](#)

Marianne C. Verberne, Robert Verpoorte, John F. Bol, Jesús Mercado-Blanco ...+2 more authors

Institutions: Leiden University, Spanish National Research Council, Utrecht University

Published on: 01 Jul 2000 - Nature Biotechnology (Nature Publishing Group)

Topics: Systemic acquired resistance, Pathogenesis-related protein, Hypersensitive response, Plant defense against herbivory and Plant disease resistance

Related papers:

- [Isochorismate synthase is required to synthesize salicylic acid for plant defence](#)
- [Requirement of Salicylic Acid for the Induction of Systemic Acquired Resistance](#)
- [A Central Role of Salicylic Acid in Plant Disease Resistance](#)
- [Salicylic Acid Induction–Deficient Mutants of Arabidopsis Express PR-2 and PR-5 and Accumulate High Levels of Camalexin after Pathogen Inoculation](#)
- [Salicylic acid : a likely endogenous signal in the resistance response of tobacco to viral infection](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/overproduction-of-salicylic-acid-in-plants-by-bacterial-1ig2smfmdg>

El artículo seleccionado no se encuentra disponible por ahora a texto completo

The requested paper is not presently available in its full-text version